
CORE PURE / COMPLEX NUMBERS

Loci and Regions in the Argand Diagram

Worked solutions 10 questions · 88 marks

Questions

Question 1	*6 marks*	2
Question 2	*6 marks*	5
Question 3	*10 marks*	8
Question 4	*12 marks*	12
Question 5	*12 marks*	15
Question 6	*12 marks*	19
Question 7	*7 marks*	23
Question 8	*7 marks*	25
Question 9	*9 marks*	29
Question 10	*7 marks*	32



1. The Argand diagram shows a circle of radius 4. The centre of the circle is the point which represents the complex number $-2+i$.

(a) Use set notation to define the locus of complex numbers, z, represented by points which lie on the circle. **[2]**

The locus L is defined by

$$L = \left\{ z \in \mathbb{C} \;\middle|\; |z-(2+i)| = |z+1-3i| \right\}$$

(b) On a copy of the Argand diagram, sketch and label the locus L. **[2]**

You are given that the locus

$$\left\{ z \in \mathbb{C} \;\middle|\; \arg(z+1-2i) = \frac{2\pi}{3},\ \mathrm{Im}(z) = 5 \right\}$$

contains only one number.

(c) Find this number. **[2]**

Solution

(a) The circle has centre $-2+i$ and radius 4.

So the locus is the set of complex numbers whose distance from $-2+i$ is 4:

$$\{z \in \mathbb{C} \mid |z-(-2+i)| = 4\} = \{z \in \mathbb{C} \mid |z+2-i| = 4\}$$

(b) Let $z = x+iy$, where $x, y \in \mathbb{R}$.

Then

$$|z-(2+i)| = |z+1-3i|$$

gives

$$|(x-2)+i(y-1)| = |(x+1)+i(y-3)|$$

Squaring both sides:

$$(x-2)^2+(y-1)^2 = (x+1)^2+(y-3)^2$$

Now expand:

$$\begin{aligned} x^2-4x+4+y^2-2y+1 &= x^2+2x+1+y^2-6y+9 \\ -4x-2y+5 &= 2x-6y+10 \\ -6x+4y-5 &= 0 \\ 6x-4y+5 &= 0 \end{aligned}$$

So

$$L:\ 6x-4y+5=0$$


or equivalently

$$y = \frac{3}{2}x+\frac{5}{4}$$

This is a straight line. Since it is the set of points equidistant from $2+i$ and $-1+3i$, it is the perpendicular bisector of the line segment joining those two points.

It passes through the midpoint of that segment,

$$\left(\frac{2+(-1)}{2}, \frac{1+3}{2}\right) = \left(\frac{1}{2}, 2\right)$$



(c) Since $\mathrm{Im}(z) = 5$, write

$$z = x+5i$$

Then

$$z + 1 - 2i = (x + 1) + 3i$$

We are given

$$\arg(z + 1 - 2i) = \frac{2\pi}{3}$$

So $(x + 1) + 3i$ lies in the second quadrant, with $x + 1 < 0$. Therefore

$$\tan \frac{2\pi}{3} = \frac{3}{x + 1}$$

Using $\tan \frac{2\pi}{3} = -\sqrt{3}$,

$$\frac{3}{x + 1} = -\sqrt{3}$$

Hence


$$x + 1 = -\sqrt{3}$$

so

$$x = -1 - \sqrt{3}$$

Therefore the required number is

$$z = -1 - \sqrt{3} + 5i$$



2. **(a)** Sketch, on the Argand diagram above, the locus of points satisfying the equation

$$|z - (2 + 4i)| = 3$$ **[2]**

(b) Sketch, also on the Argand diagram, the locus of points satisfying the equation

$$\arg(z - 1) = \frac{\pi}{4}$$ **[1]**

(c) For the complex number w, find the maximum value of $|w|$ such that


$$|w - (2 + 4i)| \leq 3 \qquad \text{and} \qquad 0 \leq \arg(w - 1) \leq \frac{\pi}{4}$$ **[3]**

Solution

(a) The locus

$$|z - (2 + 4i)| = 3$$

is a circle with centre at the point representing $2 + 4i$ and radius 3.




(b) The locus

$$\arg(z-1) = \frac{\pi}{4}$$

is the half-line starting at the point representing 1 on the real axis, making an angle of $\frac{\pi}{4}$ with the positive real axis.

In Cartesian form this is

$$y = x - 1, \qquad x > 1$$



(c) Let

$$w = x + iy$$

The condition


$$|w - (2+4i)| \leq 3$$

means that w lies inside or on the circle from part (a).

Also,

$$0 \leq \arg(w-1) \leq \frac{\pi}{4}$$

means that w lies between the half-line from 1 along the positive real axis and the half-line from part (b).



Since the circle has lowest point $(2, 1)$, every point in the circle has $y \geq 1$, so the condition $\arg(w-1) \geq 0$ is automatically satisfied.

Therefore the required region is the part of the disc lying on or below the line

$$y = x - 1$$

First find where this line meets the circle:

$$\begin{aligned}
(x-2)^2 + (y-4)^2 &= 9 \\
(x-2)^2 + (x-1-4)^2 &= 9 \\
(x-2)^2 + (x-5)^2 &= 9 \\
x^2 - 4x + 4 + x^2 - 10x + 25 &= 9 \\
2x^2 - 14x + 20 &= 0 \\
x^2 - 7x + 10 &= 0 \\
(x-2)(x-5) &= 0
\end{aligned}$$

so the intersection points are $P(2, 1)$ and $Q(5, 4)$, as marked on the diagram.

Now $|w|$ is the distance from O to the point representing w. Every point of the region lies in the disc, so $x \leq 2 + 3 = 5$. It also lies on or below the line, so $y \leq x - 1 \leq 4$. Both x and y are positive in the region, since $y \geq 1$ and $x \geq y + 1$. Hence

$$|w|^2 = x^2 + y^2 \leq 5^2 + 4^2 = 41$$

Equality holds at Q, which is in the region.

Hence the maximum value of $|w|$ is $\sqrt{41}$, attained when $w = 5 + 4i$.

3. On an Argand diagram, the locus of points satisfying $|z - 4 + 3i| = 8$ is a circle.

(a) Write down

(i) the coordinates of the centre of this circle **[1]**

(ii) the radius of this circle **[1]**

(b) On an Argand diagram, shade the set of points defined by

$$|z - 4 + 3i| \leq 8$$ **[1]**

(c) For the set of points defined in part (b), determine the maximum value of $|z|$. **[3]**

The set of points A is defined by

$$A = \left\{ z \in \mathbb{C} \;\middle|\; -\frac{\pi}{4} \leq \arg(z - 3 - 6i) \leq \frac{3\pi}{4} \right\} \bigcap \left\{ z \in \mathbb{C} \;\middle|\; |z - 4 + 3i| \leq 8 \right\}$$

(d) Determine the area of the region defined by A, giving your answer to 3 significant figures. **[4]**

Solution

(a) (i)

$$|z - 4 + 3i| = |z - (4 - 3i)|$$

So the circle has centre $(4, -3)$.


(ii) Comparing with the standard form $|z - a| = r$, the radius is 8.

(b) The inequality

$$|z - 4 + 3i| \leq 8$$

means all points whose distance from $4 - 3i$ is at most 8.

So we shade the interior and boundary of the circle with centre $(4, -3)$ and radius 8.



(c) The modulus $|z|$ is the distance from the origin O to the point representing z.

The centre C of the circle represents $4 - 3i$, so

$$OC = |4 - 3i| = \sqrt{4^2 + (-3)^2} = 5$$

The point of the region furthest from O is where the line from O through C meets the circle on the far side of C. Its distance from O is OC plus the radius.

Therefore the maximum value of $|z|$ is

$$5 + 8 = 13$$

(d) Let $z = x + iy$.

First consider the argument condition:

$$-\frac{\pi}{4} \leq \arg(z - 3 - 6i) \leq \frac{3\pi}{4}$$

Since

$$z - 3 - 6i = (x - 3) + i(y - 6)$$

the boundary arguments $-\pi/4$ and $3\pi/4$ are opposite directions, so they form a straight line through $(3, 6)$ with gradient -1.

Hence the boundary line is

$$y - 6 = -(x - 3)$$

so

$$x + y = 9$$

To decide which side is needed, take the point $(3, 7)$. Then

$$\arg\big((3 + 7i) - 3 - 6i\big) = \arg(i) = \frac{\pi}{2}$$

which is in the given range, and $(3, 7)$ satisfies $x + y = 10$.

So the argument condition is the half-plane

$$x + y \geq 9$$

The circle has centre $C = (4, -3)$ and radius 8. Since

$$4 + (-3) = 1 < 9$$

the centre lies on the opposite side of the line, so the region A is the minor segment cut off by the line $x + y = 9$.

The circle has equation $(x - 4)^2 + (y + 3)^2 = 64$. To find where the line meets it, substitute $y = 9 - x$:

$$(x - 4)^2 + (12 - x)^2 = 64$$

$$x^2 - 8x + 16 + x^2 - 24x + 144 = 64$$

$$2x^2 - 32x + 96 = 0$$


$$x^2 - 16x + 48 = 0$$

$$(x - 4)(x - 12) = 0$$

So the line meets the circle at $P = (4, 5)$ and $Q = (12, -3)$.

CP joins $(4, -3)$ to $(4, 5)$, so it is vertical, and CQ joins $(4, -3)$ to $(12, -3)$, so it is horizontal. Hence

$$\angle PCQ = \frac{\pi}{2}$$




Therefore

$$\begin{aligned}\text{Area}(A) &= \text{area of sector } PCQ - \text{area of } \triangle PCQ \\ &= \frac{1}{2}r^2\theta - \frac{1}{2}r^2\sin\theta \\ &= \frac{1}{2}\cdot 8^2\cdot\frac{\pi}{2} - \frac{1}{2}\cdot 8^2\cdot\sin\frac{\pi}{2} \\ &= 16\pi - 32\end{aligned}$$

So the area is

$$16\pi - 32 \approx 18.3$$

to 3 significant figures.



4. **(a)** The diagram above shows an Argand diagram.

(i) Write down the equation of the locus represented by the circumference of circle A. **[3]**

(ii) Write down the two inequalities that define the shaded region, not including either circumference. **[3]**

(b) (i) Draw an Argand diagram to show the region where

$$-\frac{\pi}{6} < \arg(z+2-i) < \frac{\pi}{4}$$ **[3]**

(ii) Determine whether the point $1+5i$ lies within this region. **[3]**

Solution

(a) (i) For a circle with centre z_0 and radius r, the locus is

$$|z - z_0| = r$$

Circle A has centre $-2+4i$ and radius 3, so

$$|z-(-2+4i)| = 3$$

hence the circumference of circle A is

$$|z+2-4i| = 3$$

(ii) The shaded region is the part inside both circles, and neither circumference is included.

Inside circle A:

$$|z+2-4i| < 3$$

Circle B has centre $2+6i$ and radius 4, so inside circle B:

$$|z-(2+6i)|<4$$

So the two inequalities are

$$|z+2-4i|<3 \qquad \text{and} \qquad |z-(2+6i)|<4$$

(b) (i) We have

$$z+2-i=z-(-2+i)$$


so the argument is measured from the point $-2+i$.

The region

$$-\frac{\pi}{6}<\arg(z+2-i)<\frac{\pi}{4}$$

is therefore the open sector with vertex at $-2+i$ between the rays making angles $-\frac{\pi}{6}$ and $\frac{\pi}{4}$ with the positive real axis.

Since the inequalities are strict, both boundary rays are excluded. The vertex $-2+i$ is excluded too, as $\arg 0$ is undefined.



(ii) For the point $z=1+5i$,

$$z+2-i=(1+5i)+2-i=3+4i$$

So

$$\arg(z+2-i)=\arg(3+4i)=\tan^{-1}\left(\frac{4}{3}\right)\approx 0.927$$


But

$$\frac{\pi}{4}\approx 0.785$$

Since

$$0.927>\frac{\pi}{4}$$

the argument is too large, so $1 + 5i$ does not lie in the region.



5. **(a)** Find the modulus of the complex number $2 + 2\sqrt{3}i$, giving your answer as an integer. **[2]**

(b) The locus of points, L, satisfies the equation $|z - (2 - 2\sqrt{3}i)| = 2$.

(i) Sketch the locus L on the Argand diagram above. **[3]**

(ii) The complex number w lies on L so that $-\pi < \arg w \leq \pi$. Find the greatest possible value of $\arg w$, giving your answer in terms of π. **[2]**

(c) Solve the equation $z^4 = 2+2\sqrt{3}i$, giving your answers in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$. **[5]**

Solution

(a) The modulus is


$$|2 + 2\sqrt{3}i| = \sqrt{2^2 + (2\sqrt{3})^2} = \sqrt{4 + 12} = \sqrt{16} = 4$$

(b) (i) The equation

$$|z - (2 - 2\sqrt{3}i)| = 2$$

represents all points a distance 2 from the point $2 - 2\sqrt{3}i$.

So L is a circle with centre $(2, -2\sqrt{3})$ and radius 2. The centre is a distance 2 from the imaginary axis, so the circle touches the imaginary axis at $-2\sqrt{3}i$.



(ii) Let C be the centre of the circle, so

$$C = 2 - 2\sqrt{3}i$$

First find OC:

$$OC = |2 - 2\sqrt{3}i| = \sqrt{2^2 + (-2\sqrt{3})^2} = \sqrt{4 + 12} = 4$$


Also,

$$\arg C = -\frac{\pi}{3}$$

because C is in the fourth quadrant and

$$\tan|\arg C| = \frac{2\sqrt{3}}{2} = \sqrt{3}$$

The greatest possible value of $\arg w$ occurs when the line from the origin to the circle is a tangent, touching the circle at w.



Then triangle OCW is right-angled at W, with

$$CW = 2, \qquad OC = 4$$

Let $\angle COW = \alpha$. Then

$$\sin\alpha = \frac{CW}{OC} = \frac{2}{4} = \frac{1}{2}$$

so

$$\alpha = \frac{\pi}{6}$$

Since this tangent is above the line OC,

$$\arg w = \arg C + \alpha = -\frac{\pi}{3} + \frac{\pi}{6} = -\frac{\pi}{6}$$

Therefore the greatest possible value of $\arg w$ is $-\frac{\pi}{6}$.

(c) From part (a),

$$|2 + 2\sqrt{3}i| = 4$$

and since $2 + 2\sqrt{3}i$ lies in the first quadrant,

$$\arg(2 + 2\sqrt{3}i) = \frac{\pi}{3}$$

So

$$2 + 2\sqrt{3}i = 4e^{i\pi/3}$$

We solve

$$z^4 = 4e^{i\pi/3}$$

Let $z = re^{i\theta}$. Then

$$z^4 = r^4 e^{i4\theta}$$

so

$$r^4 e^{i4\theta} = 4e^{i(\pi/3+2k\pi)} \qquad (k \in \mathbb{Z})$$

Equating moduli gives

$$r^4 = 4$$

so, with $r > 0$,

$$r = 4^{1/4} = \sqrt{2}$$

Equating arguments gives

$$4\theta = \frac{\pi}{3} + 2k\pi$$

hence

$$\theta = \frac{\pi}{12} + \frac{k\pi}{2}$$

The four distinct roots come from $k = 0, 1, 2, 3$:

$$\theta = \frac{\pi}{12}, \quad \frac{7\pi}{12}, \quad \frac{13\pi}{12}, \quad \frac{19\pi}{12}$$

Writing these in the range $-\pi < \theta \leq \pi$,

$$\theta = \frac{\pi}{12}, \quad \frac{7\pi}{12}, \quad -\frac{11\pi}{12}, \quad -\frac{5\pi}{12}$$

Therefore

$$z = \sqrt{2}e^{i\pi/12}, \qquad \sqrt{2}e^{i7\pi/12}, \qquad \sqrt{2}e^{-i11\pi/12}, \qquad \sqrt{2}e^{-i5\pi/12}$$

6. The point P represents a complex number z on an Argand diagram such that

$$|z-(1+i)| = \frac{1}{2}|z-(4+i)|$$

(a) Show that, as z varies, the locus of P is a circle, and give the coordinates of the centre and the radius of the circle. **[5]**

The point Q represents a complex number z on an Argand diagram such that

$$|z-(2+3i)| = |z+2-i|$$

(b) Sketch, on the same Argand diagram, the locus of P and the locus of Q as z varies. **[5]**

(c) On your diagram shade the region which satisfies

$$|z-(1+i)| \leq \frac{1}{2}|z-(4+i)| \quad \text{and} \quad |z-(2+3i)| \geq |z+2-i|$$

[2]

Solution

(a) Let

$$z = x + iy$$

Then

$$|z-(1+i)| = \sqrt{(x-1)^2+(y-1)^2}$$

and

$$|z-(4+i)| = \sqrt{(x-4)^2+(y-1)^2}$$

So the given condition becomes

$$\sqrt{(x-1)^2+(y-1)^2} = \frac{1}{2}\sqrt{(x-4)^2+(y-1)^2}$$

Squaring both sides,

$$4\big((x-1)^2+(y-1)^2\big) = (x-4)^2+(y-1)^2$$

Expand:

$$4\big(x^2-2x+1+y^2-2y+1\big) = x^2-8x+16+y^2-2y+1$$

$$4x^2-8x+4+4y^2-8y+4 = x^2-8x+16+y^2-2y+1$$

Bring everything to one side:

$$3x^2+3y^2-6y-9 = 0$$

$$x^2+y^2-2y-3 = 0$$

Complete the square in y:

$$x^2 + (y^2 - 2y + 1) - 4 = 0$$
$$x^2 + (y-1)^2 = 4$$

This is the equation of a circle.

Hence the locus is the circle with centre $(0, 1)$ and radius 2.

(b) For the second locus, let again $z = x + iy$.

Then

$$|z - (2 + 3i)| = \sqrt{(x-2)^2 + (y-3)^2}$$

and

$$|z + 2 - i| = |z - (-2 + i)| = \sqrt{(x+2)^2 + (y-1)^2}$$

So

$$\sqrt{(x-2)^2 + (y-3)^2} = \sqrt{(x+2)^2 + (y-1)^2}$$

Squaring both sides,

$$(x-2)^2 + (y-3)^2 = (x+2)^2 + (y-1)^2$$

Expand and simplify:

$$x^2 - 4x + 4 + y^2 - 6y + 9 = x^2 + 4x + 4 + y^2 - 2y + 1$$
$$-4x - 6y + 13 = 4x - 2y + 5$$
$$-8x - 4y + 8 = 0$$
$$2x + y - 2 = 0$$


So the locus of Q is the straight line

$$y = 2 - 2x$$

This is the perpendicular bisector of the line segment joining $(2, 3)$ and $(-2, 1)$.

On the same Argand diagram, we sketch:

- the circle $x^2 + (y-1)^2 = 4$, centre $(0, 1)$, radius 2,
- the line $y = 2 - 2x$.



(c) From part (a),

$$|z - (1 + i)| \leq \frac{1}{2}|z - (4 + i)|$$

gives

$$x^2 + (y - 1)^2 \leq 4$$

so the point lies inside or on the circle.

From part (b),

$$|z - (2 + 3i)| \geq |z + 2 - i|$$

gives


$$\begin{aligned}(x - 2)^2 + (y - 3)^2 &\geq (x + 2)^2 + (y - 1)^2 \\ -8x - 4y + 8 &\geq 0 \\ 2x + y - 2 &\leq 0 \\ y &\leq 2 - 2x\end{aligned}$$

So the point lies on or below the line $y = 2 - 2x$.

Therefore the required region is the intersection of:

- the inside of the circle $x^2 + (y - 1)^2 = 4$,
- the half-plane $y \leq 2 - 2x$.

The shaded region is shown below.


Im
(0, 1)
O
Re

7. On an Argand diagram, the point P representing the complex number z lies on the locus

$$|z-(2-i)|=3$$

The point Q representing the complex number w lies on the locus

$$|w+1-11i|=3|w+1-3i|$$

Find the exact coordinates of the points common to these two loci. **[7]**

Solution

Let a point common to both loci represent the complex number

$$u=x+yi$$

so its coordinates are (x,y). (A new letter is used because the question uses z and w for the two loci.)

For the locus of P,

$$|u-(2-i)|=3$$

Now

$$u-(2-i)=(x-2)+(y+1)i$$

so

$$|(x-2)+(y+1)i|=3$$

Using $|a+bi|=\sqrt{a^2+b^2}$,

$$(x-2)^2+(y+1)^2=9$$

For the locus of Q,

$$|u+1-11i|=3|u+1-3i|$$

Now

$$u+1-11i=(x+1)+(y-11)i, \qquad u+1-3i=(x+1)+(y-3)i$$

So

$$|(x+1)+(y-11)i|=3|(x+1)+(y-3)i|$$

Squaring both sides gives

$$(x+1)^2+(y-11)^2=9\big((x+1)^2+(y-3)^2\big)$$

Expand and simplify:

$$\begin{aligned} x^2+2x+1+y^2-22y+121 &= 9x^2+18x+9+9y^2-54y+81 \\ 0 &= 8x^2+16x+8+8y^2-32y-40 \\ 0 &= x^2+2x+y^2-4y-4 \\ 0 &= (x+1)^2+(y-2)^2-9 \end{aligned}$$

Hence the second locus is

$$(x+1)^2 + (y-2)^2 = 9$$

So the common points satisfy the simultaneous equations

$$(x-2)^2 + (y+1)^2 = 9$$

and

$$(x+1)^2 + (y-2)^2 = 9$$

Both right-hand sides equal 9, so equating the left-hand sides:

$$(x-2)^2 + (y+1)^2 = (x+1)^2 + (y-2)^2$$

$$x^2 - 4x + 4 + y^2 + 2y + 1 = x^2 + 2x + 1 + y^2 - 4y + 4$$

$$-6x + 6y = 0$$

$$y = x$$

Substitute $y = x$ into $(x-2)^2 + (y+1)^2 = 9$:

$$(x-2)^2 + (x+1)^2 = 9$$

$$x^2 - 4x + 4 + x^2 + 2x + 1 = 9$$

$$2x^2 - 2x - 4 = 0$$

$$x^2 - x - 2 = 0$$

$$(x-2)(x+1) = 0$$

So


$$x = 2 \quad \text{or} \quad x = -1$$

and since $y = x$,

$$y = 2 \quad \text{or} \quad y = -1$$

Therefore the exact coordinates of the common points are

$$(2, 2) \quad \text{and} \quad (-1, -1)$$



8. **(a)** Sketch, on the Argand diagram above, the locus of points satisfying the equation

$$|z - (3 + i)| = 1$$ **[1]**

(b) There is a unique complex number w that satisfies both

$$|w - (3 + i)| = 1 \quad \text{and} \quad \arg(w) = \alpha$$

where α is a constant such that $0 < \alpha < \pi$.

(i) Find the value of α. **[2]**

(ii) Find w and express it in the form $r(\cos\theta + i\sin\theta)$. Give each of r and θ to two significant figures. **[4]**


Solution

(a) The locus

$$|z - (3 + i)| = 1$$

is the set of points whose distance from $3 + i$ is 1.

So it is a circle with centre $(3, 1)$ and radius 1.



(b) (i) If $\arg(w) = \alpha$, then w lies on the half-line from the origin at angle α. This half-line is part of the line

$$y = mx$$

where

$$m = \tan\alpha$$

For there to be a unique point w satisfying both conditions, the half-line must touch the circle at exactly one point, so it must be a tangent.

The circle has equation $(x-3)^2 + (y-1)^2 = 1$. Substituting $y = mx$,

$$(x-3)^2 + (mx-1)^2 = 1$$

$$x^2 - 6x + 9 + m^2x^2 - 2mx + 1 = 1$$

$$(1+m^2)x^2 - (6+2m)x + 9 = 0$$

For a tangent this quadratic has a repeated root, so its discriminant is zero:

$$(6+2m)^2 - 4(1+m^2)(9) = 0$$

$$36 + 24m + 4m^2 - 36 - 36m^2 = 0$$

$$24m - 32m^2 = 0$$

$$8m(3-4m) = 0$$

So

$$m = 0 \quad \text{or} \quad m = \frac{3}{4}$$

If $m = 0$, then $\alpha = 0$ (the real axis, which touches the circle at 3), but the question states $0 < \alpha < \pi$, so this is not allowed.

Hence

$$\tan\alpha = \frac{3}{4}$$

so

$$\alpha = \arctan\left(\frac{3}{4}\right) \approx 0.644\,\text{rad}$$

(ii) Using the value of m, the tangent line is

$$y = \frac{3}{4}x$$

The circle equation is

$$(x-3)^2 + (y-1)^2 = 1$$

Substituting $y = \frac{3}{4}x$,

$$(x-3)^2 + \left(\frac{3}{4}x - 1\right)^2 = 1$$

$$x^2 - 6x + 9 + \frac{9}{16}x^2 - \frac{3}{2}x + 1 = 1$$

$$\frac{25}{16}x^2 - \frac{15}{2}x + 9 = 0$$

Multiplying by 16,

$$25x^2 - 120x + 144 = 0$$

Factorising,

$$(5x - 12)^2 = 0$$

So

$$x = \frac{12}{5}$$

Then

$$y = \frac{3}{4}\cdot\frac{12}{5} = \frac{9}{5}$$

Hence

$$w = \frac{12}{5} + \frac{9}{5}i$$

Now find its modulus:

$$|w| = \sqrt{\left(\frac{12}{5}\right)^2 + \left(\frac{9}{5}\right)^2}$$

$$= \sqrt{\frac{144}{25} + \frac{81}{25}}$$

$$= \sqrt{\frac{225}{25}}$$

$$= 3$$

Its argument is

$$\theta = \arg(w) = \alpha = \arctan\left(\frac{3}{4}\right) \approx 0.64$$

So, with $r = 3.0$ and $\theta = 0.64$ to two significant figures,

$$w = \frac{12}{5} + \frac{9}{5}i = 3.0(\cos 0.64 + i \sin 0.64)$$

9. In an Argand diagram, the point P representing the complex number w lies on the locus defined by

$$\left\{z \in \mathbb{C} \;\middle|\; \arg(z-2+i) = \frac{\pi}{6}\right\}$$

You are given that $\text{Im}(w) = 3$.

(a) Find w. **[2]**

The point P also lies on the locus defined by $\left\{z \in \mathbb{C} \;\middle|\; |z-2-11i| = k\right\}$, where k is a constant.

(b) Find the complex number represented by the other point of intersection of the loci defined by

$$\left\{z \in \mathbb{C} \;\middle|\; |z-2-11i| = k\right\} \quad \text{and} \quad \left\{z \in \mathbb{C} \;\middle|\; \arg(z-2+i) = \frac{\pi}{6}\right\}$$

[7]

Solution

(a) Let

$$w = x + 3i$$

since $\text{Im}(w) = 3$.

Then

$$w - 2 + i = (x-2) + 4i$$

Given

$$\arg(w-2+i) = \frac{\pi}{6}$$

so

$$\tan\frac{\pi}{6} = \frac{4}{x-2}$$

Hence

$$\frac{1}{\sqrt{3}} = \frac{4}{x-2}$$

so

$$x - 2 = 4\sqrt{3}$$

and therefore

$$x = 2 + 4\sqrt{3}$$

Thus

$$w = 2 + 4\sqrt{3} + 3i$$

(b) From part (a),

$$P \text{ represents } w = 2 + 4\sqrt{3} + 3i$$

The second locus is

$$|z-2-11i| = k = |z-(2+11i)|$$

which is a circle with centre $2+11i$.

Using the point P to find k,

$$\begin{aligned} k &= |w-(2+11i)| \\ &= \left|(2+4\sqrt{3}+3i)-(2+11i)\right| \\ &= \left|4\sqrt{3}-8i\right| \end{aligned}$$

So

$$\begin{aligned} k^2 &= (4\sqrt{3})^2+(-8)^2 \\ &= 48+64 \\ &= 112 \end{aligned}$$

and hence

$$k = 4\sqrt{7}$$

Now let a general point of intersection be

$$z = x+yi$$

Since

$$\arg(z-2+i) = \frac{\pi}{6}$$

the point lies on the half-line starting at $(2,-1)$ with gradient

$$\tan\frac{\pi}{6} = \frac{1}{\sqrt{3}}$$

so its equation is

$$y+1 = \frac{x-2}{\sqrt{3}}$$

that is,

$$y = \frac{x-2}{\sqrt{3}} - 1$$

The circle has equation

$$(x-2)^2+(y-11)^2 = 112$$

Substitute $y = \frac{x-2}{\sqrt{3}} - 1$:

$$(x-2)^2 + \left(\frac{x-2}{\sqrt{3}} - 12\right)^2 = 112$$

Let

$$u = x-2$$

Then

$$u^2 + \left(\frac{u}{\sqrt{3}} - 12\right)^2 = 112$$

Expanding,

$$u^2 + \left(\frac{u^2}{3} - 8\sqrt{3}\,u + 144\right) = 112$$

$$\frac{4u^2}{3} - 8\sqrt{3}\,u + 144 = 112$$

$$\frac{4u^2}{3} - 8\sqrt{3}\,u + 32 = 0$$

Multiply by 3:

$$4u^2 - 24\sqrt{3}\,u + 96 = 0$$

Divide by 4:

$$u^2 - 6\sqrt{3}\,u + 24 = 0$$

Factorising,

$$(u - 4\sqrt{3})(u - 2\sqrt{3}) = 0$$

So

$$u = 4\sqrt{3} \quad \text{or} \quad u = 2\sqrt{3}$$

Both values are positive, so both points have $x > 2$ and lie on the half-line itself, not on its extension. The value $u = 4\sqrt{3}$ gives the known point P from part (a), so the other intersection has

$$u = 2\sqrt{3}$$

Therefore

$$x = 2 + 2\sqrt{3}$$

and

$$y = \frac{2\sqrt{3}}{\sqrt{3}} - 1 = 2 - 1 = 1$$

Hence the other point represents

$$z = 2 + 2\sqrt{3} + i$$

10. The locus of points $z = x + iy$ that satisfy

$$\arg\left(\frac{z-(2+i)}{z-(6+5i)}\right) = \frac{\pi}{4}$$

is an arc of a circle C.

(a) On an Argand diagram sketch the locus of z. **[2]**

(b) Explain why the centre of C lies on the line $y = 7 - x$. **[1]**

(c) Determine the radius of C. **[2]**

(d) Determine the coordinates of the centre of C. **[2]**

Solution

(a) Let

$$A = 2 + i \quad \text{and} \quad B = 6 + 5i$$

so $A = (2, 1)$ and $B = (6, 5)$.

The condition


$$\arg\left(\frac{z-A}{z-B}\right) = \frac{\pi}{4}$$

means

$$\arg(z-A) - \arg(z-B) = \frac{\pi}{4}$$

so, if Z is the point representing z, the angle AZB is $\frac{\pi}{4}$.

Angles in the same segment are equal, so the locus is an arc of a circle through A and B. Since $\frac{\pi}{4} < \frac{\pi}{2}$, it is the major arc. It lies on the side below/right of the line segment AB, as checked in part (d). Also, A and B are excluded because the expression is undefined there.



(b) Since A and B lie on circle C, the centre of C is equidistant from A and B. Therefore it lies on the perpendicular bisector of AB.

The midpoint of AB is

$$\left(\frac{2+6}{2}, \frac{1+5}{2}\right) = (4, 3)$$

The gradient of AB is

$$\frac{5-1}{6-2} = 1$$

so the perpendicular bisector has gradient -1.

Its equation is

$$y - 3 = -1(x - 4)$$

$$y = 7 - x$$

Therefore the centre lies on the line $y = 7 - x$.

(c) Let P be the centre of C and r its radius.

The angle at the circumference is $\frac{\pi}{4}$, so the angle at the centre subtended by the same chord AB is

$$\angle APB = 2 \times \frac{\pi}{4} = \frac{\pi}{2}$$

So triangle APB is right-angled at P, with $PA = PB = r$. Also

$$|AB|^2 = (6-2)^2 + (5-1)^2 = 32$$

By Pythagoras,

$$r^2 + r^2 = 32$$

so $r^2 = 16$, giving

$$r = 4$$

(d) From part (b), the centre P lies on $y = 7 - x$, so write $P = (a,\ 7 - a)$.

From part (c), $PA = 4$, so

$$(a-2)^2 + (7-a-1)^2 = 16$$

$$a^2 - 4a + 4 + a^2 - 12a + 36 = 16$$

$$2a^2 - 16a + 24 = 0$$

$$a^2 - 8a + 12 = 0$$

$$(a-2)(a-6) = 0$$

Hence the two possible centres are $(2, 5)$ and $(6, 1)$.

To decide which is correct, test a point on one of the two circles. The point $z = 10 + i$ lies on the circle centred at $(6, 1)$, diametrically opposite A:

$$\begin{aligned}\arg\left(\frac{(10+i)-(2+i)}{(10+i)-(6+5i)}\right) &= \arg\left(\frac{8}{4-4i}\right)\\ &= \arg\left(\frac{8(4+4i)}{(4-4i)(4+4i)}\right)\\ &= \arg(1+i)\\ &= \frac{\pi}{4}\end{aligned}$$

So $10 + i$ lies on the locus. But

$$\text{distance from } (10, 1) \text{ to } (2, 5) = \sqrt{(8)^2 + (-4)^2} = \sqrt{80} \neq 4$$

so it does not lie on the circle centred at $(2, 5)$. This also confirms that the arc lies below/right of AB.

Therefore the centre of C is $(6, 1)$.