

Convex Analysis

Daniel Mao

Contents

1	Affine Sets	1
1.1	Definitions	1
2	Relative Topology	3
2.1	Definitions	3
2.2	Basic Properties	3
2.3	Arithmetic Properties	4
3	Convex Sets	5
3.1	Definitions	5
3.2	Arithmetic Properties of Convex Sets	6
3.3	Topological Properties of Convex Sets	6
3.4	The Convex Hull Operator	9
3.5	The Closed Convex Hull Operator	11
3.6	Stability of Convexity	11
3.7	Examples of Convex Sets	16
3.8	The Carathéodory Theorem	16
4	Geometric Objects	19
4.1	Definitions	19
4.2	Properties	19
5	Cones (in Analysis)	21
5.1	Definitions	21
5.2	The cone Operator	23
5.3	Other Properties	24
6	Tangent Cones and Normal Cones	27
6.1	Definitions	27
6.2	Basic Properties	27

6.3	Arithmetic Properties	32
6.4	Other Properties	33
7	Dual Cones and Polar Cones	35
7.1	Definitions	35
7.2	Properties	35
8	Extreme Point and Face	37
8.1	Definitions	37
8.2	Properties of Extreme Points	38
8.3	Face	38
8.4	The Krein-Milman Theorem	39
9	Projection	41
9.1	Definitions	41
9.2	Properties of the Projection Operator	42
9.3	Characterization	44
10	Separation	45
10.1	Definitions	45
10.2	Main Results	45
11	Convex Functions	49
11.1	Preliminaries	49
11.2	The Indicator Function	49
11.3	Definitions	52
11.4	Basic Properties	54
11.5	Differentiable Convex Functions	54
11.6	Convexity and Lipschitz-ness	57
11.7	Stability of Convexity	58
11.8	Examples	61
12	More Convex Functions	65
12.1	Strictly Convex	65
12.2	Strongly Convex	65
12.3	Quasiconvex	67
13	Support	69
13.1	Definitions	69
13.2	Properties	69
13.3	Supporting Hyperplane	72

14 Conjugacy	73
14.1 Definition and Examples	73
14.2 Basic Properties	74
14.3 Double Conjugate	74
14.4 Conjugates and Sub-Differentials	76
15 The Proximal Operator	79
15.1 Definitions	79
15.2 Examples	79
15.3 Basic Properties	79
15.4 Prox Calculus Rules	81
15.5 The Second Prox Theorem	81
15.6 Moreau Decomposition	83
16 Ellipsoids	85
16.1 Properties	85

Chapter 1

Affine Sets

1.1 Definitions

DEFINITION (Affine Combination). Let S be a set in \mathbb{E} . We define an **affine combination** of S to be a point x in the space of the form

$$x = \sum_{i=1}^n \lambda_i v_i$$

where (1) $n \in \mathbb{N}$, (2) $v_i \in S$ for all i , (3) $\lambda_i \in \mathbb{R}$ for all i , and (4) $\sum_{i=1}^n \lambda_i = 1$.

DEFINITION (Affine Span). Let S be a set in \mathbb{E} . We define the **affine span** of S , denoted by $\text{affspan}(S)$, to be the set of all affine combinations of S .

DEFINITION (Affine Set). Let S be a set in \mathbb{E} . We say that S is an **affine set** if $S = \text{aff}(S)$.

DEFINITION (Affine Hull). Let S be a set in \mathbb{E} . We define the **affine hull** of S , denoted by $\text{affhull}(S)$, to be the smallest affine set containing S .

Chapter 2

Relative Topology

2.1 Definitions

DEFINITION (Relative Interior). Let \mathbb{E} be some Euclidean space. Let S be a set in the space. We define the **relative interior** of S , denoted by $\text{ri}(S)$, to be the interior of S for the topology relative to the affine hull $\text{aff}(S)$. i.e., the set given by

$$\text{ri}(S) := \{x \in \text{aff}(S) : \exists r > 0, \text{ball}(x, r) \cap \text{aff}(S) \subseteq S\}.$$

A quick result. For a singleton set S , $\text{ri}(S) = S = \text{cl}(S)$.

2.2 Basic Properties

PROPOSITION 2.2.1. For any set S , we have $\text{ri}(S) \subseteq S$.

REMARK. The relative interior operator is not monotonic.

EXAMPLE 2.2.1. Consider \mathbb{R} with the usual topology and sets $\{0\}$ and $[0, 1]$. Then $\text{ri}(\{0\}) = \{0\}$ and $\text{ri}([0, 1]) = (0, 1)$.

PROPOSITION 2.2.2. Let S be a set in some Euclidean space \mathbb{E} . Then if $\text{int}(S) \neq \emptyset$, $\text{ri}(S) = \text{int}(S)$.

Proof.

It suffices to show that $\text{aff}(S) = \mathbb{R}^n$.

Since $\text{int}(S) \neq \emptyset$, $\exists x \in \text{int}(S)$.

Since $x \in \text{int}(S)$, $\exists r > 0$, $\text{ball}(x, r) \subseteq S$.

$\mathbb{E} = \text{aff}(\text{ball}(x, r)) \subseteq \text{aff}(S) \subseteq \mathbb{E}$.

This shows $\text{aff}(S) = \mathbb{E}$. ■

2.3 Arithmetic Properties

PROPOSITION 2.3.1. Let C_1 and C_2 be convex subsets of \mathbb{E} . Let λ_1 and λ_2 be scalars in \mathbb{R} . Then

$$\text{ri}(\lambda_1 C_1 + \lambda_2 C_2) = \lambda_1 \text{ri}(C_1) + \lambda_2 \text{ri}(C_2).$$

PROPOSITION 2.3.2. Let C_1 be a convex set in \mathbb{E}_1 . Let C_2 be a convex set in \mathbb{E}_2 . Then

$$\text{ri}(C_1 \oplus C_2) = \text{ri}(C_1) \oplus \text{ri}(C_2).$$

Chapter 3

Convex Sets

3.1 Definitions

DEFINITION (Convex Combination). Let \mathcal{V} be a vector space over field \mathbb{F} . Let S be a subset of \mathcal{V} . We define a **convex combination** of S to be a point x in \mathcal{V} of the form

$$x = \sum_{i=1}^n \lambda_i v_i$$

where (1) $n \in \mathbb{N}$, (2) $v_1, \dots, v_n \in S$, (3) $\lambda_1, \dots, \lambda_n \in \mathbb{R}_+$, and (4) $\sum_{i=1}^n \lambda_i = 1$.

DEFINITION (Convex Span). Let \mathbb{E} be a Euclidean space. Let S be a subset of \mathbb{E} . We define a **convex span** of S , denoted by $\text{convspan}(S)$, to be the set of all convex combinations of S .

DEFINITION (Convex). Let \mathbb{E} be a Euclidean space. Let S be a subset of \mathbb{E} . We say that S is **convex** if $S = \text{convspan}(S)$, or equivalently, if

$$\forall x, y \in S, \forall \alpha, \beta \in [0, 1] : \alpha + \beta = 1, \quad \alpha x + \beta y \in S.$$

DEFINITION (Pointed). Let \mathbb{E} be a Euclidean space. Let S be a subset of \mathbb{E} . We

say that S is **pointed** if S contains no line.

3.2 Arithmetic Properties of Convex Sets

PROPOSITION 3.2.1. Let C be a convex set. Let λ_1 and λ_2 be in \mathbb{R}_+ . Then

$$(\lambda_1 + \lambda_2)C = \lambda_1 C + \lambda_2 C.$$

Proof.

The case where any of λ_1 and λ_2 is 0 is trivial. I will assume that $\lambda_1, \lambda_2 > 0$.

For one direction, let x be an arbitrary point in $(\lambda_1 + \lambda_2)C$.

Since $x \in (\lambda_1 + \lambda_2)C$, $\exists c \in C, x = (\lambda_1 + \lambda_2)c$.

Since $\begin{cases} (\lambda_1 + \lambda_2)c = \lambda_1 c + \lambda_2 c \\ x = (\lambda_1 + \lambda_2)c \end{cases}$, we get $x = \lambda_1 c + \lambda_2 c$.

Since $\begin{cases} x = \lambda_1 c + \lambda_2 c \\ \lambda_1 c \in \lambda_1 C \\ \lambda_2 c \in \lambda_2 C \end{cases}$, we get $x \in \lambda_1 C + \lambda_2 C$.

Since $x \in \lambda_1 C + \lambda_2 C$ for any $x \in (\lambda_1 + \lambda_2)C$, $\lambda_1 C + \lambda_2 C \subseteq (\lambda_1 + \lambda_2)C$.

For the reverse direction, let x be an arbitrary point in $\lambda_1 C + \lambda_2 C$.

Since $x \in \lambda_1 C + \lambda_2 C$, $\exists c_1, c_2 \in C, x = \lambda_1 c_1 + \lambda_2 c_2$.

Define scalars $\mu_1 := \frac{\lambda_1}{\lambda_1 + \lambda_2}$ and $\mu_2 := \frac{\lambda_2}{\lambda_1 + \lambda_2}$.

Then $x = (\lambda_1 + \lambda_2)c$.

Since $\lambda_1, \lambda_2 > 0$, $\mu_1, \mu_2 \in [0, 1]$.

Define point $c := \mu_1 c_1 + \mu_2 c_2$.

Since $\begin{cases} c = \mu_1 c_1 + \mu_2 c_2 \\ c_1, c_2 \in C \\ \mu_1, \mu_2 \in [0, 1] \\ \mu_1 + \mu_2 = 1 \\ C \text{ is convex} \end{cases}$, we get $c \in C$.

Since $x = (\lambda_1 + \lambda_2)c$ and $c \in C$, $x \in (\lambda_1 + \lambda_2)C$.

Since $x \in (\lambda_1 + \lambda_2)C$ for any $x \in \lambda_1 C + \lambda_2 C$, $(\lambda_1 + \lambda_2)C \subseteq \lambda_1 C + \lambda_2 C$.

Since $\lambda_1 C + \lambda_2 C \subseteq (\lambda_1 + \lambda_2)C$ and $(\lambda_1 + \lambda_2)C \subseteq \lambda_1 C + \lambda_2 C$, $(\lambda_1 + \lambda_2)C = \lambda_1 C + \lambda_2 C$ ■

3.3 Topological Properties of Convex Sets

THEOREM 3.1. Let C be a convex set such that $\text{int}(C) \neq \emptyset$. Then

- (1) $\text{int}(C) = \text{int}(\text{cl}(C))$, and
- (2) $\text{cl}(C) = \text{cl}(\text{int}(C))$.

Proof of (1). $\text{int}(C) \subseteq \text{int}(\text{cl}(C))$ is clear. For $\text{int}(\text{cl}(C)) \subseteq \text{int}(C)$, let x be an arbitrary point in $\text{int}(\text{cl}(C))$.

Since $x \in \text{int}(\text{cl}(C))$,

$$\exists r > 0 \text{ such that } \text{ball}(x, r) \subseteq \text{cl}(C).$$

Since $\text{int}(C) \neq \emptyset$, pick $y \in \text{int}(C)$.

Define a scalar λ by

$$\lambda := \frac{r}{2\|x - y\|}.$$

Define a point z by

$$z := x + \lambda(x - y).$$

Since $\lambda = \frac{r}{2\|x - y\|}$ and $z = x + \lambda(x - y)$,

$$\begin{aligned} & \|z - x\| \\ &= \|x + \lambda(x - y) - x\| \\ &= \|\lambda(x - y)\| \\ &= \lambda\|x - y\| \\ &= \frac{r}{2\|x - y\|} \|x - y\| \\ &= \frac{r}{2} \\ &< r. \end{aligned}$$

That is,

$$\|z - x\| < r.$$

So $z \in \text{ball}(x, r)$. It follows that $z \in \text{cl}(C)$.

Since $z = x + \lambda(x - y)$, rearranging this yields

$$x = \frac{1}{1 + \lambda}z + \frac{\lambda}{1 + \lambda}y.$$

$$\text{Since } \begin{cases} x = \frac{1}{1+\lambda}z + \frac{\lambda}{1+\lambda}y \\ z \in \text{cl}(C) \\ y \in \text{int}(C) \\ \frac{1}{1+\lambda}, \frac{\lambda}{1+\lambda} \in (0, 1) \\ \frac{1}{1+\lambda} + \frac{\lambda}{1+\lambda} = 1 \end{cases}, \text{ by the lemma, we get}$$

$$x \in \text{int}(C).$$

Since $\forall x \in \text{int}(\text{cl}(C)), x \in \text{int}(C)$, we get $\text{int}(\text{cl}(C)) \subseteq \text{int}(C)$. ■

Proof of (2). $\text{cl}(\text{int}(C)) \subseteq \text{cl}(C)$ is clear. For $\text{cl}(C) \subseteq \text{cl}(\text{int}(C))$, let x be an arbitrary point in $\text{cl}(C)$.

Since $\text{int}(C) \neq \emptyset$, pick $y \in \text{int}(C)$.

Let $\lambda \in [0, 1)$.

Define a point z by

$$z(\lambda) := \lambda x + (1 - \lambda)y.$$

$$\text{Since } \begin{cases} z(\lambda) := \lambda x + (1 - \lambda)y \\ x \in \text{cl}(C) \\ y \in \text{int}(C) \\ \lambda \in [0, 1) \end{cases}, \text{ by the lemma, we get}$$

$$z(\lambda) \in \text{int}(C).$$

$$\text{Since } \begin{cases} z(\lambda) \in \text{int}(C) \\ \lim_{\lambda \rightarrow 1} z(\lambda) = x \end{cases}, \text{ we get}$$

$$x \in \text{cl}(\text{int}(C)).$$

Since $\forall x \in \text{cl}(C), x \in \text{cl}(\text{int}(C))$, we get $\text{cl}(C) \subseteq \text{cl}(\text{int}(C))$. ■

PROPOSITION 3.3.1. Let C be a convex set. Then

- (1) $\text{aff}(\text{ri}(C)) = \text{aff}(C) = \text{aff}(\text{cl}(C))$,
- (2) $\text{ri}(\text{ri}(C)) = \text{ri}(C) = \text{ri}(\text{cl}(C))$, and
- (3) $\text{cl}(\text{ri}(C)) = \text{cl}(C) = \text{cl}(\text{cl}(C))$.

PROPOSITION 3.3.2. Let C be a convex set. Then

$$C \neq \emptyset \iff \text{ri}(C) \neq \emptyset.$$

Proof. Forward Direction: Assume that $C \neq \emptyset$. I will show that $\text{ri}(C) \neq \emptyset$. Since $C \neq \emptyset$, $\text{aff}(C) \neq \emptyset$. Since C is convex, $\text{aff}(C) = \text{aff}(\text{ri}(C))$. Since $\begin{cases} \text{aff}(C) \neq \emptyset \\ \text{aff}(C) = \text{aff}(\text{ri}(C)) \end{cases}$, we get

$$\text{aff}(\text{ri}(C)) \neq \emptyset.$$

Since $\text{aff}(\text{ri}(C)) \neq \emptyset$, we get $\text{ri}(C) \neq \emptyset$.

Backward Direction: Assume that $\text{ri}(C) \neq \emptyset$. I will show that $C \neq \emptyset$. Since $\text{ri}(C) \neq \emptyset$ and $\text{ri}(C) \subseteq C$, we get $C \neq \emptyset$. ■

3.4 The Convex Hull Operator

(bug)

DEFINITION (Convex Hull). Let S be a set in \mathbb{E} . We define the **convex hull** of S , denoted by $\text{convhull}(S)$, to be the smallest convex set containing S .

PROPOSITION 3.4.1. For any set S , $\text{convspan}(S) = \text{convhull}(S)$. They will both be denoted by $\text{conv}(S)$ from now on.

Proof. Forward Direction: I will show that $\text{convspan}(S) \subseteq \text{convhull}(S)$. Let x be an arbitrary element of $\text{convspan}(S)$. We are to prove that $x \in \text{convhull}(S)$. Let C be an arbitrary convex set containing S . Since x is a convex combination of S , x is also a convex combination of C . Since x is a convex combination of C and C is convex, $x \in C$. Since x is in any convex set containing S , $x \in \text{convhull}(S)$. Since $x \in \text{convhull}(S)$ for any $x \in \text{convspan}(S)$, $\text{convspan}(S) \subseteq \text{convhull}(S)$.

Backward Direction: I will show that $\text{convhull}(S) \subseteq \text{convspan}(S)$.

proof incomplete. ■

not finished

PROPOSITION 3.4.2 (The Convex Hull Operator). Let \mathbb{E} be a Euclidean space.

(1) Expansive

$$\forall S \subseteq \mathbb{E}, \quad S \subseteq \text{conv}(S).$$

(2) Monotonic Increasing

$$\forall S_1, S_2 \subseteq \mathbb{E} : S_1 \subseteq S_2, \quad \text{conv}(S_1) \subseteq \text{conv}(S_2).$$

(3) Idempotent

$$\forall S \subseteq \mathbb{E}, \quad \text{conv}(\text{conv}(S)) = \text{conv}(S).$$

PROPOSITION 3.4.3 (Bounded). The convex hull of a bounded set is bounded.

PROPOSITION 3.4.4 (Open). The convex hull of an open set is open.

Proof. Let \mathcal{V} be a topological vector space. Let G be an open subset of \mathcal{V} . I will show that $\text{conv}(G)$ is open. Let $\sum_{i=1}^n \lambda_i x_i$ be an arbitrary convex combination of elements of G . Let $i_0 \in \{1..n\}$ be such that $\lambda_{i_0} \neq 0$. Then

$$\sum_{i=1}^n \lambda_i x_i \in \sum_{i \neq i_0} \lambda_i x_i + i_0 G \subseteq \text{conv}(G).$$

So

$$\text{conv}(G) = \bigcup \left\{ \sum_{i \neq i_0} \lambda_i x_i + i_0 G \right\}.$$

Note that the function $f(x) := \sum_{i \neq i_0} \lambda_i x_i + i_0 x$ is a homeomorphism. So $\text{conv}(G)$ is a union of open sets and hence open. ■

REMARK (Closed). The convex hull of a closed set need not be closed.

EXAMPLE 3.4.1. The set $S := \{(x, y) \in \mathbb{R}^2 : y \geq \frac{1}{1+x^2}\}$ is closed. However, $\text{conv}(S) = \{(x, y) \in \mathbb{R}^2 : y > 0\}$ is open.

PROPOSITION 3.4.5 (Compact). The convex hull of a compact set is compact.

3.5 The Closed Convex Hull Operator

DEFINITION (Closed Convex Hull). Let S be a set in some Euclidean space. We define the **closed convex hull** of S , denoted by $\overline{\text{conv}}(S)$, to be the smallest closed convex containing S .

PROPOSITION 3.5.1. The closed convex hull is the closure of the convex hull.

PROPOSITION 3.5.2. A closed convex hull does not distinguish a set from its closure. i.e., for any set S , we have $\overline{\text{conv}}(S) = \overline{\text{conv}}(\text{cl}(S))$.

PROPOSITION 3.5.3. If S is bounded, then the closure operation and the convex hull operation commute. i.e., $\text{conv}(\text{cl}(S)) = \text{cl}(\text{conv}(S))$.

REMARK. The closure operation and the convex hull operation do not commute in general.

3.6 Stability of Convexity

PROPOSITION 3.6.1 (Intersection). Convexity is stable under intersection. i.e., the intersection of any collection of convex sets is convex.

Proof. Let $\{C_i\}_{i \in I}$ be an arbitrary collection of convex sets where I is an index set and C_i is convex for any $i \in I$. Let C denote their intersection. If $C = \emptyset$, then we are done. Else, let x and y be two arbitrary points in C . Let λ be an arbitrary number in $(0, 1)$. Define a point $z := \lambda x + (1 - \lambda)y$. Since $x \in C$ and $C = \bigcap_{i \in I} C_i$, we get $x \in C_i$ for any $i \in I$. Since $y \in C$ and $C = \bigcap_{i \in I} C_i$, we get $y \in C_i$ for any $i \in I$. Let i be an arbitrary index in I . Since $x \in C_i$ and $y \in C_i$ and $\lambda \in (0, 1)$ and C_i is convex and $z = \lambda x + (1 - \lambda)y$, we get $z \in C_i$. Since $z \in C_i$ for any $i \in I$ and $C = \bigcap_{i \in I} C_i$, we get $z \in C$. Since

$$\forall x, y \in C, \forall \lambda \in (0, 1), \quad \lambda x + (1 - \lambda)y \in C,$$

by definition of convex sets, we get C is convex. ■

PROPOSITION 3.6.2 (Affine Map). Convexity is stable under affine mapping. i.e., the affine image of a convex set is convex.

PROPOSITION 3.6.3 (Linear Combinations). Convexity is stable under linear combinations. i.e., if C_1 and C_2 are convex sets and λ_1 and λ_2 are real numbers, then the set C defined as

$$C := \lambda_1 C_1 + \lambda_2 C_2$$

is convex.

Proof. If $C_1 = \emptyset$ or $C_2 = \emptyset$, then $\lambda_1 C_1 + \lambda_2 C_2 = \emptyset$ and we are done. Now assume that $C_1, C_2 \neq \emptyset$. Then $C = \lambda_1 C_1 + \lambda_2 C_2 \neq \emptyset$. Let x and y be arbitrary points in C .

Since $x \in C$, $\exists x_1 \in C_1, x_2 \in C_2$ such that $x = \lambda_1 x_1 + \lambda_2 x_2$.

Since $y \in C$, $\exists y_1 \in C_1, y_2 \in C_2$ such that $y = \lambda_1 y_1 + \lambda_2 y_2$.

Let $\lambda \in [0, 1]$ be arbitrary. Define a point z as $z := \lambda x + (1 - \lambda)y$. Then

$$\begin{aligned} z &= \lambda x + (1 - \lambda)y \\ &= \lambda(\lambda_1 x_1 + \lambda_2 x_2) + (1 - \lambda)(\lambda_1 y_1 + \lambda_2 y_2) \\ &= \lambda_1(\lambda x_1 + (1 - \lambda)y_1) + \lambda_2(\lambda x_2 + (1 - \lambda)y_2). \end{aligned}$$

Since $x_1, y_1 \in C_1$, $\lambda \in [0, 1]$ and C_1 is convex, we get $\lambda x_1 + (1 - \lambda)y_1 \in C_1$.

Since $x_2, y_2 \in C_2$, $\lambda \in [0, 1]$ and C_2 is convex, we get $\lambda x_2 + (1 - \lambda)y_2 \in C_2$.

So $z = \lambda_1(\lambda x_1 + (1 - \lambda)y_1) + \lambda_2(\lambda x_2 + (1 - \lambda)y_2) \in \lambda_1 C_1 + \lambda_2 C_2$.

That is, $\forall x \in C, \forall y \in C, \forall \lambda \in [0, 1]$, we have $\lambda x + (1 - \lambda)y \in C$.

So by definition, C is convex. ■

COROLLARY 3.1. The Minkowski sum of two convex sets is convex.

LEMMA 3.1. Let C be a convex set in \mathbb{E} . Let $x \in \text{int}(C)$. Let $y \in \text{cl}(C)$. Then

$$\forall \lambda \in (0, 1], \quad \lambda x + (1 - \lambda)y \in C.$$

Proof.

Since $x \in \text{int}(S)$, there exists some radius r_x such that $\text{ball}(x, r_x) \subseteq S$.

Define $r_z := \lambda r_x$.

Let z' be an arbitrary point in $\text{ball}(z, r_z)$.

Define $x' := \frac{1}{\lambda}(z' - (1 - \lambda)y)$.

Notice

$$\begin{aligned} & \|x - x'\| \\ &= \frac{1}{|\lambda|} \|\lambda x - \lambda x'\| \\ &= \frac{1}{|\lambda|} \|(z - (1 - \lambda)y) - (z' - (1 - \lambda)y)\| \\ &= \frac{1}{|\lambda|} \|z - z'\| \\ &\leq \frac{1}{|\lambda|} r_z, \text{ since } z' \in \text{ball}(z, r_z) \\ &= \frac{1}{|\lambda|} \lambda r_x \\ &= r_x. \end{aligned}$$

That is,

$$\|x - x'\| \leq r_x.$$

So $x' \in \text{ball}(x, r_x)$.

Since $x' \in \text{ball}(x, r_x)$ and $\text{ball}(x, r_x) \subseteq S$, we get $x' \in S$.

$$\text{Since } \begin{cases} z' = \lambda x' + (1 - \lambda)y \\ x', y \in S \\ \lambda \in (0, 1] \\ S \text{ is convex} \end{cases}, \text{ we get } z' \in S.$$

Since $z' \in S$ for any $z' \in \text{ball}(z, r_z)$, $\text{ball}(z, r_z) \subseteq S$.

Since there exists some radius r_z such that $\text{ball}(z, r_z) \subseteq S$, $z \in \text{int}(S)$.

Alternative Expressing:

Define $B := \text{ball}(0, 1)$.

$$\begin{aligned}
 & (1 - \lambda)x + \lambda y + \varepsilon B \\
 & \subseteq (1 - \lambda)x + \lambda(C + \varepsilon B) + \varepsilon B \\
 & = (1 - \lambda)x + \lambda C + \lambda \varepsilon B + \varepsilon B \\
 & = (1 - \lambda)x + (1 + \lambda)\varepsilon B + \lambda C \\
 & = (1 - \lambda)\left(x + \frac{1 + \lambda}{1 - \lambda}\varepsilon B\right) + \lambda C \\
 & \subseteq (1 - \lambda)C + \lambda C \\
 & = C.
 \end{aligned}$$

■

LEMMA 3.2. Let C be a convex set in \mathbb{E} . Let $x \in \text{ri}(C)$. Let $y \in \text{cl}(C)$. Then

$$\forall \lambda \in (0, 1], \quad \lambda x + (1 - \lambda)y \in C.$$

Proof.

Case 1. $\text{int}(C) \neq \emptyset$.

Then $\text{int}(C) = \text{ri}(C)$.

Since $x \in \text{int}(C)$ and $y \in \text{cl}(C)$, $\forall t \in (0, 1]$, $z := tx + (1 - t)y \in C$.

Case 2. $\text{int}(C) = \emptyset$.

Now $\dim(C) < d$.

Say $\dim(C) = l$.

Apply case 1 in \mathbb{R}^l .

■

PROPOSITION 3.6.4 (Interior). Convexity is stable under interior. i.e., the interior of a convex set is convex.

Proof. Let S be a convex set in \mathbb{E} . If $\text{int}(S) = \emptyset$, then we are done. Else: let x and y be two arbitrary points in $\text{int}(S)$. Let λ be an arbitrary number in $(0, 1)$. Define a point z by $z := \lambda x + (1 - \lambda)y$. Since $x, y \in \text{int}(S)$ and $\lambda \in (0, 1)$, by the lemma, we get $z \in \text{int}(S)$. Since

$$\forall x, y \in \text{int}(S), \forall \lambda \in (0, 1), \quad \lambda x + (1 - \lambda)y \in \text{int}(S),$$

we get $\text{int}(S)$ is convex.

■

PROPOSITION 3.6.5 (Relative Interior). Convexity is stable under relative interior. i.e., the relative interior of a convex set is convex.

Proof. Let S be a convex set in \mathbb{E} . If $\text{ri}(S) = \emptyset$, then we are done. Else: let x and y be two arbitrary points in $\text{ri}(S)$. Let λ be an arbitrary number in $(0, 1)$. Define a point z by $z := \lambda x + (1 - \lambda)y$. Since $x, y \in \text{ri}(S)$ and $\lambda \in (0, 1)$, by the lemma, we get $z \in \text{ri}(S)$. Since

$$\forall x, y \in \text{ri}(S), \forall \lambda \in (0, 1), \quad \lambda x + (1 - \lambda)y \in \text{ri}(S),$$

we get $\text{ri}(S)$ is convex. ■

PROPOSITION 3.6.6 (Closure). Convexity is stable under closure. i.e., the closure of a convex set is convex.

Proof Approach 1.

Let $x, y \in \text{cl}(C)$.

Let $t \in [0, 1]$.

Since $x \in \text{cl}(C)$, $\exists \{x_i\}_{i \in \mathbb{N}} \subseteq C, \lim_{i \rightarrow \infty} x_i = x$.

Since $y \in \text{cl}(C)$, $\exists \{y_i\}_{i \in \mathbb{N}} \subseteq C, \lim_{i \rightarrow \infty} y_i = y$.

Since $\lim_{i \rightarrow \infty} x_i = x$ and $\lim_{i \rightarrow \infty} y_i = y$, $\lim_{i \rightarrow \infty} (tx_i + (1 - t)y_i) = tx + (1 - t)y$.

Since $x_i, y_i \in C$ and C is convex, $tx_i + (1 - t)y_i \in C$.

Since $tx_i + (1 - t)y_i \in C$, $\lim_{i \rightarrow \infty} (tx_i + (1 - t)y_i) = tx + (1 - t)y$, $tx + (1 - t)y \in \text{cl}(C)$.

Since $\forall x, y \in \text{cl}(C), \forall t \in [0, 1], tx + (1 - t)y \in \text{cl}(C)$, we get $\text{cl}(C)$ is convex. ■

Proof Approach 2.

$\text{cl}(C) = \bigcap_{\varepsilon > 0} [C + \varepsilon \text{ball}(0, 1)]$. This is an intersection of linear combinations of convex sets and hence convex. ■

PROPOSITION 3.6.7 (Conical Hull). Convexity is stable under conical hull. i.e., if C is convex, then $\text{cone}(C)$ is convex.

Proof.

Let x and y be arbitrary points in $\text{cone}(C)$.

Let λ be an arbitrary number in $(0, 1)$.

Define point z as $z := \lambda x + (1 - \lambda)y$.

Since $x \in \text{cone}(C)$, $\exists x' \in C$ and $\exists \alpha > 0$ such that $x = \alpha x'$.

Since $y \in \text{cone}(C)$, $\exists y' \in C$ and $\exists \beta > 0$ such that $y = \beta y'$.

Define point z' as $z' := \frac{\lambda\alpha}{\lambda\alpha + (1-\lambda)\beta}x' + \frac{(1-\lambda)\beta}{\lambda\alpha + (1-\lambda)\beta}y'$.

Since $x', y' \in C$ and $\frac{\lambda\alpha}{\lambda\alpha + (1-\lambda)\beta} \in (0, 1)$ and $\frac{\lambda\alpha}{\lambda\alpha + (1-\lambda)\beta} + \frac{(1-\lambda)\beta}{\lambda\alpha + (1-\lambda)\beta} = 1$ and C is convex and $z' := \frac{\lambda\alpha}{\lambda\alpha + (1-\lambda)\beta}x' + \frac{(1-\lambda)\beta}{\lambda\alpha + (1-\lambda)\beta}y'$, we get $z' \in C$.

Since $z' \in C$ and $z = (\lambda\alpha + (1 - \lambda)\beta)z'$, $z \in \text{cone}(C)$.

That is, $\lambda x + (1 - \lambda)y \in \text{cone}(C)$.

Since $\forall x, y \in \text{cone}(C)$, $\forall \lambda \in (0, 1)$, $\lambda x + (1 - \lambda)y \in \text{cone}(C)$, we get $\text{cone}(C)$ is convex. ■

3.7 Examples of Convex Sets

EXAMPLE 3.7.1. Let I be an index set. Let b_i for $i \in I$ be vectors in \mathbb{E} . Let β_i for $i \in I$ be reals. Then the set C given by

$$C := \{x \in \mathbb{E} : \forall i \in I, \langle x, b_i \rangle \leq \beta_i\}$$

is convex.

Proof.

Each of $C_i := \{x \in \mathbb{E} : \langle x, b_i \rangle \leq \beta_i\}$ is convex and $C = \bigcap_{i \in I} C_i$.

$$\begin{aligned} \langle z, b_i \rangle &= \langle \lambda x + (1 - \lambda)y, b_i \rangle \\ &= \lambda \langle x, b_i \rangle + (1 - \lambda) \langle y, b_i \rangle \\ &\leq \lambda \beta_i + (1 - \lambda) \beta_i \\ &= \beta_i. \end{aligned}$$
■

3.8 The Carathéodory Theorem

THEOREM 3.2 (Carathéodory). Let \mathbb{E} be some Euclidean space. Let S be some set in the space. Let x be some point in $\text{conv}(S)$. Then x can be represented as a convex

combination of at most $d + 1$ points in S . i.e., x lies in some r -simplex with vertices in S , where $r \leq d$.

Chapter 4

Geometric Objects

4.1 Definitions

DEFINITION (Hyperplane). Let \mathbb{E} be a Euclidean space over \mathbb{R} . Let H be a subset of \mathbb{E} . We say that H is a **hyperplane** if and only if H can be expressed as

$$H = \{x \in \mathbb{E} : a^\top x = b\}$$

for some $a \in \mathbb{E} \setminus \{0\}$ and $b \in \mathbb{R}$.

DEFINITION (Closed Half-Space). Let \mathbb{E} be a Euclidean space over \mathbb{R} . Let P be a subset of \mathbb{E} . We say that P is a **closed half-space** if and only if P can be expressed as

$$P = \{x \in \mathbb{E} : a^\top x \leq b\}$$

for some $a \in \mathbb{E} \setminus \{0\}$ and $b \in \mathbb{R}$.

DEFINITION (Polyhedron). Let \mathbb{E} be a Euclidean space over \mathbb{R} . Let P be a subset of \mathbb{E} . We say that P is a **polyhedron** if and only if P can be expressed as the intersection of finitely many closed half-spaces in \mathbb{E} .

4.2 Properties

PROPOSITION 4.2.1. Polyhedrons are convex.

Chapter 5

Cones (in Analysis)

5.1 Definitions

DEFINITION (Conical Combination). Let S be a set in \mathbb{E} . We define a **conical combination** of S to be a point x in the space of the form

$$x = \sum_{i=1}^n \lambda_i v_i$$

where (1) $n \in \mathbb{N}$, (2) $v_i \in S$ for all i , and (3) $\lambda_i \in \mathbb{R}_{++}$ for all i .

DEFINITION (Cone). Let S be a set in \mathbb{E} . We say that S is a **cone** if $S = \mathbb{R}_{++}S$.

DEFINITION (Conical Hull). Let S be a set in \mathbb{E} . We define the **conical hull** of S , denoted by $\text{cone}(S)$, to be the intersection of all cones containing C .

PROPOSITION 5.1.1. Let S be a set in \mathbb{E} . Then $\text{cone}(S) = \mathbb{R}_{++}S$.

Proof. **Forward Direction:** I will show that $\text{cone}(S) \subseteq \mathbb{R}_{++}S$. Since $\mathbb{R}_{++}\mathbb{R}_{++}S = \mathbb{R}_{++}S$, $\mathbb{R}_{++}S$ is a cone. Since $1 \in \mathbb{R}_{++}$, $S \subseteq \mathbb{R}_{++}S$. Since $\mathbb{R}_{++}S$ is a cone containing S and $\text{cone}(S)$ is the smallest cone containing S , we get

$$\text{cone}(S) \subseteq \mathbb{R}_{++}S.$$

Backward Direction: I will show that $\mathbb{R}_{++}S \subseteq \text{cone}(S)$. Let C be an arbitrary cone containing S . Since $S \subseteq C$, $\mathbb{R}_{++}S \subseteq \mathbb{R}_{++}C$. Since C is a cone, $\mathbb{R}_{++}C = C$. So $\mathbb{R}_{++}S \subseteq C$. Since $\mathbb{R}_{++}S \subseteq C$ for any cone C containing S , we get

$$\mathbb{R}_{++}S \subseteq \text{cone}(S).$$

■

DEFINITION (Closed Conical Hull). Let S be a set in \mathbb{E} . We define the **closed conical hull** of S , denoted by $\text{clcone}(S)$, to be the intersection of all closed cones containing S .

PROPOSITION 5.1.2. For any set S in \mathbb{E} , we have

$$\text{clcone}(S) = \text{cl}(\text{cone}(S)).$$

Proof.

For $\text{clcone}(S) \subseteq \text{cl}(\text{cone}(S))$.

Since $\text{cl}(\text{cone}(S))$ is a closed cone containing S and $\text{clcone}(S)$ is the smallest closed cone containing S , $\text{clcone}(S) \subseteq \text{cl}(\text{cone}(S))$.

For $\text{cl}(\text{cone}(S)) \subseteq \text{clcone}(S)$.

Since $S \subseteq \text{clcone}(S)$, by the monotonicity of the cone operator, $\text{cone}(S) \subseteq \text{cone}(\text{clcone}(S))$.

Since $\text{cone}(S) \subseteq \text{cone}(\text{clcone}(S))$, by the monotonicity of the closure operator, $\text{cl}(\text{cone}(S)) \subseteq \text{cl}(\text{cone}(\text{clcone}(S)))$.

Since $\text{clcone}(S)$ is a cone, $\text{cone}(\text{clcone}(S)) = \text{clcone}(S)$.

Since $\text{clcone}(S)$ is closed, $\text{cl}(\text{clcone}(S)) = \text{clcone}(S)$.

Since $\text{cone}(\text{clcone}(S)) = \text{clcone}(S)$ and $\text{cl}(\text{clcone}(S)) = \text{clcone}(S)$, we get $\text{cl}(\text{cone}(\text{clcone}(S))) = \text{clcone}(S)$.

Since $\text{cl}(\text{cone}(S)) \subseteq \text{cl}(\text{cone}(\text{clcone}(S)))$ and $\text{cl}(\text{cone}(\text{clcone}(S))) = \text{clcone}(S)$, we get $\text{cl}(\text{cone}(S)) \subseteq \text{clcone}(S)$.

Since $\text{clcone}(S) \subseteq \text{cl}(\text{cone}(S))$ and $\text{cl}(\text{cone}(S)) \subseteq \text{clcone}(S)$, we get $\text{clcone}(S) = \text{cl}(\text{cone}(S))$.

■

DEFINITION (Ray). Let \mathbb{E} be a Euclidean space. Let R be a subset of \mathbb{E} . We say that R is a **ray** if R can be expressed as

$$R = \{\alpha v : \alpha \in \mathbb{R}_+\}$$

for some $v \in \mathbb{E} \setminus \{0\}$.

DEFINITION (Extreme Ray). Let \mathbb{E} be a Euclidean space. Let K be a cone in \mathbb{E} . Let R be a ray in K . We say that R is an **extreme ray** in K if for any pair of rays R_1 and R_2 in K such that $R_1 + R_2 \supseteq R$, we have either $R_1 = R$ or $R_2 = R$ (or both).

5.2 The cone Operator

PROPOSITION 5.2.1 (The cone Operator). The cone operator has the following properties.

$$(1) \quad \forall S \subseteq \mathbb{E},$$

$$S \subseteq \text{cone}(S).$$

$$(2) \quad \forall S_1, S_2 \subseteq \mathbb{E},$$

$$S_1 \subseteq S_2 \implies \text{cone}(S_1) \subseteq \text{cone}(S_2).$$

$$(3) \quad \forall S \subseteq \mathbb{E},$$

$$\text{cone}(\text{cone}(S)) = \text{cone}(S).$$

PROPOSITION 5.2.2. The conv operator and the cone operator commute. Let S be a set in \mathbb{E} . Then

$$\text{conv}(\text{cone}(S)) = \text{cone}(\text{conv}(S)).$$

Proof.

For $\text{cone}(\text{conv}(S)) \subseteq \text{conv}(\text{cone}(S))$, let x be an arbitrary point in $\text{cone}(\text{conv}(S))$.

Since $x \in \text{cone}(\text{conv}(S))$, we get $\exists \lambda \in \mathbb{R}_+$, $\exists n \in \mathbb{N}$, $\exists v_1, \dots, v_n \in S$, $\exists \mu_1, \dots, \mu_n \in [0, 1]$, $\sum_{i=1}^n \mu_i = 1$ such that $x = \lambda \sum_{i=1}^n \mu_i v_i$.

Since $x = \lambda \sum_{i=1}^n \mu_i v_i$, $x = \sum_{i=1}^n \mu_i (\lambda v_i)$.

Since $\lambda \in \mathbb{R}_+$ and $v_i \in S$, $\lambda v_i \in \text{cone}(S)$.

Since $\lambda v_i \in \text{cone}(S)$ and $\mu_i \in [0, 1]$, $\sum_{i=1}^n \mu_i = 1$, $\sum_{i=1}^n \mu_i (\lambda v_i) \in \text{conv}(\text{cone}(S))$.

Since $\forall x \in \text{cone}(\text{conv}(S))$, $x \in \text{conv}(\text{cone}(S))$, $\text{cone}(\text{conv}(S)) \subseteq \text{conv}(\text{cone}(S))$.

For $\text{conv}(\text{cone}(S)) \subseteq \text{cone}(\text{conv}(S))$, let x be an arbitrary point in $\text{conv}(\text{cone}(S))$.

Since $x \in \text{conv}(\text{cone}(S))$, $\exists n \in \mathbb{N}$, $\exists \lambda_i \in [0, 1]$, $\sum_{i=1}^n \lambda_i = 1$, $\exists \mu_i \in \mathbb{R}_+$, $\exists v_i \in S$ such that $x = \sum_{i=1}^n \lambda_i \mu_i v_i$.

Define $\alpha := \sum_{i=1}^n \lambda_i \mu_i$.

Define $\beta_i := \lambda_i \mu_i / \alpha$.

Then $\alpha \in \mathbb{R}_+$ and $\beta_i \in [0, 1]$ and $\sum_{i=1}^n \beta_i = 1$ and $x = \alpha \sum_{i=1}^n \beta_i v_i$.

Since $\beta_i \in [0, 1]$ and $\sum_{i=1}^n \beta_i = 1$ and $v_i \in S$, we get $\sum_{i=1}^n \beta_i v_i \in \text{conv}(S)$.

Since $\alpha \in \mathbb{R}_+$ and $\sum_{i=1}^n \beta_i v_i \in \text{conv}(S)$ and $x = \alpha \sum_{i=1}^n \beta_i v_i$, we get $x \in \text{cone}(\text{conv}(S))$.

Since $\forall x \in \text{conv}(\text{cone}(S)), x \in \text{cone}(\text{conv}(S))$, we get $\text{conv}(\text{cone}(S)) \subseteq \text{cone}(\text{conv}(S))$.

Since $\text{cone}(\text{conv}(S)) \subseteq \text{conv}(\text{cone}(S))$ and $\text{conv}(\text{cone}(S)) \subseteq \text{cone}(\text{conv}(S))$, we get $\text{conv}(\text{cone}(S)) = \text{cone}(\text{conv}(S))$. ■

5.3 Other Properties

PROPOSITION 5.3.1. Let C be a convex set in \mathbb{E} . Assume $\text{int}(C) \neq \emptyset$ and $0 \in C$. Then $\text{int}(\text{cone}(C)) = \text{cone}(\text{int}(C))$.

Proof.

For one direction, let x be an arbitrary point in $\text{int}(\text{cone}(C))$. We are to prove that $x \in \text{cone}(\text{int}(C))$.

Since $x \in \text{int}(\text{cone}(C))$, $\exists r$ such that $\text{ball}(x, r) \subseteq \text{cone}(C)$.

Since $x \in \text{int}(\text{cone}(C))$, $x \in \text{cone}(C)$.

Since $x \in \text{cone}(C)$, $\exists n \in \mathbb{N}$, $\exists \lambda_1, \dots, \lambda_n > 0$, $\exists v_1, \dots, v_n \in C$ such that $x = \sum_{i=1}^n \lambda_i v_i$.

Assume for the sake of contradiction that $\exists k \in \{1, \dots, n\}$ such that $\forall r_k > 0$, $\text{ball}(v_k, r_k) \cap \mathbb{E} \setminus C \neq \emptyset$.

not finished

For the reverse direction, let x be an arbitrary point in $\text{cone}(\text{int}(C))$. We are to prove that $x \in \text{int}(\text{cone}(C))$.

Since $x \in \text{cone}(\text{int}(C))$, $\exists n \in \mathbb{N}$, $\exists \lambda_1, \dots, \lambda_n > 0$, $\exists v_1, \dots, v_n \in \text{int}(C)$ such that $x = \sum_{i=1}^n \lambda_i v_i$.

Since $v_i \in \text{int}(C)$ for each $i \in \{1, \dots, n\}$, $\exists r_i$ such that $\text{ball}(v_i, r_i) \subseteq C$.

Define $R := \min\{\lambda_i r_i\}_{i=1}^n$.

Say $R = \lambda_k r_k$ for some $k \in \{1, \dots, n\}$.

Let y be an arbitrary point in $\text{ball}(x, R)$.

Since $y \in \text{ball}(x, R)$, $\exists w$ such that $\|w\| < R$ and $y = x + w$.

$$\begin{aligned}
y &= \sum_{i=1}^n \lambda_i v_i + w \\
&= \sum_{i \neq k} \lambda_i v_i + \lambda_k v_k + w \\
&= \sum_{i \neq k} \lambda_i v_i + \lambda_k (v_k + w/\lambda_k).
\end{aligned}$$

Since $\|w\| < R$, $\|w/\lambda_k\| < R/\lambda_k = r_k$.

Since $\|w/\lambda_k\| < r_k$, $v_k + w/\lambda_k \in \text{ball}(v_k, r_k)$.

So $v_k + w/\lambda_k \in C$.

So $y \in \text{cone}(C)$.

Since $\forall y \in \text{ball}(x, R)$, $y \in \text{cone}(C)$, $\text{ball}(x, R) \subseteq \text{cone}(C)$.

Since $\exists r$ such that $\text{ball}(x, r) \subseteq \text{cone}(C)$, $x \in \text{int}(\text{cone}(C))$.

This proves $\text{cone}(\text{int}(C)) \subseteq \text{int}(\text{cone}(C))$.

■

PROPOSITION 5.3.2. Let C be a convex set in \mathbb{E} . Assume $\text{int}(C) \neq \emptyset$ and $0 \in C$. Then

$$0 \in \text{int}(C) \iff \text{cone}(C) = \mathbb{E}.$$

Proof. For one direction, assume that $0 \in \text{int}(C)$. We are to prove that $\text{cone}(C) = \mathbb{E}$. Clearly

$$\text{cone}(C) \subseteq \mathbb{E}.$$

Since $0 \in \text{int}(C)$, $\exists r > 0$ such that $\text{ball}(0, r) \subseteq C$. Since $\text{ball}(0, r) \subseteq C$, $\text{cone}(\text{ball}(0, r)) \subseteq \text{cone}(C)$. Since $\text{cone}(\text{ball}(0, r)) = \mathbb{E}$ and $\text{cone}(\text{ball}(0, r)) \subseteq \text{cone}(C)$, we get

$$\mathbb{E} \subseteq \text{cone}(C).$$

For the reverse direction, assume that $\text{cone}(C) = \mathbb{E}$. We are to prove that $0 \in \text{int}(C)$.

$$\mathbb{E} = \text{int}(\mathbb{E}) = \text{int}(\text{cone}(C)) = \text{cone}(\text{int}(C)).$$

If $0 \notin \text{int}(C)$, then $0 \notin \text{cone}(\text{int}(C))$. So $0 \in \text{int}(C)$.

■

Chapter 6

Tangent Cones and Normal Cones

6.1 Definitions

DEFINITION (Tangent Cones). Let C be a non-empty convex set in \mathbb{E} . Let x be a point in \mathbb{E} . We define the **tangent cone** to C at point x , denoted by $T_C(x)$, to be a set given by

$$T_C(x) := \begin{cases} \text{clcone}(C - x), & \text{if } x \in C, \text{ or} \\ \emptyset, & \text{if } x \notin C. \end{cases}$$

DEFINITION (Normal Cones). Let C be a non-empty convex set in \mathbb{E} . Let x be a point in \mathbb{E} . We define the **normal cone** to C at point x , denoted by $N_C(x)$, to be a set given by

$$N_C(x) := \begin{cases} \{v \in \mathbb{E} : \forall y \in C - x, \langle y, v \rangle \leq 0\}, & \text{if } x \in C, \text{ or} \\ \emptyset, & \text{if } x \notin C. \end{cases}$$

6.2 Basic Properties

PROPOSITION 6.2.1. Let C be a closed convex set in \mathbb{E} . Let x be a point in \mathbb{E} . Then $T_C(x)$ and $N_C(x)$ are closed convex cones.

Proof.

If $C = \emptyset$, then $T_C(x) = N_C(x) = \emptyset$.

If $C \neq \emptyset$ and $x \notin C$, then $T_C(x) = N_C(x) = \emptyset$.

So now I assume that $C \neq \emptyset$ and $x \in C$.

Tangent Cone is Closed:

By definition, $T_C(x) = \text{clcone}(C - x)$. So $T_C(x)$ is a closed.

Tangent Cone is Convex:

C is convex	
\Downarrow	since affine mapping preserves convexity
$C - x$ is convex	
\Downarrow	since the cone operator preserves convexity
$\text{cone}(C - x)$ is convex	
\Downarrow	since the cl operator preserves convexity
$\text{cl}(\text{cone}(C - x))$ is convex	
\Downarrow	since $\text{cl} \circ \text{cone} = \text{clcone}$
$\text{clcone}(C - x)$ is convex	

That is, $T_C(x)$ is convex.

Tangent Cone is a Cone

By definition, $T_C(x) = \text{clcone}(C - x)$. So $T_C(x)$ is a cone.

Normal Cone is Closed:

Let $\{x_i\}_{i \in \mathbb{N}}$ be an arbitrary sequence in $N_C(x)$ that converges to some point in \mathbb{E} .

Say $x_i \rightarrow x_\infty$.

Let y be an arbitrary point in $C - x$.

Since $x_i \in N_C(x)$ and $y \in C - x$, by definition of $N_C(x)$, we get $\langle x_i, y \rangle \leq 0$.

Since $\langle x_i, y \rangle \leq 0$ for any $i \in \mathbb{N}$ and $x_i \rightarrow x_\infty$, we get $\langle x_\infty, y \rangle \leq 0$.

Since $\forall y \in C - x$, $\langle x_\infty, y \rangle \leq 0$, by definition of $N_C(x)$, we get $x_\infty \in N_C(x)$.

Since any convergent sequence whose terms are in $N_C(x)$ has its limit also in $N_C(x)$, $N_C(x)$ is closed.

Normal Cone is Convex:

Let u and v be arbitrary points in $N_C(x)$.

Let λ be an arbitrary number in $(0, 1)$.

Define point z as $z := \lambda u + (1 - \lambda)v$.

Let y be an arbitrary point in $C - x$.

Since $u \in N_C(x)$, $\langle u, y \rangle \leq 0$.

Since $v \in N_C(x)$, $\langle v, y \rangle \leq 0$.

$$\begin{aligned} & \langle z, y \rangle \\ &= \langle \lambda u + (1 - \lambda)v, y \rangle \\ &= \lambda \langle u, y \rangle + (1 - \lambda) \langle v, y \rangle \\ &\leq \lambda 0 + (1 - \lambda) 0 \\ &= 0. \end{aligned}$$

That is, $\langle z, y \rangle \leq 0$.

Since $\forall y \in C - x$, $\langle z, y \rangle \leq 0$, we get $z \in N_C(x)$.

That is, $\lambda u + (1 - \lambda)v \in N_C(x)$.

Since $\forall u, v \in N_C(x)$, $\forall \lambda \in (0, 1)$, $\lambda u + (1 - \lambda)v \in N_C(x)$, we get $N_C(x)$ is convex.

Normal Cone is a Cone:

Let v be an arbitrary point in $N_C(x)$.

Let λ be an arbitrary number such that $\lambda > 0$.

Let y be an arbitrary point in $C - x$.

Since $v \in N_C(x)$, $\langle v, y \rangle \leq 0$.

Since $\langle v, y \rangle \leq 0$ and $\lambda > 0$, $\langle \lambda v, y \rangle \leq 0$.

Since $\forall y \in C - x$, $\langle \lambda v, y \rangle \leq 0$, we get $\lambda v \in N_C(x)$.

Since $\forall v \in N_C(x)$, $\forall \lambda > 0$, $\lambda v \in N_C(x)$, we get $N_C(x)$ is a cone. ■

PROPOSITION 6.2.2. Let C be a non-empty closed convex set in \mathbb{E} . Let x be a point in C . Let n be a point in \mathbb{E} . Then

$$n \in N_C(x) \iff \forall t \in T_C(x), \langle n, t \rangle \leq 0.$$

Proof.

For one direction, assume that $n \in N_C(x)$.

We are to prove that

$$\forall t \in T_C(x), \quad \langle n, t \rangle \leq 0.$$

Let t be an arbitrary point in $T_C(x)$.

Since $t \in T_C(x) = \text{cl}(\text{cone}(C - x))$,

$$\exists \{t_i\}_{i \in \mathbb{N}} \subseteq \text{cone}(C - x), \text{ such that } t_i \rightarrow t. \tag{1}$$

Since $t_i \in \text{cone}(C - x)$,

$$\forall i \in \mathbb{N}, \exists \lambda_i \in \mathbb{R}_{++}, \exists c_i \in C \text{ such that } t_i = \lambda_i(c_i - x). \quad (2)$$

Since $n \in N_C(x)$ and $c_i \in C$,

$$\langle n, c_i - x \rangle \leq 0. \quad (3)$$

Now using (2) and (3), we have

$$\begin{aligned} & \langle n, t_i \rangle \\ &= \langle n, \lambda_i(c_i - x) \rangle, & \text{since } t_i = \lambda_i(c_i - x)s \\ &= \lambda_i \langle n, c_i - x \rangle \\ &\leq \lambda_i \cdot 0, & \text{since } \langle n, c_i - x \rangle \leq 0 \\ &= 0. \end{aligned}$$

That is,

$$\forall i \in \mathbb{N}, \quad \langle n, t_i \rangle \leq 0.$$

Since $\langle n, t_i \rangle \leq 0$ for each $i \in \mathbb{N}$ and $t_i \rightarrow t$, we get

$$\langle n, t \rangle \leq 0.$$

For the reverse direction, assume that n is a vector such that

$$\forall t \in T_C(x), \quad \langle n, t \rangle \leq 0.$$

We are to prove that $n \in N_C(x)$.

Let y be an arbitrary point in $C - x$.

Since $C - x \subseteq \text{clcone}(C - x) = T_C(x)$ and $y \in C - x$, we get $y \in T_C(x)$.

Since $y \in T_C(x)$ and $\forall t \in T_C(x), \langle n, t \rangle \leq 0$, we get $\langle n, y \rangle \leq 0$.

Since $\forall y \in C - x, \langle n, y \rangle \leq 0$, we get $n \in N_C(x)$. ■

THEOREM 6.1. Let C be a closed convex set in \mathbb{E} such that $\text{int}(C) \neq \emptyset$. Let x be a point in \mathbb{E} . Then

$$x \in \text{int}(C) \iff T_C(x) = \mathbb{E} \iff N_C(x) = \{0\}.$$

Proof.

Part 1.

$x \in \text{int}(C)$ if and only if $0 \in \text{int}(C - x)$, if and only if $\text{clcone}(C - x) = \mathbb{E}$.

Part 2.

For one direction, assume that $T_C(x) = \mathbb{E}$.

We are to prove that $N_C(x) = \{0\}$.

Consider $n = 0$.

Since

$$\forall t \in T_C(x), \quad \langle 0, t \rangle = 0 \leq 0,$$

we get $0 \in N_C(x)$.

Let n be an arbitrary vector in $N_C(x)$.

By another proposition, we have

$$\begin{aligned} n &\in N_C(x) \\ \iff \forall t \in T_C(x) = \mathbb{E}, \langle n, t \rangle &\leq 0 \\ \implies \text{for } t = n, \langle n, t \rangle = \langle n, n \rangle &\leq 0 \\ \implies n &= 0. \end{aligned}$$

That is, $n \in N_C(x) \implies n = 0$.

So $N_C(x) = \{0\}$.

For the reverse direction, assume that $N_C(x) = \{0\}$.

We are to prove that $T_C(x) = \mathbb{E}$.

Clearly $T_C(x) \subseteq \mathbb{E}$.

For $\mathbb{E} \subseteq T_C(x)$, let x be an arbitrary point in \mathbb{E} .

Define $p := \text{proj}_{T_C(x)}(x)$.

Since $p = \text{proj}_{T_C(x)}(x)$,

$$\forall y \in T_C(x), \quad \langle x - p, y - p \rangle \leq 0. \quad (1)$$

Since $p = \text{proj}_{T_C(x)}(x)$, $p \in T_C(x)$.

Since $p \in T_C(x)$ and $T_C(x)$ is a cone,

$$2p \in T_C(x). \quad (2)$$

Apply (1) to $y = 2p$, we get

$$\langle x - p, 2p - p \rangle = \langle x - p, p \rangle \leq 0. \quad (3)$$

Since $T_C(x)$ is a closed cone,

$$0 \in T_C(x). \quad (4)$$

Apply (1) to $y = 0$, we get

$$\langle x - p, 0 - p \rangle = \langle x - p, -p \rangle \leq 0. \quad (5)$$

From (3) and (5), we get

$$\langle x - p, p \rangle = 0.$$

So (1) becomes

$$\forall y \in T_C(x), \quad \langle x - p, y \rangle \leq 0.$$

So $x - p \in N_C(x)$.

So $x - p = 0$.

So $x = p$.

So $x \in T_C(x)$.

Since $\forall x \in \mathbb{E}, x \in T_C(x)$, we get

$$\mathbb{E} \subseteq T_C(x).$$

■

6.3 Arithmetic Properties

PROPOSITION 6.3.1. Let C and D be convex subsets of \mathbb{E} . Let x be a point in \mathbb{E} . Then

$$N_C(x) + N_D(x) \subseteq N_{C \cap D}(x).$$

Proof.

If C or D is empty, then $N_C(x) + N_D(x) = N_{C \cap D}(x) = \emptyset$.

So now I assume that $C, D \neq \emptyset$.

If $x \notin C \cap D$, then $N_C(x) + N_D(x) = N_{C \cap D}(x) = \emptyset$.

So now I assume that $x \in C \cap D$.

Let v be an arbitrary point in $N_C(x) + N_D(x)$.

Since $v \in N_C(x) + N_D(x)$, $\exists u \in N_C(x)$, $\exists w \in N_D(x)$ such that $v = u + w$.

Since $u \in N_C(x)$, $\forall y \in C - x$, $\langle u, y \rangle \leq 0$.

Since $w \in N_D(x)$, $\forall y \in D - x$, $\langle w, y \rangle \leq 0$.

Let y be an arbitrary point in $C \cap D - x$.

Since $y \in C \cap D - x$, we get $y \in C - x$ and $y \in D - x$.

$$\begin{aligned} & \langle v, y \rangle \\ &= \langle u + w, y \rangle \\ &= \langle u, y \rangle + \langle w, y \rangle \\ &\leq 0 + 0 = 0. \end{aligned}$$

This is true for any $y \in C \cap D - x$.

So $v \in N_{C \cap D}(x)$.

This is true for any $v \in N_C(x) + N_D(x)$.

So $N_C(x) + N_D(x) \subseteq N_{C \cap D}(x)$.

■

THEOREM 6.2. Let C and D be convex sets in \mathbb{E} . Assume that $\text{ri}(C) \cap \text{ri}(D) \neq \emptyset$. Let x be a point in $C \cap D$. Then

$$N_{C \cap D}(x) = N_C(x) + N_D(x).$$

6.4 Other Properties

PROPOSITION 6.4.1. Let f be a proper, convex, and lower semi-continuous function from \mathbb{E} to \mathbb{R}^* . Let x be a point in $\text{dom}(f)$. Let u be a point in \mathbb{E} . Then $u \in \partial f(x)$ if and only if $(u, -1) \in N_{\text{epi}(f)}(x, f(x))$.

Proof.

$$\begin{aligned} u &\in \partial f(x) \\ \iff \forall y \in \mathbb{E}, f(y) &\geq f(x) + \langle u, y - x \rangle \\ \iff \forall y \in \text{dom}(f), f(y) &\geq f(x) + \langle u, y - x \rangle \\ \iff \forall (y, \beta) \in \text{epi}(f), f(x) + \langle u, y - x \rangle &\leq \beta \\ \iff \forall (y, \beta) \in \text{epi}(f), \langle (u, -1), (y - x, \beta - f(x)) \rangle &\leq 0 \\ \iff \forall (y, \beta) \in \text{epi}(f), \langle (u, -1), (y, \beta) - (x, f(x)) \rangle &\leq 0 \\ \iff (u, -1) \in N_{\text{epi}(f)}(x, f(x)). \end{aligned}$$

■

Chapter 7

Dual Cones and Polar Cones

7.1 Definitions

DEFINITION (Dual Cone). Let K be a cone in \mathbb{E} . We define the **dual cone** of K , denoted by K^* , to be the set given by

$$K^* := \{x \in \mathbb{E} : \forall k \in K, x \cdot k \geq 0\}.$$

DEFINITION (Polar Cone). Let \mathbb{E} be some Euclidean space. Let S be some set in the space. We define the **polar cone** of S , denoted by C° , to be the set given by

$$C^\circ := \{y \in \mathbb{E} : \forall x \in C, \langle y, x \rangle \leq 0\}.$$

7.2 Properties

PROPOSITION 7.2.1. If S is a linear subspace of some Euclidean space \mathbb{E} , then $S^\circ = S^\perp$.

Chapter 8

Extreme Point and Face

8.1 Definitions

DEFINITION (Extreme Points - 1). Let \mathcal{V} be a vector space. Let C be a nonempty convex set in \mathcal{V} . Let x be some point in C . We say that x is an **extreme point** of C if it does not lie strictly between any two distinct points in C .

DEFINITION (Extreme Points - 2). Let \mathbb{E} be some Euclidean space. Let C be a nonempty convex set in the space. Let x be some point in C . We say that x is an **extreme point** of C if $C \setminus \{x\}$ is convex.

PROPOSITION 8.1.1. The two definitions of extreme point are equivalent.

Proof. **Definition 1** \iff **Definition 2**:

Forward Direction: Assume that x does not lie between any two distinct points in C . I will show that $C \setminus \{x\}$ is convex. Let x_1 and x_2 be two arbitrary distinct points in $C \setminus \{x\}$. Let λ be an arbitrary number in $(0, 1)$. Define a point y as $y := \lambda x_1 + (1 - \lambda)x_2$. Since C is convex, $x_1, x_2 \in C$, and $\lambda \in (0, 1)$, we get $y \in C$. Since x does not lie between any two distinct points in C , $y \neq x$. So $y \in C \setminus \{x\}$. That is, I have proved that

$$\forall x_1, x_2 \in C \setminus \{x\}, \forall \lambda \in (0, 1), \quad y = \lambda x_1 + (1 - \lambda)x_2 \in C \setminus \{x\}.$$

By definition, $C \setminus \{x\}$ is convex.

Backward Direction: Assume that $C \setminus \{x\}$ is convex. I will show that x does not lie between any two distinct points in C . Assume for the sake of contradiction that x does lie between two distinct points in C . Say $x = \lambda x_1 + (1 - \lambda)x_2$ where $x_1, x_2 \in C$, $x_1 \neq x_2$, and $\lambda \in (0, 1)$. Clearly $x \neq x_1$ and $x \neq x_2$. So $x_1, x_2 \in C \setminus \{x\}$. Since $C \setminus \{x\}$ is convex, $x_1, x_2 \in C \setminus \{x\}$, and $\lambda \in (0, 1)$, we get $x = \lambda x_1 + (1 - \lambda)x_2 \in C \setminus \{x\}$. This leads to a contradiction. So the assumption that x lies between two distinct points in C does not hold. i.e. x does not lie between two distinct points in C . ■

8.2 Properties of Extreme Points

PROPOSITION 8.2.1. If C is nonempty, convex, and compact, then $\text{Ext}(C) \neq \emptyset$.

PROPOSITION 8.2.2. Let \mathcal{V} be a locally convex space. Let K be a nonempty, compact, and convex set in \mathcal{V} . Then $\text{Ext}(K) \neq \emptyset$.

8.3 Face

DEFINITION (Face). Let \mathcal{V} be a vector space. Let C be a nonempty convex set in \mathcal{V} . Let F be another nonempty convex set in \mathcal{V} such that $F \subseteq C$. We say that F is a **face** of C if

$$\forall x, y \in C, \forall t \in (0, 1), \quad tx + (1 - t)y \in F \implies x, y \in F.$$

Faces are generalizations of extreme points.

PROPOSITION 8.3.1 (Transitivity). If A is a face of B and B is a face of C , then A is a face of C .

DEFINITION (Extreme Points - 3). Let \mathbb{E} be some Euclidean space. Let C be a nonempty convex set in the space. Let x be some point in C . We say that x is an

extreme point of C if $\{x\}$ is a face of C .

PROPOSITION 8.3.2. This definition of extreme points is equivalent to the previous two.

PROPOSITION 8.3.3. If F is a face of C , then $\text{Ext}(F) \subseteq \text{Ext}(C)$.

8.4 The Krein-Milman Theorem

LEMMA 8.1. Let \mathcal{V} be a locally convex space. Let K be a nonempty, compact, and convex set in \mathcal{V} . Let $\rho \in \mathcal{V}^*$. Define $r := \sup\{\Re\rho(x) : x \in K\}$. Define $F := \{x \in K : \Re\rho(x) = r\}$. Then F is a nonempty compact face of K .

Proof. **Nonempty:** Since $\Re\rho$ is continuous and K is compact, $\{\Re\rho(x) : x \in K\}$ is a compact set in \mathbb{R} . So $r = \sup\{\Re\rho(x) : x \in K\}$ is attained. So $F \neq \emptyset$.

Compact: Notice $F = (\Re\rho)^{-1}(\{r\})$. Since $\Re\rho$ is continuous and $\{r\} \subseteq \mathbb{R}$ is closed, F is closed. Since F is a closed subset of K and K is compact, F is compact.

Convex: Let x and y be arbitrary elements of F . Let $t \in (0, 1)$. Since $x, y \in F$, we have $\Re\rho(x) = \Re\rho(y) = r$. So

$$\Re\rho(tx + (1-t)y) = t\Re\rho(x) + (1-t)\Re\rho(y) = tr + (1-t)r = r.$$

So $tx + (1-t)y \in F$. So F is convex.

Face: Let x and y be arbitrary elements of K . Let $t \in (0, 1)$. Suppose that $tx + (1-t)y \in F$. Since $x, y \in K$, we have $\Re\rho(x) \leq r$ and $\Re\rho(y) \leq r$. Since $tx + (1-t)y \in F$, we have

$$t\Re\rho(x) + (1-t)\Re\rho(y) = \Re\rho(tx + (1-t)y) = r.$$

So we must have $\Re\rho(x) = \Re\rho(y) = r$. So $x, y \in F$. So F is a face of K . ■

THEOREM 8.1 (Krein-Milman Theorem). A compact convex set in a locally convex space is the closed convex hull of its extreme points.

Proof. Let \mathcal{V} be a locally convex space. Let K be a nonempty, compact, and convex set in \mathcal{V} .

Forward Direction: Show that $K \subseteq \text{clconv}(\text{Ext}(K))$. Let m be an arbitrary element of K . Assume for the sake of contradiction that $m \notin \text{clconv}(\text{Ext}(K))$. By the Hahn-Banach Theorem, there is some $\tau \in \mathcal{V}^*$ and $\alpha, \beta \in \mathbb{R}$ such that $\alpha > \beta$ and

$$\forall b \in \text{clconv}(\text{Ext}(K)), \quad \Re \tau(m) \geq \alpha > \beta \geq \Re \tau(b).$$

Define $s := \sup\{\Re \tau(w) : w \in K\}$. Define $L := \{z \in K : \Re \tau(z) = s\}$. Then L is a nonempty compact face of K . So $\text{Ext}(L) \neq \emptyset$. Let e be an element of $\text{Ext}(L)$. Then $e \in \text{Ext}(L) \subseteq L$. So $\Re \tau(e) = s$. So

$$\forall b \in \text{clconv}(\text{Ext}(K)), \quad \Re \tau(e) = s \geq \Re \tau(m) \geq \alpha > \beta \geq \Re \tau(b).$$

That is, $\Re \tau(e) > \Re \tau(b)$. Since L is a face of K , $\text{Ext}(L) \subseteq \text{Ext}(K)$. Notice $e \in \text{Ext}(L) \subseteq \text{Ext}(K) \subseteq \text{clconv}(\text{Ext}(K))$. So in particular, $\Re \tau(e) > \Re \tau(e)$. This is a contradiction. So $m \in \text{clconv}(\text{Ext}(K))$. So $K \subseteq \text{clconv}(\text{Ext}(K))$.

Backward Direction: Show that $\text{clconv}(\text{Ext}(K)) \subseteq K$. Note that $\text{Ext}(K) \subseteq K$. Since K is closed and convex and $\text{Ext}(K) \subseteq K$, we get $\text{clconv}(\text{Ext}(K)) \subseteq K$. ■

Chapter 9

Projection

9.1 Definitions

DEFINITION (Projection). Let \mathcal{H} be a Hilbert space. Let S be a non-empty set in the space. Let x be a point in the space. We define the **projection** of x onto S , denoted by $\text{proj}_S(x)$, to be a point given by

$$\text{proj}_S(x) := \operatorname{argmin}_{p \in S} \|p - x\|.$$

i.e., $\text{proj}_S(x)$ is the closest point in S to x .

PROPOSITION 9.1.1 (Existence). If S is non-empty and closed, then the projection $\text{proj}_S(x)$ exists.

Proof. Define for an $n \in \mathbb{N}$ a point c_n to be a point in S that satisfies

$$\lim_{i \in \mathbb{N}} \|c_i - x\| = d_S(x) \text{ where } d_S(x) = \inf_{p \in S} \|p - x\|.$$

Since \mathcal{H} is a Hilbert space, the norm $\|\cdot\|$ on \mathcal{H} satisfies the Parallelogram Law. So

$$\begin{aligned} \|c_m - c_n\|^2 &= 2\|c_m - x\|^2 + 2\|c_n - x\|^2 - \|c_m + c_n - 2x\|^2 \\ &= 2\|c_m - x\|^2 + 2\|c_n - x\|^2 - 4\left\|\frac{c_m + c_n}{2} - x\right\|^2 \\ &\leq 2\|c_m - x\|^2 + 2\|c_n - x\|^2 - 4d_S(x)^2 \\ &\rightarrow 2d_S(x)^2 + 2d_S(x)^2 - 4d_S(x)^2 = 0. \end{aligned}$$

So the sequence $(c_i)_{i \in \mathbb{N}}$ is Cauchy. Since \mathcal{H} is a Hilbert space, it is complete. So $(c_i)_{i \in \mathbb{N}}$ converges. Since S is closed, and $(c_i)_{i \in \mathbb{N}}$ is a Cauchy sequence in S , $p := \lim_{i \in \mathbb{N}} c_i \in S$. So $\|p - x\| = \|\lim_{i \in \mathbb{N}} c_i - x\| = \lim_{i \in \mathbb{N}} \|c_i - x\| = d_S(x)$. So p is the minimizer of the distance to the point x over S . So $p = \text{proj}_S(x)$. ■

PROPOSITION 9.1.2 (Uniqueness). If S is non-empty, closed, and convex, then the projection $\text{proj}_S(x)$ is unique.

Proof. Let p denote $\text{proj}_S(x)$. Then $\|p - x\| = d_S(x)$. Let q be a point in S such that $\|q - x\| = d_S(x)$. Then by the Parallelogram Law,

$$\begin{aligned} 0 \leq \|p - q\|^2 &= 2\|x - p\|^2 + 2\|q - x\|^2 - 4\left\|x - \frac{1}{2}(p + q)\right\|^2 \\ &\leq 2d_S^2(x) + 2d_S^2(x) - 4d_S^2(x) \\ &= 0. \end{aligned}$$

This shows $\|p - q\| = 0$ and hence $p = q$. Thus the projection is unique. ■

9.2 Properties of the Projection Operator

PROPOSITION 9.2.1 (Idempotent). The projection operator is idempotent. i.e., if C is a nonempty closed convex set in \mathbb{E} , then $\text{proj}_C = \text{proj}_C \text{proj}_C$.

Proof. Let x be an arbitrary point in \mathbb{E} . By definition, $\text{proj}_C(x) \in C$. Since $\text{proj}_C(x) \in C$, the closest point in C to $\text{proj}_C(x)$ is $\text{proj}_C(x)$. So $\text{proj}_C \text{proj}_C(x) = \text{proj}_C(x)$. This is true for any $x \in \mathbb{E}$. So $\text{proj}_C = \text{proj}_C \text{proj}_C$. ■

PROPOSITION 9.2.2. Let C be a nonempty closed convex set in \mathbb{E} . Then the set of fixed points of the operator proj_C is C .

Proof. For one direction, let x be an arbitrary fixed point of proj_C . We are to prove that $x \in C$. Since x is a fixed point of proj_C , $x = \text{proj}_C(x)$. By definition of projection, $\text{proj}_C(x) \in C$. So $x = \text{proj}_C(x) \in C$.

For the reverse direction, let x be an arbitrary point in C . We are to prove that x is a fixed point of proj_C . Since $x \in C$, the closest point in C to x is x . So $x = \text{proj}_C(x)$. So x is a fixed point of proj_C . ■

PROPOSITION 9.2.3 (Linearity). Let C be a nonempty closed convex set in \mathbb{E} . Then the operator proj_C is linear if and only if C is a linear subspace.

PROPOSITION 9.2.4 (Non-expansive). The projection operator is non-expansive. i.e., if C is a nonempty closed convex set in \mathbb{E} , then $\|\text{proj}_C(x) - \text{proj}_C(y)\| \leq \|x - y\|$ for any $x, y \in \mathbb{E}$.

this is not true. I guess it will be true when C is a linear subspace.

PROPOSITION 9.2.5. Let C be a nonempty closed convex set in \mathbb{E} . Then proj_C is Lipschitz with constant 1.

Proof. Let x and y be two arbitrary points in \mathbb{E} . If $\|\text{proj}_C(x) - \text{proj}_C(y)\| = 0$, then $\|\text{proj}_C(x) - \text{proj}_C(y)\| \leq \|x - y\|$. Otherwise,

$$\begin{aligned}
 & \|\text{proj}_C(x) - \text{proj}_C(y)\|^2 \\
 &= \langle \text{proj}_C(x) - \text{proj}_C(y), \text{proj}_C(x) - \text{proj}_C(y) \rangle \\
 &= \langle \text{proj}_C(x) - \text{proj}_C(y), \text{proj}_C(x) - x \rangle \\
 &+ \langle \text{proj}_C(x) - \text{proj}_C(y), x - y \rangle \\
 &+ \langle \text{proj}_C(x) - \text{proj}_C(y), y - \text{proj}_C(y) \rangle \\
 &= \langle x - \text{proj}_C(x), \text{proj}_C(y) - \text{proj}_C(x) \rangle \\
 &+ \langle y - \text{proj}_C(y), \text{proj}_C(x) - \text{proj}_C(y) \rangle \\
 &+ \langle \text{proj}_C(x) - \text{proj}_C(y), x - y \rangle \\
 &\leq 0 + 0 + \langle \text{proj}_C(x) - \text{proj}_C(y), x - y \rangle \\
 &= \langle \text{proj}_C(x) - \text{proj}_C(y), x - y \rangle \\
 &\leq \|\text{proj}_C(x) - \text{proj}_C(y)\| \|x - y\|.
 \end{aligned}$$

That is,

$$\|\text{proj}_C(x) - \text{proj}_C(y)\|^2 \leq \|\text{proj}_C(x) - \text{proj}_C(y)\| \|x - y\|.$$

Dividing both sides by $\|\text{proj}_C(x) - \text{proj}_C(y)\|$ gives

$$\|\text{proj}_C(x) - \text{proj}_C(y)\| \leq \|x - y\|.$$

So proj_C is Lipschitz with constant 1. ■

PROPOSITION 9.2.6 (Firmly Non-expansive). Let C be a nonempty closed convex set in \mathbb{E} . Then proj_C is firmly non-expansive.

Proof. This is to prove.

$$\forall x, y \in \mathbb{E}, \quad \|\text{proj}_C(y) - \text{proj}_C(x)\|^2 \leq \langle \text{proj}_C(y) - \text{proj}_C(x), y - x \rangle.$$

$$\begin{aligned} & \|\text{proj}_C(y) - \text{proj}_C(x)\|^2 \\ &= \langle \text{proj}_C(y) - \text{proj}_C(x), \text{proj}_C(y) - \text{proj}_C(x) \rangle \\ &= \langle \text{proj}_C(y) - \text{proj}_C(x), \text{proj}_C(y) - y \rangle \\ &+ \langle \text{proj}_C(y) - \text{proj}_C(x), y - x \rangle \\ &+ \langle \text{proj}_C(y) - \text{proj}_C(x), x - \text{proj}_C(x) \rangle \\ &\leq 0 + \langle \text{proj}_C(y) - \text{proj}_C(x), y - x \rangle + 0 \\ &= \langle \text{proj}_C(y) - \text{proj}_C(x), y - x \rangle. \end{aligned}$$

■

9.3 Characterization

THEOREM 9.1 (Projection Theorem). Let C be a nonempty closed convex set in \mathbb{E} . Let x and p be points in \mathbb{E} . Then $p = \text{proj}_C(x)$ if and only if

$$\forall y \in C, \quad \langle y - p, x - p \rangle \leq 0.$$

Proof. Let y be an arbitrary point in C . Let α be an arbitrary number in $[0, 1]$. Define $y_\alpha := \alpha y + (1 - \alpha)p$. Now

$$\begin{aligned} & p = \text{proj}_C(x) \\ \iff & \forall y \in C, \forall \alpha \in [0, 1], \|x - p\|^2 \leq \|x - y_\alpha\|^2 \\ \iff & \forall y \in C, \forall \alpha \in [0, 1], \|x - p\|^2 \leq \|x - p - \alpha(y - p)\|^2 \\ \iff & \forall y \in C, \langle x - p, y - p \rangle \leq 0. \end{aligned}$$

■

Chapter 10

Separation

10.1 Definitions

DEFINITION (Separated). Let S_1 and S_2 be two sets in \mathbb{E} . We say that S_1 and S_2 are **separated** if $\exists b \in \mathbb{E} \setminus \{\vec{0}\}$ such that

$$\sup_{s_1 \in S_1} \langle s_1, b \rangle \leq \inf_{s_2 \in S_2} \langle s_2, b \rangle.$$

DEFINITION (Strongly Separated). Let S_1 and S_2 be two sets in \mathbb{E} . We say that they are **strongly separated** if the inequality holds strictly.

DEFINITION (Properly Separated). Let S_1 and S_2 be two sets in \mathbb{E} . We say that S_1 and S_2 are **properly separated** if $\exists b \in \mathbb{E}$ such that

$$\begin{aligned} \sup_{x \in S_1} \langle x, b \rangle &\leq \inf_{y \in S_2} \langle y, b \rangle, \text{ and} \\ \inf_{x \in S_1} \langle x, b \rangle &> \sup_{y \in S_2} \langle y, b \rangle. \end{aligned}$$

10.2 Main Results

PROPOSITION 10.2.1. Let C be a nonempty closed convex set in \mathbb{E} . Let x be a point in \mathbb{E} such that $x \notin C$. Then x and C are strongly separated.

Proof. Define a point p by

$$p := \text{proj}_C(x).$$

Define a point a by

$$a := x - p.$$

To prove that x is strongly separated from C , it suffices to prove that

$$\forall y \in C, \quad \langle y, a \rangle < \langle x, a \rangle.$$

Since $x \notin C$ and C is closed,

$$a \neq 0. \tag{1}$$

Let y be an arbitrary point in C . Since $p = \text{proj}_C(x)$ and $y \in C$,

$$\langle y - p, x - p \rangle \leq 0. \tag{2}$$

$$\begin{aligned} & \langle y, a \rangle \\ & < \langle y, a \rangle + \langle a, a \rangle, \text{ since } a \neq 0 \\ & = \langle y + a, a \rangle \\ & = \langle y + x - p, x - p \rangle, \text{ substitute } a = x - p \\ & = \langle y - p, x - p \rangle + \langle x, x - p \rangle \\ & \leq 0 + \langle x, x - p \rangle, \text{ since } \langle y - p, x - p \rangle \leq 0 \\ & = \langle x, x - p \rangle \\ & = \langle x, a \rangle. \end{aligned}$$

That is,

$$\forall y \in C, \quad \langle y, a \rangle < \langle x, a \rangle.$$

So x is strongly separated from C . ■

PROPOSITION 10.2.2. Let C_1 be a non-empty closed convex set in \mathbb{E} . Let C_2 be a non-empty compact convex set in \mathbb{E} . Assume that C_1 and C_2 are disjoint. Then C_1 and C_2 are strongly separated.

Proof. Since C_1 is non-empty closed and convex and C_2 is non-empty compact and convex, we get $C_1 - C_2$ is non-empty closed and convex. Since $C_1 \cap C_2 = \emptyset$, $0 \notin C_1 - C_2$. Since $C_1 - C_2$ is non-empty closed and convex and $0 \in C_1 - C_2$, 0 and $C_1 - C_2$ are strongly separated. Since 0 is strongly separated from $C_1 - C_2$,

$$\exists a \neq 0 \text{ such that } \forall c_1 \in C_1, c_2 \in C_2, \quad \langle c_1 - c_2, a \rangle < \langle 0, a \rangle.$$

That is,

$$\langle c_1, a \rangle < \langle c_2, a \rangle.$$

So C_1 and C_2 are strongly separated. ■

THEOREM 10.1. Let C_1 and C_2 be non-empty closed convex sets in \mathbb{E} . Assume that C_1 and C_2 are disjoint. Then C_1 and C_2 are separated.

Proof. For $n \in \mathbb{N}$, define

$$D_n := C_2 \cap \text{ball}(0, n).$$

Then D_n is compact for any $n \in \mathbb{N}$. Since $\left\{ \begin{array}{l} C_1 \text{ is non-empty closed and convex} \\ D_n \text{ is non-empty compact and convex} \end{array} \right.$ we get C_1 and D_n are strongly separated for any $n \in \mathbb{N}$. So

$$\forall n \in \mathbb{N}, \exists a_n \in \mathbb{E}, \|a_n\| = 1 \text{ such that } \forall c_1 \in C_1, \forall d_2 \in D_n, \quad \langle c_1, a_n \rangle < \langle d_2, a_n \rangle.$$

Since $\|a_n\| = 1$ for any $n \in \mathbb{N}$, there exists a subsequence $\{a_n\}_{n \in I}$ where I is some infinite subset of \mathbb{N} such that $\{a_n\}_{n \in I}$ converges to some point $a \in \mathbb{E}$. Let x be an arbitrary point in C_1 . Let y be an arbitrary point in C_2 . For large enough n , $y \in D_n$. Since

$$\left\{ \begin{array}{l} \langle x, a_n \rangle < \langle y, a_n \rangle \text{ for large enough } n \\ \lim_{n \in I, n \rightarrow \infty} \langle x, a_n \rangle = \langle x, a \rangle \\ \lim_{n \in I, n \rightarrow \infty} \langle y, a_n \rangle = \langle y, a \rangle \end{array} \right., \text{ we get}$$

$$\langle x, a \rangle \leq \langle y, a \rangle.$$

Since

$$\exists a \neq 0 \text{ such that } \forall x \in C_1, \forall y \in C_2, \quad \langle x, a \rangle \leq \langle y, a \rangle,$$

by definition of separated, C_1 and C_2 are separated. ■

PROPOSITION 10.2.3. Let C_1 and C_2 be non-empty convex subsets of \mathbb{E} . Then C_1 and C_2 are properly separated if and only if

$$\text{ri}(C_1) \cap \text{ri}(C_2) = \emptyset.$$



Chapter 11

Convex Functions

11.1 Preliminaries

DEFINITION (Epigraph). Let f be a function from \mathbb{E} to \mathbb{R}^* . We define the **epigraph** of f , denoted by $\text{epi}(f)$, to be the set given by

$$\text{epi}(f) := \{(x, \alpha) \in \mathbb{E} \times \mathbb{R} : f(x) \leq \alpha\}.$$

DEFINITION (Domain). Let f be a function from \mathbb{E} to \mathbb{R}^* . We define the **domain** of f , denoted by $\text{dom}(f)$, to be a set given by

$$\text{dom}(f) := \{x \in \mathbb{E} : f(x) < +\infty\}.$$

DEFINITION (Proper). Let f be a function from \mathbb{E} to \mathbb{R}^* . We say that f is **proper** if

$$\begin{aligned} \exists x \in \mathbb{E}, \quad f(x) &\neq +\infty, \text{ and} \\ \forall x \in \mathbb{E}, \quad f(x) &\neq -\infty \end{aligned}$$

11.2 The Indicator Function

DEFINITION (The Indicator Function). Let S be a subset of \mathbb{E} . We define the **indicator function** of S , denoted by δ_S , to be a function from \mathbb{E} to \mathbb{R}^* given by

$$\delta_S(x) = \begin{cases} 0, & \text{if } x \in S \\ +\infty, & \text{if } x \notin S. \end{cases}$$

PROPOSITION 11.2.1. Let S be a subset of \mathbb{E} . Then

- (1) S is non-empty if and only if δ_S is proper.
- (2) S is convex if and only if δ_S is convex.
- (3) S is closed if and only if δ_S is lower semi-continuous.

Proof of (1).

For one direction, assume that S is not empty.

We are to prove that δ_S is proper.

Since $S \neq \emptyset$, pick $p \in S$.

Since $p \in S$, $\delta_S(p) = 0$.

Since $\delta_S(p) = 0$, $\exists x_0 \in \mathbb{E}$ such that $\delta_S(x_0) \neq +\infty$.

By definition of the indicator function, it never takes $-\infty$.

Since $\exists x_0 \in \mathbb{E}$ such that $\delta_S(x_0) \neq +\infty$ and $\forall x \in \mathbb{E}$, $\delta_S(x) \neq -\infty$, we get δ_S is proper.

For the reverse direction, assume that δ_S is proper.

We are to prove that S is non-empty.

Assume for the sake of contradiction that S is empty.

Let x be an arbitrary point in \mathbb{E} .

Since $S = \emptyset$, $x \notin S$.

Since $x \notin S$, $\delta_S(x) = +\infty$.

Since $\forall x \in \mathbb{E}$, $\delta_S(x) = +\infty$, by definition of proper function, δ_S is not proper.

This contradicts to the assumption that δ_S is proper.

So the assumption that $S = \emptyset$ is false.

i.e., S is non-empty. ■

Proof of (2).

For one direction, assume that S is convex.

We are to prove that δ_S is convex.

Let x and y be arbitrary points in $\text{dom}(\delta_S)$.

By definition of indicator functions, $\text{dom}(\delta_S) = S$.

So $x, y \in S$.

Let λ be an arbitrary number in $(0, 1)$.

Define point z as $z := \lambda x + (1 - \lambda)y$.

Since $x, y \in S$ and $\lambda \in (0, 1)$ and S is convex and $z = \lambda x + (1 - \lambda)y$, we get $z \in S$.

Since $z \in S$, $\delta_S(z) = 0$.

Since $\lambda \in (0, 1)$ and $\text{range}(\delta_S) = \{0, +\infty\}$, we get $\lambda\delta_S(x) + (1 - \lambda)\delta_S(y) \geq 0$.

Since $\delta_S(z) = 0$ and $\lambda\delta_S(x) + (1 - \lambda)\delta_S(y) \geq 0$, we get $\delta_S(z) \leq \lambda\delta_S(x) + (1 - \lambda)\delta_S(y)$.

That is, $\delta_S(\lambda x + (1 - \lambda)y) \leq \lambda\delta_S(x) + (1 - \lambda)\delta_S(y)$.

Since $\forall x, y \in \text{dom}(\delta_S)$, $\forall \lambda \in (0, 1)$, $\delta_S(\lambda x + (1 - \lambda)y) \leq \lambda\delta_S(x) + (1 - \lambda)\delta_S(y)$, we get δ_S is convex.

For the reverse direction, assume that δ_S is convex.

We are to prove that S is convex.

The case where S is empty is trivial.

So now I assume $S \neq \emptyset$.

Let x and y be arbitrary points in S .

Let λ be an arbitrary number in $(0, 1)$.

Define point z as $z := \lambda x + (1 - \lambda)y$.

Since $x \in S$, $\delta_S(x) = 0$.

Since $y \in S$, $\delta_S(y) = 0$.

Since $\delta_S(x) = \delta_S(y) = 0$, we get $\lambda\delta_S(x) + (1 - \lambda)\delta_S(y) = 0$.

Since $\lambda \in (0, 1)$ and δ_S is convex, $\delta_S(z) \leq \lambda\delta_S(x) + (1 - \lambda)\delta_S(y)$.

Since $\delta_S(z) \leq \lambda\delta_S(x) + (1 - \lambda)\delta_S(y)$ and $\lambda\delta_S(x) + (1 - \lambda)\delta_S(y) = 0$, we get $\delta_S(z) \leq 0$.

By definition of the indicator function, $\delta_S(z) \geq 0$.

Since $\delta_S(z) \leq 0$ and $\delta_S(z) \geq 0$, we get $\delta_S(z) = 0$.

Since $\delta_S(z) = 0$, $z \in S$.

That is, $\lambda x + (1 - \lambda)y \in S$.

Since $\forall x, y \in S$, $\forall \lambda \in (0, 1)$, $\lambda x + (1 - \lambda)y \in S$, we get S is convex. ■

Proof of (3).

For one direction, assume that S is closed.

We are to prove that δ_S is lower semi-continuous.

Let $\{(x_i, \alpha_i)\}_{i \in \mathbb{N}}$ be an arbitrary sequence in $\text{epi}(\delta_S)$ that converges.

Say its limit is $(x_\infty, \alpha_\infty)$.

Since $(x_i, \alpha_i) \rightarrow (x_\infty, \alpha_\infty)$, $x_i \rightarrow x_\infty$.

Since $(x_i, \alpha_i) \in \text{epi}(\delta_S)$, $\delta_S(x_i) \leq \alpha_i$.

Since $\delta_S(x_i) \leq \alpha_i$ and $\alpha_i \in \mathbb{R}$, we get $\delta_S(x_i) \neq +\infty$.

Since $\delta_S(x_i) \neq +\infty$, $x_i \in S$.

Since $x_i \in S$ and $x_i \rightarrow x_\infty$ and S is closed, $x_\infty \in S$.

Since $x_\infty \in S$, $\delta_S(x_\infty) = 0$.

Since $x_i \in S$, $\delta_S(x_i) = 0$.

Since $\delta_S(x_i) = 0$ and $\delta_S(x_i) \leq \alpha_i$, $\alpha_i \geq 0$.

Since $(x_i, \alpha_i) \rightarrow (x_\infty, \alpha_\infty)$, $\alpha_i \rightarrow \alpha_\infty$.

Since $\alpha_i \geq 0$ and $\alpha \rightarrow \alpha_\infty$, $\alpha_\infty \geq 0$.

Since $\delta_S(x_\infty) = 0$ and $\alpha_\infty \geq 0$, $\delta_S(x_\infty) \leq \alpha_\infty$.

Since $\delta_S(x_\infty) \leq \alpha_\infty$, $(x_\infty, \alpha_\infty) \in \text{epi}(\delta_S)$.

Since for any convergent sequence in $\text{epi}(\delta_S)$, its limit is also in $\text{epi}(\delta_S)$, we get $\text{epi}(\delta_S)$ is closed.

For the reverse direction, assume that δ_S is lower semi-continuous.

We are to prove that S is closed.

Let $\{x_i\}_{i \in \mathbb{N}}$ be an arbitrary sequence in S that converges.

Say its limit is x_∞ .

Since $x_i \in S$, $\delta_S(x_i) = 0$.

Since $\delta_S(x_i) = 0$, $(x_i, 0) \in \text{epi}(\delta_S)$.

Since $x_i \rightarrow x_\infty$, $(x_i, 0) \rightarrow (x_\infty, 0)$.

Since $(x_i, 0) \in \text{epi}(\delta_S)$ and $(x_i, 0) \rightarrow (x_\infty, 0)$, $(x_\infty, 0) \in \text{epi}(\delta_S)$.

Since $(x_\infty, 0) \in \text{epi}(\delta_S)$, $\delta_S(x_\infty) \leq 0$.

By definition of the indicator function, $\delta_S(x_\infty) \geq 0$.

Since $\delta_S(x_\infty) \leq 0$ and $\delta_S(x_\infty) \geq 0$, we get $\delta_S(x_\infty) = 0$.

Since $\delta_S(x_\infty) = 0$, $x_\infty \in S$.

Since for any convergent sequence in S , its limit is also in S , we get S is closed. ■

11.3 Definitions

DEFINITION (Convex Function). Let f be a function from \mathbb{E} to \mathbb{R}^* . We say that f is **convex** if

$$\forall x, y \in \text{dom}(f), \forall \lambda \in [0, 1], \quad f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y).$$

DEFINITION (Convex Function). Let f be a function from \mathbb{E} to \mathbb{R}^* . We say that f is **convex** if the epigraph of f is convex.

PROPOSITION 11.3.1. The two definitions of convexity of functions are equivalent.

Proof.

The case where $\text{dom}(f), \text{epi}(f) = \emptyset$ is trivial.

So now I assume that $\text{dom}(f), \text{epi}(f) \neq \emptyset$.

For one direction, assume that $\forall x, y \in \text{dom}(f), \forall \lambda \in [0, 1]$, we have $f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$.

We are to prove that the epigraph of f is convex.

Let (x, α) and (y, β) be two arbitrary points in $\text{epi}(f)$.

Since $(x, \alpha), (y, \beta) \in \text{epi}(f)$, $x, y \in \text{dom}(f)$.

Let λ be an arbitrary number in $[0, 1]$.

Define a point $(z, \gamma) := \lambda(x, \alpha) + (1 - \lambda)(y, \beta)$.

Then $z = \lambda x + (1 - \lambda)y$ and $\gamma = \lambda\alpha + (1 - \lambda)\beta$.

Since $x, y \in \text{dom}(f)$, $\lambda \in [0, 1]$, we get $f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$.

Since $(x, \alpha) \in \text{epi}(f)$, $f(x) \leq \alpha$.

Since $(y, \beta) \in \text{epi}(f)$, $f(y) \leq \beta$.

Since $f(x) \leq \alpha$ and $f(y) \leq \beta$ and $f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$, we get $f(\lambda x + (1 - \lambda)y) \leq \lambda\alpha + (1 - \lambda)\beta$.

Since $z = \lambda x + (1 - \lambda)y$ and $\gamma = \lambda\alpha + (1 - \lambda)\beta$ and $f(\lambda x + (1 - \lambda)y) \leq \lambda\alpha + (1 - \lambda)\beta$, we get $f(z) \leq \gamma$.

Since $f(z) \leq \gamma$, $(z, \gamma) \in \text{epi}(f)$.

For the reverse direction, assume that $\text{epi}(f)$ is convex.

We are to prove that $\forall x, y \in \text{dom}(f), \forall \lambda \in [0, 1]$, we have $f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$.

Let x and y be two arbitrary points in $\text{dom}(f)$.

Let λ be an arbitrary number in $[0, 1]$.

Define $z := \lambda x + (1 - \lambda)y$.

Define $\gamma := \lambda f(x) + (1 - \lambda)f(y)$.

Since $(x, f(x)) \in \text{epi}(f)$ and $(y, f(y)) \in \text{epi}(f)$ and $\lambda \in [0, 1]$ and $\text{epi}(f)$ is convex, we get $\lambda(x, f(x)) + (1 - \lambda)(y, f(y)) \in \text{epi}(f)$.

Since $z = \lambda x + (1 - \lambda)y$ and $\gamma = \lambda f(x) + (1 - \lambda)f(y)$ and $\lambda(x, f(x)) + (1 - \lambda)(y, f(y)) \in \text{epi}(f)$, we get $(z, \gamma) \in \text{epi}(f)$.

Since $(z, \gamma) \in \text{epi}(f)$, $f(z) \leq \gamma$.

That is, $f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$.

■

11.4 Basic Properties

PROPOSITION 11.4.1 (Necessary Condition). The domain of a convex function is convex.

Proof. Follows from the fact that convexity is stable under affine transformations. Define $A((x, \alpha)) := x$. Then $\text{dom}(f) = A(\text{epi}(f))$. ■

PROPOSITION 11.4.2. The level sets of a convex function are convex.

PROPOSITION 11.4.3 (Restriction to a Line). A function $f : \mathbb{E} \rightarrow \mathbb{R}$ is convex if and only if $\forall x \in \text{dom}(f), \forall v \in \mathbb{E}$, the function $g_{x,v} : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$g_{x,v}(t) = f(x + tv)$$

is convex.

11.5 Differentiable Convex Functions

PROPOSITION 11.5.1. Let f be a proper convex function from \mathbb{E} to \mathbb{R}^* . Let $x \in \text{dom}(f)$. If f is differentiable at point x , then $\nabla(f)(x)$ is the unique subgradient of f at point x . i.e., $\partial(f)(x) = \{\nabla(f)(x)\}$. Conversely, if the subgradient $\partial(f)(x)$ of f at point x is a singleton set $\{v\}$, then f is differentiable at point x and $\nabla(f)(x) = v$.

Proof. ■

PROPOSITION 11.5.2 (First-Order Condition). Let f be a proper function from \mathbb{E} to \mathbb{R}^* . Assume that $\text{dom}(f)$ is convex and open and that f is differentiable on $\text{dom}(f)$. Then f is convex if and only if

$$\forall x, y \in \text{dom}(f), \quad f(y) - f(x) \geq \langle \nabla(f)(x), y - x \rangle.$$

i.e., the first-order approximation of f is a global under-estimator.

Proof.

Part 1.

For one direction, assume that f is convex. We are to prove that

$$\forall x, y \in \text{dom}(f), \quad f(y) - f(x) \geq \langle \nabla(f)(x), y - x \rangle.$$

Let x and y be arbitrary points in $\text{dom}(f)$. Since f is convex and differentiable at point x , $\nabla(f)(x) = \partial(f)(x)$. So $\nabla(f)(x)$ satisfies the subgradient inequality. That is,

$$f(y) - f(x) \geq \langle \nabla(f)(x), y - x \rangle.$$

Part 2.

For the reverse direction, assume that

$$\forall x, y \in \text{dom}(f), \quad f(y) - f(x) \geq \langle \nabla(f)(x), y - x \rangle.$$

We are to prove that f is convex.

Not Finished.

■

PROPOSITION 11.5.3. Let f be a proper function from \mathbb{E} to \mathbb{R}^* . Assume that $\text{dom}(f)$ is convex and open and that f is differentiable on $\text{dom}(f)$. Then f is convex if and only if

$$\forall x, y \in \text{dom}(f), \quad \langle \nabla(f)(x) - \nabla(f)(y), x - y \rangle \geq 0.$$

Proof.

Part 1.

For one direction, assume that f is convex. We are to prove that

$$\forall x, y \in \text{dom}(f), \quad \langle \nabla(f)(x) - \nabla(f)(y), x - y \rangle \geq 0.$$

Let x and y be arbitrary points in $\text{dom}(f)$. Since f is convex and differentiable at point x , $\nabla(f)(x) = \partial(f)(x)$. So $\nabla(f)(x)$ satisfies the subgradient inequality. That is,

$$f(y) - f(x) \geq \langle \nabla(f)(x), y - x \rangle. \tag{1}$$

Since f is convex and differentiable at point y , $\nabla(f)(y) = \partial(f)(y)$. So $\nabla(f)(y)$ satisfies the subgradient inequality. That is,

$$f(x) - f(y) \geq \langle \nabla(f)(y), x - y \rangle. \tag{2}$$

Take the sum of inequalities (1) and (2), we get

$$(f(y) - f(x)) + (f(x) - f(y)) \geq \langle \nabla(f)(x), y - x \rangle + \langle \nabla(f)(y), x - y \rangle$$

$$\begin{aligned}
&\implies 0 \geq -\langle \nabla(f)(x), x-y \rangle + \langle \nabla(f)(y), x-y \rangle \\
&\implies \langle \nabla(f)(x), x-y \rangle - \langle \nabla(f)(y), x-y \rangle \geq 0 \\
&\implies \langle \nabla(f)(x) - \nabla(f)(y), x-y \rangle \geq 0.
\end{aligned}$$

Part 2.

For the reverse direction, assume that

$$\forall x, y \in \text{dom}(f), \quad \langle \nabla(f)(x) - \nabla(f)(y), x-y \rangle \geq 0.$$

We are to prove that f is convex. Let x and y be arbitrary points in $\text{dom}(f)$. Define a function φ on $(0, 1)$ by

$$\varphi(\lambda) := f(\lambda x + (1-\lambda)y).$$

Notice φ is differentiable and

$$\varphi'(\lambda) = \langle \nabla(f)(\lambda x + (1-\lambda)y), x-y \rangle.$$

Let α and β be arbitrary numbers in $(0, 1)$. Assume that $\alpha < \beta$. Define two points z_α and z_β by $z_\alpha := \alpha x + (1-\alpha)y$ and $z_\beta := \beta x + (1-\beta)y$. Then

$$\begin{aligned}
&\varphi'(\beta) - \varphi'(\alpha) \\
&= \langle \nabla(f)(\beta x + (1-\beta)y), x-y \rangle - \langle \nabla(f)(\alpha x + (1-\alpha)y), x-y \rangle \\
&= \langle \nabla(f)(z_\beta), x-y \rangle - \langle \nabla(f)(z_\alpha), x-y \rangle \\
&= \langle \nabla(f)(z_\beta) - \nabla(f)(z_\alpha), x-y \rangle \\
&= \left\langle \nabla(f)(z_\beta) - \nabla(f)(z_\alpha), \frac{z_\beta - z_\alpha}{\beta - \alpha} \right\rangle \\
&= \frac{1}{\beta - \alpha} \langle \nabla(f)(z_\beta) - \nabla(f)(z_\alpha), z_\beta - z_\alpha \rangle \\
&\geq \frac{1}{\beta - \alpha} \cdot 0, \text{ by assumption} \\
&= 0.
\end{aligned}$$

That is,

$$\forall \alpha, \beta \in (0, 1), \quad \beta > \alpha \implies \varphi'(\beta) - \varphi'(\alpha) \geq 0.$$

So φ' is increasing. So φ is convex. So

$$\varphi(\lambda) \leq \lambda\varphi(1) + (1-\lambda)\varphi(0).$$

That is,

$$f(\lambda x + (1-\lambda)y) \leq \lambda f(x) + (1-\lambda)f(y).$$

By definition, f is convex. ■

PROPOSITION 11.5.4 (Second-Order Condition). A twice continuously differentiable real-valued function f defined on a convex set is convex if and only if

$$\forall x \in \text{dom}(f), \quad \nabla^2 f(x) \geq 0$$

where $\nabla^2 f(x) = \begin{bmatrix} \frac{\partial^2 f(x)}{\partial x_1 \partial x_1} & \cdots & \frac{\partial^2 f(x)}{\partial x_1 \partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 f(x)}{\partial x_n \partial x_1} & \cdots & \frac{\partial^2 f(x)}{\partial x_n \partial x_n} \end{bmatrix}$ denotes the Hessian matrix of f at x_0 .

PROPOSITION 11.5.5. Let f be a twice continuously differentiable function from \mathbb{E} to \mathbb{R} . Then f is convex if and only if $\forall x \in \mathbb{E}$, $\nabla^2 f(x)$ is positive semi-definite.

11.6 Convexity and Lipschitz-ness

THEOREM 11.1. Let f be a differentiable convex function from \mathbb{E} to \mathbb{R} . Then the following statements are equivalent.

(1) ∇f is Lipschitz with constant L .

(2) $\forall x, y \in \mathbb{E}$, we have

$$f(y) - f(x) \leq \langle \nabla f(x), y - x \rangle + \frac{L}{2} \|y - x\|^2.$$

(3) $\forall x, y \in \mathbb{E}$, we have

$$f(y) - f(x) \geq \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \|\nabla f(y) - \nabla f(x)\|^2.$$

(4) $\forall x, y \in \mathbb{E}$, we have

$$L \langle \nabla f(y) - \nabla f(x), y - x \rangle \geq \|\nabla f(y) - \nabla f(x)\|^2.$$

(1) \implies (2).

Assume that ∇f is Lipschitz with constant L .

Let x and y be two arbitrary points in \mathbb{E} .

$$f(y) - f(x)$$

$$\begin{aligned}
&= \int_0^1 \langle \nabla f(x + t(y - x)), y - x \rangle dt \\
&= \langle \nabla f(x), y - x \rangle + \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), y - x \rangle dt \\
&\leq \langle \nabla f(x), y - x \rangle + \int_0^1 \|\nabla f(x + t(y - x)) - \nabla f(x)\| \|y - x\| dt \\
&\leq \langle \nabla f(x), y - x \rangle + \int_0^1 L \|x + t(y - x) - x\| \|y - x\| dt \\
&= \langle \nabla f(x), y - x \rangle + L \|y - x\|^2 \int_0^1 t dt \\
&= \langle \nabla f(x), y - x \rangle + \frac{L}{2} \|y - x\|^2
\end{aligned}$$

That is,

$$f(y) - f(x) \leq \langle \nabla f(x), y - x \rangle + \frac{L}{2} \|y - x\|^2.$$

■

THEOREM 11.2. Let f be a twice continuously differentiable function from \mathbb{E} to \mathbb{R} . Let L be some non-negative number. Then the following statements are equivalent.

- (1) ∇f is L -Lipschitz.
- (2) $\forall x \in \mathbb{E}, \|\nabla^2 f(x)\| \leq L$.

11.7 Stability of Convexity

PROPOSITION 11.7.1 (Non-Negative Linear Combination). A non-negative linear combination of proper convex functions is again convex.

Proof. It suffices to prove that non-negative scalar multiples of convex functions are convex and sums of two convex functions are convex.

Part 1.

Let f be a proper convex function. Let $\alpha \geq 0$ be an arbitrary scalar. We are to prove that αf is convex. Notice $\text{dom}(f) = \text{dom}(\alpha f)$. Since f is proper, $\text{dom}(f) \neq \emptyset$. So $\text{dom}(\alpha f) \neq \emptyset$. Let x and y be two arbitrary points in $\text{dom}(\alpha f)$. Let λ be an arbitrary number in $(0, 1)$.

Define a point z as $z := \lambda x + (1 - \lambda)y$. Then

$$\begin{aligned} (\alpha f)(\lambda x + (1 - \lambda)y) &= \alpha f(\lambda x + (1 - \lambda)y) \\ &\leq \alpha(\lambda f(x) + (1 - \lambda)f(y)) \\ &= \lambda \alpha f(x) + (1 - \lambda) \alpha f(y) \\ &= \lambda(\alpha f)(x) + (1 - \lambda)(\alpha f)(y). \end{aligned}$$

That is,

$$\forall x, y \in \text{dom}(\alpha f), \forall \lambda \in (0, 1), \quad (\alpha f)(\lambda x + (1 - \lambda)y) \leq \lambda(\alpha f)(x) + (1 - \lambda)(\alpha f)(y).$$

So by definition, αf is convex.

Part 2.

Let f and g be proper convex functions. We are to prove that $f + g$ is convex. Notice $\text{dom}(f + g) = \text{dom}(f) \cap \text{dom}(g)$. Since f is proper, $\text{dom}(f) \neq \emptyset$. Since g is proper, $\text{dom}(g) \neq \emptyset$. So $\text{dom}(f + g) \neq \emptyset$. Let x and y be two arbitrary points in $\text{dom}(f + g)$. Let λ be an arbitrary number in $(0, 1)$. Define a point z as $z := \lambda x + (1 - \lambda)y$. Then

$$\begin{aligned} (f + g)(\lambda x + (1 - \lambda)y) &= f(\lambda x + (1 - \lambda)y) + g(\lambda x + (1 - \lambda)y) \\ &\leq \lambda f(x) + (1 - \lambda)f(y) + \lambda g(x) + (1 - \lambda)g(y) \\ &= \lambda(f(x) + g(x)) + (1 - \lambda)(f(y) + g(y)) \\ &= \lambda(f + g)(x) + (1 - \lambda)(f + g)(y). \end{aligned}$$

That is,

$$\forall x, y \in \text{dom}(f + g), \forall \lambda \in (0, 1), \quad (f + g)(\lambda x + (1 - \lambda)y) \leq \lambda(f + g)(x) + (1 - \lambda)(f + g)(y).$$

So by definition, $f + g$ is convex. ■

PROPOSITION 11.7.2 (Direct Sum). Direct sums of convex functions are convex.

Proof. Let z and w be two arbitrary points in $\text{dom}(f \oplus g)$. Let $\lambda \in (0, 1)$ be arbitrary. Say $z = x \oplus y$ and $w = u \oplus v$ where $x, u \in \mathbb{R}^m$ and $y, v \in \mathbb{R}^p$. Since $z \in \text{dom}(f \oplus g)$, $(f \oplus g)(z) \neq +\infty$. That is, $f(x) + g(y) \neq +\infty$. So neither $f(x)$ nor $g(y)$ is $+\infty$. So both $x \in \text{dom}(f)$ and $y \in \text{dom}(g)$. Similarly, we have $u \in \text{dom}(f)$ and $v \in \text{dom}(g)$. Consider the point

$$\begin{aligned} &\lambda z + (1 - \lambda)w \\ &= \lambda x \oplus y + (1 - \lambda)u \oplus v \end{aligned}$$

$$= (\lambda x + (1 - \lambda)u) \oplus (\lambda y + (1 - \lambda)v).$$

Apply $f \oplus g$ to both sides, we get

$$\begin{aligned} & (f \oplus g)(\lambda z + (1 - \lambda)w) \\ &= (f \oplus g)[(\lambda x + (1 - \lambda)u) \oplus (\lambda y + (1 - \lambda)v)] \\ &= f(\lambda x + (1 - \lambda)u) + g(\lambda y + (1 - \lambda)v). \end{aligned}$$

Since f and g are convex, we get

$$\begin{aligned} f(\lambda x + (1 - \lambda)u) &\leq \lambda f(x) + (1 - \lambda)f(u), \text{ and} \\ g(\lambda y + (1 - \lambda)v) &\leq \lambda g(y) + (1 - \lambda)g(v). \end{aligned}$$

So

$$\begin{aligned} & (f \oplus g)(\lambda z + (1 - \lambda)w) \\ &\leq \lambda f(x) + (1 - \lambda)f(u) + \lambda g(y) + (1 - \lambda)g(v) \\ &= \lambda(f(x) + g(y)) + (1 - \lambda)(f(u) + g(v)) \\ &= \lambda(f \oplus g)(x \oplus y) + (1 - \lambda)(f \oplus g)(u \oplus v) \\ &= \lambda(f \oplus g)(z) + (1 - \lambda)(f \oplus g)(w). \end{aligned}$$

That is,

$$(f \oplus g)(\lambda z + (1 - \lambda)w) \leq \lambda(f \oplus g)(z) + (1 - \lambda)(f \oplus g)(w).$$

This holds for any $z, w \in \text{dom}(f \oplus g)$ and any $\lambda \in (0, 1)$. So $(f \oplus g)$ is convex. ■

PROPOSITION 11.7.3 (Composition). The composition of a convex function with an affine function is convex. i.e., if f is convex, then $f(Ax + b)$ is convex.

Proof. Let x and y be arbitrary points in \mathbb{E} . Let λ be an arbitrary number in $(0, 1)$. Define a point z by $z := \lambda x + (1 - \lambda)y$.

$$\begin{aligned} & g(\lambda x + (1 - \lambda)y) \\ &= f(A(\lambda x + (1 - \lambda)y) + b) \\ &= f(\lambda Ax + (1 - \lambda)Ay + b), && \text{by linearity of } A \\ &= f(\lambda Ax + (1 - \lambda)Ay + \lambda b + (1 - \lambda)b), && \text{decompose } b \\ &= f(\lambda(Ax + b) + (1 - \lambda)(Ay + b)) \\ &\leq \lambda f(Ax + b) + (1 - \lambda)f(Ay + b), && \text{by convexity of } f \end{aligned}$$

$$= \lambda g(x) + (1 - \lambda)g(y).$$

That is,

$$\forall x, y \in \mathbb{E}, \forall \lambda \in (0, 1), \quad g(\lambda x + (1 - \lambda)y) \leq \lambda g(x) + (1 - \lambda)g(y).$$

So g is convex. ■

PROPOSITION 11.7.4 (Supremum). The supremum of a collection of convex functions is again convex. i.e., Let $\{f_i\}_{i \in I}$ be a collection of convex functions where I is some index set. Then the function F given by $F := \sup_{i \in I} f_i$ is convex.

Proof.

$$\begin{aligned} (x, \alpha) &\in \text{epi}(F) \\ \iff \sup_{i \in I} f_i(x) &\leq \alpha \\ \iff \forall i \in I, f_i(x) &\leq \alpha \\ \iff \forall i \in I, (x, \alpha) &\in \text{epi}(f_i) \\ \iff (x, \alpha) &\in \bigcap_{i \in I} \text{epi}(f_i). \end{aligned}$$

So $\text{epi}(F) = \bigcap_{i \in I} \text{epi}(f_i)$. Since f_i are convex, $\text{epi}(f_i)$ are convex. Since $\text{epi}(f_i)$ are convex, $\bigcap_{i \in I} \text{epi}(f_i)$ is convex. That is, $\text{epi}(F)$ is convex. Since $\text{epi}(F)$ is convex, F is convex. ■

PROPOSITION 11.7.5 (Pointwise Supremum). If $f(x, y)$ is convex in x for each y in some set \mathcal{A} , then the function g given by

$$g(x) = \sup_{y \in \mathcal{A}} f(x, y)$$

is convex.

11.8 Examples

EXAMPLE 11.8.1. Affine functions are convex.

EXAMPLE 11.8.2. Norms are convex.

Proof.

$$\begin{aligned}
 & \|\alpha x + \beta y\| \\
 & \leq \|\alpha x\| + \|\beta y\| \\
 & = |\alpha|\|x\| + |\beta|\|y\| \\
 & = \alpha\|x\| + \beta\|y\|.
 \end{aligned}$$

■

EXAMPLE 11.8.3. Square norms are convex.

Proof Approach 1. Notice $\|\cdot\|^2$ is the direct sum of m squares and squares are convex. So by CO 463 Assignment 2 Problem 3, $\|\cdot\|^2$ is convex. ■

Proof Approach 2. The domain is \mathbb{E} . Let x and y be two points in \mathbb{E} . Let λ be an arbitrary number in $(0, 1)$. Define a point z as $z := \lambda x + (1 - \lambda)y$.

$$\begin{aligned}
 & \|\lambda x + (1 - \lambda)y\|^2 \\
 & = \|\lambda x\|^2 + \|(1 - \lambda)y\|^2 + 2\langle \lambda x, (1 - \lambda)y \rangle \\
 & = \lambda^2\|x\|^2 + (1 - \lambda)^2\|y\|^2 + 2\lambda(1 - \lambda)\langle x, y \rangle \\
 & \leq \lambda^2\|x\|^2 + (1 - \lambda)^2\|y\|^2 + 2\lambda(1 - \lambda)\|x\|\|y\| \\
 & \leq \lambda(\lambda - 1)\|x\|^2 + \lambda(\lambda - 1)\|y\|^2 + 2\lambda(1 - \lambda)\|x\|\|y\| \\
 & \quad + \lambda\|x\|^2 + (1 - \lambda)\|y\|^2 \\
 & = \lambda\|x\|^2 + (1 - \lambda)\|y\|^2 \\
 & \quad + \lambda(\lambda - 1)[\|x\|^2 + \|y\|^2 - 2\|x\|\|y\|] \\
 & \leq \lambda\|x\|^2 + (1 - \lambda)\|y\|^2
 \end{aligned}$$

That is,

$$\forall x, y \in \mathbb{E}, \forall \lambda \in (0, 1), \quad \|\lambda x + (1 - \lambda)y\|^2 \leq \lambda\|x\|^2 + (1 - \lambda)\|y\|^2.$$

So by definition, $\|\cdot\|^2$ is convex. ■

EXAMPLE 11.8.4. The distance function to a convex set is convex.

EXAMPLE 11.8.5. The perspective of a convex function is convex. i.e., if $f : \mathbb{E} \rightarrow \mathbb{R}$

Chapter 12

More Convex Functions

12.1 Strictly Convex

DEFINITION (Strictly Convex). Let f be a proper function from \mathbb{E} to \mathbb{R}^* . We say that f is **strictly convex** if $\forall x, y \in \text{dom}(f)$, $\forall \lambda \in [0, 1]$, we have $f(\lambda x + (1 - \lambda)y) < \lambda f(x) + (1 - \lambda)f(y)$, except when $\lambda x + (1 - \lambda)y = x$ or y .

PROPOSITION 12.1.1. Strictly convex functions are convex.

12.2 Strongly Convex

DEFINITION (Strongly Convex). Let f be a proper function from \mathbb{E} to \mathbb{R}^* . Let β be a positive constant. We say that f is β **-strongly convex** if $\forall x, y \in \text{dom}(f)$, $\forall \lambda \in [0, 1]$, we have

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) - \frac{\beta}{2} \lambda(1 - \lambda) \|x - y\|^2.$$

PROPOSITION 12.2.1. Strongly convex functions are strictly convex.

PROPOSITION 12.2.2. Let f be a function from \mathbb{E} to \mathbb{R}^* . Then f is β -strongly convex if and only if $f - \frac{\beta}{2}\|\cdot\|^2$ is convex.

Proof. Let β be a positive constant. Let g denote $f - \frac{\beta}{2}\|\cdot\|^2$. Let x and y be arbitrary elements of \mathbb{E} . Let $\lambda \in (0, 1)$ be arbitrary.

$$\begin{aligned}
& f \text{ is } \beta\text{-strongly convex} \\
\iff & f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) \\
& \quad - \frac{\beta}{2}\lambda(1 - \lambda)\|x - y\|^2 \\
\iff & f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) \\
& \quad - \frac{\beta}{2}\lambda(1 - \lambda)(\|x\|^2 + \|y\|^2 - 2\langle x, y \rangle) \\
\iff & f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) \\
& \quad - \lambda\frac{\beta}{2}\|x\|^2 + \frac{\beta}{2}\lambda^2\|x\|^2 \\
& \quad - (1 - \lambda)\frac{\beta}{2}\|y\|^2 + \frac{\beta}{2}(1 - \lambda)^2\|y\|^2 \\
& \quad + \beta\lambda(1 - \lambda)\langle x, y \rangle \\
\iff & f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) \\
& \quad - \lambda\frac{\beta}{2}\|x\|^2 - (1 - \lambda)\frac{\beta}{2}\|y\|^2 \\
& \quad + \frac{\beta}{2}\|\lambda x\|^2 + \frac{\beta}{2}\|(1 - \lambda)y\|^2 + \beta\langle \lambda x, (1 - \lambda)y \rangle \\
\iff & f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) \\
& \quad - \lambda\frac{\beta}{2}\|x\|^2 - (1 - \lambda)\frac{\beta}{2}\|y\|^2 \\
& \quad + \frac{\beta}{2}\|\lambda x + (1 - \lambda)y\|^2 \\
\iff & g(\lambda x + (1 - \lambda)y) \leq \lambda g(x) + (1 - \lambda)g(y) \\
\iff & f - \frac{\beta}{2}\|\cdot\|^2 \text{ is } \beta \text{ convex.}
\end{aligned}$$

■

Question: Can we allow f to take on $-\infty$? Do we need f to be proper?

PROPOSITION 12.2.3. Let f and g be functions from \mathbb{E} to \mathbb{R}^* . Suppose f is

β -strongly convex for some positive constant β and g is convex. Then $f + g$ is also β -strongly convex.

Question: Can we allow f or g to take on $-\infty$? Do we need f and g to be proper?

Proof.

$$\begin{aligned}
 & f \text{ is } \beta\text{-strongly convex} \\
 \implies & f - \frac{\beta}{2} \|\cdot\|^2 \text{ is convex} \\
 \implies & f + g - \frac{\beta}{2} \|\cdot\|^2 \text{ is convex} \\
 \implies & f + g \text{ is } \beta\text{-strongly convex.}
 \end{aligned}$$

■

12.3 Quasiconvex

DEFINITION (Quasiconvex). Let $f : \mathbb{E} \rightarrow \mathbb{R}$ be a function with convex domain. We say that f is **quasiconvex** if any level set of f is convex.

PROPOSITION 12.3.1 (Jensen's Inequality for Quasiconvex Functions). Let f be a quasiconvex function. Then $\forall x, y \in \text{dom}(f)$, $\forall \alpha, \beta \in [0, 1]$ such that $\alpha + \beta = 1$,

$$f(\alpha x + \beta y) \leq \max\{f(x), f(y)\}.$$

PROPOSITION 12.3.2. A differentiable real-valued function f with convex domain is convex if and only if $\forall x, y \in \text{dom}(f)$,

$$f(y) \leq f(x) \implies \nabla f(x) \cdot (y - x) \leq 0. \quad ???$$

Not sure where did this come from but I don't think this is correct.

Chapter 13

Support

13.1 Definitions

DEFINITION (Support Function). Let S be a subset of \mathbb{E} . We define the **support function** of S , denoted by σ_S , to be a function from \mathbb{E} to \mathbb{R}^* given by

$$\sigma_S(x) = \sup_{s \in S} \langle x, s \rangle.$$

DEFINITION (Supporting Hyperplane). Let S be a set in \mathbb{E} with nonempty boundary. Let x_0 be a point in the boundary of S . We define a **supporting hyperplane** H to set S at point x_0 to be a set of the form

$$H = \{x \in \mathbb{E} : a^T x = a^T x_0\},$$

such that $a \in \mathbb{E}$ and $a \neq \vec{0}$ and $\forall x \in S, a^T x \leq a^T x_0$.

13.2 Properties

PROPOSITION 13.2.1. The support function of a non-empty set S is proper, convex, and lower semi-continuous.

Proof.

Part 1. Proper.

Define f_s to be a function from \mathbb{E} to \mathbb{R} by $f_s(x) = \langle s, x \rangle$.

These functions are linear and hence proper, convex, and lower semi-continuous.

Notice $\sigma_S = \sup_{s \in S} f_s$.

So σ_S is convex and lower semi-continuous.

Since $\sigma_S(0) = \sup_{s \in S} \langle 0, s \rangle = 0$, $\exists x_0 \in \mathbb{E}, \sigma_S(x) \neq +\infty$.

Since $\sigma_S(x) = \sup_{s \in S} \langle x, s \rangle \geq \langle x, s \rangle \neq -\infty$, $\forall x \in \mathbb{E}, \sigma_S(x) \neq -\infty$.

Since $\exists x_0 \in \mathbb{E}, \sigma_S(x) \neq +\infty$ and $\forall x \in \mathbb{E}, \sigma_S(x) \neq -\infty$, by definition, σ_S is proper. ■

PROPOSITION 13.2.2. The support function of a non-empty and bounded set is continuous.

Proof.

Let x_0 be an arbitrary point in \mathbb{E} . Let ε be an arbitrary positive number. Define $M := \sup_{y \in C} \|y\| + 1$. Since C is bounded, M is finite. Define $\delta := \varepsilon/M$. Let x be an arbitrary point such that $\|x - x_0\| < \delta$. Let y be an arbitrary point in \mathbb{E} . Then by the Cauchy Schwarz inequality, we have

$$\langle x - x_0, y \rangle \leq \|x - x_0\| \|y\|.$$

That is,

$$\langle x, y \rangle \leq \|x - x_0\| \|y\| + \langle x_0, y \rangle.$$

It follows that

$$\begin{aligned} \sup_{y \in C} \langle x, y \rangle &\leq \sup_{y \in C} (\|x - x_0\| \|y\| + \langle x_0, y \rangle) \\ &\leq \|x - x_0\| \sup_{y \in C} \|y\| + \sup_{y \in C} \langle x_0, y \rangle. \end{aligned}$$

That is,

$$\sigma_C(x) \leq \sigma_C(x_0) + \|x - x_0\| \sup_{y \in C} \|y\|.$$

By definition of δ and M ,

$$\sigma_C(x) < \sigma_C(x_0) + \varepsilon. \tag{1}$$

Similarly, reversing the role of x and x_0 , we can prove that

$$\sigma_C(x_0) < \sigma_C(x) + \varepsilon. \tag{2}$$

From (1) and (2) we get

$$|\sigma_C(x) - \sigma_C(x_0)| < \varepsilon.$$

Since $\forall \varepsilon > 0, \exists \delta > 0$ such that $|\sigma_C(x) - \sigma_C(x_0)| < \varepsilon$ whenever $\|x - x_0\| < \delta$, by definition, δ_C is continuous. ■

PROPOSITION 13.2.3. Let S be a subset of \mathbb{E} . Then $\sigma_S = \sigma_{\text{conv}(S)} = \sigma_{\text{clconv}(S)}$.

Proof.

Let x be an arbitrary point in \mathbb{E} .

$$\begin{aligned}\sigma_S(x) &= \sup \{ \langle x, s \rangle : s \in S \} \\ \sigma_{\text{conv}(S)}(x) &= \sup \{ \langle x, s \rangle : s \in \text{conv}(S) \} \\ \sigma_{\text{clconv}(S)}(x) &= \sup \{ \langle x, s \rangle : s \in \text{clconv}(S) \}.\end{aligned}$$

It is easy to see that by the linearity of inner products,

$$\text{conv} \{ \langle x, s \rangle : s \in S \} = \{ \langle x, s \rangle : s \in \text{conv}(S) \}.$$

It is easy to see that by the linearity and the continuity of inner products,

$$\text{clconv} \{ \langle x, s \rangle : s \in S \} = \{ \langle x, s \rangle : s \in \text{clconv}(S) \}.$$

It is also easy to see that for any subset A of the reals,

$$\sup(A) = \sup(\text{conv}(A)),$$

and

$$\sup(A) = \sup(\text{cl}(A)).$$

So

$$\begin{aligned}\sigma_S(x) &= \sup \{ \langle x, s \rangle : s \in S \} \\ &= \sup \text{conv} \{ \langle x, s \rangle : s \in S \} \\ &= \sup \{ \langle x, s \rangle : s \in \text{conv}(S) \} \\ &= \sigma_{\text{conv}(S)}(x).\end{aligned}$$

That is, $\sigma_S(x) = \sigma_{\text{conv}(S)}(x)$.

$$\begin{aligned}\sigma_S(x) &= \sup \{ \langle x, s \rangle : s \in S \}\end{aligned}$$

$$\begin{aligned}
&= \sup \operatorname{conv} \{ \langle x, s \rangle : s \in S \} \\
&= \sup \{ \langle x, s \rangle : s \in \operatorname{conv}(S) \} \\
&= \sup \operatorname{cl} \{ \langle x, s \rangle : s \in \operatorname{conv}(S) \} \\
&= \sup \{ \langle x, s \rangle : s \in \operatorname{cl}(\operatorname{conv}(S)) \} \\
&= \sup \{ \langle x, s \rangle : s \in \operatorname{clconv}(S) \} \\
&= \sigma_{\operatorname{clconv}(S)}(x).
\end{aligned}$$

That is, $\sigma_S(x) = \sigma_{\operatorname{clconv}(S)}(x)$.

■

13.3 Supporting Hyperplane

THEOREM 13.1 (Supporting Hyperplane Theorem). For any boundary point x_0 of a convex set C , there exists a supporting hyperplane to C at x_0 .

Chapter 14

Conjugacy

14.1 Definition and Examples

DEFINITION (Convex Conjugate (Legendre–Fenchel Transformation)). Let f be a function from \mathbb{E} to \mathbb{R}^* . We define the **convex conjugate** of f , denoted by f^* , to be a function also from \mathbb{E} to \mathbb{R}^* given by

$$f^*(x) := \sup_{y \in \mathbb{E}} \{ \langle y, x \rangle - f(y) \}.$$

EXAMPLE 14.1.1. Let S be a subset of \mathbb{E} . Then $\delta_S^* = \sigma_S$.

Proof. Recall that

$$\delta_S(x) = \begin{cases} 0, & x \in S, \\ +\infty, & x \notin S, \end{cases}$$
$$\sigma_S(x) = \sup_{y \in S} \langle x, y \rangle.$$

Now for any $x \in \mathbb{E}$,

$$\begin{aligned} \delta_S^*(x) &= \sup_{y \in S} (\langle x, y \rangle - \delta_S(y)) \\ &= \sup_{y \in S} (\langle x, y \rangle - 0) \end{aligned}$$

$$\begin{aligned}
&= \sup_{y \in S} \langle x, y \rangle \\
&= \sigma_S(x).
\end{aligned}$$

So $\delta_S^* = \sigma_S$. ■

14.2 Basic Properties

PROPOSITION 14.2.1. The convex conjugate function is convex.

Proof. If $\text{dom}(f) = \emptyset$, then one can see that $f^* \equiv -\infty$. It is a pointwise supremum of affine functions. ■

PROPOSITION 14.2.2. The convex conjugate function is lower semi-continuous.

14.3 Double Conjugate

PROPOSITION 14.3.1. Let f be any function from \mathbb{E} to \mathbb{R}^* . Then $f^{**} \leq f$.

Proof. Let x be an arbitrary point in \mathbb{E} .

$$\begin{aligned}
&f^{**}(x) \\
&= \sup_{y \in \mathbb{E}} \{ \langle y, x \rangle - f^*(y) \} \\
&= \sup_{y \in \mathbb{E}} \left\{ \langle y, x \rangle - \sup_{z \in \mathbb{E}} \{ \langle z, y \rangle - f(z) \} \right\} \\
&\leq \sup_{y \in \mathbb{E}} \left\{ \langle y, x \rangle - \{ \langle x, y \rangle - f(x) \} \right\} \\
&= \sup_{y \in \mathbb{E}} \{ f(x) \} \\
&= f(x).
\end{aligned}$$

That is, $f^{**}(x) \leq f(x)$. Since $\forall x \in \mathbb{E}$, $f^{**}(x) \leq f(x)$, we get $f^{**} \leq f$. ■

PROPOSITION 14.3.2. Let f be a proper function. Then f is convex and lower semi-continuous if and only if

$$f^{**} = f.$$

PROPOSITION 14.3.3. Let f and g be functions from \mathbb{E} to \mathbb{R}^* . Then $f \leq g$ implies $f^* \geq g^*$ and $f^{**} \leq g^{**}$.

Proof. Let x be an arbitrary point in \mathbb{E} .

$$\begin{aligned} f^*(x) &= \sup_{y \in \mathbb{E}} \{ \langle y, x \rangle - f(y) \} \\ &\geq \sup_{y \in \mathbb{E}} \{ \langle y, x \rangle - g(y) \} \\ &= g^*(x). \end{aligned}$$

That is, $f^*(x) \geq g^*(x)$. Since $\forall x \in \mathbb{E}$, $f^*(x) \geq g^*(x)$, we get $f^* \geq g^*$.

Let x be an arbitrary point in \mathbb{E} .

$$\begin{aligned} f^{**}(x) &= \sup_{y \in \mathbb{E}} \{ \langle y, x \rangle - f^*(y) \} \\ &= \sup_{y \in \mathbb{E}} \left\{ \langle y, x \rangle - \sup_{z \in \mathbb{E}} \{ \langle z, y \rangle - f(z) \} \right\} \\ &\leq \sup_{y \in \mathbb{E}} \left\{ \langle y, x \rangle - \sup_{z \in \mathbb{E}} \{ \langle z, y \rangle - g(z) \} \right\} \\ &= \sup_{y \in \mathbb{E}} \{ \langle y, x \rangle - g^*(y) \} \\ &= g^{**}(x). \end{aligned}$$

That is, $f^{**}(x) \leq g^{**}(x)$. Since $\forall x \in \mathbb{E}$, $f^{**}(x) \leq g^{**}(x)$, we get $f^{**} \leq g^{**}$. ■

PROPOSITION 14.3.4.

$$\text{epi}(f^{**}) = \text{conv}(\text{epi}(f)).$$

14.4 Conjugates and Sub-Differentials

THEOREM 14.1 (Fenchel-Young). Let f be a proper function from \mathbb{E} to \mathbb{R}^* . Then $\forall x, y \in \mathbb{E}$, we have

$$f(x) + f^*(y) \geq \langle x, y \rangle.$$

PROPOSITION 14.4.1. Let f be a proper closed convex function from \mathbb{E} to \mathbb{R}^* . Then $\forall x, y \in \mathbb{E}$,

$$y \in \partial f(x) \iff x \in \partial f^*(y) \iff f(x) + f^*(y) = \langle x, y \rangle.$$

Proof of $y \in \partial f(x) \iff x \in \partial f^(y)$.* For one direction, assume that $y \in \partial f(x)$. We are to prove that $x \in \partial f^*(y)$. Consider an arbitrary point $z \in \mathbb{E}$. Since $y \in \partial f(x)$, we get

$$\forall u \in \mathbb{E}, \quad \langle y, u - x \rangle \leq f(u) - f(x).$$

Rearranging yields

$$\forall u \in \mathbb{E}, \quad \langle y, u \rangle - f(u) \leq \langle y, x \rangle - f(x).$$

It follows that

$$\sup_{u \in \mathbb{E}} (\langle y, u \rangle - f(u)) \leq \langle y, x \rangle - f(x). \quad (1)$$

By definition of supremum, we have

$$\sup_{u \in \mathbb{E}} (\langle y, u \rangle - f(u)) \geq \langle y, x \rangle - f(x). \quad (2)$$

From (1) and (2), we get

$$\sup_{u \in \mathbb{E}} (\langle y, u \rangle - f(u)) = \langle y, x \rangle - f(x).$$

That is,

$$f^*(y) = \langle y, x \rangle - f(x).$$

Then

$$\begin{aligned} & f^*(z) - f^*(y) \\ &= \sup_{u \in \mathbb{E}} (\langle z, u \rangle - f(u)) - \sup_{u \in \mathbb{E}} (\langle y, u \rangle - f(u)) \\ &= \sup_{u \in \mathbb{E}} (\langle z, u \rangle - f(u)) - \langle y, x \rangle + f(x) \\ &\geq \langle z, x \rangle - f(x) - \langle y, x \rangle + f(x) \end{aligned}$$

$$= \langle z - y, x \rangle.$$

That is,

$$\langle x, z - y \rangle \leq f^*(z) - f^*(y).$$

So $x \in \partial f^*(y)$. This proves

$$y \in \partial f(x) \implies x \in \partial f^*(y).$$

Since $f^{**} = f$, similarly, we can prove that

$$x \in \partial f^*(y) \implies y \in \partial f(x).$$

■

PROPOSITION 14.4.2. Let f be a proper convex function from \mathbb{E} to \mathbb{R}^* . Let x be a point in \mathbb{E} . Assume that $\partial f(x) \neq \emptyset$. Then $f^{**}(x) = f(x)$.

Chapter 15

The Proximal Operator

15.1 Definitions

DEFINITION (Proximal Operator). Let f be a function from \mathbb{E} to \mathbb{R}^* . We define the **proximal operator** of f , denoted by prox_f , to be a function from \mathbb{E} to $\mathcal{P}(\mathbb{E})$ given by

$$\text{prox}_f(x) := \underset{y \in \mathbb{E}}{\text{argmin}} \left\{ f(y) + \frac{1}{2} \|y - x\|^2 \right\}.$$

15.2 Examples

EXAMPLE 15.2.1 (Soft Threshold). Let $\lambda \geq 0$. Let f be a function from \mathbb{R} to \mathbb{R} given by $f(x) := \lambda|x|$. Then

$$\text{prox}_f(x) = \begin{cases} x + \lambda, & \text{if } x < -\lambda \\ 0, & \text{if } -\lambda \leq x \leq \lambda \\ x - \lambda, & \text{if } x > \lambda. \end{cases}$$

15.3 Basic Properties

(bug)

PROPOSITION 15.3.1. Let f be a proper, convex, and lower semi-continuous function from \mathbb{E} to \mathbb{R}^* . Then $\forall x \in \mathbb{E}$, $\text{prox}_f(x)$ is a singleton set.

Proof. Let x be an arbitrary element of \mathbb{E} . Define a function $h : \mathbb{E} \rightarrow \mathbb{R}^*$ by $h(y) := \frac{1}{2}\|y-x\|^2$. Define a function $g : \mathbb{E} \rightarrow \mathbb{R}^*$ by $g(y) := f(y) + h(y)$. Then $\text{prox}_f(x) = \underset{y \in \mathbb{E}}{\text{argmin}} g(y)$. Note that h is proper, lower semi-continuous, and β -strongly convex for any $\beta \in (0, 1)$. Since f and h are proper, g is proper (why?). Since f and h are lower semi-continuous, g is lower semi-continuous. Since f is convex and h is β -strongly convex, g is β -strongly convex. Since g is proper, lower semi-continuous, and strongly convex, g has a unique minimizer (why?). So $\text{prox}_f(x)$ is a singleton set. ■

not fully understood

PROPOSITION 15.3.2. Let C be a nonempty closed convex subset of \mathbb{E} . Then prox_{δ_C} and proj_C are both singleton and $\text{prox}_{\delta_C} = \text{proj}_C$.

Proof. Since C is nonempty, convex, and closed, δ_C is proper, convex, and lower semi-continuous and hence prox_{δ_C} is singleton. Since C is nonempty, convex, and closed, proj_C is singleton. Let x and p be arbitrary elements of \mathbb{E} . Then

$$\begin{aligned}
 & p \in \text{prox}_{\delta_C}(x) \\
 \iff & p \in \underset{y \in \mathbb{E}}{\text{argmin}} \{ \delta_C(y) + \frac{1}{2}\|y-x\|^2 \} \\
 \iff & \forall y \in \mathbb{E}, \delta_C(y) + \frac{1}{2}\|y-x\|^2 \geq \delta_C(p) + \frac{1}{2}\|p-x\|^2 \\
 \iff & p \in C \text{ and } \forall y \in C, \frac{1}{2}\|y-x\|^2 \geq \frac{1}{2}\|p-x\|^2 \\
 \iff & p \in C \text{ and } \forall y \in C, \|y-x\| \geq \|p-x\| \\
 \iff & p \in \underset{y \in C}{\text{argmin}} \|y-x\| \\
 \iff & p \in \text{proj}_C(x).
 \end{aligned}$$

■

PROPOSITION 15.3.3 (Firmly Non-Expansive). Let f be a proper closed convex function. Then prox_f is firmly non-expansive.

15.4 Prox Calculus Rules

PROPOSITION 15.4.1 (Scaling and Translation).

THEOREM 15.1 (Norm Composition).

PROPOSITION 15.4.2. Let f_1, \dots, f_m be proper, convex, and lower semi-continuous functions from \mathbb{R} to \mathbb{R}^* . Define a function $f : \mathbb{R}^m \rightarrow \mathbb{R}^*$ by $f((x_i)_{i=1}^m) := \sum_{i=1}^m f_i(x_i)$. Then

$$\text{prox}_f((x_i)_{i=1}^m) = (\text{prox}_{f_i}(x_i))_{i=1}^m.$$

Proof. Since each f_i is proper, convex, and lower semi-continuous, f is proper, convex, and lower semi-continuous. Let $(x_i)_{i=1}^m$ and $(p_i)_{i=1}^m$ be arbitrary elements of \mathbb{R}^m . Then

$$(p_i)_{i=1}^m = \text{prox}_f((x_i)_{i=1}^m)$$

$$\iff$$

■

15.5 The Second Prox Theorem

PROPOSITION 15.5.1. Let f be a proper, convex, and lower semi-continuous function from \mathbb{E} to \mathbb{R}^* . Let x and p be points in \mathbb{E} . Then $p = \text{prox}_f(x)$ if and only if

$$x - p \in \partial f(p).$$

PROPOSITION 15.5.2. Let f be a proper, convex, and lower semi-continuous function from \mathbb{E} to \mathbb{R}^* . Let x and p be elements of \mathbb{E} . Then $p = \text{prox}_f(x)$ if and only if

$$\forall y \in \mathbb{E}, \quad \langle y - p, x - p \rangle \leq f(y) - f(p).$$

Proof. Forward Direction:

Assume that $p = \text{prox}_f(x)$. I will show that $\forall y \in \mathbb{E}$, $\langle y - p, x - p \rangle \leq f(y) - f(p)$. Let y be an arbitrary element of \mathbb{E} . Define for each $\lambda \in (0, 1)$ a point p_λ by $p_\lambda := \lambda y + (1 - \lambda)p$. Then

$$\begin{aligned}
& p = \text{prox}_f(x) \\
& \implies f(p) + \frac{1}{2}\|x - p\|^2 \leq f(p_\lambda) + \frac{1}{2}\|x - p_\lambda\|^2 \\
& \iff f(p) \leq f(p_\lambda) + \frac{1}{2}\|x - p_\lambda\|^2 - \frac{1}{2}\|x - p\|^2 \\
& \iff f(p) \leq f(p_\lambda) + \frac{1}{2} \left\langle \left[(x - p_\lambda) + (x - p) \right], \left[(x - p_\lambda) - (x - p) \right] \right\rangle \\
& \iff f(p) \leq f(p_\lambda) + \frac{1}{2} \left\langle \left[2x - \lambda y - (1 - \lambda)p - p \right], \left[p - \lambda y - (1 - \lambda)p \right] \right\rangle \\
& \iff f(p) \leq f(p_\lambda) + \frac{1}{2} \left\langle \left[2(x - p) + \lambda(p - y) \right], \left[\lambda(p - y) \right] \right\rangle \\
& \iff f(p) \leq f(p_\lambda) + \lambda \langle x - p, p - y \rangle + \frac{1}{2}\lambda^2\|p - y\|^2 \\
& \iff f(p) \leq f(\lambda y + (1 - \lambda)p) + \lambda \langle x - p, p - y \rangle + \frac{1}{2}\lambda^2\|p - y\|^2 \\
& \implies f(p) \leq \lambda f(y) + (1 - \lambda)f(p) + \lambda \langle x - p, p - y \rangle + \frac{1}{2}\lambda^2\|p - y\|^2 \\
& \iff \lambda \langle y - p, x - p \rangle \leq \lambda f(y) - \lambda f(p) + \frac{1}{2}\lambda^2\|p - y\|^2 \\
& \iff \langle y - p, x - p \rangle \leq f(y) - f(p) + \frac{1}{2}\lambda\|p - y\|^2 \\
& \iff \langle y - p, x - p \rangle \leq f(y) - f(p).
\end{aligned}$$

Backward Direction:

Assume that $\forall y \in \mathbb{E}$, $\langle y - p, x - p \rangle \leq f(y) - f(p)$. I will show that $p = \text{prox}_f(x)$. Let y be an arbitrary element of \mathbb{E} . Then

$$\begin{aligned}
& \langle y - p, x - p \rangle \leq f(y) - f(p) \\
& \iff f(p) + \frac{1}{2}\|x - p\|^2 \leq f(y) + \langle x - p, p - y \rangle + \frac{1}{2}\|x - p\|^2 \\
& \implies f(p) + \frac{1}{2}\|x - p\|^2 \leq f(y) + \langle x - p, p - y \rangle + \frac{1}{2}\|x - p\|^2 + \frac{1}{2}\|p - y\|^2 \\
& \iff f(p) + \frac{1}{2}\|x - p\|^2 \leq f(y) + \frac{1}{2}\|(x - p) + (p - y)\|^2 \\
& \iff f(p) + \frac{1}{2}\|x - p\|^2 \leq f(y) + \frac{1}{2}\|x - y\|^2 \\
& \implies p = \text{prox}_f(x).
\end{aligned}$$

This completes the proof. ■

15.6 Moreau Decomposition

THEOREM 15.2 (Moreau Decomposition). Let f be a proper closed convex function from \mathbb{E} to \mathbb{R}^* . Then

$$\text{prox}_f + \text{prox}_{f^*} = \text{id}.$$

Proof. Let x be an arbitrary point in \mathbb{E} . We are to prove that

$$\text{prox}_f(x) + \text{prox}_{f^*}(x) = x.$$

Let p denote $\text{prox}_f(x)$. Since f is proper convex and lower semi-continuous and $p = \text{prox}_f(x)$, we get

$$x - p \in \partial f(p).$$

Since $x - p \in \partial f(p)$, we get $p \in \partial f^*(x - p)$. It follows that $x - p = \text{prox}_{f^*}(x)$. Substitute $p = \text{prox}_f(x)$ and rearrange the equation, we get

$$\text{prox}_f(x) + \text{prox}_{f^*}(x) = x.$$

■

Chapter 16

Ellipsoids

DEFINITION (Ellipsoid). Let v be a point in some Euclidean space \mathbb{E} . We define an **ellipsoid**, centered at point v , to be a set of the form

$$\{x \in \mathbb{E} : (x - v)^T A (x - v) = 1\}$$

where A is some d by d positive definite matrix.

16.1 Properties

PROPOSITION 16.1.1. The eigenvectors of A define the principal axes of the ellipsoid.