

Functional Analysis

Daniel Mao

Contents

1	Balanced Sets	1
1.1	Definitions	1
1.2	Properties	1
1.3	Stability of Balance	2
1.4	Absorbing Sets	2
2	Normed Linear Spaces	3
2.1	Definitions	3
2.2	Properties	3
2.3	Equivalence of Norms	4
2.4	Dual Norms	5
2.5	p -norms	6
3	Inner Product Spaces	9
3.1	Inner Products	9
3.2	Inner Product Space	10
3.3	Inequalities	10
4	Orthogonality	13
4.1	Orthogonal Sets	13
4.2	Orthogonal Bases	13
4.3	Orthogonal Complements	14
4.4	Orthogonal Projection	15
4.5	Inequalities in Hilbert Spaces	15
5	Sequence Spaces	17
5.1	ℓ^p Space	17
5.2	c_0 Space and c_{00} Space	18
5.3	Hölder's Inequality	21

6	Function Spaces	23
6.1	The \mathcal{L}^p Norm	23
7	Banach Space	25
7.1	Definition	25
7.2	Properties	25
7.3	Examples of Banach Space	25
7.4	Construction of Banach Spaces	26
8	Hilbert Space	27
8.1	Definition	27
8.2	Examples of Hilbert Space	27
8.3	Properties of Hilbert Space	27
9	Operators	31
9.1	Bounded Operators	31
9.2	Examples of Bounded Operators	31
9.3	The Space of Bounded Operators	34
9.4	Invertible Bounded Operators	34
10	Dual Space	37
10.1	Definition	37
10.2	Properties	37
10.3	Examples of Dual Space	37
11	Quotient Spaces	39
11.1	Definitions	39
12	Topological Vector Space	41
12.1	Definitions	41
12.2	Examples	42
12.3	Neighborhood Improvements	42
13	Equicontinuity in Metric Spaces	43
13.1	Definitions	43
13.2	Sufficient Conditions	43
14	Adjoint Operator	45
14.1	Definitions	45
14.2	Properties of the Adjoint Operator	45
14.3	Normal Operators	46

<i>CONTENTS</i>	iii
14.4 Self-adjoint	46
15 Convolution	47
16 Coercive Functions	49
16.1 Definitions	49
16.2 Properties	49
17 Unclassified Results	51

Chapter 1

Balanced Sets

1.1 Definitions

Definition (Balanced Sets). *Let X be a vector space over field \mathbb{F} . Let S be a subset of X . We say that S is **balanced** if*

$$\forall a \in \mathbb{F} : |a| \leq 1, \quad aS \subseteq S.$$

Definition (Balanced Hull). *Let X be a vector space over field \mathbb{F} . Let S be a subset of X . We define the **balanced hull** of S , denoted by $\text{balhull}(S)$, to be the smallest balanced set containing S .*

Definition (Balanced Core). *Let X be a vector space over field \mathbb{F} . Let S be a subset of X . We define the **balanced core** of S , denoted by $\text{balcore}(S)$, to be the largest balanced set contained in S .*

1.2 Properties

Proposition 1.2.1. *Let X be a vector space over field \mathbb{F} . Let B be a balanced subset of X . Then*

$$\forall a, b \in \mathbb{F} : |a| \leq |b|, \quad aB \subseteq bB.$$

Proposition 1.2.2. *Balanced sets are path connected.*

Proposition 1.2.3 (Act on Other Properties).

- *The balanced hull of a compact set is compact.*

- *The balanced hull of a totally bounded set is totally bounded.*
- *The balanced hull of a bounded set is bounded.*

Proposition 1.2.4 (Act on Other Properties). • *The balanced core of a closed set is closed.*

Proposition 1.2.5. *Let X be a vector space over field \mathbb{F} . Let a be a scalar in field \mathbb{F} . Then*

$$a \operatorname{balhull}(S) = \operatorname{balhull}(aS).$$

1.3 Stability of Balance

Proposition 1.3.1 (Set Operations). • *The union of balanced sets is also balanced.*

- *The intersection of balanced sets is also balanced.*

Proposition 1.3.2 (Linear Mappings). • *The scalar multiple of a balanced set is also balanced.*

- *The (Minkowski) sum of two balanced sets is also balanced.*
- *The image of a balanced set under a linear operator is also balanced.*
- *The inverse image of a balanced set under a linear operator is also balanced.*

Proposition 1.3.3 (Topological Operations). *The closure of a balanced set is also balanced.*

Proposition 1.3.4. *The convex hull of a balanced set is also balanced (and also convex).*

1.4 Absorbing Sets

Definition (Absorbing Sets). *Let \mathfrak{X} be a vector space over field \mathbb{F} . Let S be a subset of X . We say that S is **absorbing** if*

$$\forall x \in \mathfrak{X}, \quad \exists r \in \mathbb{R} : r > 0, \quad \forall c \in \mathbb{F} : |c| \geq r, \quad x \in cS.$$

i.e.,

$$\bigcup_{n \in \mathbb{N}} nS = \mathfrak{X}.$$

Proposition 1.4.1. *Every absorbing set contains the origin.*

Chapter 2

Normed Linear Spaces

2.1 Definitions

Definition (Seminorm). Let \mathfrak{X} be a vector space over field \mathbb{F} . We define a **seminorm** on \mathfrak{X} , denoted by ν , to be a map from \mathfrak{X} to \mathbb{R} that satisfies the following conditions.

(1) $\forall x \in \mathfrak{X}, \quad \nu(x) \geq 0.$

(2) $\forall \lambda \in \mathbb{F}, \forall x \in \mathfrak{X}, \quad \nu(\lambda x) = \lambda \nu(x).$

(3) *Triangle Inequality.*

$$\forall x, y \in \mathfrak{X}, \quad \nu(x + y) \leq \nu(x) + \nu(y).$$

The idea behind the seminorm is that we are trying to give our vector space a notion of “length” of vectors.

Definition (Norm). Let \mathfrak{X} be a vector space over field \mathbb{F} . We define a **norm** on \mathfrak{X} , denoted by ν , to be a seminorm on \mathfrak{X} that satisfies the additional condition:

$$\forall x \in \mathfrak{X}, \quad \nu(x) = 0 \iff x = 0.$$

2.2 Properties

Proposition 2.2.1. Let $(V, \|\cdot\|_V)$ be a normed vector space over field \mathbb{F} . Then $(V, \|\cdot\|)$ is complete if and only if $(\overline{B(0,1)}, \|\cdot\|_V)$ is complete.

Proof.

For one direction, assume that $(V, \|\cdot\|)$ is complete.

We are to prove that $(\overline{B(0,1)}, \|\cdot\|_V)$ is complete.

Since $(\overline{B(0,1)}, \|\cdot\|_V)$ is a closed subspace of $(V, \|\cdot\|)$ and $(V, \|\cdot\|)$ is complete, $(\overline{B(0,1)}, \|\cdot\|_V)$ is also complete.

For the reverse direction, assume that $(\overline{B(0,1)}, \|\cdot\|_V)$ is complete.

We are to prove that $(V, \|\cdot\|_V)$ is complete.

Let $\{x_i\}_{i \in \mathbb{N}}$ be an arbitrary Cauchy sequence in $(V, \|\cdot\|_V)$.

Since $\{x_i\}_{i \in \mathbb{N}}$ is Cauchy in $(V, \|\cdot\|_V)$, $\{x_i\}_{i \in \mathbb{N}}$ is bounded in $(V, \|\cdot\|_V)$.

Let λ be a positive upper bound for $\{\|x_i\|_V\}_{i \in \mathbb{N}}$.

Since $\{x_i\}_{i \in \mathbb{N}}$ is Cauchy in $(V, \|\cdot\|_V)$, $\{x_i/\lambda\}_{i \in \mathbb{N}}$ is Cauchy in $(\overline{B(0,1)}, \|\cdot\|_V)$.

Since $\{x_i/\lambda\}_{i \in \mathbb{N}}$ is Cauchy in $(\overline{B(0,1)}, \|\cdot\|_V)$ and $(\overline{B(0,1)}, \|\cdot\|_V)$ is complete, $\{x_i/\lambda\}_{i \in \mathbb{N}}$ converges in $(\overline{B(0,1)}, \|\cdot\|_V)$.

Since $\{x_i/\lambda\}_{i \in \mathbb{N}}$ converges in $(\overline{B(0,1)}, \|\cdot\|_V)$, $\{x_i\}_{i \in \mathbb{N}}$ converges in $(V, \|\cdot\|_V)$.

Since any Cauchy sequence in $(V, \|\cdot\|_V)$ converges in $(V, \|\cdot\|_V)$, $(V, \|\cdot\|_V)$ is complete. ■

Proposition 2.2.2. *Proper subspaces of a normed linear space has empty interior.*

Proof. Let \mathfrak{X} be a normed linear space. Let \mathcal{M} be a proper subspace of \mathfrak{X} . Assume for the sake of contradiction that \mathcal{M} has non-empty interior. Then $\exists x_0 \in \mathcal{M}$ and $\exists r > 0$ such that $\text{ball}(x_0, r) \subseteq \mathcal{M}$ where $\text{ball}(x_0, r)$ denotes the open ball centered at point x_0 with radius r . Let x be an arbitrary point in \mathfrak{X} . Define a point $y(x)$ as $y(x) := x_0 + \frac{r}{2\|x\|}x$. Then $x = \frac{2\|x\|}{r}(y - x_0)$. It is easy to verify that $\|y - x_0\| = \frac{r}{2} < r$. So $y \in \text{ball}(x_0, r)$. So $y \in \mathcal{M}$. Since $y, x_0 \in \mathcal{M}$ and \mathcal{M} is a subspace, we get $\frac{2\|x\|}{r}(y - x_0) \in \mathcal{M}$. That is, $x \in \mathcal{M}$. So $\forall x \in \mathfrak{X}, x \in \mathcal{M}$. So $\mathcal{M} = \mathfrak{X}$. This contradicts to the assumption that \mathcal{M} is a proper subspace of \mathfrak{X} . So \mathcal{M} has empty interior. ■

Proposition 2.2.3. *Closed proper subspaces of a normed linear space are nowhere dense.*

Proof. Let \mathfrak{X} be a normed linear space. Let \mathcal{M} be a closed proper subspace of \mathfrak{X} . Since \mathcal{M} is closed, $\text{cl}(\mathcal{M}) = \mathcal{M}$. So $\text{cl}(\mathcal{M}) = \mathcal{M}$ is a closed proper subspace of \mathfrak{X} . Since $\text{cl}(\mathcal{M})$ is a proper subspace of \mathfrak{X} , $\text{int}(\text{cl}(\mathcal{M})) = \emptyset$. So \mathcal{M} is nowhere dense. ■

Proposition 2.2.4. *Finite dimensional subspace of a normed linear space is closed.*

Proposition 2.2.5. *Finite-dimensional normed linear spaces are complete.*

2.3 Equivalence of Norms

Definition (Equivalence of Norms). *Let \mathfrak{X} be a vector space over field \mathbb{F} . Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be two norms on V . We say that $\|\cdot\|_1$ and $\|\cdot\|_2$ are **equivalent** if*

$$\exists c_1, c_2 > 0, \quad \forall v \in \mathfrak{X}, \quad c_1\|v\|_1 \leq \|v\|_2 \leq c_2\|v\|_1.$$

Or equivalently,

$$c_1 \|v\|_2 \leq \|v\|_1 \leq c_2 \|v\|_2.$$

Proposition 2.3.1. *The equivalence of norms is an equivalence relation.*

Theorem 1. *Let V be a finite dimensional vector space over field $\mathbb{F} = \{\mathbb{R}, \mathbb{C}\}$. Then any two norms on V are equivalent.*

Proof.

Let $\|\cdot\|_p$ be an arbitrary p -norm on V and $\|\cdot\|$ be an arbitrary norm on V .

Let \mathcal{B} be the standard basis for V . Say $\mathcal{B} = \{e_1, e_2, \dots, e_n\}$.

Let v be an arbitrary vector in V .

$$\begin{aligned} \|v\| &= \left\| \sum_{i=1}^n v_i e_i \right\| \\ &\leq \sum_{i=1}^n |v_i| \|e_i\| \\ &\leq \left(\sum_{i=1}^n |v_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n \|e_i\|^{\frac{p}{p-1}} \right)^{1-\frac{1}{p}} \\ &= \left(\sum_{i=1}^n \|e_i\|^{\frac{p}{p-1}} \right)^{1-\frac{1}{p}} \|v\|_p \\ &:= c_1 \|v\|_p. \end{aligned}$$

■

Proposition 2.3.2. *Let X be a vector space. Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be two norms on X . Then $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent if and only if they generate the same metric topology.*

Proof. Convergence to 0 is equivalent under either $\|\cdot\|_1$ or $\|\cdot\|_2$. i.e., equivalent norms give rise to the same set of sequences that are convergent. Convergence of sequences defines the topology. ■

Proposition 2.3.3. *Let \mathfrak{X} be a vector space. Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be two norms on \mathfrak{X} . Let ι be the identity map from $(\mathfrak{X}, \|\cdot\|_1)$ to $(\mathfrak{X}, \|\cdot\|_2)$. Then if $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent, ι is continuous, and in fact, a homeomorphism between $(\mathfrak{X}, \|\cdot\|_1)$ and $(\mathfrak{X}, \|\cdot\|_2)$.*

2.4 Dual Norms

Definition (Dual Norm). *Let $(V, \|\cdot\|)$ be an normed vector space. We define the **dual norm** of $\|\cdot\|$, denoted by $\|\cdot\|_\circ$, to be a function given by*

$$\|v\|_\circ := \max_{\|w\|=1} v \cdot w = \max_{\|w\| \neq 0} \frac{|v \cdot w|}{\|w\|}.$$

Proposition 2.4.1. *Dual norms of norms are indeed norms.*

Proposition 2.4.2. *Let $(V, \|\cdot\|)$ be a normed vector space. Let v, w be vectors in the space. Then*

$$|v \cdot w| \leq \|v\| \cdot \|w\|.$$

2.5 p -norms

Definition (p -norm). *Let V be a finite-dimensional normed vector space over field \mathcal{F} . Let $\mathcal{B} = \{b_1, \dots, b_n\}$ be a basis for V where $n = \dim(V)$. Let v be a vector in a normed vector space. For $p \in [1, +\infty)$, we define the **p -norm** of v , denoted by $\|v\|_p$, to be the number given by*

$$\|v\|_p = \left(\sum_{i=1}^n |(v_{\mathcal{B}})_i|^p \right)^{\frac{1}{p}}.$$

Definition (Infinity Norm - 1). *Let $\mathfrak{X} = \mathbb{K}^n$ where \mathbb{K} is a field and $n \in \mathbb{N}$. We define the **infinity norm** on \mathfrak{X} , denoted by $\|\cdot\|_\infty$, to be a function given by*

$$\|v\|_\infty := \max\{|v_i|\}_{i=1}^n.$$

Definition (Infinity Norm - 2). *Let $\mathfrak{X} = \mathbb{K}^{\mathbb{N}}$. We define the **infinity norm** on \mathfrak{X} , denoted by $\|\cdot\|_\infty$, to be a function given by*

$$\|v\|_\infty := \sup_{i \in \mathbb{N}} |v_i|.$$

Definition (Infinity Norm - 3). *Let $\mathfrak{X} = \mathcal{C}([0, 1], \mathbb{C})$. We define the **infinity norm** on \mathfrak{X} , denoted by $\|\cdot\|_\infty$, to be a function given by*

$$\nu(f) := \sup_{x \in [0, 1]} |f(x)|.$$

Proposition 2.5.1. *Let $\mathfrak{X} := \mathcal{C}([0, 1], \mathbb{C})$. Let x be an arbitrary number in $[0, 1]$. Define a function ν_x on \mathfrak{X} by $\nu_x(f) := |f(x)|$. Define a function ν on \mathfrak{X} by $\nu(f) := \sup_{x \in [0, 1]} |f(x)|$. Then ν_x is a seminorm on \mathfrak{X} for each x and ν is a norm on \mathfrak{X} and we have $\nu = \sup_{x \in [0, 1]} \nu_x$.*

Proposition 2.5.2. *p -norms are indeed norms.*

Proposition 2.5.3. *For any vector v in \mathbb{R}^n , we have*

$$\lim_{p \rightarrow \infty} \|v\|_p = \|v\|_\infty.$$

i.e.,

$$\lim_{p \rightarrow \infty} \left(\sum_{i=1}^n |v_i|^p \right)^{\frac{1}{p}} = \max\{|v_i|\}_{i=1}^n.$$

Proof. Let p be an arbitrary number in $[1, +\infty)$. Let k be an arbitrary index in $\{1, \dots, n\}$. Then

$$|v_k| \leq \left(\sum_{i=1}^n |v_i|^p \right)^{1/p} = \|v\|_p.$$

So

$$\max\{|v_k|\} = \|v\|_\infty \leq \|v\|_p.$$

So

$$\lim_{p \rightarrow \infty} \|v\|_p \geq \|v\|_\infty. \quad (1)$$

On the other hand, note that

$$\left(\sum_{i=1}^n |v_i|^p \right) / \|v\|_\infty^p = \sum_{i=1}^n \left(\frac{|v_i|}{\|v\|_\infty} \right)^p$$

decreases as p increases. So it is bounded above. Say

$$\left(\sum_{i=1}^n |v_i|^p \right) / \|v\|_\infty^p \leq C$$

for some $C \in \mathbb{R}$. Then

$$\left(\sum_{i=1}^n |v_i|^p \right)^{1/p} = \|v\|_p \leq C^{1/p} \|v\|_\infty.$$

So

$$\lim_{p \rightarrow \infty} \|v\|_p \leq \lim_{p \rightarrow \infty} C^{1/p} \|v\|_\infty = \|v\|_\infty. \quad (2)$$

From (1) and (2) we get

$$\lim_{p \rightarrow \infty} \|v\|_p = \|v\|_\infty.$$

■

Proposition 2.5.4. *Let p be an arbitrary number in $[1, +\infty)$. Then the dual norm of the p -norm $\|\cdot\|_p$ is the q -norm $\|\cdot\|_q$ where q is such that satisfies*

$$\frac{1}{p} + \frac{1}{q} = 1.$$

Proposition 2.5.5. *Let p and q be numbers in $[1, +\infty]$. Let v be a vector in \mathbb{R}^n . Then if $p \leq q$,*

$$\|x\|_q \leq \|x\|_p \leq n^{\frac{1}{p} - \frac{1}{q}} \cdot \|x\|_q.$$

Proposition 2.5.6. *Let w and z be vectors in \mathbb{E}^d . Then*

$$\|w + z\|_2^2 + \|w - z\|_2^2 = 2(\|w\|_2^2 + \|z\|_2^2).$$

Chapter 3

Inner Product Spaces

3.1 Inner Products

3.1.1 Definitions

Definition (Inner Product). *Let V be a vector space over field \mathbb{F} . We define an **inner product** on V , denoted by $\langle \cdot, \cdot \rangle$, to be a scalar-valued function defined on $V \times V$ such that*

(1) *Positive Definiteness:*

$$\forall x \in V, \langle x, x \rangle \geq 0 \text{ and } \langle x, x \rangle = 0 \iff x = 0.$$

(2) *Sesqui-Linearity:*

$$\begin{aligned} \forall x, y, z, w \in V, \quad \langle x + y, z + w \rangle &= \langle x, z \rangle + \langle y, z \rangle + \langle x, w \rangle + \langle y, w \rangle, \text{ and} \\ \forall a, b \in \mathbb{F}, \forall x, y \in V, \quad \langle ax, by \rangle &= a\bar{b}\langle x, y \rangle. \end{aligned}$$

(3) *Conjugate Symmetry:*

$$\forall x, y \in V, \quad \langle x, y \rangle = \overline{\langle y, x \rangle}.$$

Definition (Induced Norm). *Let \mathfrak{X} be an inner product space over field \mathbb{K} . We define the **norm induced by $\langle \cdot, \cdot \rangle$** , denoted by $\| \cdot \|$, to be a function from \mathfrak{X} to \mathbb{R}_+ given by*

$$\|x\| := \sqrt{\langle x, x \rangle}$$

3.1.2 Examples of Inner Products

Definition (Standard Inner Product). *For $V = \mathbb{F}^n$, we define the **standard inner product** by*

$$\langle x, y \rangle := \sum_{i=1}^n x_i \bar{y}_i.$$

Definition (Frobenius Inner Product). For $V = \mathbb{F}^{n \times n}$, we define the **Frobenius inner product** by

$$\langle M_1, M_2 \rangle := \text{tr}(M_2^* M_1).$$

Definition. Let V be the space of continuous scalar-valued functions on $[0, 2\pi]$. We define the inner product on V by

$$\langle f, g \rangle := \frac{1}{2\pi} \int_0^{2\pi} f(x) \overline{g(x)} dx.$$

3.1.3 Properties

Proposition 3.1.1. Let V be a finite dimensional inner product space. Let \mathcal{B} be a basis for V . Let x and y be vectors in V . Then

$$x = y \iff \forall b \in \mathcal{B}, \quad \langle x, b \rangle = \langle y, b \rangle.$$

3.2 Inner Product Space

Definition (Inner Product Space). Let \mathfrak{X} be a vector space over field \mathbb{F} . Let $\langle \cdot, \cdot \rangle$ be an inner product on \mathfrak{X} . We define an **inner product space** to be the pair $(\mathfrak{X}, \langle \cdot, \cdot \rangle)$.

3.3 Inequalities

Theorem 2 (Minkowski).

$$\left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{p}} \leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}}$$

Proposition 3.3.1 (Cauchy-Schwarz Inequality). Let V be an inner product space. Then

$$\forall x, y \in V, \quad |\langle x, y \rangle| \leq \|x\| \cdot \|y\|$$

Proposition 3.3.2 (Triangle Inequality). Let V be an inner product space. Then

$$\forall x, y \in V, \quad \|x + y\| \leq \|x\| + \|y\|$$

Proposition 3.3.3 (Parallelogram Law). Let \mathfrak{X} be an inner product space. Then

$$\forall x, y \in \mathfrak{X}, \quad \|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2.$$

Proof.

$$\|x + y\|^2 + \|x - y\|^2 = \langle x + y, x + y \rangle + \langle x - y, x - y \rangle$$

$$\begin{aligned} &= \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle \\ &\quad + \langle x, x \rangle - \langle x, y \rangle - \langle y, x \rangle + \langle y, y \rangle \\ &= 2\langle x, x \rangle + 2\langle y, y \rangle \\ &= 2\|x\|^2 + 2\|y\|^2. \end{aligned}$$

That is,

$$\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2.$$

■

Chapter 4

Orthogonality

4.1 Orthogonal Sets

Definition (Orthogonality). *Let V be an inner product space. We say that points x and y in V are **orthogonal** if $\langle x, y \rangle = 0$.*

Definition (Orthogonal Set). *Let \mathfrak{X} be an inner product space. Let S be a subset of \mathfrak{X} . We say that S is **orthogonal** if*

$$\forall x, y \in S : x \neq y, \quad \langle x, y \rangle = 0.$$

i.e., if any two distinct vectors are orthogonal.

Definition (Orthonormal Set). *Let \mathfrak{X} be an inner product space. Let S be a set in the space. We say that S is **orthonormal** if S is orthogonal and $\forall x \in S, \|x\| = 1$ where $\|\cdot\|$ is the norm induced by the inner product.*

Proposition 4.1.1. *Orthogonal sets are linearly independent.*

4.2 Orthogonal Bases

Definition (Orthogonal Basis). *Let V be an inner product space. Let S be a set in the space. We say that S is an **orthogonal basis** for V if it is an ordered basis for V and orthogonal.*

Definition (Orthonormal Basis). *Let \mathfrak{X} be an inner product space. Let S be a set in the space. We say that S is an **orthonormal basis** for \mathfrak{X} if it is the maximal, with respect to inclusion, in the collection of all orthonormal sets in the space.*

Proposition 4.2.1. *Let V be an inner product space. Let $S = \{v_1, \dots, v_n\}$ be an orthogonal subset of V where each v_i is non-zero. Then*

$$\forall y \in \text{span}(S), \quad y = \sum_{i=1}^n \frac{\langle y, v_i \rangle}{\|v_i\|^2} v_i.$$

Theorem 3 (Gram-Schmidt Process). *Let V be an inner product space. Let $S = \{x_0, \dots, x_n\}$ be a linearly independent subset of V . Then the set $T = \{y_0, \dots, y_n\}$ given by $y_0 := x_0$ and*

$$\forall i \in \{1, \dots, n\}, \quad y_i := x_i - \sum_{j=1}^{i-1} \frac{\langle x_i, y_j \rangle}{\|y_j\|} y_j$$

is an orthogonal subset of V consisting of non-zero vectors such that $\text{span}(S) = \text{span}(S')$.

Proposition 4.2.2. *Let V be an inner product space and $S = \{v_0, v_1, \dots, v_n\}$ be an orthogonal subset of V . Then the set S' derived from the Gram-Schmidt process is exactly S .*

Theorem 4 (Parseval's Identity). *Let V be a finite-dimensional inner product space. Let $\mathcal{B} = \{v_1, \dots, v_n\}$ be an orthogonal basis for V . Then*

$$\forall x, y \in V, \quad \langle x, y \rangle = \sum_{i=1}^n \langle x, v_i \rangle \overline{\langle y, v_i \rangle}.$$

Proposition 4.2.3. *Let \mathcal{H} be a Hilbert space. Let \mathcal{E} be an orthonormal set in \mathcal{H} . Then \mathcal{E} is an orthonormal basis for \mathcal{H} if and only if*

$$\forall x \in \mathcal{H}, \quad x = \sum_{e \in \mathcal{E}} \langle x, e \rangle e.$$

4.3 Orthogonal Complements

Definition (Orthogonal Complement). *Let \mathfrak{X} be an inner product space. Let S be a non-empty subset of V . We define the **orthogonal complement** of S , denoted by S^\perp , to be a set given by*

$$S^\perp := \{x \in \mathfrak{X} : \forall s \in S, \langle x, s \rangle = 0\}.$$

i.e., the set of all points in \mathfrak{X} that are orthogonal to all vectors in S .

Proposition 4.3.1. *Let V be a finite-dimensional inner product space. Then*

$$(1) \quad V^\perp = \{0_V\}$$

$$(2) \quad \{0_V\}^\perp = V$$

Proposition 4.3.2. *Orthogonal complements are always linear subspaces.*

Proposition 4.3.3. *Let V be an inner product space and W be a subspace of V with basis β . Then a vector in V is also in W^\perp if and only if it is orthogonal to all vectors in β .*

Proposition 4.3.4 (Extension). *Let V be an n -dimensional inner product space and $S = \{v_1, v_2, \dots, v_k\}$ be an orthogonal subset of V . Then S can be extended to an orthogonal basis $B = \{v_1, v_2, \dots, v_k, v_{k+1}, \dots, v_n\}$ for V .*

Proposition 4.3.5. *Let V be an inner product space. Then*

- (1) $S \subseteq T$ implies $T^\perp \subseteq S^\perp$ for any subsets S and T of V .
- (2) $S \subseteq (S^\perp)^\perp$ for any subset S of V .

Proposition 4.3.6. *Let V be a finite-dimensional inner product space and W be a subspace of V . Then*

- (1) $W = (W^\perp)^\perp$
- (2) $V = W \oplus W^\perp$

Proposition 4.3.7. *Let V be a finite-dimensional inner product space and W_1 and W_2 be subspaces of V . Then*

- (1) $(W_1 + W_2)^\perp = W_1^\perp \cap W_2^\perp$
- (2) $(W_1 \cap W_2)^\perp = W_1^\perp + W_2^\perp$

4.4 Orthogonal Projection

Definition (Orthogonal Projection). *Let V be a vector space. Let W be a finite-dimensional subspace of V . Let x be a vector in V . We define the **orthogonal projection** of x on W , denoted by (x) , to be the vector u in W such that $x = u + v$ where v is another vector in W^\perp .*

4.5 Inequalities in Hilbert Spaces

Theorem 5 (Bessel's Inequality). *Let \mathcal{H} be a Hilbert space. Let \mathcal{E} be an orthonormal set in the space. Then*

$$\forall x \in \mathcal{H}, \quad \sum_{e \in \mathcal{E}} |\langle x, e \rangle|^2 \leq \|x\|^2.$$

Proposition 4.5.1. *Let \mathcal{H} be a Hilbert space. Let \mathcal{E} be an orthonormal set in \mathcal{H} . Let x be a point in the space. Then the net $\sum_{e \in \mathcal{E}} \langle x, e \rangle e$ converges in \mathcal{H} .*

Proof. Let \mathcal{F} be the collection of all finite subsets of \mathcal{E} , partially ordered by inclusion. Define for each $F \in \mathcal{F}$ a vector y_F as $y_F := \sum_{e \in F} \langle x, e \rangle e$. Let ε be an arbitrary positive number. Since \mathcal{E} is an orthonormal set, the set $\{e \in \mathcal{E} : \langle x, e \rangle \neq 0\}$ is countable. Let $\{e_i\}_{i \in \mathbb{N}}$ denote the set. By the Bessel's inequality, $\exists N \in \mathbb{N}$ such that $\sum_{i=N+1}^{\infty} |\langle x, e_i \rangle|^2 < \varepsilon^2$. Define a set F_0 as $F_0 := \{e_1, \dots, e_N\}$. Let F and G be arbitrary elements in \mathcal{F} such that $F_0 \leq F$ and $F_0 \leq G$. Then

$$\begin{aligned}
 \|y_F - y_G\|^2 &= \left\| \sum_{e \in F \setminus G} \langle x, e \rangle e - \sum_{e \in G \setminus F} \langle x, e \rangle e \right\|^2 \\
 &= \sum_{e \in F \cup G \setminus F \cap G} |\langle x, e \rangle|^2 \\
 &= \sum_{e \in F \cup G} |\langle x, e \rangle|^2 - \sum_{e \in F \cap G} |\langle x, e \rangle|^2 \\
 &\leq \sum_{i=N+1}^{\infty} |\langle x, e_i \rangle|^2 \\
 &< \varepsilon^2.
 \end{aligned}$$

So $\{y_F\}_{F \in \mathcal{F}}$ is Cauchy. Since \mathcal{H} is complete and $\{y_F\}_{F \in \mathcal{F}}$ is Cauchy, $\{y_F\}_{F \in \mathcal{F}}$ converges. ■

Chapter 5

Sequence Spaces

5.1 ℓ^p Space

Definition (ℓ^p Space). We define the ℓ^p space to be the set of all sequences x such that $\|x\|_p$ is finite, equipped with the p -norm $\|\cdot\|_p$.

Definition (Weighted ℓ^p Space). Let $(r_i)_{i \in \mathbb{N}}$ be a sequence of positive integers. We define the **weighted ℓ^p** space to be the set given by

$$\ell^p := \left\{ (x_i)_{i \in \mathbb{N}} \in \mathbb{K}^{\mathbb{N}} : \sum_{i \in \mathbb{N}} r_i |x_i|^p < +\infty \right\}.$$

Proposition 5.1.1. For $p \in [1, +\infty)$, $(\ell^p, \|\cdot\|_p)$ is complete.

Proof.

Let $\{x_n\}_{n \in \mathbb{N}}$ be an arbitrary Cauchy sequence in ℓ^p .

Since $\{x_n\}_{n \in \mathbb{N}}$ is Cauchy in ℓ^p , $\forall \varepsilon > 0$, $\exists N(\varepsilon) \in \mathbb{N}$ such that $\forall m, n > N$, we have $\|x_m - x_n\|_p < \varepsilon$.

Since $\|x_m - x_n\|_p < \varepsilon$ and $|x_m^{(i)} - x_n^{(i)}| \leq \|x_m - x_n\|_p$ for any $i \in \mathbb{N}$, $|x_m^{(i)} - x_n^{(i)}| < \varepsilon$ for any $i \in \mathbb{N}$.

Since for any $i \in \mathbb{N}$ and any positive number ε , there exists an integer $N(\varepsilon)$ such that for any indices $m, n > N$, we have $|x_m^{(i)} - x_n^{(i)}| < \varepsilon$, by definition, $\{x_n^{(i)}\}_{n \in \mathbb{N}}$ is Cauchy in \mathbb{F} .

Since $\{x_n^{(i)}\}_{n \in \mathbb{N}}$ is Cauchy in \mathbb{F} and \mathbb{F} is complete, $\{x_n^{(i)}\}_{n \in \mathbb{N}}$ converges.

Let $x_0^{(i)} = x_n^{(i)}$. Let $x_0 = \{x_0^{(i)}\}_{i \in \mathbb{N}}$.

$$\|x_0\|_p = \left(\sum_{i=1}^{\infty} |x_0^{(i)}|^p \right)^{\frac{1}{p}}$$

■

5.2 c_0 Space and c_{00} Space

Definition (c_0 Space). We define c_0 to be

$$c_0 := \left\{ \{x_n\}_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} : \lim_{n \rightarrow \infty} x_n = 0 \right\}.$$

Definition (c_{00} Space). We define c_{00} to be

$$c_{00} := \left\{ (x_n)_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} : \exists N \in \mathbb{N}, \forall n > N, x_n = 0 \right\}.$$

i.e., the set of all eventually zero sequences of real numbers. i.e., the set of all sequences with finite support.

Proposition 5.2.1. The c_{00} is not complete in $(\ell_1, \|\cdot\|_1)$.

Proof. Define a sequence of vectors $(\mathbf{r}_i)_{i \in \mathbb{N}}$ by $\mathbf{r}_i^j := \frac{1}{j^2}$ for $j \in \{1..i\}$ and $\mathbf{r}_i^j := 0$ for $j > i$. Then $(\mathbf{r}_i)_{i \in \mathbb{N}}$ converges to something that is not in c_{00} . ■

Proposition 5.2.2. The closure of c_{00} in the space $(\mathbb{R}^{\omega}, d_1)$ is ℓ_1 .

Proof. For one direction, we are to prove that $\text{cl}(c_{00}) \subseteq \ell_1$. Let x be an arbitrary element in $\text{cl}(c_{00})$. Since $x \in \text{cl}(c_{00})$, there exists another element $y \in c_{00}$ such that $d_1(x, y) < 1$. Let $N \in \mathbb{N}$ be such that $\forall n > N, y_n = 0$. Then

$$\begin{aligned} & d_1(x, y) < 1 \\ \iff & \sum_{n \in \mathbb{N}} |x_n - y_n| < 1 \\ \iff & \sum_{n=1}^N |x_n - y_n| + \sum_{n>N} |x_n - y_n| < 1 \\ \iff & \sum_{n=1}^N |x_n - y_n| + \sum_{n>N} |x_n| < 1 \\ \implies & \sum_{n=1}^N \big| |x_n| - |y_n| \big| + \sum_{n>N} |x_n| < 1 \\ \implies & \sum_{n=1}^N (|x_n| - |y_n|) + \sum_{n>N} |x_n| < 1 \\ \implies & \sum_{n=1}^N |x_n| - \sum_{n=1}^N |y_n| + \sum_{n>N} |x_n| < 1 \\ \iff & \sum_{n \in \mathbb{N}} |x_n| - \sum_{n=1}^N |y_n| < 1 \\ \iff & \sum_{n \in \mathbb{N}} |x_n| < 1 + \sum_{n=1}^N |y_n|. \end{aligned}$$

Since $\sum_{n \in \mathbb{N}} |x_n|$ is bounded, $x \in \ell_1$.

For the reverse direction, we are to prove that $\ell_1 \subseteq \text{cl}(c_{00})$. Let x be an arbitrary element in ℓ_1 . For $i \in \mathbb{N}$, define $x^i = \{x_j^i\}_{j \in \mathbb{N}}$ as $x_j^i = x_j$ for $j \leq i$ and $x_j^i = 0$ for $j > i$. Then $\forall i \in \mathbb{N}, x^i \in c_{00}$. Then

$$\begin{aligned} & \lim_{i \in \mathbb{N}} d_1(x^i, x) \\ &= \lim_{i \in \mathbb{N}} \sum_{j \in \mathbb{N}} |x_j^i - x_j| \\ &= \lim_{i \in \mathbb{N}} \sum_{j > i} |x_j^i - x_j| \\ &= \lim_{i \in \mathbb{N}} \sum_{j > i} |x_j| \\ &= 0. \end{aligned}$$

That is, $\lim_{i \in \mathbb{N}} d_1(x^i, x) = 0$. So $\lim_{i \in \mathbb{N}} x^i = x$. So $x \in \text{cl}(c_{00})$. ■

Proposition 5.2.3. *The closure of c_{00} in the space $(\mathbb{R}^\omega, d_\infty)$ is c_0 .*

Proof. For one direction, we are to prove that $\text{cl}(c_{00}) \subseteq c_0$. Let x be an arbitrary element in $\text{cl}(c_{00})$. Let ε be an arbitrary positive real number. Since $x \in \text{cl}(c_{00})$, there exists another element y in c_{00} such that $d_\infty(x, y) < \varepsilon$. That is, $\forall j \in \mathbb{N}, |x_j - y_j| < \varepsilon$. Since $y \in c_{00}$, $\exists N \in \mathbb{N}$ such that $\forall j > N, y_j = 0$. So $\forall j > N, |x_j| < \varepsilon$. That is,

$$\forall \varepsilon > 0, \quad \exists N \in \mathbb{N}, \quad \forall j > N, \quad |x_j| < \varepsilon.$$

By definition of convergence of limits, $\lim_{j \in \mathbb{N}} x_j = 0$. So $x \in c_0$.

For the reverse direction, we are to prove that $c_0 \subseteq \text{cl}(c_{00})$. Let x be an arbitrary element in c_0 . For $i \in \mathbb{N}$, define x^i as $x_j^i = x_j$ for $j \leq i$ and $x_j^i = 0$ for $j > i$. Then $\forall i \in \mathbb{N}, x^i \in c_{00}$. Let ε be an arbitrary positive real number. Since $x \in c_0$,

$$\exists N \in \mathbb{N}, \quad \forall j > N, \quad |x_j| < \varepsilon/2.$$

Let $i > N$. Then

$$\begin{aligned} & d_\infty(x^i, x) \\ &= \sup_{j \in \mathbb{N}} |x_j^i - x_j| \\ &= \sup_{j > i} |x_j^i - x_j| \\ &= \sup_{j > i} |x_j| \\ &\leq \varepsilon/2 < \varepsilon. \end{aligned}$$

That is,

$$\forall \varepsilon > 0, \quad \exists N \in \mathbb{N}, \quad \forall i > N, \quad d_\infty(x^i, x) < \varepsilon.$$

By definition of convergence of sequences, $\lim_{i \in \mathbb{N}} x^i = x$. So $x \in \text{cl}(c_{00})$. ■

Proposition 5.2.4. *Let $A := \{\{x_n\}_{n \in \mathbb{N}} \in c_{00} : \sum_{n \in \mathbb{N}} x_n = 0\}$. Then A is a subset of ℓ^1 and is closed in (ℓ^1, d_1) . i.e. $\text{cl}(A) = A$ in (ℓ^1, d_1) .*

Proof. Let $x = \{x^i\}_{i \in \mathbb{N}}$ be a sequence in ℓ^1 , where each $x^i = \{x_j^i\}_{j \in \mathbb{N}}$ is an element in A , that converges in (ℓ^1, d_1) . Say $\lim_{i \rightarrow \infty} x^i = x^\infty$.

First I claim that $x^\infty \in c_{00}$.

Now I claim that $\sum_{j \in \mathbb{N}} x_j^\infty = 0$. i.e. $x^\infty \in A$. Since $x^\infty \in c_{00}$,

$$\exists N \in \mathbb{N}, \quad \forall j > N, \quad x_j^\infty = 0.$$

Define $y_i := \sum_{j=1}^N x_j^i$. Define $y_\infty := \sum_{j=1}^N x_j^\infty$. It is easy to see that $\lim_{i \in \mathbb{N}} y_i = y_\infty$. Assume for the sake of contradiction that $y_\infty \neq 0$. i.e. $\{y_i\}_{i \in \mathbb{N}}$ does not converge to 0. Then

$$\exists \varepsilon_0 > 0, \quad \forall M \in \mathbb{N}, \quad \exists i_0 > M, \quad |y_{i_0} - 0| = |y_{i_0}| \geq \varepsilon_0. \quad (1)$$

Since $\lim_{i \rightarrow \infty} x^i = x^\infty$,

$$\exists M_0 \in \mathbb{N}, \quad \forall i > M_0, \quad d_1(x^i, x^\infty) < \varepsilon_0. \quad (2)$$

Consider statement (1) for a particular M, M_0 , we have

$$\exists i_0 > M_0, \quad |y_{i_0}| \geq \varepsilon_0. \quad (3)$$

That is,

$$\left| \sum_{j=1}^N x_j^{i_0} \right| \geq \varepsilon_0. \quad (3')$$

Consider statement (2) for a particular i, i_0 , we have

$$d_1(x^{i_0}, x^\infty) < \varepsilon_0. \quad (4)$$

From statement (4) we can derive:

$$\begin{aligned} & d_1(x^{i_0}, x^\infty) < \varepsilon_0 \\ \iff & \sum_{j \in \mathbb{N}} |x_j^{i_0} - x_j^\infty| < \varepsilon_0 \\ \iff & \sum_{j=1}^N |x_j^{i_0} - x_j^\infty| + \sum_{j > N} |x_j^{i_0} - x_j^\infty| < \varepsilon_0 \end{aligned}$$

$$\begin{aligned}
&\implies \sum_{j>N} |x_j^{i_0} - x_j^\infty| < \varepsilon_0 \\
&\iff \sum_{j>N} |x_j^{i_0} - 0| < \varepsilon_0 \\
&\iff \sum_{j>N} |x_j^{i_0}| < \varepsilon_0 \\
&\implies \left| \sum_{j>N} x_j^{i_0} \right| < \varepsilon_0 \\
&\implies \left| \sum_{j>N} x_j^{i_0} \right| < \varepsilon_0 \\
&\iff \left| \sum_{j \in \mathbb{N}} x_j^{i_0} - \sum_{j=1}^N x_j^{i_0} \right| < \varepsilon_0 \\
&\iff \left| 0 - \sum_{j=1}^N x_j^{i_0} \right| < \varepsilon_0 \\
&\iff \left| \sum_{j=1}^N x_j^{i_0} \right| < \varepsilon_0.
\end{aligned}$$

This contradicts to statement (3'). So the original assumption that $y_\infty \neq 0$ is false. i.e. $y_\infty = 0$. It follows that $\sum_{j \in \mathbb{N}} x_j^\infty = 0$. This completes the proof. ■

5.3 Hölder's Inequality

Theorem 6 (Hölder's Inequality). *Let $\mathfrak{X} = \mathbb{R}^n$ for some $n \in \mathbb{N}$. Let $x = (x_i)_{i=1}^n$ and $y = (y_i)_{i=1}^n$ be vectors in \mathfrak{X} . Then $\forall p, q \in (1, +\infty) : 1/p + 1/q = 1$, $\|xy\|_1 \leq \|x\|_p \|y\|_q$. i.e.,*


$$\sum_{i=1}^n |x_i y_i| \leq \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} \left(\sum_{i=1}^n |y_i|^q \right)^{1/q}.$$

Chapter 6

Function Spaces

6.1 The \mathcal{L}^p Norm

$$\|f\|_p = \left(\int_a^b |f(x)|^p dx \right)^{\frac{1}{p}}.$$

 the instructors' answer, where instructors collectively construct a single answer

In the sup norm, convergence coincides with uniform convergence. Moreover, $C[a, b]$ is complete in this norm. It is not complete in any of the L^p norms for $1 \leq p < \infty$. The completion in these norms is called $L^p(a, b)$.

[undo](#) [thanks](#) | 1

Updated 1 day ago by Kenneth Davidson

Chapter 7

Banach Space

7.1 Definition

Definition (Banach Space). Let $(\mathfrak{X}, \|\cdot\|)$ be a normed linear space. Let d be the metric induced by $\|\cdot\|$. We say that \mathfrak{X} is a **Banach space** if (\mathfrak{X}, d) is a complete metric space. i.e., we define a Banach space to be a complete normed linear space.

7.2 Properties

Proposition 7.2.1. Let $(\mathfrak{X}, \|\cdot\|)$ be a normed vector space over field \mathbb{F} . Then $(\mathfrak{X}, \|\cdot\|)$ is a Banach space if and only if every absolutely summable series in X is summable.

Proposition 7.2.2. Any Banach space with a Schauder basis has to be separable.

7.3 Examples of Banach Space

Example 7.3.1. $(\mathcal{C}([0, 1], \mathbb{F}), \|\cdot\|_\infty)$ is a Banach space.

Example 7.3.2 (Disc Algebra). Define $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$. Define $\mathcal{A}(\mathbb{D}) := \{f \in \mathcal{C}(\overline{\mathbb{D}}) : f|_{\mathbb{D}} \text{ is holomorphic}\}$. Define $\|\cdot\|_\infty$ by $\|f\|_\infty := \sup_{z \in \overline{\mathbb{D}}} |f(z)|$. Then $(\mathcal{A}(\mathbb{D}), \|\cdot\|_\infty)$ is a Banach space.

Example 7.3.3. Let (X, Ω, μ) be a measure space. Let p be a number in $[1, +\infty)$. Define

$$\mathcal{L}^p(X, \mu) := \text{span}\{f : X \rightarrow [0, +\infty] \mid f \text{ is measurable and } \int_X |f|^p < +\infty\}.$$

Define an equivalence relation on $\mathcal{L}^p(X, \mu)$ by $f \equiv g$ if and only if

$$\mu(\{x \in X : f(x) \neq g(x)\}) = 0.$$

Define a space $L^p(X, \mu) := \mathcal{L}^p(X, \mu) / \equiv$. Then $L^p(X, \mu)$ is a Banach space when equipped with the norm

$$\|[f]\|_p := \left(\int_X |f|^p \right)^{1/p}.$$

Example 7.3.4. Let $\mathcal{P}_{\mathbb{C}}[0, 1]$ denote the set of all polynomials with complex coefficients. For each $p \in [1, +\infty)$, define a norm

$$\|f\|_p := \left(\int_0^1 |f|^p \right)^{1/p}.$$

For $p = +\infty$, define a norm

$$\|f\|_{\infty} := \sup_{x \in [0, 1]} |f(x)|.$$

7.4 Construction of Banach Spaces

Definition. Let $(\mathfrak{X}, \|\cdot\|_{\mathfrak{X}})$ and $(\mathfrak{Y}, \|\cdot\|_{\mathfrak{Y}})$ be two Banach spaces over field \mathbb{K} . Let $p \in [1, +\infty)$. We define

$$\mathfrak{X} \oplus_p \mathfrak{Y} := \{(x, y) : x \in \mathfrak{X}, y \in \mathfrak{Y}\}$$

and

$$\|(x, y)\|_p := (\|x\|_{\mathfrak{X}}^p + \|y\|_{\mathfrak{Y}}^p)^{1/p}.$$

For $p = +\infty$, we define

$$\mathfrak{X} \oplus_{\infty} \mathfrak{Y} := \{(x, y) : x \in \mathfrak{X}, y \in \mathfrak{Y}\}$$

and

$$\|(x, y)\|_{\infty} := \max(\|x\|_{\mathfrak{X}}, \|y\|_{\mathfrak{Y}}).$$

- Note that the norms behave similarly to what the p norm would do.
- We can similarly define the direct sum of finitely many Banach spaces.

Proposition 7.4.1. $\|\cdot, \cdot\|_p$ is a norm on $\mathfrak{X} \oplus_p \mathfrak{Y}$.

Proposition 7.4.2. $\mathfrak{X} \oplus_p \mathfrak{Y}$ is complete with respect to $\|\cdot, \cdot\|_p$.

Chapter 8

Hilbert Space

8.1 Definition

Definition (Hilbert Space). We define a **Hilbert space**, denoted by \mathcal{H} , to be a complete inner product space.

8.2 Examples of Hilbert Space

Example 8.2.1. Let (X, μ) be a measure space. Then $L^2(X, \mu)$ is a Hilbert space with inner product given by

$$\langle f, g \rangle := \int_X f(x) \overline{g(x)} d\mu(x).$$

Example 8.2.2. $\ell^2(\mathbb{K}) = \{(x_i)_{i \in \mathbb{N}} \in \mathbb{K}^\infty : \sum_{i \in \mathbb{N}} |x_i|^2 < +\infty\}$ is a Hilbert space with inner product given by

$$\langle (x_i)_{i \in \mathbb{N}}, (y_i)_{i \in \mathbb{N}} \rangle := \sum_{i \in \mathbb{N}} x_i \overline{y_i}.$$

8.3 Properties of Hilbert Space

Proposition 8.3.1. Let \mathcal{H} be a Hilbert space. Let S be a non-empty set in the space. Then $S^{\perp\perp} = \text{clspan}(S)$.

Proof. For one direction, we are to prove that $\text{clspan}(S) \subseteq S^{\perp\perp}$.

For the reverse direction, we are to prove that $S^{\perp\perp} \subseteq \text{clspan}(S)$. Assume for the sake of contradiction that $\exists x \in S^{\perp\perp}$ with $x \neq 0$ such that $x \notin \text{clspan}(S)$. Say $x = m_1 + m_2$ for some $m_1 \in \text{clspan}(S)$ and some $m_2 \in \text{clspan}(S)^\perp$. Note that $\text{clspan}(S)^\perp = S^\perp$. So $m_2 \in S^\perp$. Since $x \in S^{\perp\perp}$ and $m_2 \in S^\perp$, we should have $\langle x, m_2 \rangle = 0$. However,

$$\langle x, m_2 \rangle = \langle m_1 + m_2, m_2 \rangle$$

$$\begin{aligned}
&= \langle m_1, m_2 \rangle + \langle m_2, m_2 \rangle \\
&= 0 + \langle m_2, m_2 \rangle \\
&> 0, \text{ since } m_2 \neq 0.
\end{aligned}$$

This leads to a contradiction. So $S^{\perp\perp} \subseteq \text{clspan}(S)$. ■

Theorem 7 (The Riesz Representation Theorem). *Let \mathcal{H} be a Hilbert space over field $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$. Suppose that $\mathcal{H} \neq \{0\}$. Then for any $\varphi \in \mathcal{H}^*$, $\exists y \in \mathcal{H}$ such that*

$$\forall x \in \mathcal{H}, \quad \varphi(x) = \langle x, y \rangle.$$

Proof. Define for each $y \in \mathcal{H}$ a function $\beta_y \in \mathcal{H}^*$ by $\beta_y(x) := \langle x, y \rangle$. We are to prove that $\mathcal{H}^* = \{\beta_y : y \in \mathcal{H}\}$. It is easy to verify that each β_y is linear and bounded. So $\forall y \in \mathcal{H}$, $\beta_y \in \mathcal{H}^*$. i.e., $\{\beta_y : y \in \mathcal{H}\} \subseteq \mathcal{H}^*$. Define a map Θ from \mathcal{H} to \mathcal{H}^* as $\Theta(y) := \beta_y$. It is easy to verify that Θ is linear.

$$\begin{aligned}
\|\Theta(y)\| &= \|\beta_y\| = \sup\{\beta_y(x) : \|x\| = 1\} \\
&= \sup\{\langle x, y \rangle : \|x\| = 1\} \\
&\leq \sup\{\|x\|\|y\| : \|x\| = 1\} \\
&= \|y\|.
\end{aligned}$$

That is, $\|\Theta(y)\| \leq \|y\|$. So $\|\Theta\| \leq 1$. On the other hand, consider an arbitrary point $y_0 \in \mathcal{H}$ with $y_0 \neq 0$:

$$\begin{aligned}
\|\Theta\| &= \sup \left\{ \frac{\|\Theta(y)\|}{\|y\|} : y \neq 0 \right\} \\
&\geq \frac{\|\Theta(y)\|}{\|y\|} \Big|_{y=y_0} \\
&= \frac{\|\Theta(y_0)\|}{\|y_0\|} \\
&= \frac{\|\beta_{y_0}\|}{\|y_0\|} \\
&= \frac{1}{\|y_0\|} \sup \left\{ \frac{\|\beta_{y_0}(y)\|}{\|y\|} : y \neq 0 \right\} \\
&\geq \frac{1}{\|y_0\|} \frac{\|\beta_{y_0}(y_0)\|}{\|y_0\|} \\
&\geq \frac{1}{\|y_0\|} \frac{\|y_0\|^2}{\|y_0\|} \\
&= 1.
\end{aligned}$$

That is, $\|\Theta\| \geq 1$. So $\|\Theta\| = 1$. So Θ is isometric. It immediately follows that Θ is injective. Now it remains to prove that Θ is surjective. Let $\varphi \in \mathcal{H}^*$. If $\varphi = 0$, then $\varphi = \Theta(0)$ and

we are done. Otherwise, let $\mathcal{M} := \ker(\varphi)$. Then we have $\text{codim } \mathcal{M} = \dim \mathcal{M}^\perp = 1$. Take $e \in \mathcal{M}^\perp$ such that $\|e\| = 1$. Let P denote the orthogonal projection onto \mathcal{M} . Then $1 - P$ is the orthogonal projection onto \mathcal{M}^\perp .

$$x = Px + (1 - P)x = Px + \langle x, e \rangle e.$$

So for $x \in \mathcal{H}$,

$$\varphi(x) = \varphi(Px) + \langle x, e \rangle \varphi(e) = \langle x, \overline{\varphi(e)} e \rangle = \beta_y(x)$$

where $y := \overline{\varphi(e)}e$. Hence $\varphi = \beta_y$. So Θ is surjective. This completes the proof. ■

Chapter 9

Operators

9.1 Bounded Operators

Definition (Bounded Operator). *Let \mathfrak{X} and \mathfrak{Y} be normed linear spaces. Let T be a linear map from \mathfrak{X} to \mathfrak{Y} . We say that T is a **bounded operator** if*

$$\exists k \in \mathbb{R}, \quad \forall x \in \mathfrak{X}, \quad \|Tx\|_{\mathfrak{Y}} \leq k\|x\|_{\mathfrak{X}}.$$

Definition (Operator Norm). *Let \mathfrak{X} and \mathfrak{Y} be normed linear spaces. Let T be a bounded operator from \mathfrak{X} to \mathfrak{Y} . We define the **operator norm** of T , denoted by $\|T\|$, to be the number given by*

$$\|T\| := \inf\{k \in \mathbb{R} : \forall x \in \mathfrak{X}, \|Tx\|_{\mathfrak{Y}} \leq k\|x\|_{\mathfrak{X}}\}.$$

Proposition 9.1.1.

$$\|T\| = \sup\{\|Tx\|_{\mathfrak{Y}} : x \in \mathfrak{X}, \|x\|_{\mathfrak{X}} = 1\}.$$

Proposition 9.1.2. *Let X and Y be normed linear spaces. Let T be a linear map from X to Y . Then T is bounded if and only if T is continuous.*

9.2 Examples of Bounded Operators

Example 9.2.1 (The Multiplication Operator). *Let $\mathfrak{X} = (\mathcal{C}([0, 1], \mathbb{C}), \|\cdot\|_{\infty})$. Let f be a function in \mathfrak{X} . We define the **multiplication operator** on \mathfrak{X} , w.r.t. f , denoted by M_f , as*

$$M_f(g) = fg.$$

Then M_f is bounded and $\|M_f\| = \|f\|_{\infty}$.

Proof. Let g be an arbitrary function in \mathfrak{X} . Then

$$\|M_f g\|_{\infty} = \|fg\|_{\infty}$$

$$\begin{aligned}
&= \sup_{x \in [0,1]} |f(x)g(x)| \\
&= \sup_{x \in [0,1]} |f(x)| |g(x)| \\
&\leq \sup_{x \in [0,1]} |f(x)| \sup_{x \in [0,1]} |g(x)| \\
&= \|f\|_\infty \|g\|_\infty.
\end{aligned}$$

That is, $\|M_f g\|_\infty \leq \|f\|_\infty \|g\|_\infty$. So $\|f\|_\infty$ is an element of the set $S = \{k \in \mathbb{R} : \forall g \in \mathfrak{X}, \|M_f g\|_\infty \leq k \|g\|_\infty\}$. So $\|M_f\| = \inf(S) \leq \|f\|_\infty$. Consider g_0 given by $g_0(x) = 1$. Then g_0 in \mathfrak{X} . Then

$$\|M_f g_0\|_\infty = \|f g_0\|_\infty = \|f\|_\infty = \|f\|_\infty \|g_0\|_\infty.$$

Let k be an arbitrary element in S . Assume for the sake of contradiction that $k < \|f\|_\infty$. Then

$$\begin{aligned}
\|f\|_\infty \|g_0\|_\infty &= \|M_f g_0\|_\infty \\
&\leq k \|g_0\|_\infty \\
&< \|f\|_\infty \|g_0\|_\infty.
\end{aligned}$$

This leads to a contradiction. So $\forall k \in S, k \geq \|f\|_\infty$. So $\|f\|_\infty$ is a lower bound for the set S . So $\|M_f\| = \inf(S) \geq \|f\|_\infty$. Since $\|M_f\| \leq \|f\|_\infty$ and $\|M_f\| \geq \|f\|_\infty$, we get $\|M_f\| = \|f\|_\infty$. ■

Example 9.2.2 (The Volterra Operator). Let $\mathfrak{X} = (\mathcal{C}([0, 1], \mathbb{C}), \|\cdot\|_\infty)$. Define

$$Vf := x \mapsto \int_0^x f(t) dt.$$

Then the Volterra Operator is bounded and $\|V\| \leq 1$.

Proof. Let f be an arbitrary function in \mathfrak{X} with $\|f\|_\infty = 1$. Then $\forall x \in [0, 1]$,

$$\begin{aligned}
|Vf(x)| &= \left| \int_0^x f(t) dt \right| \\
&\leq \int_0^x |f(t)| dt \\
&\leq \int_0^x \sup_{t \in [0,1]} |f(t)| dt \\
&= \int_0^x \|f\|_\infty dt \\
&= \int_0^x 1 dt \\
&= x.
\end{aligned}$$

That is, $\forall x \in [0, 1]$, $|Vf(x)| \leq 1$. So $\|Vf\|_\infty \leq 1$. Since $\forall f \in \mathfrak{X} : \|f\|_\infty = 1$, $\|Vf\|_\infty \leq 1$, we get $\|V\| \leq 1$. ■

Example 9.2.3 (The Diagonal Operator). Let $\mathfrak{X} = \ell^2(\mathbb{N})$. Let

$$D = \begin{bmatrix} d_1 & & & \\ & d_2 & & \\ & & d_3 & \\ & & & \ddots \end{bmatrix}.$$

Then D is bounded if and only if $(d_i)_{i \in \mathbb{N}}$ is bounded and $\|D\| = \|(d_i)_{i \in \mathbb{N}}\|_\infty$.

Proof. Case 1.

$$\begin{aligned} \|Dx\|_2^2 &= \sum_{i \in \mathbb{N}} |d_i x_i|^2 \\ &\leq \sum_{i \in \mathbb{N}} \|(d_j)_{j \in \mathbb{N}}\|_\infty |x_i|^2 \\ &= \|(d_j)_{j \in \mathbb{N}}\|_\infty \sum_{i \in \mathbb{N}} |x_i|^2 \\ &= \|(d_j)_{j \in \mathbb{N}}\|_\infty \|x\|_2^2. \end{aligned}$$

Case 2.

If $(d_i)_{i \in \mathbb{N}} \notin \ell^\infty$, $\exists (d_{n_i})_{i \in \mathbb{N}} \rightarrow \infty$.

$$\begin{aligned} \|De_{n_i}\|_2 &= \|d_{n_i} e_{n_i}\|_2 \\ &= |d_{n_i}| \|e_{n_i}\|_2 \\ &= |d_{n_i}|. \end{aligned}$$

So $\|D\| \geq \|De_{n_i}\|_2 \rightarrow \infty$. ■

Example 9.2.4 (Weighted Shifts).

- Let $\mathcal{H} = \ell^2_{\mathbb{N}}$. Let $(w_n)_{n \in \mathbb{N}} \in \ell^\infty_{\mathbb{N}}$. We define an *unilateral forward weighted shift* W on \mathcal{H} as

$$W(x_n) := (0, w_1 x_1, w_2 x_2, w_3 x_3, \dots).$$

i.e.,

$$W = \begin{bmatrix} 0 & & & & \\ w_1 & 0 & & & \\ & w_2 & 0 & & \\ & & w_3 & 0 & \\ & & & \ddots & \ddots \end{bmatrix}.$$

Then W is bounded and $\|W\| = \sup\{|w_n| : n \in \mathbb{N}\}$.

- Let $\mathcal{H} = \ell_{\mathbb{N}}^2$. Let $(v_n)_{n \in \mathbb{N}} \in \ell_{\mathbb{N}}^\infty$. We define an **unilateral backward weighted shift** V on \mathcal{H} as

$$V(x_n) := (v_1 x_2, v_2 x_3, v_3 x_4, \dots).$$

Then V is bounded and $\|V\| = \sup\{|v_n| : n \in \mathbb{N}\}$.

- Let $\mathcal{H} = \ell_{\mathbb{Z}}^2$. Let $(u_n)_{n \in \mathbb{Z}} \in \ell_{\mathbb{Z}}^\infty$. We define a **bilateral weighted shift** U on \mathcal{H} as

$$U(x_n) := (u_{n-1} x_{n-1})_{n \in \mathbb{Z}}.$$

Then U is bounded and $\|U\| = \sup\{|u_n| : n \in \mathbb{Z}\}$.

Example 9.2.5 (The Composition Operators). Let $\mathfrak{X} = \mathcal{C}([0, 1], \mathbb{C})$. Let $\varphi \in \mathcal{C}([0, 1], [0, 1])$. We define the **composition operator** on \mathfrak{X} , denoted by C_φ as

$$C_\varphi(f) := f \circ \varphi.$$

Then C_φ is contractive.

Proof.

$$\begin{aligned} \|C_\varphi(f)\| &= \sup_{x \in [0, 1]} |(f \circ \varphi)(x)| \\ &\leq \|f\|_\infty. \end{aligned}$$

■

9.3 The Space of Bounded Operators

Proposition 9.3.1. Let \mathfrak{X} and \mathfrak{Y} be normed linear spaces. Then $\mathcal{B}(\mathfrak{X}, \mathfrak{Y})$ is a vector space and the operator norm is a norm on $\mathcal{B}(\mathfrak{X}, \mathfrak{Y})$.

Proposition 9.3.2. Let \mathfrak{X} and \mathfrak{Y} be normed linear spaces. Let $\mathcal{B}(\mathfrak{X}, \mathfrak{Y})$ be the space of bounded linear operators from \mathfrak{X} to \mathfrak{Y} . Then if \mathfrak{Y} is complete, $\mathcal{B}(\mathfrak{X}, \mathfrak{Y})$ is complete.

Proposition 9.3.3. Let \mathfrak{X} and \mathfrak{Y} be normed linear spaces. Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be two equivalent norms on $\mathcal{B}(\mathfrak{X}, \mathfrak{Y})$. Then $T \in \mathcal{B}(\mathfrak{X}, \mathfrak{Y}, \|\cdot\|_1)$ if and only if $T \in \mathcal{B}(\mathfrak{X}, \mathfrak{Y}, \|\cdot\|_2)$.

9.4 Invertible Bounded Operators

Proposition 9.4.1. Let $(\mathfrak{X}, \|\cdot\|_1)$ be a Banach space. Let $S \in \mathcal{B}(\mathfrak{X})$ be a bounded linear map that is invertible. Define a norm $\|\cdot\|_2$ on \mathfrak{X} as

$$\|x\|_2 := \|Sx\|_1.$$

Then $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent.

Proof. On one hand, since S is bounded, $\exists c_1$ such that $\forall x \in \mathfrak{X}$, $\|Sx\|_1 \leq c_1\|x\|_1$. That is, $\|x\|_2 \leq c_1\|x\|_1$.

On the other hand, since S is invertible, S^{-1} exists and is also bounded. Since S^{-1} is bounded, $\exists c_2$ such that $\forall x \in \mathfrak{X}$, $\|S^{-1}x\|_1 \leq c_2\|x\|_1$. Consider $x = Sx$, we get $\forall x \in \mathfrak{X}$, $\|S^{-1}Sx\|_1 \leq c_2\|Sx\|_1$. That is, $\|x\|_1 \leq c_2\|x\|_2$.

So $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent. ■

Proposition 9.4.2. *Let $(\mathfrak{X}, \|\cdot\|)$ be a Banach space. Let S be a map in $\mathcal{B}(\mathfrak{X})$ that is invertible. Then*

$$\|S^{-1}\| = (\inf\{\|Sx\| : \|x\| = 1\})^{-1}.$$

Proof.

$$\begin{aligned} \|S^{-1}\| &= \sup\{\|S^{-1}x\| : \|x\| = 1\} \\ &= \sup\left\{\frac{\|S^{-1}x\|}{\|x\|} : \|x\| \neq 0\right\} \\ &= \sup\left\{\frac{\|S^{-1}Sx\|}{\|Sx\|} : \|x\| \neq 0\right\} \\ &= \sup\left\{\frac{\|x\|}{\|Sx\|} : \|x\| \neq 0\right\} \\ &= \sup\left\{\frac{1}{\|Sx\|} : \|x\| = 1\right\} \\ &= (\inf\{\|Sx\| : \|x\| = 1\})^{-1}. \end{aligned}$$

That is,

$$\|S^{-1}\| = (\inf\{\|Sx\| : \|x\| = 1\})^{-1}.$$
■

Chapter 10

Dual Space

10.1 Definition

Definition ((Topological) Dual Space). *Let \mathfrak{X} be a normed linear space over field \mathbb{K} . We define the **(topological) dual space** of \mathfrak{X} , denoted by \mathfrak{X}^* , to be the space $\mathcal{B}(\mathfrak{X}, \mathbb{K})$.*

Definition (Linear Functionals). *We call the elements of \mathfrak{X}^* **linear functionals**.*

Proposition 10.1.1. *Let X be a normed linear space. Then there exists a contractive map from X to its double dual X^{**} .*

10.2 Properties

10.3 Examples of Dual Space

Example 10.3.1. $c_0(\mathbb{N})^*$ is isometrically isomorphic to $\ell^1(\mathbb{N})$.

Example 10.3.2. $c_0(\mathbb{N})^*$ is isometrically isomorphic to $\ell^\infty(\mathbb{N})$.

Chapter 11

Quotient Spaces

11.1 Definitions

Definition. Let \mathfrak{V} be a vector space. Let \mathfrak{W} be a subspace of \mathfrak{V} . We define a **quotient space**, denoted by $\mathfrak{V}/\mathfrak{W}$, to be a set $\{v + \mathfrak{W} : v \in \mathfrak{V}\}$ with operations

$$(v_1 + \mathfrak{W}) + (v_2 + \mathfrak{W}) := (v_1 + v_2) + \mathfrak{W} \text{ and}$$

$$\kappa(v + \mathfrak{W}) := (\kappa v) + \mathfrak{W}.$$

Definition (Quotient Map). Let $(\mathfrak{X}, \|\cdot\|)$ be a normed linear space. Let \mathfrak{M} be a linear manifold in \mathfrak{X} . We define the **quotient map** on \mathfrak{X} with respect to \mathfrak{M} , denoted by $q_{\mathfrak{M}}$, to be a function from \mathfrak{X} to $\mathfrak{X}/\mathfrak{M}$ given by

$$q_{\mathfrak{M}}(x) := x + \mathfrak{M}$$

Proposition 11.1.1. Quotient maps are contractive. i.e.,

$$\forall x \in \mathfrak{X}, \quad \|x + \mathfrak{M}\|_{\mathfrak{X}/\mathfrak{M}} \leq \|x\|_{\mathfrak{X}}.$$

Proposition 11.1.2. Let \mathfrak{X} be a normed linear space. Let \mathfrak{M} be a closed subspace of \mathfrak{X} . Let q be the canonical quotient map from \mathfrak{X} to $\mathfrak{X}/\mathfrak{M}$. Then

- q is a continuous map. i.e.,

$$\forall \text{ open set } W \subseteq \mathfrak{X}/\mathfrak{M}, \quad q^{-1}(W) \text{ is open in } \mathfrak{X}.$$

- q is an open map. i.e.,

$$\forall \text{ open set } G \subseteq \mathfrak{X}, \quad q(G) \text{ is open in } \mathfrak{X}/\mathfrak{M}.$$

Proof. Since q is contractive, q is continuous and hence (1). ■

Definition (Seminorm on Quotient Spaces). *Let $(\mathfrak{X}, \|\cdot\|)$ be a normed linear space. Let \mathfrak{M} be a linear manifold in \mathfrak{X} . We define a **seminorm** on $\mathfrak{X}/\mathfrak{M}$ to be a function from $\mathfrak{X}/\mathfrak{M}$ to \mathbb{R} given by*

$$p(x + \mathfrak{M}) := \inf\{\|x + m\| : m \in \mathfrak{M}\}.$$

Proposition 11.1.3. *Seminorms on quotient spaces are indeed seminorms.*

Proposition 11.1.4. *A seminorm on a quotient space $\mathfrak{X}/\mathfrak{M}$ is a norm if and only if \mathfrak{M} is closed.*

Chapter 12

Topological Vector Space

12.1 Definitions

Definition (Compatible). Let \mathcal{W} be a vector space over field \mathbb{K} . Let \mathcal{T} be a topology on \mathcal{W} . We say that \mathcal{T} is **compatible** with the vector space structure on \mathcal{W} if the addition and scalar multiplication operations on \mathcal{W} are continuous.

Definition (Topological Vector Space). We define a **topological vector space** to be a vector space with a compatible Hausdorff topology.

Proposition 12.1.1 (Stability under Linear Combinations). Let X be a normed vector space over \mathbb{F} . Let K be a compact set in the space. Let C be a closed set in the space. Then $\forall \alpha, \beta \in \mathbb{F}$, the set S given by $S := \alpha K + \beta C$ is closed.

Proof. The case where $\beta = 0$ is trivial. I will assume $\beta \neq 0$. Let $\alpha, \beta \in \mathbb{F}$ be arbitrary. Let $\{s_i\}_{i \in \mathbb{N}}$ be an arbitrary sequence in S that converges. Say the limit is s_∞ . Since $s_i \in S$ for any $i \in \mathbb{N}$ and $S = \alpha K + \beta C$, $s_i = \alpha k_i + \beta c_i$ for some $k_i \in K$ and some $c_i \in C$, for any $i \in \mathbb{N}$. Since $\{k_i\}_{i \in \mathbb{N}}$ is a sequence in K and K is compact, there exists a convergent subsequence $\{k_i\}_{i \in I}$ of $\{k_i\}_{i \in \mathbb{N}}$ in K . Say $\{k_i\}_{i \in I}$ converges to $k_\infty \in K$. Since $\{s_i\}_{i \in \mathbb{N}}$ converges to s_∞ , $\{s_i\}_{i \in I}$ also converges to s_∞ . Since $s_i = \alpha k_i + \beta c_i$, $c_i = \beta^{-1}(s_i - \alpha k_i)$. Define $c_\infty := \beta^{-1}(s_\infty - \alpha k_\infty)$. Since $\{s_i\}_{i \in I}$ converges to s_∞ and $\{k_i\}_{i \in I}$ converges to k_∞ and $c_i = \beta^{-1}(s_i - \alpha k_i)$, $\{c_i\}_{i \in I}$ converges to c_∞ . Since $\{c_i\}_{i \in I}$ is a sequence in C and converges to c_∞ and C is closed, $c_\infty \in C$. Since $s_\infty = \alpha k_\infty + \beta c_\infty$ and $k_\infty \in K$ and $c_\infty \in C$, $s_\infty \in \alpha K + \beta C$. Since for any sequence in S that converges, the limit is also in S , S is closed. ■

Remark. The sum of two closed sets may not be closed.

Proof. **Counter-example 1**

Consider $A := \{n : n \in \mathbb{N}\}$ and $B := \{n + \frac{1}{n} : n \in \mathbb{N}\}$.
 (<https://math.stackexchange.com/questions/124130/sum-of-two-closed-sets-in-mathbb-r-is-closed>)
 Their sum contains the sequence $\{\frac{1}{n}\}_{n \in \mathbb{N}}$ but does not contain 0.

Counter-example 2

Consider $A := \mathbb{R} \times \{0\}$ and $B := \{(x, y) \in \mathbb{R}^2 : x, y > 0, xy \geq 1\}$. Their sum is $\mathbb{R} \times \mathbb{R}_{++}$. ■

12.2 Examples

Example 12.2.1. Let \mathfrak{X} be a normed linear space. Then \mathfrak{X} is a topological vector space with the topology induced by the norm.

Proposition 12.2.1. Normed linear spaces are Hausdorff.

Example 12.2.2. Let \mathfrak{X} be a Banach space. Let \mathfrak{X}^* denote the dual space of \mathfrak{X} . Let τ_* denote the weak topology on \mathfrak{X}^* induced by elements of \mathfrak{X} as

$$\lim_{\alpha} x_{\alpha}^* = x^* \iff \forall x \in \mathfrak{X}, \lim_{\alpha} x_{\alpha}^*(x) = x^*(x).$$

Then (\mathfrak{X}^*, τ_*) is a topological vector space.

12.3 Neighborhood Improvements

Proposition 12.3.1. Let (\mathcal{V}, τ) be a topological vector space. Let $U \in \mathcal{U}_0$ be a neighborhood of 0 in \mathcal{V} . Then

- $\exists N \in \mathcal{U}_0$ such that $N + N \subseteq U$.
- $\exists M \in \mathcal{U}_0$ and $\exists \varepsilon > 0$ such that $\forall 0 < |k| < \varepsilon$, we have $kM \subseteq U$.
-

Chapter 13

Equicontinuity in Metric Spaces

13.1 Definitions

Definition ((Pointwise) Equicontinuity). Let (X, d_X) and (Y, d_Y) be metric spaces. Let \mathcal{F} be a collection of functions from X to Y . Let x_0 be a point in X . We say that \mathcal{F} is **(pointwise) equicontinuous** at point x_0 if for any positive number ε , there exists some number $\delta(x_0, \varepsilon)$ such that for any function f in \mathcal{F} and any point x in X , we have

$$d_Y(f(x), f(x_0)) < \varepsilon$$

whenever $d_X(x, x_0) < \delta(x_0, \varepsilon)$ is satisfied.

Definition (Uniform Equicontinuity). Let (X, d_X) and (Y, d_Y) be metric spaces. Let \mathcal{F} be a collection of functions from X to Y . We say that \mathcal{F} is **uniformly equicontinuous** if for any positive number ε , there exists some number $\delta(\varepsilon)$ such that for any function f in \mathcal{F} and any points x_1 and x_2 in X , we have

$$d_Y(f(x_1), f(x_2)) < \varepsilon$$

whenever $d_X(x_1, x_2) < \delta(\varepsilon)$ is satisfied.

13.2 Sufficient Conditions

Proposition 13.2.1. The closure of an equicontinuous family of functions is equicontinuous.

Proof.

Let (X, d_X) and (Y, d_Y) be metric spaces.

Let \mathcal{F} be an equicontinuous family of functions from X to Y .

We are to prove that $cl(\mathcal{F})$ is equicontinuous.

Let x_0 be an arbitrary point in X .

Let ε be an arbitrary positive number.

Since \mathcal{F} is equicontinuous at point x_0 , there exists some $\delta(x_0, \varepsilon)$ such that for any function f in \mathcal{F} and any point x in X such that $d_X(x, x_0) < \delta(x_0, \varepsilon)$, we have $d_Y(f(x), f(x_0)) < \varepsilon/3$.

Let f be an arbitrary function in $cl(\mathcal{F})$.

Let x be an arbitrary point in X such that $d_X(x, x_0) < \delta(x_0, \varepsilon)$.

Since $f \in cl(\mathcal{F})$, there exists some function $f_0 \in \mathcal{F}$ such that $d_\infty(f, f_0) < \varepsilon/3$.

Since $d_\infty(f, f_0) < \varepsilon/3$, $d_Y(f(x), f_0(x)) < \varepsilon/3$ and $d_Y(f(x_0), f_0(x_0)) < \varepsilon/3$.

Since $f_0 \in \mathcal{F}$ and $d_X(x, x_0) < \delta(x_0, \varepsilon)$, $d_Y(f_0(x), f_0(x_0)) < \varepsilon/3$.

Since $d_Y(f(x), f_0(x)) < \varepsilon/3$ and $d_Y(f(x_0), f_0(x_0)) < \varepsilon/3$ and $d_Y(f_0(x), f_0(x_0)) < \varepsilon/3$,
 $d_Y(f(x), f(x_0)) < \varepsilon$.

Since for any positive number ε , there exists some $\delta(x_0, \varepsilon)$ such that for any function f in $cl(\mathcal{F})$ and any point x in X such that $d_X(x, x_0) < \delta(x_0, \varepsilon)$, we have $d_Y(f(x), f(x_0)) < \varepsilon$, by definition of equicontinuous, $cl(\mathcal{F})$ is equicontinuous at point x_0 .

Since $cl(\mathcal{F})$ is equicontinuous at point x_0 for any point x_0 in X , $cl(\mathcal{F})$ is equicontinuous. ■

Chapter 14

Adjoint Operator

14.1 Definitions

Definition (Adjoint Matrix). *Let A be an $m \times n$ matrix. We define the **adjoint** of A , denoted by A^* , to be an $n \times m$ matrix given by*

$$(A^*)_{ij} := \overline{(A)_{ji}}.$$

Definition (Adjoint Operator). *Let V and W be inner product spaces. Let T be a linear map from V to W . We define the **adjoint** of T , denoted by T^* , to be a map from W to V such that*

$$\forall x \in V, \forall y \in W, \quad \langle T(x), y \rangle_W = \langle x, T^*(y) \rangle_V.$$

Proposition 14.1.1 (Existence). *Let V be a finite-dimensional inner product space and T be a linear operator on V . Then the adjoint of T exists.*

Proposition 14.1.2 (Uniqueness). *Let V be an inner product space and T be a linear operator on V . Then the adjoint of T is unique, provided that it exists.*

14.2 Properties of the Adjoint Operator

Proposition 14.2.1. *Let V be an inner product space. Then*

- (1) $(I_V)^* = I_V$ where I_V is the identity operator on V .
- (2) $T^{**} = T$ for any linear operator T on V .

Proposition 14.2.2. *Let V be an inner product space and T be a linear operator on V . Then T^* is also linear.*

Proposition 14.2.3. *Let V be an inner product space. Then*

(1) *For any linear operators T and U ,*

$$(T + U)^* = T^* + U^*.$$

(2) *For any linear operator T ,*

$$(cT)^* = \bar{c} \cdot T^*.$$

(3) *For any linear operators T and U ,*

$$(TU)^* = U^*T^*.$$

Proposition 14.2.4. *Let V be a finite-dimensional inner product space and T be a linear operator on V . Then if T is invertible, T^* is also invertible.*

Proposition 14.2.5. *Let V be an inner product space and T be an invertible linear operator on V . Then $(T^{-1})^* = (T^*)^{-1}$.*

14.3 Normal Operators

Definition (Normal). *Let V be an inner product space and T be a linear operator on V . We say that T is **normal** if $TT^* = T^*T$.*

14.4 Self-adjoint

Chapter 15

Convolution

Definition (Convolution). *Let f and g be functions from \mathbb{R} to \mathbb{R} . We define the **convolution** of f and g , denoted by $f * g$, to be a function on \mathbb{R} given by*

$$(f * g)(t) := \int_{-\infty}^{+\infty} f(\tau)g(t - \tau)dt.$$

Chapter 16

Coercive Functions

16.1 Definitions

Definition (Coercive). Let f be a function from \mathbb{R}^d to \mathbb{R}^* . We say that f is *coercive* if $\lim_{\|x\| \rightarrow \infty} f(x) = +\infty$.

16.2 Properties

Proposition 16.2.1. Let f be a proper lower semi-continuous function from \mathbb{R}^d to \mathbb{R}^* . Let K be a compact set in \mathbb{R}^d . Assume $K \cap \text{dom}(f) \neq \emptyset$. Then f attains its minimum over K .

Proof.

Define $m := \inf_{x \in K} f(x)$.

Since $m = \inf_{x \in K} f(x)$, there exists a sequence $\{x_i\}_{i \in \mathbb{N}}$ in K such that $\lim_{i \rightarrow \infty} f(x_i) = m$.

Since K is compact and $\{x_i\}_{i \in \mathbb{N}} \subseteq K$, there exists a convergent subsequence $\{x_i\}_{i \in I}$ in K where I is an infinite subset of \mathbb{N} .

Say the limit is x_∞ where $x_\infty \in K$.

Since $\lim_{i \rightarrow \infty} f(x_i) = m$, we get $\lim_{i \in I, i \rightarrow \infty} f(x_i) = m$.

Since $\lim_{i \in I, i \rightarrow \infty} f(x_i) = m$, we get $\liminf_{i \in I, i \rightarrow \infty} f(x_i) = m$.

Since f is lower semi-continuous and $\lim_{i \in I, i \rightarrow \infty} x_i = x_\infty$, we get $f(x_\infty) \leq \liminf_{i \in I, i \rightarrow \infty} f(x_i)$.

That is, $f(x_\infty) \leq m$.

Since $m = \inf_{x \in K} f(x)$, we have $\forall x \in K, f(x) \geq m$.

In particular, $f(x_\infty) \geq m$.

Since $f(x_\infty) \geq m$ and $f(x_\infty) \leq m$, $f(x_\infty) = m$.

Since f is proper, $f(x_\infty) = m \neq -\infty$.

So f attains its minimum at point x_∞ .

■

Proposition 16.2.2. *Let f be a proper, lower semi-continuous, and coercive function from \mathbb{R}^d to \mathbb{R}^* . Let C be a closed subset of \mathbb{R}^d . Assume $C \cap \text{dom}(f) \neq \emptyset$. Then f attains its minimum over C .*

Proof.

Since $C \cap \text{dom}(f) \neq \emptyset$, take $x \in C \cap \text{dom}(f)$.

Since f is coercive, $\exists R$ such that $\forall y, \|y\| > R$, we have $f(y) \geq f(x)$.

Since $x \in C \cap \text{dom}(f)$ and $\forall y, \|y\| > R$, we have $f(y) \geq f(x)$, the set of minimizers of f over C is the same as the set of minimizers of f over $C \cap \text{ball}[0, R]$.

Since C and $\text{ball}[0, R]$ are both closed, $C \cap \text{ball}[0, R]$ is closed.

Since $\text{ball}[0, R]$ is bounded, $C \cap \text{ball}[0, R]$ is bounded.

Since $C \cap \text{ball}[0, R]$ is closed and bounded, by the Heine-Borel Theorem, $C \cap \text{ball}[0, R]$ is compact.

Since f is proper and lower semi-continuous and $C \cap \text{ball}[0, R]$ is compact, f attains its minimum over $C \cap \text{ball}[0, R]$.

So f attains its minimum over C .

■

Chapter 17

Unclassified Results

Proposition 17.0.1. *Let (X, d) be a compact metric space. Let $L(X)$ be the set of all Lipschitz functions from X to \mathbb{R} . Let $C(X)$ be the set of all continuous functions from X to \mathbb{R} . Then $L(X)$ is dense in $C(X)$.*

Proposition 17.0.2. *Let $(V, \|\cdot\|)$ be a normed vector space. Let S be a subset of V . Let p be a vector in V . Then we have the followings.*

$$(1) \ p + \text{int}(S) = \text{int}(p + S),$$

$$(2) \ p + \text{cl}(S) = \text{cl}(p + S).$$

Proof.

Proof of (1).

For one direction, let x be an arbitrary point in the set $(p + \text{int}(S))$.

We are to prove that $x \in \text{int}(p + S)$.

Since $x \in (p + \text{int}(S))$, $(x - p) \in \text{int}(S)$.

Since $(x - p) \in \text{int}(S)$, by definition of interior, there exists a radius r such that

$$B(x - p, r) \subseteq S.$$

It follows that $B(x, r) \subseteq p + S$.

Since there exists a radius r such that $B(x, r) \subseteq p + S$, by definition of interior,

$$x \in \text{int}(p + S).$$

For the reverse direction, let x be an arbitrary point in $\text{int}(p + S)$.

We are to prove that $x \in p + \text{int}(S)$.

Since $x \in \text{int}(p + S)$, by definition of interior, there exists a radius r such that

$$B(x, r) \subseteq (p + S).$$

It follows that $B(x - p, r) \subseteq S$.

Since there exists a radius r such that $B(x - p, r) \subseteq S$, by definition of interior,

$$(x - p) \in \text{int}(S).$$

Since $(x - p) \in \text{int}(S)$, we get $x \in (p + \text{int}(S))$.

Proof of (2).

For one direction, let x be an arbitrary point in the set $(p + \text{cl}(S))$.

We are to prove that $x \in \text{cl}(p + S)$.

Since $x \in (p + \text{cl}(S))$, we get $(x - p) \in \text{cl}(S)$.

Since $(x - p) \in \text{cl}(S)$, by definition of closure, for any radius r , we have

$$B(x - p, r) \cap S \neq \emptyset.$$

It follows that $B(x, r) \cap (p + S) \neq \emptyset$.

Since for any radius r , $B(x, r) \cap (p + S) \neq \emptyset$, by definition of closure, we get

$$x \in \text{cl}(p + S).$$

For the reverse direction, let x be an arbitrary point in $\text{cl}(p + S)$.

We are to prove that $x \in (p + \text{cl}(S))$.

Since $x \in \text{cl}(p + S)$, by definition of closure, for any radius r , we have

$$B(x, r) \cap (p + S) \neq \emptyset.$$

It follows that $B(x - p, r) \cap S \neq \emptyset$.

Since for any radius r , $B(x - p, r) \cap S \neq \emptyset$, by definition of closure, we get

$$(x - p) \in \text{cl}(S).$$

Since $(x - p) \in \text{cl}(S)$, we get $x \in (p + \text{cl}(S))$.

■

Proposition 17.0.3. *Let $(V, \|\cdot\|)$ be a normed vector space. Let S be a subset of V . Let λ be a non-zero real number. Then*

$$(1) \lambda \text{int}(S) = \text{int}(\lambda S).$$

$$(2) \lambda \text{cl}(S) = \text{cl}(\lambda S).$$

Proof of (1). For one direction, let x be an arbitrary point in $\lambda \text{int}(S)$.

We are to prove that $x \in \text{int}(\lambda S)$.

Since $x \in \lambda \text{int}(S)$, we get $x/\lambda \in \text{int}(S)$.

Since $x/\lambda \in \text{int}(S)$, by definition of interior, there exists a radius r such that

$$B(x/\lambda, r) \subseteq S.$$

Let y be an arbitrary point in $B(x, \lambda r)$.

Since $y \in B(x, \lambda r)$, we get $\|y - x\| \leq \lambda r$.

Since $\|y - x\| \leq \lambda r$, we get $\|y/\lambda - x/\lambda\| \leq r$.

Since $\|y/\lambda - x/\lambda\| \leq r$, we get $y/\lambda \in B(x/\lambda, r)$.

Since $y/\lambda \in B(x/\lambda, r)$ and $B(x/\lambda, r) \subseteq S$, we get $y/\lambda \in S$.

Since $y/\lambda \in S$, we get $y \in \lambda S$.

Since any point in $B(x, \lambda r)$ is also in λS , we get $B(x, \lambda r) \subseteq \lambda S$.

Since there exists a radius r such that $B(x, \lambda r) \subseteq \lambda S$, by definition of interior, we get

$$x \in \text{int}(\lambda S).$$

For the reverse direction,

■