

# Functional Analysis

Daniel Mao



# Contents

<b>1</b>	<b>Balanced Sets</b>	<b>1</b>
1.1	Definitions . . . . .	1
1.2	Properties . . . . .	1
1.3	Stability of Balance . . . . .	2
1.4	Absorbing Sets . . . . .	2
<b>2</b>	<b>Inner Product Space</b>	<b>3</b>
2.1	Inner Products . . . . .	3
2.2	Inequalities . . . . .	4
2.3	Orthogonality . . . . .	5
<b>3</b>	<b>Normed Vector Spaces</b>	<b>7</b>
3.1	Definitions . . . . .	7
3.2	Properties . . . . .	7
3.3	Equivalence of Norms . . . . .	8
3.4	Dual Norms . . . . .	9
3.5	$p$ -norms . . . . .	9
3.6	Banach Spaces . . . . .	11
<b>4</b>	<b>Topological Vector Spaces</b>	<b>13</b>
4.1	Definitions . . . . .	13
4.2	Topological Vector Spaces . . . . .	13
4.3	Neighborhoods . . . . .	14
<b>5</b>	<b>Sequence Spaces</b>	<b>15</b>
5.1	$\ell_p$ Space . . . . .	15
5.2	$c_0$ Space and $c_{00}$ Space . . . . .	16
<b>6</b>	<b>Function Spaces</b>	<b>21</b>
6.1	The $\mathcal{L}^p$ Norm . . . . .	21

<b>7 Hilbert Space</b>	<b>23</b>
7.1 Hilbert Spaces . . . . .	23
<b>8 Equicontinuity in Metric Spaces</b>	<b>25</b>
8.1 Definitions . . . . .	25
8.2 Sufficient Conditions . . . . .	25
<b>9 Operators</b>	<b>27</b>
9.1 Bounded Operators . . . . .	27
9.2 Space of Bounded Operators . . . . .	28
9.3 Dual Spaces . . . . .	28
<b>10 Adjoint Operator</b>	<b>29</b>
10.1 Definitions . . . . .	29
10.2 Properties of the Adjoint Operator . . . . .	29
10.3 Normal Operators . . . . .	30
10.4 Self-adjoint . . . . .	30
<b>11 Convolution</b>	<b>31</b>
<b>12 Coercive Functions</b>	<b>33</b>
12.1 Definitions . . . . .	33
12.2 Properties . . . . .	33
<b>13 Unclassified Results</b>	<b>35</b>

# 1

## Balanced Sets

### 1.1 Definitions

**Definition** (Balanced Sets). *Let  $X$  be a vector space over field  $\mathbb{F}$ . Let  $S$  be a subset of  $X$ . We say that  $S$  is **balanced** if*

$$\forall a \in \mathbb{F} : |a| \leq 1, \quad aS \subseteq S.$$

**Definition** (Balanced Hull). *Let  $X$  be a vector space over field  $\mathbb{F}$ . Let  $S$  be a subset of  $X$ . We define the **balanced hull** of  $S$ , denoted by  $\text{balhull}(S)$ , to be the smallest balanced set containing  $S$ .*

**Definition** (Balanced Core). *Let  $X$  be a vector space over field  $\mathbb{F}$ . Let  $S$  be a subset of  $X$ . We define the **balanced core** of  $S$ , denoted by  $\text{balcore}(S)$ , to be the largest balanced set contained in  $S$ .*

### 1.2 Properties

**Proposition 1.2.1.** *Let  $X$  be a vector space over field  $\mathbb{F}$ . Let  $B$  be a balanced subset of  $X$ . Then*

$$\forall a, b \in \mathbb{F} : |a| \leq |b|, \quad aB \subseteq bB.$$

**Proposition 1.2.2.** *Balanced sets are path connected.*

**Proposition 1.2.3** (Act on Other Properties). • *The balanced hull of a compact set is compact.*

- *The balanced hull of a totally bounded set is totally bounded.*
- *The balanced hull of a bounded set is bounded.*

**Proposition 1.2.4** (Act on Other Properties). • *The balanced core of a closed set is closed.*

**Proposition 1.2.5.** *Let  $X$  be a vector space over field  $\mathbb{F}$ . Let  $a$  be a scalar in field  $\mathbb{F}$ . Then*

$$a \operatorname{balhull}(S) = \operatorname{balhull}(aS).$$

### 1.3 Stability of Balance

**Proposition 1.3.1** (Set Operations). • *The union of balanced sets is also balanced.*

- *The intersection of balanced sets is also balanced.*

**Proposition 1.3.2** (Linear Mappings). • *The scalar multiple of a balanced set is also balanced.*

- *The (Minkowski) sum of two balanced sets is also balanced.*
- *The image of a balanced set under a linear operator is also balanced.*
- *The inverse image of a balanced set under a linear operator is also balanced.*

**Proposition 1.3.3** (Topological Operations). *The closure of a balanced set is also balanced.*

**Proposition 1.3.4.** *The convex hull of a balanced set is also balanced (and also convex).*

### 1.4 Absorbing Sets

**Definition** (Absorbing Sets). *Let  $X$  be a vector space over field  $\mathbb{F}$ . Let  $S$  be a subset of  $X$ . We say that  $S$  is **absorbing** if*

$$\forall x \in X, \quad \exists r \in \mathbb{R} : r > 0, \quad \forall c \in \mathbb{F} : |c| \geq r, \quad x \in cA.$$

**Proposition 1.4.1.** *Every absorbing set contains the origin.*

## 2

# Inner Product Space

## 2.1 Inner Products

### 2.1.1 Definitions

**Definition** (Inner Product). *Let  $V$  be a vector space over field  $\mathbb{F}$ . We define an **inner product** on  $V$ , denoted by  $\langle \cdot, \cdot \rangle$ , to be a scalar-valued function defined on  $V \times V$  such that*

(1) *Positive Definiteness*

$$\begin{aligned}\forall x, y \in V, \quad \langle x, x \rangle &\geq 0, \text{ and} \\ \forall x \in V, \quad \langle x, x \rangle = 0 &\iff x = O_V.\end{aligned}$$

(2) *Sesqui-Linearity*

$$\begin{aligned}\forall x, y, z, w \in V, \quad \langle x + y, z + w \rangle &= \langle x, z \rangle + \langle y, z \rangle + \langle x, w \rangle + \langle y, w \rangle, \text{ and} \\ \forall a, b \in \mathbb{F}, \forall x, y \in V, \quad \langle ax, by \rangle &= a\bar{b}\langle x, y \rangle.\end{aligned}$$

(3) *Conjugate Symmetry*

$$\forall x, y \in V, \quad \langle x, y \rangle = \overline{\langle y, x \rangle}.$$

**Definition** (Norm). *Let  $V$  be an inner product space over field  $\mathbb{F}$ . We define the **norm**, denoted by  $\|\cdot\|$ , to be a function from  $V$  to  $\mathbb{R}_+$  given by*

$$\|x\| := \sqrt{\langle x, x \rangle}$$

**Definition** (Orthogonal Vectors). *Let  $V$  be an inner product space. Let  $x$  and  $y$  be vectors in  $V$ . We say that  $x$  and  $y$  are **orthogonal** if  $\langle x, y \rangle = 0$ .*

**Definition** (Orthogonal Sets). *Let  $S$  be a subset of  $V$ . We say that  $S$  is **orthogonal** if*

$$\forall x, y \in S, \quad \langle x, y \rangle = 0.$$

### 2.1.2 Examples

**Definition** (Standard Inner Product). For  $V = \mathbb{F}^n$ , we define the **standard inner product** by

$$\langle x, y \rangle := \sum_{i=1}^n x_i \overline{y_i}.$$

**Definition** (Frobenius Inner Product). For  $V = \mathbb{F}^{n \times n}$ , we define the **Frobenius inner product** by

$$\langle M_1, M_2 \rangle := \text{tr}(M_2^* M_1).$$

**Definition.** Let  $V$  be the space of continuous scalar-valued functions on  $[0, 2\pi]$ . We define the inner product on  $V$  by

$$\langle f, g \rangle := \frac{1}{2\pi} \int_0^{2\pi} f(x) \overline{g(x)} dx.$$

### 2.1.3 Properties

**Proposition 2.1.1.** Let  $V$  be a finite dimensional inner product space. Let  $\mathcal{B}$  be a basis for  $V$ . Let  $x$  and  $y$  be vectors in  $V$ . Then

$$x = y \iff \forall b \in \mathcal{B}, \quad \langle x, b \rangle = \langle y, b \rangle.$$

## 2.2 Inequalities

**Theorem 1** (Minkowski).

$$\left( \sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{p}} \leq \left( \sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left( \sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}}$$

**Proposition 2.2.1** (Cauchy-Schwarz Inequality). Let  $V$  be an inner product space. Then

$$\forall x, y \in V, \quad |\langle x, y \rangle| \leq \|x\| \cdot \|y\|$$

**Proposition 2.2.2** (Triangle Inequality). Let  $V$  be an inner product space. Then

$$\forall x, y \in V, \quad \|x + y\| \leq \|x\| + \|y\|$$

**Proposition 2.2.3** (Parallelogram Law). Let  $V$  be an inner product space. Then

$$\forall x, y \in V, \quad \|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2$$



## 2.3 Orthogonality

### 2.3.1 Orthogonal Sets

**Definition** (Orthogonality). *Let  $V$  be an inner product space. We say that points  $x$  and  $y$  in  $V$  are **orthogonal** if  $\langle x, y \rangle = 0$ .*

**Definition** (Orthogonal Sets). *Let  $V$  be an inner product space and  $S$  be a subset of  $V$ . We say that  $S$  is **orthogonal** if any two vectors in  $S$  are orthogonal.*

**Proposition 2.3.1.** *Orthogonal sets are linearly independent.*

### 2.3.2 Orthogonal Bases

**Definition** (Orthogonal Basis). *Let  $V$  be an inner product space and  $S$  be a subset of  $V$ . We say that  $S$  is an **orthogonal basis** for  $V$  if it is an ordered basis for  $V$  and orthogonal.*

**Proposition 2.3.2.** *Let  $V$  be an inner product space. Let  $S = \{v_1, \dots, v_n\}$  be an orthogonal subset of  $V$  where each  $v_i$  is non-zero. Then*

$$\forall y \in \text{span}(S), \quad y = \sum_{i=1}^n \frac{\langle y, v_i \rangle}{\|v_i\|^2} v_i.$$

**Theorem 2** (Gram-Schmidt Process). *Let  $V$  be an inner product space. Let  $S = \{v_0, \dots, v_n\}$  be a linearly independent subset of  $V$ . Then the set  $S' = \{v'_0, \dots, v'_n\}$  given by  $v'_0 := v_0$  and*

$$\forall i \in \{1, \dots, n\}, \quad v'_i := v_i - \sum_{j=1}^{i-1} \frac{\langle v_i, v'_j \rangle}{\|v'_j\|^2} v'_j$$

*is an orthogonal subset of  $V$  consisting of non-zero vectors. Furthermore, we have  $\text{span}(S') = \text{span}(S)$ .*

**Proposition 2.3.3.** *Let  $V$  be an inner product space and  $S = \{v_0, v_1, \dots, v_n\}$  be an orthogonal subset of  $V$ . Then the set  $S'$  derived from the Gram-Schmidt process is exactly  $S$ .*

**Theorem 3** (Parseval's Identity). *Let  $V$  be a finite-dimensional inner product space. Let  $B = \{v_1, \dots, v_n\}$  be an orthogonal basis for  $V$ . Then*

$$\forall x, y \in V, \quad \langle x, y \rangle = \sum_{i=1}^n \langle x, v_i \rangle \overline{\langle y, v_i \rangle}.$$

**Theorem 4** (Bessel's Inequality). *Let  $V$  be a finite-dimensional inner product space. Let  $B = \{v_1, \dots, v_n\}$  be an orthogonal subset for  $V$ . Then*

$$\forall x \in V, \quad \|x\|^2 \geq \sum_{i=1}^n |\langle x, v_i \rangle|^2.$$

### 2.3.3 Orthogonal Complements

**Definition** (Orthogonal Complement). *Let  $V$  be an inner product space and  $S$  be a non-empty subset of  $V$ . We define the **orthogonal complement** of  $S$ , denoted by  $S^\perp$ , to be the set of all points in  $V$  that are orthogonal to all vectors in  $S$ .*

**Proposition 2.3.4.** *Let  $V$  be a finite-dimensional inner product space. Then*

$$(1) V^\perp = \{O_V\}$$

$$(2) \{O_V\}^\perp = V$$

**Proposition 2.3.5.** *Orthogonal complements are always linear subspaces.*

**Proposition 2.3.6.** *Let  $V$  be an inner product space and  $W$  be a subspace of  $V$  with basis  $\beta$ . Then a vector in  $V$  is also in  $W^\perp$  if and only if it is orthogonal to all vectors in  $\beta$ .*

**Proposition 2.3.7** (Extension). *Let  $V$  be an  $n$ -dimensional inner product space and  $S = \{v_1, v_2, \dots, v_k\}$  be an orthogonal subset of  $V$ . Then  $S$  can be extended to an orthogonal basis  $B = \{v_1, v_2, \dots, v_k, v_{k+1}, \dots, v_n\}$  for  $V$ .*

### 2.3.4 Properties of the Orthogonal Complement Operator

**Proposition 2.3.8.** *Let  $V$  be an inner product space. Then*

$$(1) S \subseteq T \text{ implies } T^\perp \subseteq S^\perp \text{ for any subsets } S \text{ and } T \text{ of } V.$$

$$(2) S \subseteq (S^\perp)^\perp \text{ for any subset } S \text{ of } V.$$

**Proposition 2.3.9.** *Let  $V$  be a finite-dimensional inner product space and  $W$  be a subspace of  $V$ . Then*

$$(1) W = (W^\perp)^\perp$$

$$(2) V = W \oplus W^\perp$$

**Proposition 2.3.10.** *Let  $V$  be a finite-dimensional inner product space and  $W_1$  and  $W_2$  be subspaces of  $V$ . Then*

$$(1) (W_1 + W_2)^\perp = W_1^\perp \cap W_2^\perp$$

$$(2) (W_1 \cap W_2)^\perp = W_1^\perp + W_2^\perp$$

### 2.3.5 Orthogonal Projection

**Definition** (Orthogonal Projection). *Let  $V$  be a vector space. Let  $W$  be a finite-dimensional subspace of  $V$ . Let  $x$  be a vector in  $V$ . We define the **orthogonal projection** of  $x$  on  $W$ , denoted by  $(x)$ , to be the vector  $u$  in  $W$  such that  $x = u + v$  where  $v$  is another vector in  $W^\perp$ .*

# 3

## Normed Vector Spaces

### 3.1 Definitions

**Definition (Norm).** Let  $X$  be a vector space over field  $\mathbb{F}$ . We define a **norm** on  $X$ , denoted by  $\nu$ , to be a map from  $X$  to  $\mathbb{R}$  that satisfies the following conditions.

- (1)  $\forall x \in X, \quad \nu(x) = 0 \iff x = 0.$
- (2)  $\forall x \in X, \quad \nu(x) \geq 0.$
- (3)  $\forall \lambda \in \mathbb{F}, \forall x \in X, \quad \nu(\lambda x) = \lambda \nu(x).$
- (4)  $\forall x, y \in X, \quad \nu(x + y) \leq \nu(x) + \nu(y).$

**Definition (Semi-Norm).** Let  $X$  be a vector space over field  $\mathbb{F}$ . We define a **semi-norm** on  $X$ , denoted by  $\nu$ , to be a map from  $X$  to  $\mathbb{R}$  that satisfies the following conditions.

- (1)  $\forall x \in X, \quad \nu(x) \geq 0.$
- (2)  $\forall \lambda \in \mathbb{F}, \forall x \in X, \quad \nu(\lambda x) = \lambda \nu(x).$
- (3)  $\forall x, y \in X, \quad \nu(x + y) \leq \nu(x) + \nu(y).$

### 3.2 Properties

**Proposition 3.2.1.** Let  $(V, \|\cdot\|_V)$  be a normed vector space over field  $\mathbb{F}$ . Then  $(V, \|\cdot\|)$  is complete if and only if  $(\overline{B(0,1)}, \|\cdot\|_V)$  is complete.

*Proof.*

For one direction, assume that  $(V, \|\cdot\|)$  is complete.

We are to prove that  $(\overline{B(0,1)}, \|\cdot\|_V)$  is complete.

Since  $(\overline{B(0,1)}, \|\cdot\|_V)$  is a closed subspace of  $(V, \|\cdot\|)$  and  $(V, \|\cdot\|)$  is complete,  $(\overline{B(0,1)}, \|\cdot\|_V)$  is also complete.

For the reverse direction, assume that  $(\overline{B(0,1)}, \|\cdot\|_V)$  is complete.

We are to prove that  $(V, \|\cdot\|_V)$  is complete.

Let  $\{x_i\}_{i \in \mathbb{N}}$  be an arbitrary Cauchy sequence in  $(V, \|\cdot\|_V)$ .

Since  $\{x_i\}_{i \in \mathbb{N}}$  is Cauchy in  $(V, \|\cdot\|_V)$ ,  $\{x_i\}_{i \in \mathbb{N}}$  is bounded in  $(V, \|\cdot\|_V)$ .

Let  $\lambda$  be a positive upper bound for  $\{\|x_i\|_V\}_{i \in \mathbb{N}}$ .

Since  $\{x_i\}_{i \in \mathbb{N}}$  is Cauchy in  $(V, \|\cdot\|_V)$ ,  $\{x_i/\lambda\}_{i \in \mathbb{N}}$  is Cauchy in  $(\overline{B(0,1)}, \|\cdot\|_V)$ .

Since  $\{x_i/\lambda\}_{i \in \mathbb{N}}$  is Cauchy in  $(\overline{B(0,1)}, \|\cdot\|_V)$  and  $(\overline{B(0,1)}, \|\cdot\|_V)$  is complete,  $\{x_i/\lambda\}_{i \in \mathbb{N}}$  converges in  $(\overline{B(0,1)}, \|\cdot\|_V)$ .

Since  $\{x_i/\lambda\}_{i \in \mathbb{N}}$  converges in  $(\overline{B(0,1)}, \|\cdot\|_V)$ ,  $\{x_i\}_{i \in \mathbb{N}}$  converges in  $(V, \|\cdot\|_V)$ .

Since any Cauchy sequence in  $(V, \|\cdot\|_V)$  converges in  $(V, \|\cdot\|_V)$ ,  $(V, \|\cdot\|_V)$  is complete. ■

### 3.3 Equivalence of Norms

**Definition** (Equivalence of Norms). *Let  $V$  be a vector space over field  $\mathbb{F}$ . Let  $\|\cdot\|_1$  and  $\|\cdot\|_2$  be two norms on  $V$ . We say that  $\|\cdot\|_1$  and  $\|\cdot\|_2$  are **equivalent** if there exist positive constants  $c_1$  and  $c_2$  such that for any vector  $v$  in  $V$ ,*

$$c_1\|v\|_1 \leq \|v\|_2 \leq c_2\|v\|_1.$$

*Or equivalently,*

$$c_1\|v\|_2 \leq \|v\|_1 \leq c_2\|v\|_2.$$

**Proposition 3.3.1.** *Equivalence of norms is an equivalence relation.*

**Theorem 5.** *Let  $V$  be a finite dimensional vector space over field  $\mathbb{F} = \{\mathbb{R}, \mathbb{C}\}$ . Then any two norms on  $V$  are equivalent.*

*Proof.*

Let  $\|\cdot\|_p$  be an arbitrary  $p$ -norm on  $V$  and  $\|\cdot\|$  be an arbitrary norm on  $V$ .

Let  $\mathcal{B}$  be the standard basis for  $V$ . Say  $\mathcal{B} = \{e_1, e_2, \dots, e_n\}$ .

Let  $v$  be an arbitrary vector in  $V$ .

$$\begin{aligned}
\|v\| &= \left\| \sum_{i=1}^n v_i e_i \right\| \\
&\leq \sum_{i=1}^n |v_i| \|e_i\| \\
&\leq \left( \sum_{i=1}^n |v_i|^p \right)^{\frac{1}{p}} \left( \sum_{i=1}^n \|e_i\|^{\frac{p}{p-1}} \right)^{1-\frac{1}{p}} \\
&= \left( \sum_{i=1}^n \|e_i\|^{\frac{p}{p-1}} \right)^{1-\frac{1}{p}} \|v\|_p \\
&:= c_1 \|v\|_p.
\end{aligned}$$

■

**Proposition 3.3.2.** *Let  $X$  be a vector space. Let  $\|\cdot\|_1$  and  $\|\cdot\|_2$  be two norms on  $X$ . Then  $\|\cdot\|_1$  and  $\|\cdot\|_2$  are equivalent if and only if they generate the same metric topology.*

### 3.4 Dual Norms

**Definition (Dual Norm).** *Let  $(V, \|\cdot\|)$  be a normed vector space. We define the **dual norm** of  $\|\cdot\|$ , denoted by  $\|\cdot\|_\circ$ , to be a function given by*

$$\|v\|_\circ := \max_{\|w\|=1} v \cdot w = \max_{\|w\| \neq 0} \frac{|v \cdot w|}{\|w\|}.$$

**Proposition 3.4.1.** *The dual norms of norms are indeed norms.*

**Proposition 3.4.2.** *Let  $(V, \|\cdot\|)$  be a normed vector space. Let  $v, w$  be vectors in the space. Then*

$$|v \cdot w| \leq \|v\| \cdot \|w\|_\circ.$$

### 3.5 $p$ -norms

**Definition ( $p$ -norm).** *Let  $V$  be a finite-dimensional normed vector space over field  $\mathcal{F}$ . Let  $\mathcal{B} = \{b_1, \dots, b_n\}$  be a basis for  $V$  where  $n = \dim(V)$ . Let  $v$  be a vector in a normed vector space. For  $p \in [1, +\infty)$ , we define the  **$p$ -norm** of  $v$ , denoted by  $\|v\|_p$ , to be the number given by*

$$\|v\|_p = \left( \sum_{i=1}^n |(v_{\mathcal{B}})_i|^p \right)^{\frac{1}{p}}.$$

**Definition** (Infinity Norm). *Let  $v$  be a vector in a normed vector space. We define the **infinity norm** of  $v$ , denoted by  $\|v\|_\infty$ , to be the number given by*

$$\|v\|_\infty := \max\{|v_i|\}_{i=1}^n.$$

**Proposition 3.5.1.**  *$p$ -norms are indeed norms.*

**Proposition 3.5.2.** *For any vector  $v$  in  $\mathbb{R}^n$ , we have*

$$\lim_{p \rightarrow \infty} \|v\|_p = \|v\|_\infty.$$

*i.e.,*

$$\lim_{p \rightarrow \infty} \left( \sum_{i=1}^n |v_i|^p \right)^{\frac{1}{p}} = \max\{|v_i|\}_{i=1}^n.$$

*Proof.* Let  $p$  be an arbitrary number in  $[1, +\infty)$ . Let  $k$  be an arbitrary index in  $\{1, \dots, n\}$ . Then

$$|v_k| \leq \left( \sum_{i=1}^n |v_i|^p \right)^{1/p} = \|v\|_p.$$

So

$$\max\{|v_k|\} = \|v\|_\infty \leq \|v\|_p.$$

So

$$\lim_{p \rightarrow \infty} \|v\|_p \geq \|v\|_\infty. \quad (1)$$

On the other hand, note that

$$\left( \sum_{i=1}^n |v_i|^p \right) / \|v\|_\infty^p = \sum_{i=1}^n \left( \frac{|v_i|}{\|v\|_\infty} \right)^p$$

decreases as  $p$  increases. So it is bounded above. Say

$$\left( \sum_{i=1}^n |v_i|^p \right) / \|v\|_\infty^p \leq C$$

for some  $C \in \mathbb{R}$ . Then

$$\left( \sum_{i=1}^n |v_i|^p \right)^{1/p} = \|v\|_p \leq C^{1/p} \|v\|_\infty.$$

So

$$\lim_{p \rightarrow \infty} \|v\|_p \leq \lim_{p \rightarrow \infty} C^{1/p} \|v\|_\infty = \|v\|_\infty. \quad (2)$$

From (1) and (2) we get

$$\lim_{p \rightarrow \infty} \|v\|_p = \|v\|_\infty.$$

■

**Proposition 3.5.3.** *Let  $p$  be an arbitrary number in  $[1, +\infty)$ . Then the dual norm of the  $p$ -norm  $\|\cdot\|_p$  is the  $q$ -norm  $\|\cdot\|_q$  where  $q$  is such that satisfies*

$$\frac{1}{p} + \frac{1}{q} = 1.$$

**Proposition 3.5.4.** *Let  $p$  and  $q$  be numbers in  $[1, +\infty]$ . Let  $v$  be a vector in  $\mathbb{R}^n$ . Then if  $p \leq q$ ,*

$$\|x\|_q \leq \|x\|_p \leq n^{\frac{1}{p} - \frac{1}{q}} \cdot \|x\|_q.$$

**Proposition 3.5.5.** *Let  $w$  and  $z$  be vectors in  $\mathbb{E}^d$ . Then*

$$\|w + z\|_2^2 + \|w - z\|_2^2 = 2(\|w\|_2^2 + \|z\|_2^2).$$

## 3.6 Banach Spaces

**Definition** (Banach Space). *We define a **Banach space** to be a complete normed vector spaces.*

**Example 3.6.1.**  $(\mathcal{C}([0, 1], \mathbb{F}), \|\cdot\|_\infty)$  is a Banach space.

**Example 3.6.2** (Disc Algebra). *Define  $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ . Define  $\mathcal{A}(\mathbb{D}) := \{f \in \mathcal{C}(\overline{\mathbb{D}}) : f|_{\mathbb{D}} \text{ is holomorphic}\}$ . Define  $\|\cdot\|_\infty$  by  $\|f\|_\infty := \sup_{z \in \overline{\mathbb{D}}} |f(z)|$ . Then  $(\mathcal{A}(\mathbb{D}), \|\cdot\|_\infty)$  is a Banach space.*

**Proposition 3.6.1.** *Let  $(X, \|\cdot\|)$  be a normed vector space over field  $\mathbb{F}$ . Then  $(X, \|\cdot\|)$  is a Banach space if and only if every absolutely summable series in  $X$  is summable.*





# 4

## Topological Vector Spaces

### 4.1 Definitions

### 4.2 Topological Vector Spaces

**Definition** (Vector Topology). *Let  $X$  be a vector space over a topological field  $\mathbb{K}$ . We define a **vector topology** on  $X$  to be a topology on  $X$  such that vector addition and scalar multiplication are continuous.*

**Proposition 4.2.1** (Stability under Linear Combinations). *Let  $X$  be a normed vector space over  $\mathbb{F}$ . Let  $K$  be a compact set in the space. Let  $C$  be a closed set in the space. Then  $\forall \alpha, \beta \in \mathbb{F}, S := \alpha K + \beta C$  is closed.*

*Proof.*

The case where  $\beta = 0$  is trivial. I will assume  $\beta \neq 0$ .

Let  $\alpha, \beta \in \mathbb{F}$  be arbitrary.

Let  $\{s_i\}_{i \in \mathbb{N}}$  be an arbitrary sequence in  $S$  that converges.

Say the limit is  $s_\infty$ .

Since  $s_i \in S$  for any  $i \in \mathbb{N}$  and  $S = \alpha K + \beta C$ ,  $s_i = \alpha k_i + \beta c_i$  for some  $k_i \in K$  and some  $c_i \in C$ , for any  $i \in \mathbb{N}$ .

Since  $\{k_i\}_{i \in \mathbb{N}}$  is a sequence in  $K$  and  $K$  is compact, there exists a convergent subsequence  $\{k_i\}_{i \in I}$  of  $\{k_i\}_{i \in \mathbb{N}}$  in  $K$ .

Say  $\{k_i\}_{i \in I}$  converges to  $k_\infty \in K$ .

Since  $\{s_i\}_{i \in \mathbb{N}}$  converges to  $s_\infty$ ,  $\{s_i\}_{i \in I}$  also converges to  $s_\infty$ .

Since  $s_i = \alpha k_i + \beta c_i$ ,  $c_i = \beta^{-1}(s_i - \alpha k_i)$ .

Define  $c_\infty := \beta^{-1}(s_\infty - \alpha k_\infty)$

Since  $\{s_i\}_{i \in I}$  converges to  $s_\infty$  and  $\{k_i\}_{i \in I}$  converges to  $k_\infty$  and  $c_i = \beta^{-1}(s_i - \alpha k_i)$ ,  $\{c_i\}_{i \in I}$  converges to  $c_\infty$ .

Since  $\{c_i\}_{i \in I}$  is a sequence in  $C$  and converges to  $c_\infty$  and  $C$  is closed,  $c_\infty \in C$ .

Since  $s_\infty = \alpha k_\infty + \beta c_\infty$  and  $k_\infty \in K$  and  $c_\infty \in C$ ,  $s_\infty \in \alpha K + \beta C$ .

Since for any sequence in  $S$  that converges, the limit is also in  $S$ ,  $S$  is closed. ■

**Remark.** *The sum of two closed sets may not be closed.*

*Proof.*

**Counter-example 1**

Consider  $A := \{n : n \in \mathbb{N}\}$  and  $B := \{n + \frac{1}{n} : n \in \mathbb{N}\}$ .

(<https://math.stackexchange.com/questions/124130/sum-of-two-closed-sets-in-mathbb-r-is-closed>)

Their sum contains the sequence  $\{\frac{1}{n}\}_{n \in \mathbb{N}}$  but does not contain 0.

**Counter-example 2**

Consider  $A := \mathbb{R} \times \{0\}$  and  $B := \{(x, y) \in \mathbb{R}^2 : x, y > 0, xy \geq 1\}$ .

Their sum is  $\mathbb{R} \times \mathbb{R}_{++}$ . ■

## 4.3 Neighborhoods

# 5

## Sequence Spaces

### 5.1 $\ell_p$ Space

**Definition** ( $\ell_p^{(n)}$  Space). We define the  $\ell_p^{(n)}$  space to be the set of all sequences  $\{x_i\}_{i=1}^{i=n}$  such that

**Definition** ( $\ell_p$  Space). We define the  $\ell_p$  space to be the set of all sequences  $x$  such that  $\|x\|_p$  is finite, equipped with the  $p$ -norm  $\|\cdot\|_p$ .

**Proposition 5.1.1.** For  $p \in [1, +\infty)$ ,  $(\ell_p, \|\cdot\|_p)$  is complete.

*Proof.*

Let  $\{x_n\}_{n \in \mathbb{N}}$  be an arbitrary Cauchy sequence in  $\ell_p$ .

Since  $\{x_n\}_{n \in \mathbb{N}}$  is Cauchy in  $\ell_p$ ,  $\forall \varepsilon > 0$ ,  $\exists N(\varepsilon) \in \mathbb{N}$  such that  $\forall m, n > N$ , we have  $\|x_m - x_n\|_p < \varepsilon$ .

Since  $\|x_m - x_n\|_p < \varepsilon$  and  $|x_m^{(i)} - x_n^{(i)}| \leq \|x_m - x_n\|_p$  for any  $i \in \mathbb{N}$ ,  $|x_m^{(i)} - x_n^{(i)}| < \varepsilon$  for any  $i \in \mathbb{N}$ .

Since for any  $i \in \mathbb{N}$  and any positive number  $\varepsilon$ , there exists an integer  $N(\varepsilon)$  such that for any indices  $m, n > N$ , we have  $|x_m^{(i)} - x_n^{(i)}| < \varepsilon$ , by definition,  $\{x_n^{(i)}\}_{n \in \mathbb{N}}$  is Cauchy in  $\mathbb{F}$ .

Since  $\{x_n^{(i)}\}_{n \in \mathbb{N}}$  is Cauchy in  $\mathbb{F}$  and  $\mathbb{F}$  is complete,  $\{x_n^{(i)}\}_{n \in \mathbb{N}}$  converges.

Let  $x_0^{(i)} = x_n^{(i)}$ . Let  $x_0 = \{x_0^{(i)}\}_{i \in \mathbb{N}}$ .

$$\|x_0\|_p = \left( \sum_{i=1}^{\infty} |x_0^{(i)}|^p \right)^{\frac{1}{p}}$$

■

## 5.2 $c_0$ Space and $c_{00}$ Space

**Definition** ( $c_0$  Space). We define  $c_0$  to be

$$c_0 := \left\{ \{x_n\}_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} : \lim_{n \rightarrow \infty} x_n = 0 \right\}.$$

**Definition** ( $c_{00}$  Space). We define  $c_{00}$  to be

$$c_{00} := \left\{ \{x_n\}_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} : \exists N \in \mathbb{N}, \forall n > N, x_n = 0 \right\}.$$

i.e. the set of all eventually zero sequences of real numbers.

**Proposition 5.2.1.** The closure of  $c_{00}$  in the space  $(\mathbb{R}^{\omega}, d_1)$  is  $\ell_1$ .

*Proof.* For one direction, we are to prove that  $\text{cl}(c_{00}) \subseteq \ell_1$ . Let  $x$  be an arbitrary element in  $\text{cl}(c_{00})$ . Since  $x \in \text{cl}(c_{00})$ , there exists another element  $y \in c_{00}$  such that  $d_1(x, y) < 1$ . Let  $N \in \mathbb{N}$  be such that  $\forall n > N, y_n = 0$ . Then

$$\begin{aligned} d_1(x, y) &< 1 \\ \iff \sum_{n \in \mathbb{N}} |x_n - y_n| &< 1 \\ \iff \sum_{n=1}^N |x_n - y_n| + \sum_{n>N} |x_n - y_n| &< 1 \\ \iff \sum_{n=1}^N |x_n - y_n| + \sum_{n>N} |x_n| &< 1 \\ \implies \sum_{n=1}^N ||x_n| - |y_n|| + \sum_{n>N} |x_n| &< 1 \\ \implies \sum_{n=1}^N (|x_n| - |y_n|) + \sum_{n>N} |x_n| &< 1 \\ \implies \sum_{n=1}^N |x_n| - \sum_{n=1}^N |y_n| + \sum_{n>N} |x_n| &< 1 \\ \iff \sum_{n \in \mathbb{N}} |x_n| - \sum_{n=1}^N |y_n| &< 1 \\ \iff \sum_{n \in \mathbb{N}} |x_n| &< 1 + \sum_{n=1}^N |y_n|. \end{aligned}$$

Since  $\sum_{n \in \mathbb{N}} |x_n|$  is bounded,  $x \in \ell_1$ .

For the reverse direction, we are to prove that  $\ell_1 \subseteq \text{cl}(c_{00})$ . Let  $x$  be an arbitrary element in  $\ell_1$ . For  $i \in \mathbb{N}$ , define  $x^i = \{x_j^i\}_{j \in \mathbb{N}}$  as  $x_j^i = x_j$  for  $j \leq i$  and  $x_j^i = 0$  for  $j > i$ . Then

$\forall i \in \mathbb{N}, x^i \in c_{00}$ . Then

$$\begin{aligned}
 & \lim_{i \in \mathbb{N}} d_1(x^i, x) \\
 &= \lim_{i \in \mathbb{N}} \sum_{j \in \mathbb{N}} |x_j^i - x_j| \\
 &= \lim_{i \in \mathbb{N}} \sum_{j > i} |x_j^i - x_j| \\
 &= \lim_{i \in \mathbb{N}} \sum_{j > i} |x_j| \\
 &= 0.
 \end{aligned}$$

That is,  $\lim_{i \in \mathbb{N}} d_1(x^i, x) = 0$ . So  $\lim_{i \in \mathbb{N}} x^i = x$ . So  $x \in \text{cl}(c_{00})$ . ■

**Proposition 5.2.2.** *The closure of  $c_{00}$  in the space  $(\mathbb{R}^\omega, d_\infty)$  is  $c_0$ .*

*Proof.* For one direction, we are to prove that  $\text{cl}(c_{00}) \subseteq c_0$ . Let  $x$  be an arbitrary element in  $\text{cl}(c_{00})$ . Let  $\varepsilon$  be an arbitrary positive real number. Since  $x \in \text{cl}(c_{00})$ , there exists another element  $y$  in  $c_{00}$  such that  $d_\infty(x, y) < \varepsilon$ . That is,  $\forall j \in \mathbb{N}, |x_j - y_j| < \varepsilon$ . Since  $y \in c_{00}$ ,  $\exists N \in \mathbb{N}$  such that  $\forall j > N, y_j = 0$ . So  $\forall j > N, |x_j| < \varepsilon$ . That is,

$$\forall \varepsilon > 0, \quad \exists N \in \mathbb{N}, \quad \forall j > N, \quad |x_j| < \varepsilon.$$

By definition of convergence of limits,  $\lim_{j \in \mathbb{N}} x_j = 0$ . So  $x \in c_0$ .

For the reverse direction, we are to prove that  $c_0 \subseteq \text{cl}(c_{00})$ . Let  $x$  be an arbitrary element in  $c_0$ . For  $i \in \mathbb{N}$ , define  $x^i$  as  $x_j^i = x_j$  for  $j \leq i$  and  $x_j^i = 0$  for  $j > i$ . Then  $\forall i \in \mathbb{N}, x^i \in c_{00}$ . Let  $\varepsilon$  be an arbitrary positive real number. Since  $x \in c_0$ ,

$$\exists N \in \mathbb{N}, \quad \forall j > N, \quad |x_j| < \varepsilon/2.$$

Let  $i > N$ . Then

$$\begin{aligned}
 & d_\infty(x^i, x) \\
 &= \sup_{j \in \mathbb{N}} |x_j^i - x_j| \\
 &= \sup_{j > i} |x_j^i - x_j| \\
 &= \sup_{j > i} |x_j| \\
 &\leq \varepsilon/2 < \varepsilon.
 \end{aligned}$$

That is,

$$\forall \varepsilon > 0, \quad \exists N \in \mathbb{N}, \quad \forall i > N, \quad d_\infty(x^i, x) < \varepsilon.$$

By definition of convergence of sequences,  $\lim_{i \in \mathbb{N}} x^i = x$ . So  $x \in \text{cl}(c_{00})$ . ■

**Proposition 5.2.3.** *Let  $A := \{\{x_n\}_{n \in \mathbb{N}} \in c_{00} : \sum_{n \in \mathbb{N}} x_n = 0\}$ . Then  $A$  is a subset of  $\ell^1$  and is closed in  $(\ell^1, d_1)$ . i.e.  $\text{cl}(A) = A$  in  $(\ell^1, d_1)$ .*

*Proof.* Let  $x = \{x^i\}_{i \in \mathbb{N}}$  be a sequence in  $\ell^1$ , where each  $x^i = \{x_j^i\}_{j \in \mathbb{N}}$  is an element in  $A$ , that converges in  $(\ell^1, d_1)$ . Say  $\lim_{i \rightarrow \infty} x^i = x^\infty$ .

First I claim that  $x^\infty \in c_{00}$ .

Now I claim that  $\sum_{j \in \mathbb{N}} x_j^\infty = 0$ . i.e.  $x^\infty \in A$ . Since  $x^\infty \in c_{00}$ ,

$$\exists N \in \mathbb{N}, \quad \forall j > N, \quad x_j^\infty = 0.$$

Define  $y_i := \sum_{j=1}^N x_j^i$ . Define  $y_\infty := \sum_{j=1}^N x_j^\infty$ . It is easy to see that  $\lim_{i \in \mathbb{N}} y_i = y_\infty$ . Assume for the sake of contradiction that  $y_\infty \neq 0$ . i.e.  $\{y_i\}_{i \in \mathbb{N}}$  does not converge to 0. Then

$$\exists \varepsilon_0 > 0, \quad \forall M \in \mathbb{N}, \quad \exists i_0 > M, \quad |y_{i_0} - 0| = |y_{i_0}| \geq \varepsilon_0. \quad (1)$$

Since  $\lim_{i \rightarrow \infty} x^i = x^\infty$ ,

$$\exists M_0 \in \mathbb{N}, \quad \forall i > M_0, \quad d_1(x^i, x^\infty) < \varepsilon_0. \quad (2)$$

Consider statement (1) for a particular  $M$ ,  $M_0$ , we have

$$\exists i_0 > M_0, \quad |y_{i_0}| \geq \varepsilon_0. \quad (3)$$

That is,

$$\left| \sum_{j=1}^N x_j^{i_0} \right| \geq \varepsilon_0. \quad (3')$$

Consider statement (2) for a particular  $i$ ,  $i_0$ , we have

$$d_1(x^{i_0}, x^\infty) < \varepsilon_0. \quad (4)$$

From statement (4) we can derive:

$$\begin{aligned}
& d_1(x^{i_0}, x^\infty) < \varepsilon_0 \\
\iff & \sum_{j \in \mathbb{N}} |x_j^{i_0} - x_j^\infty| < \varepsilon_0 \\
\iff & \sum_{j=1}^N |x_j^{i_0} - x_j^\infty| + \sum_{j>N} |x_j^{i_0} - x_j^\infty| < \varepsilon_0 \\
\implies & \sum_{j>N} |x_j^{i_0} - x_j^\infty| < \varepsilon_0 \\
\iff & \sum_{j>N} |x_j^{i_0} - 0| < \varepsilon_0 \\
\iff & \sum_{j>N} |x_j^{i_0}| < \varepsilon_0 \\
\implies & \left| \sum_{j>N} x_j^{i_0} \right| < \varepsilon_0 \\
\implies & \left| \sum_{j>N} x_j^{i_0} \right| < \varepsilon_0 \\
\iff & \left| \sum_{j \in \mathbb{N}} x_j^{i_0} - \sum_{j=1}^N x_j^{i_0} \right| < \varepsilon_0 \\
\iff & \left| 0 - \sum_{j=1}^N x_j^{i_0} \right| < \varepsilon_0 \\
\iff & \left| \sum_{j=1}^N x_j^{i_0} \right| < \varepsilon_0.
\end{aligned}$$

This contradicts to statement (3'). So the original assumption that  $y_\infty \neq 0$  is false. i.e.  $y_\infty = 0$ . It follows that  $\sum_{j \in \mathbb{N}} x_j^\infty = 0$ . This completes the proof. ■






## 6

# Function Spaces

### 6.1 The $\mathcal{L}^p$ Norm

$$\|f\|_p = \left( \int_a^b |f(x)|^p dx \right)^{\frac{1}{p}}.$$

 **the instructors' answer**, where instructors collectively construct a single answer

In the sup norm, convergence coincides with uniform convergence. Moreover,  $C[a, b]$  is complete in this norm. It is not complete in any of the  $L^p$  norms for  $1 \leq p < \infty$ . The completion in these norms is called  $L^p(a, b)$ .

[undo](#) [thanks](#) | 1

Updated 1 day ago by Kenneth Davidson



# 7

## Hilbert Space

### 7.1 Hilbert Spaces

**Definition** (Hilbert Space). *We define a **Hilbert space** to be a complete inner product space.*

**Example 7.1.1.**  $\ell^2$  is a Hilbert space.



# 8

## Equicontinuity in Metric Spaces

### 8.1 Definitions

**Definition** ((Pointwise) Equicontinuity). Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces. Let  $\mathcal{F}$  be a collection of functions from  $X$  to  $Y$ . Let  $x_0$  be a point in  $X$ . We say that  $\mathcal{F}$  is **(pointwise) equicontinuous** at point  $x_0$  if for any positive number  $\varepsilon$ , there exists some number  $\delta(x_0, \varepsilon)$  such that for any function  $f$  in  $\mathcal{F}$  and any point  $x$  in  $X$ , we have

$$d_Y(f(x), f(x_0)) < \varepsilon$$

whenever  $d_X(x, x_0) < \delta(x_0, \varepsilon)$  is satisfied.

**Definition** (Uniform Equicontinuity). Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces. Let  $\mathcal{F}$  be a collection of functions from  $X$  to  $Y$ . We say that  $\mathcal{F}$  is **uniformly equicontinuous** if for any positive number  $\varepsilon$ , there exists some number  $\delta(\varepsilon)$  such that for any function  $f$  in  $\mathcal{F}$  and any points  $x_1$  and  $x_2$  in  $X$ , we have

$$d_Y(f(x_1), f(x_2)) < \varepsilon$$

whenever  $d_X(x_1, x_2) < \delta(\varepsilon)$  is satisfied.

### 8.2 Sufficient Conditions

**Proposition 8.2.1.** *The closure of an equicontinuous family of functions is equicontinuous.*

*Proof.*

Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces.

Let  $\mathcal{F}$  be an equicontinuous family of functions from  $X$  to  $Y$ .

We are to prove that  $cl(\mathcal{F})$  is equicontinuous.

Let  $x_0$  be an arbitrary point in  $X$ .

Let  $\varepsilon$  be an arbitrary positive number.

Since  $\mathcal{F}$  is equicontinuous at point  $x_0$ , there exists some  $\delta(x_0, \varepsilon)$  such that for any function  $f$  in  $\mathcal{F}$  and any point  $x$  in  $X$  such that  $d_X(x, x_0) < \delta(x_0, \varepsilon)$ , we have  $d_Y(f(x), f(x_0)) < \varepsilon/3$ .

Let  $f$  be an arbitrary function in  $cl(\mathcal{F})$ .

Let  $x$  be an arbitrary point in  $X$  such that  $d_X(x, x_0) < \delta(x_0, \varepsilon)$ .

Since  $f \in cl(\mathcal{F})$ , there exists some function  $f_0 \in \mathcal{F}$  such that  $d_\infty(f, f_0) < \varepsilon/3$ .

Since  $d_\infty(f, f_0) < \varepsilon/3$ ,  $d_Y(f(x), f_0(x)) < \varepsilon/3$  and  $d_Y(f(x_0), f_0(x_0)) < \varepsilon/3$ .

Since  $f_0 \in \mathcal{F}$  and  $d_X(x, x_0) < \delta(x_0, \varepsilon)$ ,  $d_Y(f_0(x), f_0(x_0)) < \varepsilon/3$ .

Since  $d_Y(f(x), f_0(x)) < \varepsilon/3$  and  $d_Y(f(x_0), f_0(x_0)) < \varepsilon/3$  and  $d_Y(f_0(x), f_0(x_0)) < \varepsilon/3$ ,  $d_Y(f(x), f(x_0)) < \varepsilon$ .

Since for any positive number  $\varepsilon$ , there exists some  $\delta(x_0, \varepsilon)$  such that for any function  $f$  in  $cl(\mathcal{F})$  and any point  $x$  in  $X$  such that  $d_X(x, x_0) < \delta(x_0, \varepsilon)$ , we have  $d_Y(f(x), f(x_0)) < \varepsilon$ , by definition of equicontinuous,  $cl(\mathcal{F})$  is equicontinuous at point  $x_0$ .

Since  $cl(\mathcal{F})$  is equicontinuous at point  $x_0$  for any point  $x_0$  in  $X$ ,  $cl(\mathcal{F})$  is equicontinuous. ■

# 9

## Operators

### 9.1 Bounded Operators

**Definition** (Bounded Operator). *Let  $X$  and  $Y$  be normed linear spaces. Let  $T$  be a linear map from  $X$  to  $Y$ . We say that  $T$  is a **bounded operator** if*

$$\exists k \in \mathbb{R}, \quad \forall x \in X, \quad \|Tx\|_Y \leq k\|x\|_X.$$

**Definition** (Operator Norm). *Let  $X$  and  $Y$  be normed linear spaces. Let  $T$  be a bounded operator from  $X$  to  $Y$ . We define the **operator norm** of  $T$ , denoted by  $\|T\|$ , to be the number given by*

$$\|T\| := \inf\{k \in \mathbb{R} : \forall x \in X, \|Tx\|_Y \leq k\|x\|_X\}.$$

**Proposition 9.1.1.** *Let  $X$  and  $Y$  be normed linear spaces. Let  $T$  be a linear map from  $X$  to  $Y$ . Then  $T$  is bounded if and only if  $T$  is continuous.*

**Example 9.1.1** (Multiplication Operator). *Let  $X = (\mathcal{C}([0, 1], \mathbb{C}), \|\cdot\|_\infty)$ . Let  $f$  be a function in  $X$ . We define the **multiplication operator** on  $X$ , w.r.t.  $f$ , denoted by  $M_f$ , as*

$$M_f(g) = fg.$$

*Then  $M_f$  is bounded and  $\|M_f\| = \|f\|_\infty$ .*

*Proof.* Let  $g$  be an arbitrary function in  $X$ . Then

$$\|M_f g\|_\infty = \|fg\|_\infty \leq \|f\|_\infty \|g\|_\infty.$$

So  $\|f\|_\infty$  is an element of the set  $S = \{k \in \mathbb{R} : \forall x \in X, \|Tx\|_Y \leq k\|x\|_X\}$ . So  $\|M_f\| \leq \|f\|_\infty$ . Consider  $g_0$  given by  $g_0(x) = 1$ . Then

$$\|M_f g_0\|_\infty = \|fg_0\|_\infty = \|f\|_\infty = \|f\|_\infty \|g_0\|_\infty.$$

For any  $k \in S$ , if  $k < \|f\|_\infty$ ,  $k \notin S$ . So  $\forall k \in S$ ,  $k \geq \|f\|_\infty$ . So  $\|f\|_\infty$  is a lower bound for the set  $S$ . So  $\|M_f\| \geq \|f\|_\infty$ . Since  $\|M_f\| \leq \|f\|_\infty$  and  $\|M_f\| \geq \|f\|_\infty$ , we get  $\|M_f\| = \|f\|_\infty$ . ■

**Example 9.1.2** (Weighted Shifts).

- Let  $\mathcal{H} = \ell_{\mathbb{N}}^2$ . Let  $(w_n)_{n \in \mathbb{N}} \in \ell_{\mathbb{N}}^\infty$ . We define an **unilateral forward weighted shift**  $W$  on  $\mathcal{H}$  as

$$W(x_n) := (0, w_1x_1, w_2x_2, w_3x_3, \dots).$$

Then  $W$  is bounded and  $\|W\| = \sup\{|w_n| : n \in \mathbb{N}\}$ .

- Let  $\mathcal{H} = \ell_{\mathbb{N}}^2$ . Let  $(v_n)_{n \in \mathbb{N}} \in \ell_{\mathbb{N}}^\infty$ . We define an **unilateral backward weighted shift**  $V$  on  $\mathcal{H}$  as

$$V(x_n) := (v_1x_2, v_2x_3, v_3x_4, \dots).$$

Then  $V$  is bounded and  $\|V\| = \sup\{|v_n| : n \in \mathbb{N}\}$ .

- Let  $\mathcal{H} = \ell_{\mathbb{Z}}^2$ . Let  $(u_n)_{n \in \mathbb{Z}} \in \ell_{\mathbb{Z}}^\infty$ . We define a **bilateral weighted shift**  $U$  on  $\mathcal{H}$  as

$$U(x_n) := (u_{n-1}x_{n-1})_{n \in \mathbb{Z}}.$$

Then  $U$  is bounded and  $\|U\| = \sup\{|u_n| : n \in \mathbb{Z}\}$ .

## 9.2 Space of Bounded Operators

**Proposition 9.2.1.** Let  $X$  and  $Y$  be normed linear spaces. Let  $\mathcal{B}(X, Y)$  be the space of bounded linear operators from  $X$  to  $Y$ . Then if  $Y$  is complete,  $\mathcal{B}(X, Y)$  is complete.

## 9.3 Dual Spaces

**Definition** (Dual Space). Let  $X$  be a normed linear space over field  $\mathbb{K}$ . We define the **dual** of  $X$ , denoted by  $X^*$ , to be the space  $\mathcal{B}(X, \mathbb{K})$ .

**Proposition 9.3.1.** Let  $X$  be a normed linear space. Then there exists a contractive map from  $X$  to its double dual  $X^{**}$ .



# 10

## Adjoint Operator

### 10.1 Definitions

**Definition** (Adjoint Matrix). *Let  $A$  be an  $m \times n$  matrix. We define the **adjoint** of  $A$ , denoted by  $A^*$ , to be an  $n \times m$  matrix given by*

$$(A^*)_{ij} := \overline{(A)_{ji}}.$$

**Definition** (Adjoint Operator). *Let  $V$  and  $W$  be inner product spaces. Let  $T$  be a linear map from  $V$  to  $W$ . We define the **adjoint** of  $T$ , denoted by  $T^*$ , to be a map from  $W$  to  $V$  such that*

$$\forall x \in V, \forall y \in W, \quad \langle T(x), y \rangle_W = \langle x, T^*(y) \rangle_V.$$

**Proposition 10.1.1** (Existence). *Let  $V$  be a finite-dimensional inner product space and  $T$  be a linear operator on  $V$ . Then the adjoint of  $T$  exists.*

**Proposition 10.1.2** (Uniqueness). *Let  $V$  be an inner product space and  $T$  be a linear operator on  $V$ . Then the adjoint of  $T$  is unique, provided that it exists.*

### 10.2 Properties of the Adjoint Operator

**Proposition 10.2.1.** *Let  $V$  be an inner product space. Then*

- (1)  $(I_V)^* = I_V$  where  $I_V$  is the identity operator on  $V$ .
- (2)  $T^{**} = T$  for any linear operator  $T$  on  $V$ .

**Proposition 10.2.2.** *Let  $V$  be an inner product space and  $T$  be a linear operator on  $V$ . Then  $T^*$  is also linear.*

**Proposition 10.2.3.** *Let  $V$  be an inner product space. Then*

(1) *For any linear operators  $T$  and  $U$ ,*

$$(T + U)^* = T^* + U^*.$$

(2) *For any linear operator  $T$ ,*

$$(cT)^* = \bar{c} \cdot T^*.$$

(3) *For any linear operator  $T$  and  $U$ ,*

$$(TU)^* = U^*T^*.$$

**Proposition 10.2.4.** *Let  $V$  be a finite-dimensional inner product space and  $T$  be a linear operator on  $V$ . Then if  $T$  is invertible,  $T^*$  is also invertible.*

**Proposition 10.2.5.** *Let  $V$  be an inner product space and  $T$  be an invertible linear operator on  $V$ . Then  $(T^{-1})^* = (T^*)^{-1}$ .*

### 10.3 Normal Operators

**Definition (Normal).** *Let  $V$  be an inner product space and  $T$  be a linear operator on  $V$ . We say that  $T$  is **normal** if  $TT^* = T^*T$ .*

### 10.4 Self-adjoint

# 11

## Convolution

**Definition** (Convolution). *Let  $f$  and  $g$  be functions from  $\mathbb{R}$  to  $\mathbb{R}$ . We define the **convolution** of  $f$  and  $g$ , denoted by  $f * g$ , to be a function on  $\mathbb{R}$  given by*

$$(f * g)(t) := \int_{-\infty}^{+\infty} f(\tau)g(t - \tau)dt.$$



# 12

## Coercive Functions

### 12.1 Definitions

**Definition (Coercive).** Let  $f$  be a function from  $\mathbb{R}^d$  to  $\mathbb{R}^*$ . We say that  $f$  is *coercive* if  $\lim_{\|x\| \rightarrow \infty} f(x) = +\infty$ .

### 12.2 Properties

**Proposition 12.2.1.** Let  $f$  be a proper lower semi-continuous function from  $\mathbb{R}^d$  to  $\mathbb{R}^*$ . Let  $K$  be a compact set in  $\mathbb{R}^d$ . Assume  $K \cap \text{dom}(f) \neq \emptyset$ . Then  $f$  attains its minimum over  $K$ .

*Proof.*

Define  $m := \inf_{x \in K} f(x)$ .

Since  $m = \inf_{x \in K} f(x)$ , there exists a sequence  $\{x_i\}_{i \in \mathbb{N}}$  in  $K$  such that  $\lim_{i \rightarrow \infty} f(x_i) = m$ .

Since  $K$  is compact and  $\{x_i\}_{i \in \mathbb{N}} \subseteq K$ , there exists a convergent subsequence  $\{x_i\}_{i \in I}$  in  $K$  where  $I$  is an infinite subset of  $\mathbb{N}$ .

Say the limit is  $x_\infty$  where  $x_\infty \in K$ .

Since  $\lim_{i \rightarrow \infty} f(x_i) = m$ , we get  $\lim_{i \in I, i \rightarrow \infty} f(x_i) = m$ .

Since  $\lim_{i \in I, i \rightarrow \infty} f(x_i) = m$ , we get  $\liminf_{i \in I, i \rightarrow \infty} f(x_i) = m$ .

Since  $f$  is lower semi-continuous and  $\lim_{i \in I, i \rightarrow \infty} x_i = x_\infty$ , we get  $f(x_\infty) \leq \liminf_{i \in I, i \rightarrow \infty} f(x_i)$ .

That is,  $f(x_\infty) \leq m$ .

Since  $m = \inf_{x \in K} f(x)$ , we have  $\forall x \in K, f(x) \geq m$ .

In particular,  $f(x_\infty) \geq m$ .

Since  $f(x_\infty) \geq m$  and  $f(x_\infty) \leq m$ ,  $f(x_\infty) = m$ .

Since  $f$  is proper,  $f(x_\infty) = m \neq -\infty$ .

So  $f$  attains its minimum at point  $x_\infty$ .

■

**Proposition 12.2.2.** *Let  $f$  be a proper, lower semi-continuous, and coercive function from  $\mathbb{R}^d$  to  $\mathbb{R}^*$ . Let  $C$  be a closed subset of  $\mathbb{R}^d$ . Assume  $C \cap \text{dom}(f) \neq \emptyset$ . Then  $f$  attains its minimum over  $C$ .*

*Proof.*

Since  $C \cap \text{dom}(f) \neq \emptyset$ , take  $x \in C \cap \text{dom}(f)$ .

Since  $f$  is coercive,  $\exists R$  such that  $\forall y, \|y\| > R$ , we have  $f(y) \geq f(x)$ .

Since  $x \in C \cap \text{dom}(f)$  and  $\forall y, \|y\| > R$ , we have  $f(y) \geq f(x)$ , the set of minimizers of  $f$  over  $C$  is the same as the set of minimizers of  $f$  over  $C \cap \text{ball}[0, R]$ .

Since  $C$  and  $\text{ball}[0, R]$  are both closed,  $C \cap \text{ball}[0, R]$  is closed.

Since  $\text{ball}[0, R]$  is bounded,  $C \cap \text{ball}[0, R]$  is bounded.

Since  $C \cap \text{ball}[0, R]$  is closed and bounded, by the Heine-Borel Theorem,  $C \cap \text{ball}[0, R]$  is compact.

Since  $f$  is proper and lower semi-continuous and  $C \cap \text{ball}[0, R]$  is compact,  $f$  attains its minimum over  $C \cap \text{ball}[0, R]$ .

So  $f$  attains its minimum over  $C$ . ■

# 13

## Unclassified Results

**Proposition 13.0.1.** *Let  $(X, d)$  be a compact metric space. Let  $L(X)$  be the set of all Lipschitz functions from  $X$  to  $\mathbb{R}$ . Let  $C(X)$  be the set of all continuous functions from  $X$  to  $\mathbb{R}$ . Then  $L(X)$  is dense in  $C(X)$ .*

**Proposition 13.0.2.** *Let  $(V, \|\cdot\|)$  be a normed vector space. Let  $S$  be a subset of  $V$ . Let  $p$  be a vector in  $V$ . Then we have the followings.*

$$(1) \ p + \text{int}(S) = \text{int}(p + S),$$

$$(2) \ p + \text{cl}(S) = \text{cl}(p + S).$$

*Proof.*

**Proof of (1).**

For one direction, let  $x$  be an arbitrary point in the set  $(p + \text{int}(S))$ .

We are to prove that  $x \in \text{int}(p + S)$ .

Since  $x \in (p + \text{int}(S))$ ,  $(x - p) \in \text{int}(S)$ .

Since  $(x - p) \in \text{int}(S)$ , by definition of interior, there exists a radius  $r$  such that

$$B(x - p, r) \subseteq S.$$

It follows that  $B(x, r) \subseteq p + S$ .

Since there exists a radius  $r$  such that  $B(x, r) \subseteq p + S$ , by definition of interior,

$$x \in \text{int}(p + S).$$

For the reverse direction, let  $x$  be an arbitrary point in  $\text{int}(p + S)$ .

We are to prove that  $x \in p + \text{int}(S)$ .

Since  $x \in \text{int}(p + S)$ , by definition of interior, there exists a radius  $r$  such that

$$B(x, r) \subseteq (p + S).$$

It follows that  $B(x - p, r) \subseteq S$ .

Since there exists a radius  $r$  such that  $B(x - p, r) \subseteq S$ , by definition of interior,

$$(x - p) \in \text{int}(S).$$

Since  $(x - p) \in \text{int}(S)$ , we get  $x \in (p + \text{int}(S))$ .

**Proof of (2).**

For one direction, let  $x$  be an arbitrary point in the set  $(p + \text{cl}(S))$ .

We are to prove that  $x \in \text{cl}(p + S)$ .

Since  $x \in (p + \text{cl}(S))$ , we get  $(x - p) \in \text{cl}(S)$ .

Since  $(x - p) \in \text{cl}(S)$ , by definition of closure, for any radius  $r$ , we have

$$B(x - p, r) \cap S \neq \emptyset.$$

It follows that  $B(x, r) \cap (p + S) \neq \emptyset$ .

Since for any radius  $r$ ,  $B(x, r) \cap (p + S) \neq \emptyset$ , by definition of closure, we get

$$x \in \text{cl}(p + S).$$

For the reverse direction, let  $x$  be an arbitrary point in  $\text{cl}(p + S)$ .

We are to prove that  $x \in (p + \text{cl}(S))$ .

Since  $x \in \text{cl}(p + S)$ , by definition of closure, for any radius  $r$ , we have

$$B(x, r) \cap (p + S) \neq \emptyset.$$

It follows that  $B(x - p, r) \cap S \neq \emptyset$ .

Since for any radius  $r$ ,  $B(x - p, r) \cap S \neq \emptyset$ , by definition of closure, we get

$$(x - p) \in \text{cl}(S).$$

Since  $(x - p) \in \text{cl}(S)$ , we get  $x \in (p + \text{cl}(S))$ . ■

**Proposition 13.0.3.** *Let  $(V, \|\cdot\|)$  be a normed vector space. Let  $S$  be a subset of  $V$ . Let  $\lambda$  be a non-zero real number. Then*

$$(1) \lambda \text{int}(S) = \text{int}(\lambda S).$$

$$(2) \lambda \text{cl}(S) = \text{cl}(\lambda S).$$

*Proof.*

**Proof of (1).**

For one direction, let  $x$  be an arbitrary point in  $\lambda \text{int}(S)$ .

We are to prove that  $x \in \text{int}(\lambda S)$ .



Since  $x \in \lambda \operatorname{int}(S)$ , we get  $x/\lambda \in \operatorname{int}(S)$ .

Since  $x/\lambda \in \operatorname{int}(S)$ , by definition of interior, there exists a radius  $r$  such that

$$B(x/\lambda, r) \subseteq S.$$

Let  $y$  be an arbitrary point in  $B(x, \lambda r)$ .

Since  $y \in B(x, \lambda r)$ , we get  $\|y - x\| \leq \lambda r$ .

Since  $\|y - x\| \leq \lambda r$ , we get  $\|y/\lambda - x/\lambda\| \leq r$ .

Since  $\|y/\lambda - x/\lambda\| \leq r$ , we get  $y/\lambda \in B(x/\lambda, r)$ .

Since  $y/\lambda \in B(x/\lambda, r)$  and  $B(x/\lambda, r) \subseteq S$ , we get  $y/\lambda \in S$ .

Since  $y/\lambda \in S$ , we get  $y \in \lambda S$ .

Since any point in  $B(x, \lambda r)$  is also in  $\lambda S$ , we get  $B(x, \lambda r) \subseteq \lambda S$ .

Since there exists a radius  $r$  such that  $B(x, \lambda r) \subseteq \lambda S$ , by definition of interior, we get

$$x \in \operatorname{int}(\lambda S).$$

For the reverse direction,

■