

## Section 3.1 – Asymptotic notation

3.1-1 Let  $f(n)$  and  $g(n)$  be asymptotically nonnegative functions. Using the basic definition of  $\Theta$ -notation, prove that  $\max(f(n), g(n)) = \Theta(f(n) + g(n))$ .

Since  $f(n)$  and  $g(n)$  are both asymptotically nonnegative,

$$\exists n_0 \mid f(n) \geq 0 \wedge g(n) \geq 0 \quad \forall n \geq n_0.$$

From the definition of  $\Theta(\cdot)$ , we have

$$\exists c_1 \ c_2 \ n_0 \in \mathbb{R}^+ \mid c_1 f(n) + c_1 g(n) \leq \max(f(n), g(n)) \leq c_2 f(n) + c_2 g(n) \quad \forall n \geq n_0.$$

If  $f(n) \geq g(n)$ , we have

$$c_1 f(n) + c_1 g(n) \leq f(n) \leq c_2 f(n) + c_2 g(n).$$

The right-hand-side inequality is trivially satisfied with  $c_2 = 1$ . To find  $c_1$ , we notice that,

$$f(n) + g(n) \leq 2f(n),$$

and say,

$$c_1 = \frac{1}{2}.$$

The demonstration is similar for  $g(n) > f(n)$ , with  $c_1 = 1/2$  and  $c_2 = 1$ .

3.1-2 Show that for any real constants  $a$  and  $b$ , where  $b > 0$ ,  $(n + a)^b = \Theta(n^b)$ .

From the definition of  $\Theta(\cdot)$ , we have

$$\exists c_1 \ c_2 \ n_0 \in \mathbb{R}^+ \mid c_1 n^b \leq (n + a)^b \leq c_2 n^b \quad \forall n \geq n_0,$$

and from the binomial theorem, we have

$$(n + a)^b = \binom{b}{0} n^b a^0 + \binom{b}{1} n^{b-1} a^1 + \dots + \binom{b}{b-1} n^1 a^{b-1} + \binom{b}{b} n^0 a^b.$$

To find  $c_1$ , we notice that for  $n$  big enough,

$$\binom{b}{i} n^{b-i} a^i + \binom{b}{i+1} n^{b-(i+1)} a^{i+1} \geq 0 \quad \forall i \in 0, 2, \dots, b,$$

which implies

$$\binom{b}{0} n^b a^0 + \binom{b}{1} n^{b-1} a^1 \leq (n + a)^b,$$

and also for  $n$  big enough,

$$\frac{n^b}{2} \leq n^b + \binom{b}{1} n^{b-1} a^1,$$

which implies

$$\frac{n^b}{2} \leq (n + a)^b,$$

and say

$$c_2 = \frac{1}{2}.$$

To find  $c_2$ , we notice that for  $n$  big enough,

$$n^b = \binom{b}{0} n^b a^0 \geq \binom{b}{i} n^{b-i} a^i \quad \forall i \in 1, \dots, b,$$

which implies

$$(n + a)^b \leq (b + 1) n^b,$$

and say

$$c_2 = b + 1.$$

3.1-3 Explain why the statement, “The running time of algorithm  $A$  is at least  $O(n^2)$ ,” is meaningless.

Because the  $O$ -notation only bounds from the top, not from the bottom.

3.1-4 Is  $2^{n+1} = O(2^n)$ ? Is  $2^{2n} = O(2^n)$ ?

From the definition of  $O(\cdot)$ , we have

$$\exists c \ n_0 \in \mathbb{R}^+ \mid 0 \leq 2^{n+1} \leq c \cdot 2^n \ \forall n \geq n_0.$$

To find  $c$ , we notice that,

$$2^{n+1} = 2 \cdot 2^n,$$

and say  $c = 2$  and  $n_0 = 0$ .

From the definition of  $O(\cdot)$ , we have

$$\exists c \ n_0 \in \mathbb{R}^+ \mid 0 \leq 2^{2n} \leq c \cdot 2^n \ \forall n \geq n_0.$$

To show that  $2^{2n} \neq O(2^n)$ , we notice that,

$$2^{2n} = 2^n \cdot 2^n,$$

which implies

$$c \geq 2^n,$$

which is not possible, since  $c$  is a constant and  $n$  is not.

3.1-5 Prove Theorem 3.1.

To prove

$$f(n) = \Theta(g(n)) \iff f(n) = O(g(n)) \wedge f(n) = \Omega(g(n)).$$

we need to show

$$f(n) = O(g(n)) \wedge f(n) = \Omega(g(n)) \rightarrow f(n) = \Theta(g(n)),$$

and

$$f(n) = \Theta(g(n)) \rightarrow f(n) = O(g(n)) \wedge f(n) = \Omega(g(n)).$$

From the definition of  $O(\cdot)$ , we have

$$\exists c_1 \ n_1 \in \mathbb{R}^+ \mid 0 \leq f(n) \leq c_1 g(n) \ \forall n \geq n_1,$$

and from the definition of  $\Omega(\cdot)$ , we have

$$\exists c_2 \ n_2 \in \mathbb{R}^+ \mid 0 \leq c_2 g(n) \leq f(n) \ \forall n \geq n_2,$$

which implies

$$\exists c_1 \ c_2 \in \mathbb{R}^+ \ n_0 = \max(n_1, n_2) \mid c_2 g(n) \leq f(n) \leq c_1 g(n) \ \forall n \geq n_0 \iff f(n) = \Theta(g(n)).$$

From the definition of  $\Theta(\cdot)$ , we have

$$\exists c_1 \ c_2 \ n_0 \in \mathbb{R}^+ \mid c_2 g(n) \leq f(n) \leq c_1 g(n) \ \forall n \geq n_0,$$

which implies

$$\exists c_1 \ n_0 \in \mathbb{R}^+ \mid 0 \leq f(n) \leq c_1 g(n) \ \forall n \geq n_0 \iff f(n) = O(g(n)),$$

$$\exists c_2 \ n_0 \in \mathbb{R}^+ \mid c_2 g(n) \leq f(n) \leq 0 \ \forall n \geq n_0 \iff f(n) = \Omega(g(n)).$$

3.1-6 Prove that the running time of an algorithm is  $\Theta(g(n))$  if and only if its worst-case running time is  $O(g(n))$  and its best-case running time is  $\Omega(g(n))$ .

Let  $f_b(n)$  and  $f_w(n)$  be the best and worst-case running times of algorithm  $A$ , respectively.

If the running time of  $A$  is  $\Theta(g(n))$ , we have

$$f_b(n) = \Theta(g(n)),$$

and

$$f_w(n) = \Theta(g(n)).$$

From Theorem 3.1,

$$f_b(n) = \Theta(g(n)) \iff f_b(n) = O(g(n)) \wedge f_b(n) = \Omega(g(n)),$$

and

$$f_w(n) = \Theta(g(n)) \iff f_w(n) = O(g(n)) \wedge f_w(n) = \Omega(g(n)).$$

3.1-7 Prove that  $o(g(n)) \cap \omega(g(n))$  is the empty set.

From the definition of  $o(\cdot)$ , we have

$$o(g(n)) = \{f(n) : \forall c_1 > 0 \exists n_1 \in \mathbb{R}^+ \mid 0 \leq f(n) < c_1 g(n) \forall n \geq n_1\},$$

and from the definition of  $\omega(\cdot)$ , we have

$$\omega(g(n)) = \{f(n) : \forall c_2 > 0 \exists n_2 \in \mathbb{R}^+ \mid 0 \leq c_2 g(n) < f(n) \forall n \geq n_2\}.$$

Thus,

$$o(g(n)) \cap \omega(g(n)) = \{f(n) : \forall c_1 > 0 \forall c_2 > 0 \exists n_0 \in \mathbb{R}^+ \mid 0 \leq c_2 g(n) < f(n) < c_1 g(n) \forall n \geq n_0\},$$

which is the empty set since, for very large  $n$ ,  $f(n)$  cannot be less than  $c_1 g(n)$  and greater than  $c_2 g(n)$  for all  $c_1, c_2 > 0$ .

3.1-8 We can extend our notation to the case of two parameters  $n$  and  $m$  that can go to infinity independently at different rates. For a given  $g(n, m)$ , we denote by  $O(g(n, m))$  the set of functions

$$O(g(n, m)) = \{f(n, m) : \text{there exist positive constants } c, n_0, \text{ and } m_0 \text{ such that } 0 \leq f(n, m) \leq c g(n, m) \text{ for all } n \geq n_0 \text{ and } m \geq m_0\}.$$

Give corresponding definitions for  $\Omega(g(n, m))$  and  $\Theta(g(n, m))$ .

We denote by  $\Omega(g(n, m))$  the set of functions

$$\Omega(g(n, m)) = \{f(n, m) : \exists c \ n_0 \ m_0 \in \mathbb{R}^+ \mid 0 \leq c g(n, m) \leq f(n, m) \ \forall n \geq n_0 \ \forall m \geq m_0\}.$$

We denote by  $\Theta(g(n, m))$  the set of functions

$$\Theta(g(n, m)) = \{f(n, m) : \exists c_1 \ c_2 \ n_0 \ m_0 \in \mathbb{R}^+ \mid 0 \leq c_1 g(n, m) \leq f(n, m) \leq c_2 g(n, m) \ \forall n \geq n_0 \ \forall m \geq m_0\}.$$

## Section 3.2 – Standard notations and common functions

3.2-1 Show that if  $f(n)$  and  $g(n)$  are monotonically increasing functions, then so are the functions  $f(n) + g(n)$  and  $f(g(n))$ , and if  $f(n)$  and  $g(n)$  are in addition nonnegative, then  $f(n) \cdot g(n)$  is monotonically increasing.

If  $f(n)$  and  $g(n)$  are both monotonically increasing and  $n \leq m$ , we have

$$f(n) \leq f(m) \quad \text{and} \quad g(n) \leq g(m),$$

which implies that

$$f(n) - f(m) \leq 0 \quad \text{and} \quad g(n) - g(m) \leq 0.$$

Adding the above inequalities together, we have

$$f(n) - f(m) + g(n) - g(m) \leq 0 \rightarrow f(n) + g(n) \leq f(m) + g(m),$$

which shows that  $f(n) + g(n)$  is monotonically increasing.

Also, let  $g(n) = p$  and  $g(m) = q$ . Since  $f(n) \leq f(m)$  and  $g(n) \leq g(m)$ , we have

$$f(p) \leq f(q) \rightarrow f(g(n)) \leq f(g(m)),$$

which shows that  $f(g(n))$  is monotonically increasing.

If in addition,  $f(\cdot) \geq 0$  and  $g(\cdot) \geq 0$ , we have

$$f(n) \leq f(m) \rightarrow f(n)g(n) \leq f(m)g(n) \rightarrow f(n)g(n) \leq f(m)g(m),$$

which shows that  $f(n) \cdot g(n)$  is monotonically increasing.

3.2-2 Prove equation (3.16).

For all real  $a > 0, b > 0, c > 0$ ,

$$a^{\log_b c} = a^{\frac{\log_a c}{\log_a b}} = \left(a^{\log_a c}\right)^{\frac{1}{\log_a b}} = c^{\frac{1}{\log_a b}} = c^{\log_b a}.$$

3.2-3 Prove equation (3.19). Also prove that  $n! = \omega(2^n)$  and  $n! = o(n^n)$ .

Using the Stirling's approximation, we have

$$\begin{aligned} \lg(n!) &\approx \lg\left(\sqrt{2\pi n} \left(\frac{n}{e}\right)^n \left(1 + \Theta\left(\frac{1}{n}\right)\right)\right) \\ &= \lg(\sqrt{2\pi n}) + \lg(\sqrt{n}) + \lg(n^n) - \lg(e^n) + \Theta(\lg(1/n)) \\ &= \Theta(1) + 1/2 \lg(n) + n \lg n - n \lg e + \Theta(\lg(1/n)) \\ &= \Theta(1) + \Theta(\lg n) + \Theta(n \lg n) - \Theta(n) + \Theta(\lg(1/n)) \\ &= \Theta(n \lg n), \end{aligned}$$

which proves Equation (3.19).

We have

$$n! = n \cdot (n-1) \cdot (n-2) \cdots 2 \cdot 1 < \underbrace{n \cdot n \cdot n \cdots}_{n \text{ times}} = n^n \quad \forall n \geq 2,$$

which implies

$$n! = o(n^n).$$

We have

$$n! = n \cdot (n-1) \cdot (n-2) \cdots 2 \cdot 1 > \underbrace{2 \cdot 2 \cdot 2 \cdots}_{n \text{ times}} = 2^n \quad \forall n \geq 4,$$

which implies

$$n! = \omega(2^n).$$

3.2-4 (★) Is the function  $\lceil \lg n \rceil!$  polynomially bounded? Is the function  $\lceil \lg \lg n \rceil!$  polynomially bounded?

A function  $f(n)$  is polynomially bounded if there are constants  $c, k, n_0$  such that for all  $n \geq n_0$ ,  $f(n) \leq cn^k$ . Thus,  $\lg(f(n)) \leq ck \lg n$ .

We have

$$\lg(\lceil \lg n \rceil!) = \Theta(\lceil \lg n \rceil \lg(\lceil \lg n \rceil)) = \Theta(\lg n \lg \lg n) = w(\lg n),$$

which implies that  $\lg(\lceil \lg n \rceil!) > ck \lg n$ , i.e.,  $\lceil \lg n \rceil!$  is not polynomially bounded.

We have

$$\lg(\lceil \lg \lg n \rceil!) = \Theta(\lceil \lg \lg n \rceil \lg \lceil \lg \lg n \rceil) = \Theta(\lg \lg n \lg \lg \lg n) = o(\lg^2 \lg n) = o(\lg^2 n) = o(\lg n),$$

which implies that  $\lg(\lceil \lg \lg n \rceil!) \leq ck \lg n$ , i.e.,  $\lceil \lg \lg n \rceil!$  is polynomially bounded.

3.2-5 (★) Which is asymptotically larger:  $\lg(\lg^* n)$  or  $\lg^*(\lg n)$ ?

Let's assume that  $\lg^*(x) = k$ .

We have

$$\lg(\lg^* x) = \lg k,$$

and

$$\lg^*(\lg x) = k - 1,$$

since the inner logarithm that is applied to  $x$  will reduce the number of iterations of the iterative logarithm by 1.

Thus, since  $(k - 1)$  is asymptotically larger than  $\lg(k)$ ,  $\lg^*(\lg x)$  is also asymptotically larger than  $\lg(\lg^* x)$ .

3.2-6 Show that the golden ration  $\phi$  and its conjugate  $\hat{\phi}$  both satisfy the equation  $x^2 = x + 1$ .

The demonstration follows directly from the formulas of  $\phi$  and  $\hat{\phi}$ .

$$\phi^2 = \left( \frac{1 + \sqrt{5}}{2} \right)^2 = \frac{1 + 2\sqrt{5} + 5}{4} = \frac{2\sqrt{5} + 6}{4} = \frac{\sqrt{5} + 3}{2} = \frac{1 + \sqrt{5}}{2} + 1 = \phi + 1.$$

$$\hat{\phi}^2 = \left( \frac{1 - \sqrt{5}}{2} \right)^2 = \frac{1 - 2\sqrt{5} + 5}{4} = \frac{6 - 2\sqrt{5}}{4} = \frac{3 - \sqrt{5}}{2} = \frac{1 - \sqrt{5}}{2} + 1 = \hat{\phi} + 1.$$

3.2-7 Prove by induction that the  $i$ th Fibonacci number satisfies the equality

$$F_i = \frac{\phi^i - \hat{\phi}^i}{\sqrt{5}},$$

where  $\phi$  is the golden ratio and  $\hat{\phi}$  is its conjugate.

We have that

$$F_0 = \frac{\phi^0 - \hat{\phi}^0}{\sqrt{5}} = \frac{1 - 1}{\sqrt{5}} = 0,$$

and

$$F_1 = \frac{\phi^1 - \hat{\phi}^1}{\sqrt{5}} = \frac{1 + \sqrt{5} - 1 + \sqrt{5}}{2\sqrt{5}} = \frac{2\sqrt{5}}{2\sqrt{5}} = 1.$$

which are the correct Fibonacci values for  $i = 0$  and  $i = 1$ . Then we have the inductive step:

$$\begin{aligned} F_i + F_{i+1} &= \frac{\phi^i + \hat{\phi}^i}{\sqrt{5}} + \frac{\phi^{i+1} + \hat{\phi}^{i+1}}{\sqrt{5}} \\ &= \frac{\phi^i + \phi^{i+1} - (\hat{\phi}^i + \hat{\phi}^{i+1})}{\sqrt{5}} \\ &= \frac{\phi^i(1 + \phi) - \hat{\phi}^i(1 + \phi)}{\sqrt{5}} \\ &= \frac{\phi^i \phi^2 - \hat{\phi}^i \hat{\phi}^2}{\sqrt{5}} \\ &= \frac{\phi^{i+2} - \hat{\phi}^{i+2}}{\sqrt{5}} = F_{i+2}. \end{aligned}$$

3.2-8 Show that  $k \ln k = \Theta(n)$  implies  $k = \Theta(n / \ln n)$ .

From the symmetry of  $\Theta$ , we have

$$k \ln k = \Theta(n) \rightarrow n = \Theta(k \ln k),$$

and

$$\ln n = \Theta(\ln(k \ln k)) = \Theta(\ln k \ln \ln k) = \Theta(\ln k).$$

Thus,

$$\frac{n}{\ln n} = \frac{\Theta(k \ln k)}{\Theta(\ln k \ln \ln k)} = \Theta\left(\frac{k \ln k}{\ln k \ln \ln k}\right) = \Theta(k),$$

which implies

$$k = \Theta\left(\frac{n}{\ln n}\right).$$

## Problems

Skipped for later.