

## ECON 706: PROBLEM SET 1

DANIEL PFEFFER

**Problem 1.** Prove that for a weakly stationary process,  $|\gamma(\tau)| \leq \gamma(0)$  for all  $\tau$ .

*Solution.* Let  $\{y_t\}$  be a weakly stationary process, and without loss of generality assume the mean of  $y_t$  is zero for all  $t$ . Then, by the Cauchy-Schwartz inequality,

$$|\gamma(\tau)| = E y_t y_{t-\tau} \leq \sqrt{E y_t^2 E y_{t-\tau}^2} \leq \gamma(0).$$

□

**Problem 2.** Consider an AR(2) process

- (i) Characterize the conditions for stationarity.

*Solution.* Write the AR(2) process as  $(1 - \phi_1 L - \phi_2 L^2)y_t = \varepsilon_t$  where  $\varepsilon_t \sim WN(0, \sigma^2)$ , and consider the characteristic polynomial  $\phi(z) = 1 - \phi_1 z - \phi_2 z^2$ . Then the  $\{y_t\}$  process is non-explosive if the characteristic equation

$$z^2 - \phi_1 z - \phi_2 = 0$$

has root pairs  $(z_1, z_2)$  outside the locus  $\{z : |z| > 1\}$  called the unit circle:

$$\left| \frac{-\phi_1 \pm \sqrt{\phi_1^2 + 4\phi_2}}{2} \right| > 1.$$

From this restriction on the roots of  $\phi(z) = 0$  we can deduce restrictions on the autoregressive parameters. Substituting these roots into the characteristic polynomial for the coefficients gives the factorization

$$\phi(z) = (1 - z_1^{-1}z)(1 - z_2^{-1}z).$$

Here, it is identical to the case that  $\phi(z_1) = \phi(z_2) = 0$ . Next rewrite the AR(2) process with the lag operator using this factorization:  $(1 - z_1^{-1}L)(1 - z_2^{-1}L)y_t = \varepsilon_t$ , which implies that  $\phi_1 = z_1^{-1} + z_2^{-1}$  and  $\phi_2 = -(z_1 z_2)^{-1}$ . Then, together with the restriction that  $|z_1| > 1$  and  $|z_2| > 1$ , we obtain the values for which  $\{y_t\}$  is non-explosive:

$$\phi_1 + \phi_2 < 1, \quad \phi_2 - \phi_1 < 1, \quad |\phi_2| < 1,$$

a convex set in parameter space. □

- (ii) Assuming stationarity, derive its Wold representation, i.e., the coefficients  $\psi_i$  for  $i = 1, 2, \dots$

*Solution.* To obtain the Wold representation of an AR(2) process, invert the autoregressive lag polynomial using the partial fraction decomposition:

$$\frac{1}{(1 - \lambda_1 L)(1 - \lambda_2 L)} = \frac{c_1}{(1 - \lambda_1 L)} + \frac{c_2}{(1 - \lambda_2 L)} = \frac{c_1(1 - \lambda_2 L) + c_2(1 - \lambda_1 L)}{(1 - \lambda_1 L)(1 - \lambda_2 L)}$$

where  $c_1$  and  $c_2$  are coefficients to be determined. We require that  $c_1 + c_2 = 1$  and  $\lambda_2 c_1 + \lambda_1 c_2 = 0$ . Hence, the solution is

$$c_1 = \frac{\lambda_1}{\lambda_1 - \lambda_2}, \quad c_2 = \frac{-\lambda_2}{\lambda_1 - \lambda_2}.$$

Substituting this solution into the partial fraction decomposition gives

$$\frac{1}{(1 - \lambda_1 L)(1 - \lambda_2 L)} = \frac{\lambda_1}{(\lambda_1 - \lambda_2)(1 - \lambda_1 L)} + \frac{\lambda_2}{(\lambda_2 - \lambda_1)(1 - \lambda_2 L)}.$$

Now apply this expression for the inverted lag polynomial operator to our AR(2) process to obtain the Wold representation:

$$\begin{aligned} y_t &= \frac{\lambda_1}{\lambda_1 - \lambda_2} \sum_{i=0}^{\infty} \lambda_1^i \varepsilon_{t-i} + \frac{\lambda_2}{\lambda_2 - \lambda_1} \sum_{i=0}^{\infty} \lambda_2^i \varepsilon_{t-i} \\ &= \sum_{i=0}^{\infty} \left( \frac{\lambda_1}{\lambda_1 - \lambda_2} \lambda_1^i + \frac{\lambda_2}{\lambda_2 - \lambda_1} \lambda_2^i \right) \varepsilon_{t-i} \\ &= \sum_{i=0}^{\infty} (c_1 \lambda_1^i + c_2 \lambda_2^i) \varepsilon_{t-i} = \sum_{i=0}^{\infty} \psi_i \varepsilon_{t-i}, \end{aligned}$$

where  $\psi_i = c_1 \lambda_1^i + c_2 \lambda_2^i$ , for  $i = 0, 1, 2, \dots$ . □

(iii) Verify that  $\psi_0 = 1$ .

*Solution.* To verify that  $\psi_0 = 1$ , first note that  $\psi_0 = c_1 \lambda_1^0 + c_2 + \lambda_2^0 = c_1 + c_2$ , and then substitute in the expressions for  $c_1$  and  $c_2$  derived in (b) to obtain

$$\psi_0 = \frac{\lambda_1}{\lambda_1 - \lambda_2} + \frac{-\lambda_2}{\lambda_1 - \lambda_2} = \frac{\lambda_1 - \lambda_2}{\lambda_1 - \lambda_2} = 1,$$

the desired result. □

(iv) Verify that the square summability condition holds, i.e.,  $\sum_{i=0}^{\infty} \psi_i^2 < \infty$ .

*Solution.* Write the infinite sum

$$\sum_{i=0}^{\infty} \psi_i^2 = \sum_{i=0}^{\infty} (c_1 \lambda_1^i + c_2 \lambda_2^i)^2 = c_1^2 \sum_{i=0}^{\infty} \lambda_1^{2i} + c_2^2 \sum_{i=0}^{\infty} \lambda_2^{2i} + c_1 c_2 \sum_{i=0}^{\infty} (\lambda_1 \lambda_2)^{2i}.$$

We know from the stationarity conditions derived in (i) that  $\lambda_1$  and  $\lambda_2$  have modulus less than unity, which implies that each term is a convergent geometric series. Hence  $\sum_{i=0}^{\infty} \psi_i^2 < \infty$ . □

**Problem 3.** Is the following process stationary?

$$y_t = 0.5y_{t-1} + 0.9y_{t-2} - 0.1y_{t-3} + 0.3y_{t-4} + 0.5\varepsilon_{t-1} + \varepsilon_t \quad (1)$$

*Solution.* First rewrite the process in lag operator notation as

$$(1 - 0.5L - 0.9L^2 + 0.1L^3 - 0.3L^4)y_t = (1 + 0.5L)\varepsilon_t.$$

Notice that we can factor autoregressive operator and moving average operator as

$$\begin{aligned} \phi(L) &= (1 - 0.5L - 0.9L^2 + 0.1L^3 - 0.3L^4) = -0.1(L + 2)(3L^3 - 7L^2 + 5L - 5) \\ \theta(L) &= (1 + 0.5L) = 0.5(L + 2). \end{aligned}$$

which have a common factor  $(L + 2)$  and therefore a redundant parameter. So we can rewrite this process as an AR(3) model of the form  $-0.1(3L^3 - 7L^2 + 5L - 5)y_t = 0.5\varepsilon_t$ . This process is stationary if the roots of the characteristic polynomial

$$\phi(z) = -0.3z^3 + 0.7z^2 - 0.5z + 0.5.$$

are outside the unit circle. Factoring the characteristic equation as

$$-0.3(z - 1.91744)(z^2 - 0.415894z + 0.869215) = 0$$

immediately gives the real root  $z_1 = 1.91744$ , which is outside the unit circle. To obtain the other two roots, use the quadratic equation

$$z_2, z_3 = \frac{0.415894 \pm \sqrt{(-0.415894)^2 + 4(0.869215)}}{2}$$

which gives the complex roots (that occur in a complex conjugate pair) as  $z_2 = 0.207947 + 0.908831i$  and  $z_3 = 0.207947 - 0.908831i$ . Their modulus is  $0.932317 < 1$ , and so the process is not stationary.  $\square$

**Problem 4.** Consider

$$\begin{aligned} x_t &= \phi_1 x_{t-1} + \phi_2 x_{t-2} + \phi_3 x_{t-3} + \varepsilon_t \\ y_t &= \theta_1 \varepsilon_1 + \theta_2 \varepsilon_{t-2} + \varepsilon_t \\ z_t &= \phi_1 z_{t-1} + \phi_2 z_{t-2} + \phi_3 z_{t-3} + \theta_1 \varepsilon_t + \theta_2 \varepsilon_{t-2} + \varepsilon_t. \end{aligned}$$

Take  $(\phi_1, \phi_2, \phi_3, \theta_1, \theta_2) = (0.6, 0.2, 0.1, 0.5, -0.2)$  and assume  $\varepsilon_t \sim N(0, 1)$ .

- (i) Starting from arbitrary initial values, simulate a plot of a series of  $T = 10000$  realizations for each process.

*Solution.* Below is the code used to simulate the processes and plot the resulting series. Figure 1 contains simulated series for the  $x_t$ ,  $y_t$  and  $z_t$  processes, respectively.

---

```
import numpy as np
import matplotlib.pyplot as plt
from statsmodels.graphics.tsaplots import plot_acf, plot_pacf

# Simulate the series
np.random.seed(42)
T = 10000
eps = np.random.normal(0,1,size=T)
x, y, z = np.empty_like(eps), np.empty_like(eps), np.empty_like(eps)
for t in range(T):
    x[t] = 0.6*x[t-1] + 0.2*x[t-2] + 0.1*x[t-3] + eps[t]
    y[t] = 0.5*eps[t-1] - 0.2*eps[t-2] + eps[t]
    z[t] = 0.6*z[t-1] + 0.2*z[t-2] + 0.1*z[t-3] \
          + 0.5*eps[t-1] - 0.2*eps[t-2] + eps[t]

# Plot results
plt.figure(figsize=(12,4))

plt.subplot(131)
plt.plot(x)
```




FIGURE 1

```
plt.ylabel('$x_t$')

plt.subplot(132)
plt.plot(y)
plt.ylabel('$y_t$')

plt.subplot(133)
plt.plot(z)
plt.ylabel('$z_t$')

plt.tight_layout()
plt.show()
```

□

- (ii) Compute and plot the (empirical) ACF and PACF for each process. Discuss.

*Solution.* Below is the code used to plot the ACF and PACF for each process. Figure 2 contains the sample autocorrelation and partial autocorrelation plots for the  $x_t$ ,  $y_t$  and  $z_t$  processes, respectively. The first row of Figure 2 (AR(3)) displays a geometrically decaying autocorrelation function and a partial autocorrelation function that becomes zero after 3 lags. The second row of Figure 2 (MA(2)) has an autocorrelation function that becomes zero after two lags and partial autocorrelation function that tends to zero only gradually. The third row of Figure 2 (ARMA(3,2)) has an autocorrelation function resembling an AR(3) process since it decays geometrically and a partial autocorrelation function that resembles an MA(2) process because it converges to zero gradually.

```
fig, ax = plt.subplots(3,2,figsize=(12,8))

plot_acf(x, ax=ax[0,0])
plot_pacf(x, ax=ax[0,1])

plot_acf(y, ax=ax[1,0], title='')
plot_pacf(y, ax=ax[1,1], title='')

plot_acf(z, ax=ax[2,0], title='')
plot_pacf(z, ax=ax[2,1], title='')
```




FIGURE 2

```
plt.tight_layout()
plt.show()
```

□