

Computational geophysics

Equations: 1D heat equation

$$\rho c_v \partial_t T = \partial_x (K \partial_x T) + f$$

ρ : density $\rho = \rho(x)$
 c_v : specific heat $c_v = c_v(x)$
 K : conductivity $K = K(x)$

homogeneous materials $\rho = \text{const}$, $c_v = \text{const}$,
 $K = \text{const}$:

$$\partial_t T = \underbrace{\frac{K}{\rho c_v}}_{D = \text{const}} \partial_x^2 T$$

D : diffusivity

$$\partial_t T = D \partial_x^2 T \quad \text{diffusion equation}$$

1D wave equation

$$\rho \partial_t^2 s = \partial_x (\mu \partial_x s) + f$$

μ : (shear) modulus $\mu = \mu(x)$
 ρ : density $\rho = \rho(x)$
 s : displacement $s(x, t)$

homogeneous material $\rho = \text{const}$, $\mu = \text{const}$

$$\partial_t^2 s = \underbrace{\frac{\mu}{\rho}}_{c^2 = \text{const}} \partial_x^2 s$$

$$\partial_t^2 s = c^2 \partial_x^2 s$$

methods: $\left. \begin{array}{l} \text{finite-difference} \\ \text{pseudo-spectral} \end{array} \right\} \text{strong form} \left\{ \begin{array}{l} \text{low-order} \\ \text{infinite order} \end{array} \right.$

$\left. \begin{array}{l} \text{finite-element} \\ \text{spectral-element} \end{array} \right\} \text{weak form} \left\{ \begin{array}{l} \text{low-order} \\ \text{high-order} \end{array} \right.$


time schemes: implicit

explicit — conditionally
stable
(CFL condition)

missing points / details:

- finite-difference :

- staggered-grids
- rotated
- curvilinear



- absorbing boundary conditions :


- PML
- Sommerfeld
- Clayton-Engquist

- other methods :

- finite-volume method ← low-order

- discontinuous Galerkin method ← high-order

- Lattice - Boltzmann method



- particle-in-cell ← non-linear equations
Lagrangian