

Numerical approximations of harmonic 1-forms on real loci of Calabi-Yau manifolds

Daniel Platt (Imperial College London)
Loughborough University, 20 March 2024

Abstract: For applications in differential geometry and string theory one would like to construct Calabi-Yau manifolds of complex dimension three with the following property: it should contain a real submanifold of real dimension three that admits a harmonic nowhere vanishing 1-form. Many examples are expected to exist, but none have been proven to exist. The problem is that there is no explicit formula for the Calabi-Yau metric which makes it hard to write down the “harmonic” equation, let alone solve it. In the talk I will present numerical approximations of the Calabi-Yau metric, and numerical approximations of harmonic 1-forms, obtained by neural networks. This suggests some conjectural examples of harmonic, nowhere vanishing 1-forms. I will also show some proven non-examples, and explain the main long-term motivation for this numerical work, which is to numerically verifiably prove that there exists a genuine solution to the harmonic equation near the approximate solutions. This is work in progress, joint with Michael Douglas and Yidi Qi.

Background I: Calabi-Yau manifolds

- ▶ Calabi conjecture (Yau's theorem):
If (Y, g, J, ω) Kähler, complex dim n with:

$\Omega \in \Omega^{n,0}(Y)$ parallel and nowhere 0 s.t.

then ex. $\phi \in C^\infty(Y)$ s.t. $\omega_{CY} = \omega + i\partial\bar{\partial}\phi$ has $\omega_{CY}^n = \Omega \wedge \bar{\Omega}$
(\Rightarrow induced metric g_{CY} is Ricci-flat)

- ▶ Example: Fermat quintic

$$Y := \{z = [z_0 : \cdots : z_4] \in \mathbb{CP}^4 : z_0^5 + \cdots + z_4^5 = 0\}$$

has $\Omega \in \Omega^{n,0}(Y) \Rightarrow g_{CY}$ exists

Background I: Calabi-Yau manifolds

- ▶ Calabi conjecture (Yau's theorem):
If (Y, g, J, ω) Kähler, complex dim n with:

$\Omega \in \Omega^{n,0}(Y)$ parallel and nowhere 0 s.t.

then ex. $\phi \in C^\infty(Y)$ s.t. $\omega_{CY} = \omega + i\partial\bar{\partial}\phi$ has $\omega_{CY}^n = \Omega \wedge \bar{\Omega}$
(\Rightarrow induced metric g_{CY} is Ricci-flat)


- ▶ Example: Fermat quintic

$$Y := \{z = [z_0 : \cdots : z_4] \in \mathbb{CP}^4 : z_0^5 + \cdots + z_4^5 = 0\}$$

has $\Omega \in \Omega^{n,0}(Y) \Rightarrow g_{CY}$ exists

Background II: Harmonic 1-forms on Calabi-Yaus

- ▶ Let Y be Calabi-Yau 3-fold with **Calabi-Yau metric** g_{CY}
- ▶ $\sigma : Y \rightarrow Y$ anti-holomorphic involution, $L := \text{fix}(\sigma)$
example: quintic with real coefficients in \mathbb{CP}^4 and $\sigma([z_0 : \cdots : z_4]) = [\bar{z}_0 : \cdots : \bar{z}_4]$
- ▶ $S^1 \times Y$ has dimension 7 and holonomy $SU(3)$. Problem: want holonomy G_2
- ▶ Define $\hat{\sigma} : S^1 \times Y \rightarrow S^1 \times Y$ as $(x, y) \mapsto (-x, \sigma(y))$



Theorem ([Joyce and Karigiannis, 2017])


If there exists $\lambda \in \Omega^1(L)$ **harmonic w.r.t. $g_{CY}|_L$ that is nowhere 0**, then there exists a resolution $N^7 \rightarrow (S^1 \times Y)/\langle \hat{\sigma} \rangle$ with holonomy equal to G_2 .

- ▶ **Goal: check if such a 1-form exists**
- ▶ First Betti number \rightarrow harmonic 1-forms. Nowhere 0? Must **know the metric!**


Background II: Harmonic 1-forms on Calabi-Yaus

- ▶ Let Y be Calabi-Yau 3-fold with **Calabi-Yau metric** g_{CY}
- ▶ $\sigma : Y \rightarrow Y$ anti-holomorphic involution, $L := \text{fix}(\sigma)$
example: quintic with real coefficients in \mathbb{CP}^4 and $\sigma([z_0 : \cdots : z_4]) = [\bar{z}_0 : \cdots : \bar{z}_4]$
- ▶ $S^1 \times Y$ has dimension 7 and holonomy $SU(3)$. Problem: want holonomy G_2
- ▶ Define $\hat{\sigma} : S^1 \times Y \rightarrow S^1 \times Y$ as $(x, y) \mapsto (-x, \sigma(y))$


$(S^1 \times Y)/\langle \hat{\sigma} \rangle$



$\{0, \frac{1}{2}\} \times L$



N^7



Theorem ([Joyce and Karigiannis, 2017])


If there exists $\lambda \in \Omega^1(L)$ **harmonic w.r.t. $g_{CY}|_L$ that is nowhere 0**, then there exists a resolution $N^7 \rightarrow (S^1 \times Y)/\langle \hat{\sigma} \rangle$ with holonomy equal to G_2 .

- ▶ **Goal: check if such a 1-form exists**
- ▶ First Betti number \rightarrow harmonic 1-forms. Nowhere 0? Must **know the metric!**


Background II: Harmonic 1-forms on Calabi-Yaus

- ▶ Let Y be Calabi-Yau 3-fold with **Calabi-Yau metric** g_{CY}
- ▶ $\sigma : Y \rightarrow Y$ anti-holomorphic involution, $L := \text{fix}(\sigma)$
example: quintic with real coefficients in \mathbb{CP}^4 and $\sigma([z_0 : \cdots : z_4]) = [\bar{z}_0 : \cdots : \bar{z}_4]$
- ▶ $S^1 \times Y$ has dimension 7 and holonomy $SU(3)$. Problem: want holonomy G_2
- ▶ Define $\hat{\sigma} : S^1 \times Y \rightarrow S^1 \times Y$ as $(x, y) \mapsto (-x, \sigma(y))$


$$(S^1 \times Y)/\langle \hat{\sigma} \rangle$$



$$\{0, \frac{1}{2}\} \times L$$



$$N^7$$



Theorem ([Joyce and Karigiannis, 2017])


If there exists $\lambda \in \Omega^1(L)$ **harmonic w.r.t. $g_{CY}|_L$ that is nowhere 0**, then there exists a resolution $N^7 \rightarrow (S^1 \times Y)/\langle \hat{\sigma} \rangle$ with holonomy equal to G_2 .

- ▶ **Goal: check if such a 1-form exists**
- ▶ First Betti number \rightarrow harmonic 1-forms. Nowhere 0? Must **know the metric!**


Background II: Harmonic 1-forms on Calabi-Yaus

- ▶ Let Y be Calabi-Yau 3-fold with **Calabi-Yau metric** g_{CY}
- ▶ $\sigma : Y \rightarrow Y$ anti-holomorphic involution, $L := \text{fix}(\sigma)$
example: quintic with real coefficients in \mathbb{CP}^4 and $\sigma([z_0 : \cdots : z_4]) = [\bar{z}_0 : \cdots : \bar{z}_4]$
- ▶ $S^1 \times Y$ has dimension 7 and holonomy $SU(3)$. Problem: want holonomy G_2
- ▶ Define $\hat{\sigma} : S^1 \times Y \rightarrow S^1 \times Y$ as $(x, y) \mapsto (-x, \sigma(y))$


$(S^1 \times Y)/\langle \hat{\sigma} \rangle$



$\{0, \frac{1}{2}\} \times L$



N^7



Theorem ([Joyce and Karigiannis, 2017])


If there exists $\lambda \in \Omega^1(L)$ **harmonic w.r.t. $g_{CY}|_L$ that is nowhere 0**, then there exists a resolution $N^7 \rightarrow (S^1 \times Y)/\langle \hat{\sigma} \rangle$ with holonomy equal to G_2 .

- ▶ **Goal: check if such a 1-form exists**
- ▶ First Betti number \rightarrow harmonic 1-forms. Nowhere 0? Must **know the metric!**


Background II: Harmonic 1-forms on Calabi-Yaus

- ▶ Let Y be Calabi-Yau 3-fold with **Calabi-Yau metric** g_{CY}
- ▶ $\sigma : Y \rightarrow Y$ anti-holomorphic involution, $L := \text{fix}(\sigma)$
example: quintic with real coefficients in \mathbb{CP}^4 and $\sigma([z_0 : \cdots : z_4]) = [\bar{z}_0 : \cdots : \bar{z}_4]$
- ▶ $S^1 \times Y$ has dimension 7 and holonomy $SU(3)$. Problem: want holonomy G_2
- ▶ Define $\hat{\sigma} : S^1 \times Y \rightarrow S^1 \times Y$ as $(x, y) \mapsto (-x, \sigma(y))$


$(S^1 \times Y)/\langle \hat{\sigma} \rangle$



$\{0, \frac{1}{2}\} \times L$



N^7



Theorem ([Joyce and Karigiannis, 2017])

If there exists $\lambda \in \Omega^1(L)$ **harmonic w.r.t. $g_{CY}|_L$ that is nowhere 0**, then there exists a resolution $N^7 \rightarrow (S^1 \times Y)/\langle \hat{\sigma} \rangle$ with holonomy equal to G_2 .

- ▶ **Goal: check if such a 1-form exists**
- ▶ First Betti number \rightarrow harmonic 1-forms. Nowhere 0? Must **know the metric!**

Non-example 1: Fermat Quintic

► $Y := \{z = [z_0 : \cdots : z_4] \in \mathbb{CP}^4 : z_0^5 + \cdots + z_4^5 = 0\}$

► $\sigma([z_0 : \cdots : z_4]) = [\overline{z_0} : \cdots : \overline{z_4}]$



$$\mathbb{RP}^3 \xrightarrow{\cong} L = \text{fix}(\sigma) = \{x = [x_0 : \cdots : x_4] \in \mathbb{RP}^4 : x_0^5 + \cdots + x_4^5 = 0\}$$

$$[x_0 : \cdots : x_4] \mapsto \left[x_0 : \cdots : x_4 : -\sqrt[5]{x_0^5 + \cdots + x_4^5} \right]$$

► $b^1(\mathbb{RP}^3) = 0 \Rightarrow$ no harmonic 1-form on L

Non-example 1: Fermat Quintic

- ▶ $Y := \{z = [z_0 : \cdots : z_4] \in \mathbb{CP}^4 : z_0^5 + \cdots + z_4^5 = 0\}$
- ▶ $\sigma([z_0 : \cdots : z_4]) = [\overline{z_0} : \cdots : \overline{z_4}]$



$$\mathbb{RP}^3 \xrightarrow{\cong} L = \text{fix}(\sigma) = \{x = [x_0 : \cdots : x_4] \in \mathbb{RP}^4 : x_0^5 + \cdots + x_4^5 = 0\}$$

$$[x_0 : \cdots : x_4] \mapsto \left[x_0 : \cdots : x_4 : -\sqrt[5]{x_0^5 + \cdots + x_4^5} \right]$$

- ▶ $b^1(\mathbb{RP}^3) = 0 \Rightarrow$ no harmonic 1-form on L

Non-example 1: Fermat Quintic

- ▶ $Y := \{z = [z_0 : \cdots : z_4] \in \mathbb{CP}^4 : z_0^5 + \cdots + z_4^5 = 0\}$
- ▶ $\sigma([z_0 : \cdots : z_4]) = [\overline{z_0} : \cdots : \overline{z_4}]$
- ▶

$$\mathbb{RP}^3 \xrightarrow{\sim} L = \text{fix}(\sigma) = \{x = [x_0 : \cdots : x_4] \in \mathbb{RP}^4 : x_0^5 + \cdots + x_4^5 = 0\}$$

$$[x_0 : \cdots : x_4] \mapsto \left[x_0 : \cdots : x_4 : -\sqrt[5]{x_0^5 + \cdots + x_4^5} \right]$$

- ▶ $b^1(\mathbb{RP}^3) = 0 \Rightarrow$ no harmonic 1-form on L

Non-example 1: Fermat Quintic

- ▶ $Y := \{z = [z_0 : \cdots : z_4] \in \mathbb{CP}^4 : z_0^5 + \cdots + z_4^5 = 0\}$
- ▶ $\sigma([z_0 : \cdots : z_4]) = [\overline{z_0} : \cdots : \overline{z_4}]$
- ▶

$$\mathbb{RP}^3 \xrightarrow{\cong} L = \text{fix}(\sigma) = \{x = [x_0 : \cdots : x_4] \in \mathbb{RP}^4 : x_0^5 + \cdots + x_4^5 = 0\}$$


$$[x_0 : \cdots : x_4] \mapsto \left[x_0 : \cdots : x_4 : -\sqrt[5]{x_0^5 + \cdots + x_4^5} \right]$$

- ▶ $b^1(\mathbb{RP}^3) = 0 \Rightarrow$ no harmonic 1-form on L

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1:0:0:0:0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma: \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)




- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fano?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)




- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fano?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)

$\text{sing}(g)$ $\text{fix}(\sigma)$




- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fano's?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)

$\text{sing}(g)$ $\text{fix}(\sigma)$




- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fano's?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)

$\text{sing}(g)$ $\text{fix}(\sigma)$





- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fano?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)

$\text{sing}(g)$ $\text{fix}(\sigma)$




- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fano's?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)

$\text{sing}(g)$ $\text{fix}(\sigma)$




- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fano's?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)

$\text{sing}(g)$ $\text{fix}(\sigma)$





- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fano?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)

$\text{sing}(g)$ $\text{fix}(\sigma)$




- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fanos?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)

$\text{sing}(g)$ $\text{fix}(\sigma)$




- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fanos?

Non-Example 2: small complex structure limit

► Another quintic in \mathbb{CP}^4 :

1. [Krasnov, 2009] $f_{\pm} = (x_0(x_1^2 + x_2^2 + x_3^2 - x_4^2) - (x_1^3 + x_2^3 + x_3^3 - \frac{1}{2}x_4^3) \pm \epsilon x_0^3)$
smoothing of **ordinary double point** $(1 : 0 : 0 : 0 : 0)$
Has $Z(f_+) \cong \mathbb{RP}^3$, $Z(f_-) \cong \mathbb{RP}^3 \# S^1 \times S^2$
2. $v := (x_0^2 + \dots + x_4^2)$ and $g = v \cdot f_-$ has $Z_{\mathbb{R}}(g) = Z_{\mathbb{R}}(f_-) \subset \mathbb{RP}^4$ so
 $\sigma : \mathbb{CP}^4 \rightarrow \mathbb{CP}^4$, $x \mapsto \bar{x}$ has $b^1(\text{fix}(\sigma)|_{Z(g)}) = 1$
3. Take smoothing $g_{\epsilon} := g + \epsilon \xi$, where ξ generic poly
4. But: ex. incompressible $S^2 \Rightarrow$ [Jaco, 1980] $Z_{\mathbb{R}}(g_{\epsilon}) \cong Z_{\mathbb{R}}(g)$ is **no fibration over S^1**
 \Rightarrow [Tischler, 1970] **no closed nowhere zero 1-form** \Rightarrow any harmonic 1-form has zeros
(even number of zeros by Poincaré–Hopf theorem and $\chi(Z_{\mathbb{R}}(g_{\epsilon})) = 0$)

$\text{sing}(g)$ $\text{fix}(\sigma)$



- [Tian and Yau, 1990] Calabi-Yau metrics on $Z(f_-) \setminus \text{sing}(g)$ and $Z(v) \setminus \text{sing}(g)$
- [Sun and Zhang, 2019] glue these to metric on smoothing $Z(g_{\epsilon})$
- More non-examples from other cubics. Examples from other Fanos?

Conjectural example 3: quadric intersect quartic

► Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. Circle $c_{aff} = x_1^2 + x_2^2 - 1$, quartic $q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$

Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$



2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$

$\text{sing}(Z(c, q_\epsilon))$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Conjectural example 3: quadric intersect quartic

► Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. **Circle $c_{aff} = x_1^2 + x_2^2 - 1$** , quartic **$q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$**

Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$



2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$

$\text{sing}(Z(c, q_\epsilon))$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Conjectural example 3: quadric intersect quartic

- Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. **Circle $c_{aff} = x_1^2 + x_2^2 - 1$** , **quartic $q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$**

Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$



2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$

$\text{sing}(Z(c, q_\epsilon))$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Conjectural example 3: quadric intersect quartic

- Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. **Circle $c_{aff} = x_1^2 + x_2^2 - 1$** , **quartic $q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$**

Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$



2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$

$\text{sing}(Z(c, q_\epsilon))$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Conjectural example 3: quadric intersect quartic

- Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. **Circle $c_{aff} = x_1^2 + x_2^2 - 1$** , **quartic $q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$**



Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$

2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Conjectural example 3: quadric intersect quartic

- Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. **Circle $c_{aff} = x_1^2 + x_2^2 - 1$** , **quartic $q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$**



Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$

2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Conjectural example 3: quadric intersect quartic

- Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. Circle $c_{aff} = x_1^2 + x_2^2 - 1$, quartic $q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$

Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$



2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$

$\text{sing}(Z(c, q_\epsilon))$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Conjectural example 3: quadric intersect quartic

- Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. Circle $c_{aff} = x_1^2 + x_2^2 - 1$, quartic $q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$

Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$


2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$

$\text{sing}(Z(c, q_\epsilon))$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Conjectural example 3: quadric intersect quartic

- Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. Circle $c_{aff} = x_1^2 + x_2^2 - 1$, quartic $q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$

Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$



2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$

$\text{sing}(Z(c, q_\epsilon))$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Conjectural example 3: quadric intersect quartic

- Construction of **quadric intersect quartic in \mathbb{CP}^5** , also Calabi-Yau

1. Circle $c_{aff} = x_1^2 + x_2^2 - 1$, quartic $q_{aff} = x_3^4 + x_4^4 + x_5^4 - 1$

Projectivise: $c = -x_0^2 + x_1^2 + x_2^2$ and $q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$


2. c and q have **$SO(2)$ -symmetry**:

$$[x_0 : x_1 : x_2 : x_3 : x_4 : x_5] \mapsto [x_0 : \cos(t)x_1 - \sin(t)x_2 : \sin(t)x_1 + \cos(t)x_2 : x_3 : x_4 : x_5]$$

Generic smoothings c_δ and q_ϵ of c and q

3. $\mathbb{RP}^5 \supset Z_{\mathbb{R}}(c, q_\epsilon) \cong S^1 \times S^2$ smooth, $Z(c, q_\epsilon) \subset \mathbb{CP}^5$ singular
 $\text{sing}(Z(c, q_\epsilon)) = \text{sing}(c) \cap Z(q_\epsilon)$

$\text{sing}(Z(c, q_\epsilon))$



- Fantasy: $Z(c, q_\epsilon)$ has singular Calabi-Yau metric $g_0 \Rightarrow$ if q_ϵ is $SO(2)$ -invariant, then **Infinitesimal $SO(2)$ -action** gives Killing field for $g_0 \Rightarrow$ (Ricci-flat) parallel vector field \Rightarrow parallel 1-form for $g_0 \Rightarrow$ nearly parallel 1-form for g_δ

Numerical Calabi-Yau metrics

- ▶ Holomorphic volume form locally $\Omega = dz^1 \wedge dz^2 \wedge dz^3 \rightsquigarrow \text{vol}_\Omega := \Omega \wedge \overline{\Omega} \in \Omega^6(Y)$
- ▶ Ample line bundle $L \rightarrow Y$ and $k \in \mathbb{N}$ such that $L^{\otimes k}$ very ample
Example: $Y \subset \mathbb{CP}^4$ quintic, $(\mathcal{O}(1)|_Y)^{\otimes k}$
- ▶ $s_1, \dots, s_N \in H^0(L^{\otimes k})$ basis of holomorphic sections
 \Rightarrow embedding $s = (s_1, \dots, s_N) : Y \rightarrow \mathbb{CP}^{N-1}$
- ▶ h positive definite Hermitian form on $H^0(L^{\otimes k}) \rightsquigarrow$ some Fubini-Study metric
Kähler potential: $K = \log \sum_{i,j} h_{i,j} s^i \bar{s}^j$. Volume form: $\omega_h^3 = \text{vol}_h \in \Omega^6(Y)$.

If $\frac{\text{vol}_h}{\text{vol}_\Omega} = 1$, then Ricci-flat

- ▶ [Donaldson, 2009]: choose h cleverly to minimise $\int_Y \left(\frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right)^2$

	$\left \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right $	Comment
[Donaldson, 2009]	10^{-2}	$n = 2$, needs symmetries
[Headrick and Nassar, 2013]	10^{-14}	$n = 3$, needs symmetries
[Larfors et al., 2022]	10^{-2}	$n = 3$, not C^0 , complete intersections+torics
[Douglas et al., 2022]+ours	10^{-4}	$n = 3$, quintics+complete intersections

Numerical Calabi-Yau metrics

- ▶ Holomorphic volume form locally $\Omega = dz^1 \wedge dz^2 \wedge dz^3 \rightsquigarrow \text{vol}_\Omega := \Omega \wedge \overline{\Omega} \in \Omega^6(Y)$
- ▶ Ample **line bundle** $L \rightarrow Y$ and $k \in \mathbb{N}$ such that $L^{\otimes k}$ very ample
Example: $Y \subset \mathbb{CP}^4$ quintic, $(\mathcal{O}(1)|_Y)^{\otimes k}$
- ▶ $s_1, \dots, s_N \in H^0(L^{\otimes k})$ basis of holomorphic sections
 \Rightarrow embedding $s = (s_1, \dots, s_N) : Y \rightarrow \mathbb{CP}^{N-1}$
- ▶ h positive definite Hermitian form on $H^0(L^{\otimes k}) \rightsquigarrow$ some Fubini-Study metric
Kähler potential: $K = \log \sum_{i,j} h_{i,j} s^i \bar{s}^j$. Volume form: $\omega_h^3 = \text{vol}_h \in \Omega^6(Y)$.

If $\frac{\text{vol}_h}{\text{vol}_\Omega} = 1$, then Ricci-flat

- ▶ [Donaldson, 2009]: choose h cleverly to minimise $\int_Y \left(\frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right)^2$

	$\left \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right $	Comment
[Donaldson, 2009]	10^{-2}	$n = 2$, needs symmetries
[Headrick and Nassar, 2013]	10^{-14}	$n = 3$, needs symmetries
[Larfors et al., 2022]	10^{-2}	$n = 3$, not C^0 , complete intersections+torics
[Douglas et al., 2022]+ours	10^{-4}	$n = 3$, quintics+complete intersections

Numerical Calabi-Yau metrics

- ▶ Holomorphic volume form locally $\Omega = dz^1 \wedge dz^2 \wedge dz^3 \rightsquigarrow \text{vol}_\Omega := \Omega \wedge \overline{\Omega} \in \Omega^6(Y)$
- ▶ Ample **line bundle** $L \rightarrow Y$ and $k \in \mathbb{N}$ such that $L^{\otimes k}$ very ample
Example: $Y \subset \mathbb{CP}^4$ quintic, $(\mathcal{O}(1)|_Y)^{\otimes k}$
- ▶ $s_1, \dots, s_N \in H^0(L^{\otimes k})$ basis of holomorphic sections
 \Rightarrow embedding $s = (s_1, \dots, s_N) : Y \rightarrow \mathbb{CP}^{N-1}$
- ▶ h positive definite Hermitian form on $H^0(L^{\otimes k}) \rightsquigarrow$ some Fubini-Study metric
Kähler potential: $K = \log \sum_{i,j} h_{i,j} s^i \bar{s}^j$. Volume form: $\omega_h^3 = \text{vol}_h \in \Omega^6(Y)$.

If $\frac{\text{vol}_h}{\text{vol}_\Omega} = 1$, then Ricci-flat

- ▶ [Donaldson, 2009]: choose h cleverly to minimise $\int_Y \left(\frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right)^2$

	$\left \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right $	Comment
[Donaldson, 2009]	10^{-2}	$n = 2$, needs symmetries
[Headrick and Nassar, 2013]	10^{-14}	$n = 3$, needs symmetries
[Larfors et al., 2022]	10^{-2}	$n = 3$, not C^0 , complete intersections+torics
[Douglas et al., 2022] + ours	10^{-4}	$n = 3$, quintics + complete intersections

Numerical Calabi-Yau metrics

- ▶ Holomorphic volume form locally $\Omega = dz^1 \wedge dz^2 \wedge dz^3 \rightsquigarrow \text{vol}_\Omega := \Omega \wedge \overline{\Omega} \in \Omega^6(Y)$
- ▶ Ample **line bundle** $L \rightarrow Y$ and $k \in \mathbb{N}$ such that $L^{\otimes k}$ very ample
Example: $Y \subset \mathbb{CP}^4$ quintic, $(\mathcal{O}(1)|_Y)^{\otimes k}$
- ▶ $s_1, \dots, s_N \in H^0(L^{\otimes k})$ basis of holomorphic sections
 \Rightarrow embedding $s = (s_1, \dots, s_N) : Y \rightarrow \mathbb{CP}^{N-1}$
- ▶ h positive definite Hermitian form on $H^0(L^{\otimes k}) \rightsquigarrow$ some Fubini-Study metric
Kähler potential: $K = \log \sum_{i,j} h_{i,j} s^i \bar{s}^j$. Volume form: $\omega_h^3 = \text{vol}_h \in \Omega^6(Y)$.

If $\frac{\text{vol}_h}{\text{vol}_\Omega} = 1$, then Ricci-flat

- ▶ [Donaldson, 2009]: choose h cleverly to minimise $\int_Y \left(\frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right)^2$

	$\left \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right $	Comment
[Donaldson, 2009]	10^{-2}	$n = 2$, needs symmetries
[Headrick and Nassar, 2013]	10^{-14}	$n = 3$, needs symmetries
[Larfors et al., 2022]	10^{-2}	$n = 3$, not C^0 , complete intersections+torics
[Douglas et al., 2022]+ours	10^{-4}	$n = 3$, quintics+complete intersections

Numerical Calabi-Yau metrics

- ▶ Holomorphic volume form locally $\Omega = dz^1 \wedge dz^2 \wedge dz^3 \rightsquigarrow \text{vol}_\Omega := \Omega \wedge \bar{\Omega} \in \Omega^6(Y)$
- ▶ Ample **line bundle** $L \rightarrow Y$ and $k \in \mathbb{N}$ such that $L^{\otimes k}$ very ample
Example: $Y \subset \mathbb{CP}^4$ quintic, $(\mathcal{O}(1)|_Y)^{\otimes k}$
- ▶ $s_1, \dots, s_N \in H^0(L^{\otimes k})$ basis of holomorphic sections
 \Rightarrow embedding $s = (s_1, \dots, s_N) : Y \rightarrow \mathbb{CP}^{N-1}$
- ▶ h positive definite Hermitian form on $H^0(L^{\otimes k}) \rightsquigarrow$ some Fubini-Study metric
Kähler potential: $K = \log \sum_{i,j} h_{i,j} s^i \bar{s}^j$. Volume form: $\omega_h^3 = \text{vol}_h \in \Omega^6(Y)$.

If $\frac{\text{vol}_h}{\text{vol}_\Omega} = 1$, then Ricci-flat

- ▶ [Donaldson, 2009]: choose h cleverly to minimise $\int_Y \left(\frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right)^2$

	$\left \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right $	Comment
[Donaldson, 2009]	10^{-2}	$n = 2$, needs symmetries
[Headrick and Nassar, 2013]	10^{-14}	$n = 3$, needs symmetries
[Larfors et al., 2022]	10^{-2}	$n = 3$, not C^0 , complete intersections+torics
[Douglas et al., 2022] + ours	10^{-4}	$n = 3$, quintics + complete intersections

Numerical Calabi-Yau metrics

- ▶ Holomorphic volume form locally $\Omega = dz^1 \wedge dz^2 \wedge dz^3 \rightsquigarrow \text{vol}_\Omega := \Omega \wedge \bar{\Omega} \in \Omega^6(Y)$
- ▶ Ample **line bundle** $L \rightarrow Y$ and $k \in \mathbb{N}$ such that $L^{\otimes k}$ very ample
Example: $Y \subset \mathbb{CP}^4$ quintic, $(\mathcal{O}(1)|_Y)^{\otimes k}$
- ▶ $s_1, \dots, s_N \in H^0(L^{\otimes k})$ basis of holomorphic sections
 \Rightarrow embedding $s = (s_1, \dots, s_N) : Y \rightarrow \mathbb{CP}^{N-1}$
- ▶ h positive definite Hermitian form on $H^0(L^{\otimes k}) \rightsquigarrow$ some Fubini-Study metric
Kähler potential: $K = \log \sum_{i,j} h_{i,j} s^i \bar{s}^j$. Volume form: $\omega_h^3 = \text{vol}_h \in \Omega^6(Y)$.

If $\frac{\text{vol}_h}{\text{vol}_\Omega} = 1$, then Ricci-flat

- ▶ [Donaldson, 2009]: choose h cleverly to minimise $\int_Y \left(\frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right)^2$

	$\left \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right $	Comment
[Donaldson, 2009]	10^{-2}	$n = 2$, needs symmetries
[Headrick and Nassar, 2013]	10^{-14}	$n = 3$, needs symmetries
[Larfors et al., 2022]	10^{-2}	$n = 3$, not C^0 , complete intersections+torics
[Douglas et al., 2022] + ours	10^{-4}	$n = 3$, quintics + complete intersections

Numerical Calabi-Yau metrics

- ▶ Holomorphic volume form locally $\Omega = dz^1 \wedge dz^2 \wedge dz^3 \rightsquigarrow \text{vol}_\Omega := \Omega \wedge \bar{\Omega} \in \Omega^6(Y)$
- ▶ Ample **line bundle** $L \rightarrow Y$ and $k \in \mathbb{N}$ such that $L^{\otimes k}$ very ample
Example: $Y \subset \mathbb{CP}^4$ quintic, $(\mathcal{O}(1)|_Y)^{\otimes k}$
- ▶ $s_1, \dots, s_N \in H^0(L^{\otimes k})$ basis of holomorphic sections
 \Rightarrow embedding $s = (s_1, \dots, s_N) : Y \rightarrow \mathbb{CP}^{N-1}$
- ▶ h positive definite Hermitian form on $H^0(L^{\otimes k}) \rightsquigarrow$ some Fubini-Study metric
Kähler potential: $K = \log \sum_{i,j} h_{i,j} s^i \bar{s}^j$. Volume form: $\omega_h^3 = \text{vol}_h \in \Omega^6(Y)$.

If $\frac{\text{vol}_h}{\text{vol}_\Omega} = 1$, then Ricci-flat

- ▶ [Donaldson, 2009]: choose h cleverly to minimise $\int_Y \left(\frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right)^2$

	$\left \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right $	Comment
[Donaldson, 2009]	10^{-2}	$n = 2$, needs symmetries
[Headrick and Nassar, 2013]	10^{-14}	$n = 3$, needs symmetries
[Larfors et al., 2022]	10^{-2}	$n = 3$, not C^0 , complete intersections+torics
[Douglas et al., 2022]+ours	10^{-4}	$n = 3$, quintics+complete intersections

Numerical Calabi-Yau metrics

- ▶ Holomorphic volume form locally $\Omega = dz^1 \wedge dz^2 \wedge dz^3 \rightsquigarrow \text{vol}_\Omega := \Omega \wedge \bar{\Omega} \in \Omega^6(Y)$
- ▶ Ample **line bundle** $L \rightarrow Y$ and $k \in \mathbb{N}$ such that $L^{\otimes k}$ very ample
Example: $Y \subset \mathbb{CP}^4$ quintic, $(\mathcal{O}(1)|_Y)^{\otimes k}$
- ▶ $s_1, \dots, s_N \in H^0(L^{\otimes k})$ basis of holomorphic sections
 \Rightarrow embedding $s = (s_1, \dots, s_N) : Y \rightarrow \mathbb{CP}^{N-1}$
- ▶ h positive definite Hermitian form on $H^0(L^{\otimes k}) \rightsquigarrow$ some Fubini-Study metric
Kähler potential: $K = \log \sum_{i,j} h_{i,j} s^i \bar{s}^j$. Volume form: $\omega_h^3 = \text{vol}_h \in \Omega^6(Y)$.

If $\frac{\text{vol}_h}{\text{vol}_\Omega} = 1$, then Ricci-flat

- ▶ [Donaldson, 2009]: choose h cleverly to minimise $\int_Y \left(\frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right)^2$

	$\left\ \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right\ $	Comment
[Donaldson, 2009]	10^{-2}	$n = 2$, needs symmetries
[Headrick and Nassar, 2013]	10^{-14}	$n = 3$, needs symmetries
[Larfors et al., 2022]	10^{-2}	$n = 3$, not C^0 , complete intersections+torics
[Douglas et al., 2022] +ours	10^{-4}	$n = 3$, quintics+complete intersections

Numerical harmonic 1-forms

Example: quintic $X := Z(f) \subset \mathbb{CP}^4$, $L := \text{fix}(\sigma) \subset X$

- ▶ $\xi_1, \dots, \xi_{10} \in \Omega^1(\mathbb{RP}^4)$ closed 1-forms s.t. $T_x^* \mathbb{RP}^4 = (\xi_1(x), \dots, \xi_{10}(x)) \forall x$
- ▶ p_1, \dots, p_N polys, p_i degree d_i ; for $\alpha = (\alpha_{1,1}, \dots, \alpha_{10,N}) \in \mathbb{R}^{10N}$ let

$$\lambda_\alpha(x) = \sum_{\substack{i=1, \dots, N \\ j=1, \dots, 10}} \alpha_{i,j} \frac{p_i(x)}{|x|^{d_i}} \xi_j|_L(x) \in \Omega^1(L)$$

For $x_1, \dots, x_{100000} \in X$ find $\min_{\alpha \text{ s.t. } \|\lambda_\alpha\|_{L^2}=1} \int_{x_1, \dots, x_{100000}} |d\lambda_\alpha| + |d^*\lambda_\alpha|$

- ▶ Stone-Weierstrass \Rightarrow best approximations converge to harmonic form as $N \rightarrow \infty$
- ▶ Ansatz for p_i : $A_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{n_i+1}$ linear, $\text{sq} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ square each coordinate

$$p(x_0, \dots, x_4) = A_k \circ \dots \circ \text{sq} \circ A_2 \circ \text{sq} \circ A_1(x_0, \dots, x_4)$$

- ▶ Equivalent: neural network with activation function $x \mapsto x^2$
- ▶ Approximate metric $\frac{i}{2} \partial \bar{\partial} K$ smooth+explicit \Rightarrow explicitly compute $(|d\lambda_\alpha|(x_i) + |d^*\lambda_\alpha|(x_i)) / \sqrt{|\lambda(x_1)|^2 + \dots + |\lambda(x_{100000})|^2}$
 \Rightarrow minimise with tensorflow

Numerical harmonic 1-forms

Example: quintic $X := Z(f) \subset \mathbb{CP}^4$, $L := \text{fix}(\sigma) \subset X$

- ▶ $\xi_1, \dots, \xi_{10} \in \Omega^1(\mathbb{RP}^4)$ closed 1-forms s.t. $T_x^* \mathbb{RP}^4 = (\xi_1(x), \dots, \xi_{10}(x)) \forall x$
- ▶ p_1, \dots, p_N polys, p_i degree d_i ; for $\alpha = (\alpha_{1,1}, \dots, \alpha_{10,N}) \in \mathbb{R}^{10N}$ let

$$\lambda_\alpha(x) = \sum_{\substack{i=1, \dots, N \\ j=1, \dots, 10}} \alpha_{i,j} \frac{p_i(x)}{|x|^{d_i}} \xi_j|_L(x) \in \Omega^1(L)$$

For $x_1, \dots, x_{100000} \in X$ find $\min_{\alpha \text{ s.t. } \|\lambda_\alpha\|_{L^2}=1} \int_{x_1, \dots, x_{100000}} |d\lambda_\alpha| + |d^*\lambda_\alpha|$

- ▶ Stone-Weierstrass \Rightarrow best approximations converge to harmonic form as $N \rightarrow \infty$
- ▶ Ansatz for p_i : $A_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{n_i+1}$ linear, $\text{sq} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ square each coordinate

$$p(x_0, \dots, x_4) = A_k \circ \dots \circ \text{sq} \circ A_2 \circ \text{sq} \circ A_1(x_0, \dots, x_4)$$

- ▶ Equivalent: neural network with activation function $x \mapsto x^2$
- ▶ Approximate metric $\frac{i}{2} \partial \bar{\partial} K$ smooth+explicit \Rightarrow explicitly compute $(|d\lambda_\alpha|(x_i) + |d^*\lambda_\alpha|(x_i)) / \sqrt{|\lambda(x_1)|^2 + \dots + |\lambda(x_{100000})|^2}$
 \Rightarrow minimise with tensorflow

Numerical harmonic 1-forms

Example: quintic $X := Z(f) \subset \mathbb{CP}^4$, $L := \text{fix}(\sigma) \subset X$

- ▶ $\xi_1, \dots, \xi_{10} \in \Omega^1(\mathbb{RP}^4)$ closed 1-forms s.t. $T_x^* \mathbb{RP}^4 = (\xi_1(x), \dots, \xi_{10}(x)) \forall x$
- ▶ p_1, \dots, p_N polys, p_i degree d_i ; for $\alpha = (\alpha_{1,1}, \dots, \alpha_{10,N}) \in \mathbb{R}^{10N}$ let

$$\lambda_\alpha(x) = \sum_{\substack{i=1, \dots, N \\ j=1, \dots, 10}} \alpha_{i,j} \frac{p_i(x)}{|x|^{d_i}} \xi_j|_L(x) \in \Omega^1(L)$$

For $x_1, \dots, x_{100000} \in X$ find $\min_{\alpha \text{ s.t. } \|\lambda_\alpha\|_{L^2}=1} \int_{x_1, \dots, x_{100000}} |d\lambda_\alpha| + |d^*\lambda_\alpha|$

- ▶ Stone-Weierstrass \Rightarrow best approximations converge to harmonic form as $N \rightarrow \infty$
- ▶ Ansatz for p_i : $A_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{n_i+1}$ linear, $\text{sq} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ square each coordinate

$$p(x_0, \dots, x_4) = A_k \circ \dots \circ \text{sq} \circ A_2 \circ \text{sq} \circ A_1(x_0, \dots, x_4)$$

- ▶ Equivalent: neural network with activation function $x \mapsto x^2$
- ▶ Approximate metric $\frac{i}{2} \partial \bar{\partial} K$ smooth+explicit \Rightarrow explicitly compute $(|d\lambda_\alpha|(x_i) + |d^*\lambda_\alpha|(x_i)) / \sqrt{|\lambda(x_1)|^2 + \dots + |\lambda(x_{100000})|^2}$
 \Rightarrow minimise with tensorflow

Numerical harmonic 1-forms

Example: quintic $X := Z(f) \subset \mathbb{CP}^4$, $L := \text{fix}(\sigma) \subset X$

- ▶ $\xi_1, \dots, \xi_{10} \in \Omega^1(\mathbb{RP}^4)$ closed 1-forms s.t. $T_x^* \mathbb{RP}^4 = (\xi_1(x), \dots, \xi_{10}(x)) \forall x$
- ▶ p_1, \dots, p_N polys, p_i degree d_i ; for $\alpha = (\alpha_{1,1}, \dots, \alpha_{10,N}) \in \mathbb{R}^{10N}$ let

$$\lambda_\alpha(x) = \sum_{\substack{i=1, \dots, N \\ j=1, \dots, 10}} \alpha_{i,j} \frac{p_i(x)}{|x|^{d_i}} \xi_j|_L(x) \in \Omega^1(L)$$

For $x_1, \dots, x_{100000} \in X$ find $\min_{\alpha \text{ s.t. } \|\lambda_\alpha\|_{L^2}=1} \int_{x_1, \dots, x_{100000}} |d\lambda_\alpha| + |d^*\lambda_\alpha|$

- ▶ Stone-Weierstrass \Rightarrow best approximations converge to harmonic form as $N \rightarrow \infty$
- ▶ Ansatz for p_i : $A_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{n_i+1}$ linear, $\text{sq} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ square each coordinate

$$p(x_0, \dots, x_4) = A_k \circ \dots \circ \text{sq} \circ A_2 \circ \text{sq} \circ A_1(x_0, \dots, x_4)$$

- ▶ Equivalent: neural network with activation function $x \mapsto x^2$
- ▶ Approximate metric $\frac{i}{2} \partial \bar{\partial} K$ smooth+explicit \Rightarrow explicitly compute $(|d\lambda_\alpha|(x_i) + |d^*\lambda_\alpha|(x_i)) / \sqrt{|\lambda(x_1)|^2 + \dots + |\lambda(x_{100000})|^2}$
 \Rightarrow minimise with tensorflow

Numerical harmonic 1-forms

Example: quintic $X := Z(f) \subset \mathbb{CP}^4$, $L := \text{fix}(\sigma) \subset X$

- ▶ $\xi_1, \dots, \xi_{10} \in \Omega^1(\mathbb{RP}^4)$ closed 1-forms s.t. $T_x^* \mathbb{RP}^4 = (\xi_1(x), \dots, \xi_{10}(x)) \forall x$
- ▶ p_1, \dots, p_N polys, p_i degree d_i ; for $\alpha = (\alpha_{1,1}, \dots, \alpha_{10,N}) \in \mathbb{R}^{10N}$ let

$$\lambda_\alpha(x) = \sum_{\substack{i=1, \dots, N \\ j=1, \dots, 10}} \alpha_{i,j} \frac{p_i(x)}{|x|^{d_i}} \xi_j|_L(x) \in \Omega^1(L)$$

For $x_1, \dots, x_{100000} \in X$ find $\min_{\alpha \text{ s.t. } \|\lambda_\alpha\|_{L^2}=1} \int_{x_1, \dots, x_{100000}} |d\lambda_\alpha| + |d^*\lambda_\alpha|$

- ▶ Stone-Weierstrass \Rightarrow best approximations converge to harmonic form as $N \rightarrow \infty$
- ▶ Ansatz for p_i : $A_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{n_i+1}$ linear, $\text{sq} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ square each coordinate

$$p(x_0, \dots, x_4) = A_k \circ \dots \circ \text{sq} \circ A_2 \circ \text{sq} \circ A_1(x_0, \dots, x_4)$$

- ▶ Equivalent: neural network with activation function $x \mapsto x^2$
- ▶ Approximate metric $\frac{i}{2} \partial \bar{\partial} K$ smooth+explicit \Rightarrow explicitly compute $(|d\lambda_\alpha|(x_i) + |d^*\lambda_\alpha|(x_i)) / \sqrt{|\lambda(x_1)|^2 + \dots + |\lambda(x_{100000})|^2}$
 \Rightarrow minimise with tensorflow

Numerical harmonic 1-forms

Example: quintic $X := Z(f) \subset \mathbb{CP}^4$, $L := \text{fix}(\sigma) \subset X$

- ▶ $\xi_1, \dots, \xi_{10} \in \Omega^1(\mathbb{RP}^4)$ closed 1-forms s.t. $T_x^* \mathbb{RP}^4 = (\xi_1(x), \dots, \xi_{10}(x)) \forall x$
- ▶ p_1, \dots, p_N polys, p_i degree d_i ; for $\alpha = (\alpha_{1,1}, \dots, \alpha_{10,N}) \in \mathbb{R}^{10N}$ let

$$\lambda_\alpha(x) = \sum_{\substack{i=1, \dots, N \\ j=1, \dots, 10}} \alpha_{i,j} \frac{p_i(x)}{|x|^{d_i}} \xi_j|_L(x) \in \Omega^1(L)$$

For $x_1, \dots, x_{100000} \in X$ find $\min_{\alpha \text{ s.t. } \|\lambda_\alpha\|_{L^2}=1} \int_{x_1, \dots, x_{100000}} |d\lambda_\alpha| + |d^*\lambda_\alpha|$

- ▶ Stone-Weierstrass \Rightarrow best approximations converge to harmonic form as $N \rightarrow \infty$
- ▶ Ansatz for p_i : $A_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{n_i+1}$ linear, $\text{sq} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ square each coordinate

$$p(x_0, \dots, x_4) = A_k \circ \dots \circ \text{sq} \circ A_2 \circ \text{sq} \circ A_1(x_0, \dots, x_4)$$

- ▶ Equivalent: neural network with activation function $x \mapsto x^2$
- ▶ Approximate metric $\frac{i}{2} \partial \bar{\partial} K$ smooth+explicit \Rightarrow explicitly compute $(|d\lambda_\alpha|(x_i) + |d^*\lambda_\alpha|(x_i)) / \sqrt{|\lambda(x_1)|^2 + \dots + |\lambda(x_{100000})|^2}$
 \Rightarrow minimise with tensorflow

Numerical harmonic 1-forms

Example: quintic $X := Z(f) \subset \mathbb{CP}^4$, $L := \text{fix}(\sigma) \subset X$

- ▶ $\xi_1, \dots, \xi_{10} \in \Omega^1(\mathbb{RP}^4)$ closed 1-forms s.t. $T_x^* \mathbb{RP}^4 = (\xi_1(x), \dots, \xi_{10}(x)) \forall x$
- ▶ p_1, \dots, p_N polys, p_i degree d_i ; for $\alpha = (\alpha_{1,1}, \dots, \alpha_{10,N}) \in \mathbb{R}^{10N}$ let

$$\lambda_\alpha(x) = \sum_{\substack{i=1, \dots, N \\ j=1, \dots, 10}} \alpha_{i,j} \frac{p_i(x)}{|x|^{d_i}} \xi_j|_L(x) \in \Omega^1(L)$$

For $x_1, \dots, x_{100000} \in X$ find $\min_{\alpha \text{ s.t. } \|\lambda_\alpha\|_{L^2}=1} \int_{x_1, \dots, x_{100000}} |d\lambda_\alpha| + |d^*\lambda_\alpha|$

- ▶ Stone-Weierstrass \Rightarrow best approximations converge to harmonic form as $N \rightarrow \infty$
- ▶ Ansatz for p_i : $A_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{n_i+1}$ linear, $\text{sq} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ square each coordinate

$$p(x_0, \dots, x_4) = A_k \circ \dots \circ \text{sq} \circ A_2 \circ \text{sq} \circ A_1(x_0, \dots, x_4)$$

- ▶ Equivalent: neural network with activation function $x \mapsto x^2$
- ▶ Approximate metric $\frac{i}{2} \partial \bar{\partial} K$ smooth+explicit \Rightarrow explicitly compute $(|d\lambda_\alpha|(x_i) + |d^*\lambda_\alpha|(x_i)) / \sqrt{|\lambda(x_1)|^2 + \dots + |\lambda(x_{100000})|^2}$
 \Rightarrow minimise with tensorflow

Numerical harmonic 1-forms

Example: quintic $X := Z(f) \subset \mathbb{CP}^4$, $L := \text{fix}(\sigma) \subset X$

- ▶ $\xi_1, \dots, \xi_{10} \in \Omega^1(\mathbb{RP}^4)$ closed 1-forms s.t. $T_x^* \mathbb{RP}^4 = (\xi_1(x), \dots, \xi_{10}(x)) \forall x$
- ▶ p_1, \dots, p_N polys, p_i degree d_i ; for $\alpha = (\alpha_{1,1}, \dots, \alpha_{10,N}) \in \mathbb{R}^{10N}$ let

$$\lambda_\alpha(x) = \sum_{\substack{i=1, \dots, N \\ j=1, \dots, 10}} \alpha_{i,j} \frac{p_i(x)}{|x|^{d_i}} \xi_j|_L(x) \in \Omega^1(L)$$

For $x_1, \dots, x_{100000} \in X$ find $\min_{\alpha \text{ s.t. } \|\lambda_\alpha\|_{L^2}=1} \int_{x_1, \dots, x_{100000}} |d\lambda_\alpha| + |d^* \lambda_\alpha|$

- ▶ Stone-Weierstrass \Rightarrow best approximations converge to harmonic form as $N \rightarrow \infty$
- ▶ Ansatz for p_i : $A_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{n_{i+1}}$ linear, $\text{sq} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ square each coordinate

$$p(x_0, \dots, x_4) = A_k \circ \dots \circ \text{sq} \circ A_2 \circ \text{sq} \circ A_1(x_0, \dots, x_4)$$

- ▶ Equivalent: neural network with activation function $x \mapsto x^2$
- ▶ Approximate metric $\frac{i}{2} \partial \bar{\partial} K$ smooth+explicit \Rightarrow explicitly compute $(|d\lambda_\alpha|(x_i) + |d^* \lambda_\alpha|(x_i)) / \sqrt{|\lambda(x_1)|^2 + \dots + |\lambda(x_{100000})|^2}$
 \Rightarrow minimise with tensorflow

Numerical harmonic 1-forms

Example: quintic $X := Z(f) \subset \mathbb{CP}^4$, $L := \text{fix}(\sigma) \subset X$

- ▶ $\xi_1, \dots, \xi_{10} \in \Omega^1(\mathbb{RP}^4)$ closed 1-forms s.t. $T_x^* \mathbb{RP}^4 = (\xi_1(x), \dots, \xi_{10}(x)) \forall x$
- ▶ p_1, \dots, p_N polys, p_i degree d_i ; for $\alpha = (\alpha_{1,1}, \dots, \alpha_{10,N}) \in \mathbb{R}^{10N}$ let

$$\lambda_\alpha(x) = \sum_{\substack{i=1, \dots, N \\ j=1, \dots, 10}} \alpha_{i,j} \frac{p_i(x)}{|x|^{d_i}} \xi_j|_L(x) \in \Omega^1(L)$$

For $x_1, \dots, x_{100000} \in X$ find $\min_{\alpha \text{ s.t. } \|\lambda_\alpha\|_{L^2}=1} \int_{x_1, \dots, x_{100000}} |d\lambda_\alpha| + |d^* \lambda_\alpha|$

- ▶ Stone-Weierstrass \Rightarrow best approximations converge to harmonic form as $N \rightarrow \infty$
- ▶ Ansatz for p_i : $A_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{n_i+1}$ linear, $\text{sq} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ square each coordinate

$$p(x_0, \dots, x_4) = A_k \circ \dots \circ \text{sq} \circ A_2 \circ \text{sq} \circ A_1(x_0, \dots, x_4)$$

- ▶ Equivalent: neural network with activation function $x \mapsto x^2$
- ▶ Approximate metric $\frac{i}{2} \partial \bar{\partial} K$ smooth+explicit \Rightarrow explicitly compute $(|d\lambda_\alpha|(x_i) + |d^* \lambda_\alpha|(x_i)) / \sqrt{|\lambda(x_1)|^2 + \dots + |\lambda(x_{100000})|^2}$
 \Rightarrow minimise with tensorflow


Experimental results: 1-forms and their zeros

1. **Fermat:** non-example 1; no harmonic 1-form.
2. **Random Quintic:** non-example 2; harmonic 1-form must have zeros
3. **CICY1:** conjectural example 3; large perturbation $\epsilon = \frac{1}{4}$, harmonic 1-form may have zeros
4. **CICY2:** conjectural example 3; small perturbation $\epsilon = \frac{1}{100}$, conjecture no zeros

y-axis: $\min |\lambda|$

x-axis:



$$\frac{\|d\lambda\|_{L^1} + \|d^*\lambda\|_{L^1}}{\|\lambda\|_{L^2}}$$



Experimental results on quintic

- ▶ $g = v \cdot f_-$ **singular quintic** from before, $\xi = 0.84x_0^5 + \dots$ **random quintic**
- ▶ Find $\epsilon > 0$ such that $g_\epsilon := g + \epsilon\xi$ has $Z_{\mathbb{R}}(g_\epsilon)$ diffeo to $Z_{\mathbb{R}}(g)$
 - ▶ $U \subset \mathbb{RP}^4$ nbhd of $Z_{\mathbb{R}}(g)$
 - ▶ $k := \min_U |Dg| > 0$, $M := \min_{\mathbb{RP}^4 \setminus U} |g| > 0$
 - ▶ if $\|\epsilon_0 \xi\|_{C^0} < M$ and $\|D\epsilon_0 \xi\|_{C^0} < k$, then $Z_{\mathbb{R}}(g_\epsilon)$ smooth for all $0 < \epsilon < \epsilon_0$
 - ▶ $\Rightarrow Z_{\mathbb{R}}(g_\epsilon)$ diffeo for all $0 < \epsilon < \epsilon_0$ (for us $\epsilon_0 = 0.00195503$)

Train Loss Curve for the CY metric





Average of $|\text{vol}_h / \text{vol}_\Omega - 1|$ while
iteratively improving vol_h

$|\text{vol}_h / \text{vol}_\Omega(x) - 1|$ over
 $\max\{v(x)/\|x\|^2, f_-(x)/\|x\|^3\}$

Experimental results on quintic

- ▶ $g = v \cdot f_-$ singular quintic from before, $\xi = 0.84x_0^5 + \dots$ random quintic
- ▶ Find $\epsilon > 0$ such that $g_\epsilon := g + \epsilon\xi$ has $Z_{\mathbb{R}}(g_\epsilon)$ diffeo to $Z_{\mathbb{R}}(g)$
 - ▶ $U \subset \mathbb{RP}^4$ nbhd of $Z_{\mathbb{R}}(g)$
 - ▶ $k := \min_U |Dg| > 0$, $M := \min_{\mathbb{RP}^4 \setminus U} |g| > 0$
 - ▶ if $\|\epsilon_0 \xi\|_{C^0} < M$ and $\|D\epsilon_0 \xi\|_{C^0} < k$, then $Z_{\mathbb{R}}(g_\epsilon)$ smooth for all $0 < \epsilon < \epsilon_0$
 - ▶ $\Rightarrow Z_{\mathbb{R}}(g_\epsilon)$ diffeo for all $0 < \epsilon < \epsilon_0$ (for us $\epsilon_0 = 0.00195503$)

Train Loss Curve for the CY metric




Average of $|\text{vol}_h / \text{vol}_\Omega - 1|$ while
iteratively improving vol_h

$|\text{vol}_h / \text{vol}_\Omega(x) - 1|$ over
 $\max\{v(x)/\|x\|^2, f_-(x)/\|x\|^3\}$


Experimental results on quintic

- ▶ $g = v \cdot f_-$ singular quintic from before, $\xi = 0.84x_0^5 + \dots$ random quintic
- ▶ Find $\epsilon > 0$ such that $g_\epsilon := g + \epsilon\xi$ has $Z_{\mathbb{R}}(g_\epsilon)$ diffeo to $Z_{\mathbb{R}}(g)$
 - ▶ $U \subset \mathbb{RP}^4$ nbhd of $Z_{\mathbb{R}}(g)$
 - ▶ $k := \min_U |Dg| > 0$, $M := \min_{\mathbb{RP}^4 \setminus U} |g| > 0$
 - ▶ if $\|\epsilon_0 \xi\|_{C^0} < M$ and $\|D\epsilon_0 \xi\|_{C^0} < k$, then $Z_{\mathbb{R}}(g_\epsilon)$ smooth for all $0 < \epsilon < \epsilon_0$
 - ▶ $\Rightarrow Z_{\mathbb{R}}(g_\epsilon)$ diffeo for all $0 < \epsilon < \epsilon_0$ (for us $\epsilon_0 = 0.00195503$)

Train Loss Curve for the CY metric




Average of $|\text{vol}_h / \text{vol}_\Omega - 1|$ while
iteratively improving vol_h



$|\text{vol}_h / \text{vol}_\Omega(x) - 1|$ over
 $\max\{v(x)/\|x\|^2, f_-(x)/\|x\|^3\}$


Experimental results on quintic

Neck formation



$$\max_{v \in T_{\|v\|_{FS}=1} \|v\|_h} \text{ over } \max\{v(x)/\|x\|^2, f_-(x)/\|x\|^3\}$$

1-form has even number of zeros




k -medoid clustering loss of 500 points with smallest $|\omega|(x)$ over number of clusters (heuristic: "elbow" $k = 4$ is optimal number of clusters)


Experimental results on quadric \cap quartic

$$\epsilon = \frac{1}{4}$$

Train Loss Curve for the CY metric




↑ Training loss




$$\epsilon = \frac{1}{100}$$

Train Loss Curve for the CY metric




↓ Loss over distance from singularity




Experimental results on quadric \cap quartic

$$\epsilon = \frac{1}{4}$$



$$\epsilon = \frac{1}{100}$$




↑ Metric stretching over distance from singularity

Experimental results on quadric \cap quartic

$$c = -x_0^2 + x_1^2 + x_2^2 \text{ and } q = -x_0^4 + x_3^4 + x_4^4 + x_5^4$$

Set $x_0 = 1$ and $x_3 = x_4 = 0 \leadsto \{(x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 = 1\} \times \{\pm 1\}$

1-form restricted to this



Bonus motivation

Proposition

For all $\epsilon > 0$ there exists $\delta > 0$ such that

$$\left\| \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right\|_{L_1^p} < \delta \Rightarrow \|g_{\text{approx}} - g_{\text{CY}}\|_{L_1^p} < \epsilon.$$

Proposition

For all $\mu > 0$ there exists $\epsilon > 0$ such that the following is true:

for $\lambda \in \Omega^1(L^3)$ such that $\Delta_{\text{approx}} \lambda = 0$ and $\|\lambda\|_{L^2, g_{\text{approx}}} = 1$ and $\min_L |\lambda| > \mu$ let $\tilde{\lambda} \in [\lambda]$ be the unique Δ_{CY} -harmonic 1-form. Then:

$$\|g_{\text{approx}} - g_{\text{CY}}\|_{L_1^p} < \epsilon \Rightarrow |\tilde{\lambda} - \lambda|(x) < \frac{\mu}{2} \Rightarrow |\tilde{\lambda}|(x) > \frac{\mu}{2} \text{ for all } x \in L.$$

- Find: g_{approx} with $\left\| \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right\|_{L_1^p} < \delta$, λ with $\Delta_{\text{approx}} \lambda = 0$ and $\min_L |\lambda| > \mu$
- \Rightarrow there exists nowhere vanishing g_{CY} -harmonic 1-form on L

Bonus motivation

Proposition

For all $\epsilon > 0$ there exists $\delta > 0$ such that

$$\left\| \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right\|_{L_1^p} < \delta \Rightarrow \|g_{\text{approx}} - g_{CY}\|_{L_1^p} < \epsilon.$$

Proposition

For all $\mu > 0$ there exists $\epsilon > 0$ such that the following is true:

for $\lambda \in \Omega^1(L^3)$ such that $\Delta_{\text{approx}} \lambda = 0$ and $\|\lambda\|_{L^2, g_{\text{approx}}} = 1$ and $\min_L |\lambda| > \mu$ let $\tilde{\lambda} \in [\lambda]$ be the unique Δ_{CY} -harmonic 1-form. Then:

$$\|g_{\text{approx}} - g_{CY}\|_{L_1^p} < \epsilon \Rightarrow |\tilde{\lambda} - \lambda|(x) < \frac{\mu}{2} \Rightarrow |\tilde{\lambda}|(x) > \frac{\mu}{2} \text{ for all } x \in L.$$

► Find: g_{approx} with $\left\| \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right\|_{L_1^p} < \delta$, λ with $\Delta_{\text{approx}} \lambda = 0$ and $\min_L |\lambda| > \mu$

► \Rightarrow there exists nowhere vanishing g_{CY} -harmonic 1-form on L

Bonus motivation

Proposition

For all $\epsilon > 0$ there exists $\delta > 0$ such that

$$\left\| \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right\|_{L_1^p} < \delta \Rightarrow \|g_{\text{approx}} - g_{CY}\|_{L_1^p} < \epsilon.$$

Proposition

For all $\mu > 0$ there exists $\epsilon > 0$ such that the following is true:





for $\lambda \in \Omega^1(L^3)$ such that $\Delta_{\text{approx}} \lambda = 0$ and $\|\lambda\|_{L^2, g_{\text{approx}}} = 1$ and $\min_L |\lambda| > \mu$ let $\tilde{\lambda} \in [\lambda]$ be the unique Δ_{CY} -harmonic 1-form. Then:

$$\|g_{\text{approx}} - g_{CY}\|_{L_1^p} < \epsilon \Rightarrow |\tilde{\lambda} - \lambda|(x) < \frac{\mu}{2} \Rightarrow |\tilde{\lambda}|(x) > \frac{\mu}{2} \text{ for all } x \in L.$$


- Find: g_{approx} with $\left\| \frac{\text{vol}_h}{\text{vol}_\Omega} - 1 \right\|_{L_1^p} < \delta$, λ with $\Delta_{\text{approx}} \lambda = 0$ and $\min_L |\lambda| > \mu$
- \Rightarrow there exists nowhere vanishing g_{CY} -harmonic 1-form on L

Thank you for the attention!

References I

-  Donaldson, S. K. (2009).
Some numerical results in complex differential geometry.
Pure Appl. Math. Q., 5(2, Special Issue: In honor of Friedrich Hirzebruch. Part 1):571–618.
-  Douglas, M., Lakshminarasimhan, S., and Qi, Y. (2022).
Numerical calabi-yau metrics from holomorphic networks.
In *Mathematical and Scientific Machine Learning*, pages 223–252. PMLR.
-  Headrick, M. and Nassar, A. (2013).
Energy functionals for calabi-yau metrics.
In *Journal of Physics: Conference Series*, volume 462, page 012019. IOP Publishing.
-  Jaco, W. H. (1980).
Lectures on three-manifold topology.
Number 43. American Mathematical Soc.

References II

-  Joyce, D. and Karigiannis, S. (2017).
A new construction of compact G_2 -manifolds by gluing families of Eguchi-Hanson spaces.
ArXiv e-prints.
-  Krasnov, V. A. (2009).
On the topological classification of real three-dimensional cubics.
Mat. Zametki, 85(6):886–893.
-  Larfors, M., Lukas, A., Ruehle, F., and Schneider, R. (2022).
Numerical metrics for complete intersection and kreuzer–skarke calabi–yau manifolds.
Machine Learning: Science and Technology, 3(3):035014.
-  Sun, S. and Zhang, R. (2019).
Complex structure degenerations and collapsing of calabi-yau metrics.
arXiv preprint arXiv:1906.03368.



Tian, G. and Yau, S.-T. (1990).

Complete kähler manifolds with zero ricci curvature. i.

Journal of the American Mathematical Society, 3(3):579–609.



Tischler, D. (1970).

On fibering certain foliated manifolds over S^1 .

Topology, 9(2):153–154.