



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Continued Monitoring



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT lay the person down** — the upright position is essential for breathing.
✕ **Do NOT leave them alone** during an attack.
✕ **Do NOT give controller medications** (inhaled corticosteroids) for acute relief — no immediate effect.
✕ **Do NOT delay calling 101** if no inhaler is available.
✕ **Do NOT ignore a 'silent chest'** — absence of wheezing during obvious distress = life-threatening.

Equipment needed:


— Rescue inhaler (Ventolin / Salbutamol)
— Spacer device (merkhav) if available

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Inhaler	משאף	mash-AF
Spacer	מרחב	mer-KHAV
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
Asthma	אסתמה	AST-ma
Asthma attack	התקף אסתמה	het-KEF AST-ma
Wheezing	צפצופים	tsif-tsu-FIM
Chest tightness	לחץ בחזה	LA-khats ba-kha-ZE
Puffs (doses)	שאיפות	she-i-FOT
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Can you breathe?	אתה יכול לנשום?	a-TA ya-KHOL lin-SHOM?
Where is the inhaler?	איפה המשאף?	EI-fo ha-mash-AF?

Source: Magen David Adom (MDA) — MDA ALS Protocol Manual 2022
URL: https://www.mdais.org/101/first-aid
Publication date: 2022-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Personal reference only — not medical advice
Daniel Rosehill · share with attribution


CHECK responsiveness and breathing (max 10 s).
Gasping is NOT normal breathing.
Unresponsive and not breathing normally?
N
Recovery position if unconscious but breathing.
Provide appropriate first aid.
CALL 101 immediately. Use speakerphone.
Dispatcher can unlock nearest smart stand AED.
BEGIN CPR immediately.
100–120 compressions/min 5–6 cm depth. Full recoil.
Another person available to retrieve AED?
N
If AED is nearby, retrieve quickly. Otherwise continue CPR.
Y
SEND them for AED. Continue CPR.
LOCATE AED: Eifo-Defi app, MDA smart stands, public buildings.

AED Pad Placement & Shock Delivery



Continued AED / CPR Cycles



Reference Notes

DO NOT:

- ✕ **Do NOT use AED on a conscious, breathing person** – AEDs are only for cardiac arrest.
- ✕ **Do NOT apply pads in water** – move to dry ground and dry the chest first.
- ✕ **Do NOT use near flammable gases** – risk of fire or explosion.
- ✕ **Do NOT place pads directly over a pacemaker** – place at least 2.5 cm away.
- ✕ **Do NOT place pads over medication patches** – remove and wipe skin first.
- ✕ **Do NOT touch patient during analysis or shock delivery** – electrocution risk.
- ✕ **Do NOT turn off AED or disconnect pads** – even if person regains consciousness.
- ✕ **Do NOT delay defibrillation** – survival drops 7–10% per minute of delay.
- ✕ **Do NOT interrupt compressions for more than 10 seconds**.
- ✕ **Do NOT let pads touch or overlap** on the chest.
- ✕ **Do NOT use AED on a breathing, pulsed unconscious person** – use recovery position instead.

Equipment needed:

— AED (from MDA smart stands, public locations, or Eifo-Defi app)
— Razor (for shaving chest hair -- often in AED kits)
— Towel or cloth (for drying the chest)
— Scissors (for cutting clothing -- often in AED kits)
— Phone (to call 101 and use speakerphone)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
Stand clear!	התרחקו!	hit-rakh-KU!
Cardiac arrest	דום לב	dom lev
Chest compressions	עיסויי חזה	i-su-YEI kha-ZE
Shock	הלם / שוק	HE-lem / shok
Not breathing	לא נושם	lo no-SHEM
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
AED	דפיברילטור חיצוני	de-fib-ri-LA-tor khi-tso-NI
Smart stand	עמדה חכמה	am-DA kha-kha-MA

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — AED Operation Protocol / AHA 2025 Guidelines
URL: https://www.mdais.org/kursim/defibrillator
Publication date: 2025-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


👁️ RECOGNISE signs: rash, swelling, breathing difficulty, weak pulse, confusion.
Signs of anaphylaxis? (breathing, circulation, or consciousness affected)
N
👁️ Mild reaction only. Monitor for worsening. Call 101 if deteriorates.
📱 CALL 101 immediately. Speakerphone on. Identify allergen if possible.
REMOVE offending agent. Scrape bee stinger sideways. Stop any infusion.
🌬️ MANAGE airway. Clear mouth. Loosen clothing around neck and chest.
Is the patient conscious?
Y
🧑 Breathing difficulty: SIT upright. Shock/hypotension: LIE down, elevate legs.
🌬️ Breathing?
N
Y
N
PLACE in recovery position. Proceed to EpiPen.
❤️ BEGIN CPR immediately.

EpiPen & Monitoring



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT delay calling 101** – anaphylaxis is life-threatening.
✕ **Do NOT delay epinephrine** – it is the only effective first-line treatment. Over 50% of deaths occur within 60 minutes.
✕ **Do NOT substitute antihistamines for epinephrine** – antihistamines are not effective against anaphylaxis.
✕ **Do NOT make a hypotensive patient stand or sit up suddenly** – this can cause cardiac arrest.
✕ **Do NOT leave the patient unattended** after epinephrine – symptoms may recur (biphasic reaction).
✕ **Do NOT assume safety after symptom resolution** – hospital evaluation is mandatory.
✕ **Do NOT waste time removing clothing** – EpiPen can be administered through trousers.
✕ **Do NOT inject EpiPen into a vein** – intramuscular outer thigh only.
✕ **Do NOT give food or drink** to a patient experiencing anaphylaxis.

Equipment needed:

— Epinephrine auto-injector (EpiPen) -- adult 0.3 mg
— Phone to call MDA at 101

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Allergic reaction	תגובה אלרגית	tgu-VA a-LER-git
Anaphylaxis	אנפילקסיס	a-na-fi-LAK-sis
EpiPen	אפיפן	E-pi-pen
Swelling	נפיחות	ne-fi-KHUT
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
Rash / hives	פריחה / סרפדת	pri-KHA / sar-PE-det
Injection	זריקה	zri-KA
Thigh	ירך	ya-REKH
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you allergic?	יש לך אלרגיה?	yesh le-KHA a-ler-GI-ya?
Where is the EpiPen?	איפה האפיפן?	EI-fo ha-E-pi-pen?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — ALS Protocol Manual April 2024
URL: https://www.mdais.org/101/allergy
Publication date: 2024-04-01
Imported: 2026-03-15 · **Last verified:** 2026-03-15


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Secondary Assessment & Monitoring



Reference Notes

DO NOT:

✕ **Do NOT approach** until security forces confirm the scene is safe – secondary devices are a known tactic.
✕ **Do NOT remove embedded shrapnel or debris** – may cause uncontrollable bleeding.
✕ **Do NOT remove a tourniquet** once applied – only medical professionals should decide.
✕ **Do NOT remove clothing stuck to burns** – cut around it.
✕ **Do NOT let the victim move or walk** until MDA arrives.
✕ **Do NOT use head-tilt for airway management** in blast/trauma – use jaw-thrust to protect cervical spine.
✕ **Do NOT administer epinephrine in hypovolemic trauma** – blood vessels are already constricted from shock.
✕ **Do NOT assume only one injury site** – blast victims have injuries from multiple mechanisms.

Equipment needed:

— Phone to call 101 (MDA)
— Sterile gauze or clean cloth for direct pressure
— Israeli bandage (emergency bandage)
— Gauze for wound packing
— Tourniquet (CAT preferred; improvised as last resort)
— Non-adherent dressings for burns
— Clean water for cooling burns
— Blanket for warmth

Personal reference only — not medical advice


Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Explosion	פיצוץ	pi-TSUTS
Shrapnel	רסיסים	rsi-SIM
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Bandage	תחבושת	takh-BO-shet
Bleeding	דימום	di-MUM
Pressure	לחץ	LA-khats
Wound	פצע	PE-tsa
Burns	כוויות	kvi-YOT
Shock	הלם	HE-lem
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you injured?	אתה פצוע?	a-TA pa-TSU-a?
Where does it hurt?	איפה כואב?	EI-fo ko-EV?

Source: Magen David Adom (MDA) — First Aid Guidelines: Blast and Shrapnel Injuries
URL: https://www.mdais.org/media/4954/mda-first-aid-guidelines-en.pdf
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Personal reference only — not medical advice
Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Chemical & Electrical Burns / Monitoring



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply ice or ice-cold water** – extreme cold causes additional tissue and blood vessel damage.
✕ **Do NOT apply toothpaste, coffee, egg, yogurt, butter, or home remedies** – these trap heat and increase infection risk.
✕ **Do NOT rupture or pop blisters** – the fluid protects underlying tissue during healing.
✕ **Do NOT use fat-based creams or ointments** on fresh burns.
✕ **Do NOT remove clothing stuck to the burn** – cut around it instead.
✕ **Do NOT attempt chemical neutralisation** – do not apply acid to base burn or vice versa.
✕ **Do NOT approach electrical burn casualty** until power source is confirmed disconnected.

Equipment needed:

— Cool running water source
— Cling film (food wrap) or clean plastic bag
— Clean non-fluffy dressings or bandages
— Scissors (to cut clothing around stuck areas)
— Blanket (to keep patient warm)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Burn	כוויה	kvi-YA
Water	מים	MA-yim
Cold water	מים קרים	MA-yim ka-RIM
Blister	שלפוחית	shal-pu-KHIT
Bandage	תחבושת	takh-BO-shet
Cling film	ניילון נצמד	NAI-lon nits-MAD
Chemical	כימיקל	khi-mi-KAL
Electricity	חשמל	KHASH-mal
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Does it hurt?	כואב לך?	ko-EV le-KHA?
Where is the burn?	איפה הכוויה?	EI-fo ha-kvi-YA?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Burns First Aid Guidelines
URL: https://www.mdais.org/101/burns
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-15


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Unconscious Choking & Self-Treatment



Self-Treatment if Choking Alone

LEAN over a firm object (chair back, countertop edge).
PRESS upper abdomen forcefully against the object.
Repeat until the object is expelled.
📱 **CALL** 101 as soon as you can speak.

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT perform abdominal thrusts on a person who can cough and speak** – you may worsen the obstruction.
✕ **Do NOT perform back blows on an adult** – MDA/Israeli protocol explicitly describes this as incorrect and potentially harmful for adults.
✕ **Do NOT perform blind finger sweeps** – only remove an object if you can clearly see it.
✕ **Do NOT give water or liquid** to a choking person.
✕ **Do NOT continue abdominal thrusts on an unconscious person** – transition to CPR.
✕ **Do NOT use abdominal thrusts on pregnant or very obese individuals** – use chest thrusts instead.

Note: Israeli Protocol Divergence


MDA/Israeli protocol **does not recommend back blows** for adults. This differs from AHA (2025) which recommends alternating 5 back blows and 5 abdominal thrusts, and from ERC (2025) which recommends 5 back blows first. Follow MDA guidance when in Israel.

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Choking	חנק	KHE-nek
Foreign body	גוף זר	guf zar
Can you breathe?	אתה יכול לנשום?	a-TA ya-KHOL lin-SHOM?
Can you speak?	אתה יכול לדבר?	a-TA ya-KHOL le-da-BER?
Cough	שיעול	shi-UL
Stomach / abdomen	בטן	BE-ten
Chest	חזה	kha-ZE
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you choking?	אתה נחנק?	a-TA nekh-NAK?

Source: Magen David Adom (MDA) — How to Treat Choking from a Foreign Body
URL: https://www.mdais.org/101/asphyxia
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution


CHECK responsiveness. Call out, shake shoulders. "Are you okay?"
Is the person responsive?
Y
ASSESS condition. Provide appropriate first aid. Call 101 if needed.
N
CHECK breathing (max 10 seconds). Look, listen, feel.
Breathing normally? (Gasping is NOT normal)
Y
Recovery position.
Call 101.
Monitor breathing.
CALL 101 immediately. Use speakerphone. Request AED.
Call 101 yourself (speakerphone). Retrieve AED if nearby. Begin compressions.
N
Another person available?
Y
DIRECT them to call 101 and retrieve nearest AED. Begin compressions.
POSITION supine on firm surface. Heel of hand on centre of chest.

Compressions, Breaths & AED



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT delay calling 101** – early activation is critical for survival.
- ✕ **Do NOT spend more than 10 seconds checking breathing** – delays reduce survival.
- ✕ **Do NOT interrupt compressions for more than 10 seconds** for any reason.
- ✕ **Do NOT lean on the chest between compressions** – allow full recoil.
- ✕ **Do NOT compress deeper than 6 cm** to avoid internal injury.
- ✕ **Do NOT over-ventilate** – 1 second per breath, just enough for visible chest rise.
- ✕ **Do NOT touch the person during AED analysis or shock delivery**.
- ✕ **Do NOT delay AED use** – apply as soon as it arrives.
- ✕ **Do NOT wait for AED to re-analyse** before resuming compressions after shock.
- ✕ **Do NOT move the person unnecessarily** – perform CPR where they are unless unsafe.
- ✕ **Do NOT assume gasping is normal breathing** – it is a sign of cardiac arrest.

Equipment needed:

— AED (Automated External Defibrillator)
— Barrier device or pocket mask (for rescue breaths, if available)
— Phone (to call 101 and use speakerphone for dispatcher-guided CPR)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
CPR	החייאה	ha-kha-ya-A
Cardiac arrest	דום לב	dom lev
Chest compressions	עיסויי חזה	i-su-YEI kha-ZE
Rescue breaths	הנשמות	han-sha-MOT
Are you okay?	אתה בסדר?	a-TA be-SE-der?
Not breathing	לא נושם	lo no-SHEM
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
Stand clear!	התרחקו!	hit-rakh-KU!
Recovery position	תנוחת התאוששות	tnu-KHAT hit-o-she-SHUT
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Call an ambulance!	הזמינו אמבולנס!	haz-MI-nu am-bu-LANS!

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — CPR Protocol / AHA 2025 Guidelines
URL: https://www.mdais.org/101/cpr-hands-only
Publication date: 2025-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-15


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Heat Stroke Emergency



Reference Notes

DO NOT:

✕ **Do NOT give large quantities of water quickly** – rapid fluid replacement disrupts electrolyte balance.
✕ **Do NOT give sugary drinks, coffee, or alcohol**.
✕ **Do NOT wet the head of a heat stroke patient** – choking risk if consciousness is altered.
✕ **Do NOT ignore the key distinction**: dehydration = continued sweating; heat stroke = ABSENT sweating with high body temperature.
✕ **Do NOT delay calling 101 for heat stroke** – it is a medical emergency requiring hospital treatment.
✕ **Do NOT give fluids to an unconscious person**.

Equipment needed:


— Water or ORS (Oral Rehydration Solution)
— Fan or method of creating air flow
— Wet cloths or towels for cooling
— Ice packs (for heat stroke)
— Straw (for controlled sipping)

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Dehydration	התייבשות	hit-yab-SHUT
Heat stroke	מכת חום	ma-KAT khom
Water	מים	MA-yim
Shade	צל	tsel
Dizzy	סחרחורת	skhar-kho-RET
Sweating	הזעה	haz-a-A
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Drink water	שתה מים	shte MA-yim
Are you thirsty?	אתה צמא?	a-TA tsa-ME?
Come to the shade	בוא לצל	bo la-TSEL


Source: Magen David Adom (MDA) — Dehydration and Heat-Related Illness First Aid
URL: https://www.mdais.org/101/first-aid
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution




Daniel Rosehill · share with attribution

Unconscious Hypoglycemia



Hyperglycemia Pathway



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT administer insulin** — insulin dosing requires medical expertise. Leave to MDA paramedics.
✕ **Do NOT give food or drink to an unconscious patient** — risk of choking and aspiration.
✕ **Do NOT give diet drinks or sugar-free foods** for hypoglycemia — they will not raise blood sugar.
✕ **Do NOT give sugary foods high in fat** (chocolate bars, ice cream) as first-line treatment — fat slows absorption.
✕ **Do NOT withhold sugar when in doubt** — giving sugar to a hypo patient is life-saving; giving sugar to a hyper patient will not cause significant immediate harm.
✕ **Do NOT leave the patient unattended** — condition can deteriorate rapidly.
✕ **Do NOT delay calling 101** if unconscious, seizing, or not improving after sugar.
✕ **Do NOT disconnect an insulin pump roughly** — disconnect gently to avoid damage.

Equipment needed:

— Phone to call 101 (MDA) or 1221 (United Hatzalah)
— Glucometer and test strips (if available)
— Rapidly absorbed sugar (glucose tablets, glucogel, juice, sugar cubes, honey)
— Complex carbohydrate for follow-up (bread, crackers)
— Glucagon injection kit (if available and trained)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Diabetes	סוכרת	su-KE-ret
Low blood sugar	סוכר נמוך	SU-kar na-MUKH
High blood sugar	סוכר גבוה	SU-kar ga-VO-ah
Sugar	סוכר	SU-kar
Insulin	אינסולין	in-su-LIN
Glucometer	מד סוכר	mad SU-kar
Juice	מיץ	mitz
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you diabetic?	אתה סוכרתי?	a-TA su-kar-TI?
Did you eat today?	אכלת היום?	a-KHAL-ta ha-YOM?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — How to Treat a Diabetes Patient
URL: https://www.mdais.org/101/diabetes
Publication date: 2026-03-15
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT enter the water** unless you are a trained swimmer — untrained rescuers frequently become victims.
✕ **Do NOT attempt to remove water from the lungs** — MDA states this is impossible, dangerous, and wastes critical time.
✕ **Do NOT perform the Heimlich manoeuvre** or turn the person upside down to "drain" water.
✕ **Do NOT use an AED on a wet patient** or wet surface — dry the chest and move to dry area first.
✕ **Do NOT allow a near-drowning victim to refuse hospital transport** — secondary drowning and electrolyte imbalances can be fatal hours later.
✕ **Do NOT stop CPR** until the person breathes on their own or MDA takes over.
✕ **Do NOT delay calling 101** — call immediately, even before attempting rescue if alone.

Equipment needed:

— AED (defibrillator) — commonly at lifeguard stations and public facilities
— Reaching aids (pole, rope, flotation device) for water rescue
— Dry blankets or clothing to keep victim warm after extraction

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Drowning	טביעה	tvi-A
Water	מים	MA-yim
Swimming pool	בריכה	bre-KHA
Beach	חוף	khof
Lifeguard	מציל	ma-TSIL
He is not breathing	הוא לא נושם	hu lo no-SHEM
CPR	החייאה	ha-kha-ya-A
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
Ambulance	אמבולנס	am-bu-LANS
Help!	הצילו!	ha-tsi-LU!
Call an ambulance	תזמין אמבולנס	taz-MIN am-bu-LANS
Recovery position	תנוחת התאוששות	tnu-KHAT hit-o-she-SHUT

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — Drowning First Aid Guidance
URL: https://www.mdais.org/101/drowning
Publication date: 2026-03-15
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


⚠️ STOP — Ensure scene safety. Is victim still in contact with source?
⚡ High-voltage incident? (power lines, substations, industrial equipment)
Y
MAINTAIN 10m distance. CALL 101 (MDA), 102 (Fire), 103 (IEC). Wait for professionals.
N
USE dry non-conducting object (wooden handle, dry rope). Stand on dry insulating surface.
N
Can you disconnect power at mains or unplug the device?
Y
SWITCH off mains or unplug appliance. Confirm power is off.
📱 CALL 101 (MDA). Report location, casualties, whether power disconnected.
🌬️ Scene confirmed safe. Is the victim responsive?
Y
KEEP still. Proceed to burns assessment (Page 2).
N
❤️ BEGIN CPR. 30:2 ratio at 100–120/ min. Use AED if available (dry chest first).
N
🌬️ Victim breathing normally (not just gasping)?
Y
🧑 PLACE in recovery position. Monitor continuously.

Burns, Spine, and Shock Assessment



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT touch the victim** while they may still be in contact with the electrical source — chain electrocution can kill the rescuer.
✕ **Do NOT use wet or damp objects** to separate the victim from the source.
✕ **Do NOT approach high-voltage incidents** — maintain at least 10 metres distance. Call 101, 102, and 103 (IEC).
✕ **Do NOT apply water to burns** while the victim is still connected to the electrical source.
✕ **Do NOT apply creams, ointments, or home remedies** to electrical burns.
✕ **Do NOT assume the injury is minor** based on small visible burns — internal damage along the current pathway may be severe.
✕ **Do NOT give food or drink** to an electrocution victim.
✕ **Do NOT perform CPR on a conscious victim.**

Equipment needed:

— Dry non-conducting object (wooden broom handle, dry rope, dry rubber mat)
— Dry insulating surface to stand on (dry wood, dry rubber mat, newspapers)
— AED (automated external defibrillator)
— Cool running water source (for burn cooling after scene is safe)
— Cling film or clean non-fluffy dressings (for burn covering)
— Blanket (for shock prevention)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Electrocution	התחשמלות	hit-khash-me-LUT
Electricity	חשמל	KHASH-mal
Electric shock	מכת חשמל	ma-KAT KHASH-mal
Power line	קו מתח	kav ME-takh
High voltage	מתח גבוה	ME-takh ga-VO-ah
Circuit breaker	מפסק חשמל	maf-SEK KHASH-mal
Burns	כוויות	kvi-YOT
He is unconscious	הוא מחוסר הכרה	hu me-khu-SAR ha-ka-RA
CPR	החייאה	ha-kha-ya-A
Ambulance	אמבולנס	am-bu-LANS
Fire department	כבאות	ka-ba-UT
Electric company	חברת חשמל	khev-RAT KHASH-mal

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Electrocution First Aid Guidelines
URL: https://www.mdais.org/101/electrocution
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Blunt Trauma / Laceration



Reference Notes

DO NOT:

✕ **Do NOT rub the eye** when a foreign body is present.
✕ **Do NOT attempt to remove an embedded or penetrating object.**
✕ **Do NOT apply pressure** to an injured eye or embedded object.
✕ **Do NOT wash the eye** if there is a laceration to the eyeball.
✕ **Do NOT delay flushing for chemical splashes** — start immediately.
✕ **Do NOT press a cold compress onto the eyeball** for blunt trauma.

Equipment needed:


— Clean running water source for flushing
— Sterile eye pads or clean cloth
— Cold compress (for blunt trauma)
— Cup or cone (to protect embedded objects)
— Bandages to loosely cover both eyes if needed

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Eye	עין	A-yin
Eye injury	פציעת עין	ptsi-AT A-yin
Chemical burn	כוויה כימית	kvi-YA khi-MIT
Foreign body	גוף זר	guf ZAR
I cannot see	אני לא רואה	a-NI lo ro-E
It hurts	כואב לי	ko-EV li
Water	מים	MA-yim
Flush / rinse	שטיפה	shti-FA
Bandage	תחבושת	takh-BO-shet
Emergency room	חדר מיון	KHE-der mi-YUN
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM

Source: Magen David Adom (MDA) — Eye Injury First Aid
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-03-16
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution


📱 CALL 101 (MDA). Keep on speaker for guidance.
Signs of fracture? (deformity, severe pain, inability to move limb, crepitus)
N
Likely sprain. See Page 2: RICE.
Y
Fracture pathway. Continue below.
Bone visible or protruding through the skin? (open fracture)
Y
Open fracture. CONTROL bleeding: press around (not on) bone. COVER wound with sterile dressing. Monitor for shock.
N
IMMOBILISE the limb in position found. Minimum movement, maximum stability.
Arm or upper limb?
N
PAD both sides of leg with pillows or rolled blankets to prevent movement.
Y
SUPPORT with improvised sling (shirt, scarf). Attach to chest.
APPLY ice wrapped in cloth (15-min intervals). REMOVE rings, watches, jewellery from affected limb.
👁 MONITOR for shock (pale skin, rapid breathing). Keep warm until MDA arrives.

Sprain — RICE Protocol



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT attempt to straighten or realign a broken limb** — this may tear arteries and nerves.
✕ **Do NOT push a protruding bone back in** during an open fracture.
✕ **Do NOT apply ice directly to skin** — always wrap in cloth first.
✕ **Do NOT apply heat** to a fresh fracture or sprain — heat increases swelling.
✕ **Do NOT allow the patient to walk** on a suspected leg fracture.
✕ **Do NOT remove embedded objects** from a wound associated with a fracture.

Equipment needed:

— Triangular bandage or improvised sling (shirt, scarf)
— Ice pack or cooling spray (wrapped in cloth)
— Elastic bandage (for compression of sprains)
— Pillows, blankets, or rolled towels (for immobilisation and elevation)
— Sterile dressings or clean cloth (for open fracture wounds)
— Tourniquet (for severe bleeding in open fractures)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Fracture	שבר	SHE-ver
Broken bone	עצם שבורה	E-tsem shvu-RA
Sprain	נקע	NE-ka
Arm	יד / זרוע	yad / zro-A
Leg	רגל	RE-gel
It hurts	כואב לי	ko-EV li
I cannot move it	אני לא יכול להזיז	a-NI lo ya-KHOL le-ha-ZIZ
Sling	מתלה	mit-LE
Ice	קרח	KE-rakh
Bandage	תחבושת	takh-BO-shet
Ambulance	אמבולנס	am-bu-LANS
Where does it hurt?	איפה כואב לך?	EI-fo ko-EV le-KHA?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Fractures and Sprains First Aid Guidelines
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Bleeding, Assessment, and Monitoring



Reference Notes

DO NOT:

- ✕ **Do NOT move the patient** unless there is immediate danger to life (fire, active threat, collapse).
- ✕ **Do NOT twist, flex, or extend the neck.**
- ✕ **Do NOT attempt to sit the patient up.**
- ✕ **Do NOT use head-tilt chin-lift** — use jaw thrust only.
- ✕ **Do NOT remove a helmet** without proper training and a second person.
- ✕ **Do NOT give food or drink.**
- ✕ **Do NOT leave the patient alone.**
- ✕ **Do NOT apply direct pressure to a suspected skull fracture.**
- ✕ **Do NOT plug ears or nose** if blood or clear fluid is draining.
- ✕ **Do NOT pull the patient by arms or legs.**
- ✕ **Do NOT extract from a vehicle** unless the vehicle is on fire or there is another immediate threat.
- ✕ **Do NOT give oral fluids** even if dehydrated — risk of vomiting and aspiration.
- ✕ **Do NOT attempt to straighten the head** if the patient resists or reports pain — stabilise in position found.

Equipment needed:

— Disposable gloves
— Clean dressings or bandages for wound care
— Sterile dressing for skull fracture coverage
— Cervical collar (if available and properly sized)
— Blanket or thermal protection
— Tourniquet (CAT) for associated limb haemorrhage
— High-visibility vest (if responding at a road accident)


Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Head injury	פגיעת ראש	pgi-AT rosh
Spinal injury	פגיעת עמוד שדרה	pgi-AT a-MUD shid-RA
Concussion	זעזוע מוח	ziz-U-a MO-akh
Neck	צוואר	tsa-VAR
Spine	עמוד שדרה	a-MUD shid-RA
Do not move	לא לזוז	lo la-ZUZ
Can you feel your legs?	אתה מרגיש את הרגליים?	a-TA mar-GISH et ha-rag-LA-yim?
Can you move your fingers?	אתה יכול להזיז אצבעות?	a-TA ya-KHOL le-ha-ZIZ ets-ba-OT?
Helmet	קסדה	kas-DA
Recovery position	תנוחת התאוששות	tnu-KHAT hit-o-she-SHUT
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM

Source: Magen David Adom (MDA) — BLS Guide 2016 / ALS Professional Manual 2024
URL: https://www.mdais.org/101/first-aid
Publication date: 2024-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice
Daniel Rosehill · share with attribution


📱 CALL 101 immediately. State suspected heart attack. Stay on line for guidance.
🧑 STOP all physical activity. POSITION half-sitting with back supported, knees slightly bent. LOOSEN tight clothing.
💊 Patient allergic to aspirin?
Y
Skip aspirin. Inform MDA team about allergy.
N
💊 GIVE 100–300 mg aspirin. Patient must CHEW (not swallow whole). Standard Israeli tablet = 100 mg.
Patient has their own prescribed nitroglycerin?
Y
HELP patient take their nitroglycerin as prescribed.
👁️ MONITOR continuously. Stay with patient at all times. Send bystander to fetch AED.
🌬️ Patient loses consciousness and stops breathing normally?
N
👁️ Continue monitoring. Keep half-sitting. Keep calm.
Y
Cardiac arrest. See Page 2: CPR / AED.

CPR and AED — Cardiac Arrest



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT allow the patient to walk, exert, or dismiss symptoms.**
✕ **Do NOT give food or beverages.**
✕ **Do NOT leave the patient alone** (except briefly to call for help).
✕ **Do NOT give aspirin if the patient is allergic.**
✕ **Do NOT give nitroglycerin** not prescribed to this specific patient.
✕ **Do NOT wait to see if symptoms resolve** — every minute of delay increases irreversible heart damage.
✕ **Do NOT drive the patient to hospital** — call 101 and wait for ambulance. MDA has cardiac monitoring equipment.
✕ **Do NOT substitute paracetamol (Acamol) for aspirin** — only aspirin provides the antiplatelet effect.
✕ **Do NOT give aspirin as a whole tablet** — it must be chewed for rapid absorption.

Equipment needed:

— Aspirin tablets (100 mg, 1–3 tablets for 100–300 mg total)
— AED (Automated External Defibrillator) — if available
— Phone (to call 101 MDA and optionally 1221 United Hatzalah)
— Nitroglycerin (only if prescribed to the patient)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Heart attack	התקף לב	het-KEF lev
Chest pain	כאב בחזה	ke-EV ba-kha-ZE
Heart	לב	lev
Aspirin	אספירין	as-pi-RIN
Chew	ללעוס	lil-OS
Nitroglycerin	ניטרוגליצרין	nit-ro-gli-tse-RIN
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
CPR	החייאה	ha-kha-ya-A
He is not breathing	הוא לא נושם	hu lo no-SHEM
Are you allergic to aspirin?	יש לך אלרגיה לאספירין?	yesh le-KHA a-ler-GI-ya le-as-pi-RIN?
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — Heart Attack Protocol / ALS Protocol 2024 / BLS Guide 2016
URL: https://www.mdais.org/101/mi-signs
Publication date: 2024-04-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice


Daniel Rosehill · share with attribution


🌡️ ASSESS condition. Heat exhaustion or heat stroke?
Confusion, hot dry skin, temp > 40 C, seizures, loss of consciousness?
Y
N
📱 CALL 101 immediately. Life-threatening emergency.
🧑 MOVE to shade or air-conditioned area.
REMOVE unnecessary clothing.
💦 COOL aggressively: wet towels on neck, armpits, groin. Spray cool water. Fan vigorously.
Heat stroke pathway continues on Page 2.
Heat exhaustion pathway. MOVE to shaded or air-conditioned location.
🧑 LAY person down. Elevate legs slightly.
REMOVE excess clothing.
💦 COOL: wet towels, spray cool water, fan.
Heat exhaustion pathway continues on Page 2.

Heat Stroke — Continued




Heat Exhaustion — Continued



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Continued Monitoring — Heat Stroke



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT administer fever-reducing medications** (paracetamol, ibuprofen) — antipyretics do not work for environmental hyperthermia and may cause harm.
✕ **Do NOT give fluids** to a person who is not fully conscious or who cannot swallow safely.
✕ **Do NOT delay calling 101** for suspected heat stroke.
✕ **Do NOT ignore early signs of heat exhaustion** — it can progress to life-threatening heat stroke.
✕ **Do NOT leave the person unattended** while waiting for emergency services.

Equipment needed:


— Wet towels or cloths
— Cool/cold water (for spraying, pouring, and drinking)
— Cold packs or ice packs (if available)
— Fan or other means of air circulation (if available)

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Heatstroke	מכת חום	ma-KAT KHOM
Heat exhaustion	תשישות חום	tshi-SHUT KHOM
Shade	צל	TSEL
Water	מים	MA-yim
Confused	מבולבל	me-vul-BAL
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Cool down	לקרר	le-ka-RER
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Can you drink?	אתה יכול לשתות?	a-TA ya-KHOL lish-TOT?
Move to shade	בוא לצל	BO la-TSEL

Source: Magen David Adom (MDA) — MDA First Aid Guidelines, Heat-Related Illness
URL: https://www.mdais.org/101/first-aid
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT rub extremities** — friction can cause tissue damage and cardiac complications.
✕ **Do NOT use direct intense heat** (hot water baths, heating pads directly on skin) — rapid rewarming can cause dangerous cardiac arrhythmias.
✕ **Do NOT give alcohol** — it dilates blood vessels and accelerates heat loss.
✕ **Do NOT give caffeine** — it worsens hypothermia.
✕ **Do NOT warm extremities first** — cold blood rushing to the heart can trigger cardiac arrest (afterdrop).
✕ **Do NOT apply vigorous external rewarming** — gradual rewarming is essential.

Equipment needed:


— Blankets or thermal blankets (space blankets)
— Dry clothing
— Heat packs with cloth barrier
— Warm (not hot) beverages (for conscious patients only)

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Hypothermia	היפותרמיה	hi-po-TER-mi-ya
Cold exposure	חשיפה לקור	kha-shi-FA la-KOR
Shivering	רעד	RA-ad
Blanket	שמיכה	smi-KHA
Warm drink	משקה חם	mash-KE KHAM
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נושם	no-SHEM
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you cold?	קר לך?	KAR le-KHA?
Cover him/her	תכסו אותו/אותה	tekh-SU o-TO / o-TA


Source: Magen David Adom (MDA) — MDA Emergency Medicine Manual 2024
URL: https://www.mdais.org/101/first-aid
Publication date: 2024-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Personal reference only — not medical advice
Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Ingestion and Inhalation Pathways



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT induce vomiting** unless specifically directed by the Poison Information Center (04-7771900) or a physician.
✕ **Do NOT give food or drink** unless directed by medical professionals (exception: water/milk for conscious patients who ingested caustic substances).
✕ **Do NOT attempt to neutralise the poison** (e.g., giving acid for base ingestion or vice versa).
✕ **Do NOT use home remedies** (salt water, mustard, raw eggs, etc.).
✕ **Do NOT give activated charcoal** without medical direction.
✕ **Do NOT enter a confined space** with potential toxic fumes without proper protection.
✕ **Do NOT delay calling for help** — even mild symptoms can deteriorate rapidly.
✕ **Do NOT administer antidotes** — this is a hospital-level intervention only.
✕ **Do NOT leave the patient alone.**

Equipment needed:


— Gloves (to protect against contamination)
— Running water source (for skin or eye decontamination)

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Poisoning	הרעלה	har-a-LA
Poison	רעל	RA-al
Swallowed	בלע	ba-LA
Vomiting	הקאה	ha-ka-A
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נושם	no-SHEM
Cleaning product	חומר ניקוי	KHO-mer ni-KUI
Poison Center	מרכז הרעלות	mer-KAZ har-a-LOT
Ambulance	אמבולנס	am-bu-LANS
What did they swallow?	מה הוא בלע?	MA HU ba-LA?
How much?	כמה?	KA-ma?


Source: Magen David Adom (MDA) — Poisoning First Aid Guidance
URL: https://www.mdais.org/101/poisoningn
Poison Center: 04-7771900 (Rambam Health Care Campus, Haifa — 24/7)
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Placing in Recovery Position



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT place in recovery position if the person is not breathing** — begin CPR instead.
✕ **Do NOT move a person with suspected spinal injury** unless they are in immediate danger.
✕ **Do NOT leave the person unattended** — monitor breathing continuously.
✕ **Do NOT place on their back if unconscious and breathing** — risk of airway obstruction and aspiration.
✕ **Do NOT forget to call 101** — an unconscious person always requires professional medical evaluation.

Equipment needed:


— Blanket (to keep the person warm)
— Phone to call MDA at 101

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Recovery position	תנוחת התאוששות	ta-nu-KHAT hit-o-she-SHUT
Lateral position	תנוחת צד	ta-nu-KHAT TSAD
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נושם	no-SHEM
Airway	נתיב אוויר	na-TIV a-VIR
Are you OK?	אתה בסדר?	a-TA be-SE-der?
He/she is not responding	הוא/היא לא מגיב/ה	HU/HI LO ma-GIV/ma-gi-VA
Roll onto side	הפוך על הצד	ha-FOKH al ha-TSAD
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM

Source: Magen David Adom (MDA) — MDA Emergency Medicine Manual 2024; BLS Guide 2016
URL: https://www.mdais.org/101/first-aid
Publication date: 2024-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply a tourniquet** — it traps venom locally and causes tissue damage.
✕ **Do NOT suck the venom out** — this is ineffective and potentially harmful.
✕ **Do NOT cut the sting area** — increases infection risk and does not remove venom.
✕ **Do NOT apply ice or cool the sting area** — Israeli Ministry of Health (2026) explicitly advises against cooling.
✕ **Do NOT give the victim food or drink** — risks venom dispersal and may complicate hospital treatment.
✕ **Do NOT allow the victim to walk or move the stung limb** — movement accelerates venom circulation.

Equipment needed:


— Phone to call 101 (MDA) and optionally 04-7771900 (Poison Center)
— Phone or camera to photograph the scorpion if safe to do so

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Scorpion	עקרב	ak-RAV
Scorpion sting	עקיצת עקרב	a-ki-TSAT ak-RAV
Deathstalker (yellow scorpion)	עקרב צהוב	ak-RAV tsa-HOV
Venom	ארס	E-res
Antivenom	נוגדן	nog-DAN
Swelling	נפיחות	ne-fi-KHUT
Pain	כאב	KE-ev
Poison Center	מרכז הרעלות	mer-KAZ har-a-LOT
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Where were you stung?	איפה נעקצת?	EI-fo ne-ek-TSATS-ta?
Stay still	תישאר במקום	ti-sha-ER ba-ma-KOM


Source: Magen David Adom (MDA) — First Aid for Scorpion Stings; Israeli Ministry of Health
URL: https://www.mdais.org/101/first-aid
Poison Center: 04-7771900 (Rambam Health Care Campus, Haifa — 24/7)
Note: All scorpion sting victims must be observed in ED for minimum 6 hours.
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Post-Seizure Assessment



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT restrain the person** or try to hold them down — you cannot stop a seizure by force, and restraining can cause fractures or dislocations.
✕ **Do NOT put anything in the person's mouth** — no spoons, wallets, sticks, or fingers. The person cannot swallow their tongue. Objects risk breaking teeth.
✕ **Do NOT attempt mouth-to-mouth during the seizure** — people usually resume breathing once the seizure ends.
✕ **Do NOT pour water on the person** or attempt to "wake them up".
✕ **Do NOT offer food, drinks, or medication by mouth** until the person is fully alert and able to swallow safely.
✕ **Do NOT crowd around the person** — give them space and privacy. Ask bystanders to step back.
✕ **Do NOT move the person** unless they are in immediate danger (road, ledge, fire, water).

Equipment needed:

— Soft padding for under the head (jacket, towel, cushion, blanket)
— Watch or phone to time the seizure duration
— Prescribed rescue medication if available (Buccolam, Stesolid, or Rivotril — prescription only)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Seizure	התקף	het-KEF
Epilepsy	אפילפסיה	e-pi-LEP-si-ya
Convulsions	פרכוסים	pir-ku-SIM
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נושם	no-SHEM
Recovery position	תנוחת התאוששות	ta-nu-KHAT hit-o-she-SHUT
How long?	כמה זמן?	KA-ma ZMAN?
First time?	פעם ראשונה?	PA-am ri-sho-NA?
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Epilepsy Association	האגודה לאפילפסיה	ha-a-gu-DA le-e-pi-LEP-si-ya

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — First Aid Provider Course 2026; ALS Manual 2024
URL: https://www.mdais.org/101/first-aid
Israeli Epilepsy Association: 02-5000283
Note: Status epilepticus = seizure > 5 minutes = medical emergency.
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


⚠️ ENSURE scene safety. Wear gloves if available.
📱 CALL 101 immediately.
Embedded or impaled object in the wound?
Y
STABILISE object in place with padding. Apply pressure around it, not on it. Skip to shock monitoring.
N
👐 APPLY strong, direct pressure to the wound with gloved hand or sterile gauze.
If dressing soaks through: ADD layers on top. Never remove original.
Direct pressure controlled bleeding?
Y
MAINTAIN pressure. ELEVATE limb above heart level if possible. Monitor for shock.
N
APPLY Israeli Emergency Bandage (pressure dressing). Pad over wound, wrap through pressure bar, secure.
Israeli bandage controlled bleeding?
Y
MAINTAIN dressing. Monitor for shock.
N
Escalate: see Page 2 (tourniquet / wound packing).

Escalation — Tourniquet / Wound Packing



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT remove an embedded or impaled object** from the wound.
- ✕ **Do NOT lift or remove a blood-soaked dressing** to check the wound — add layers on top.
- ✕ **Do NOT remove or loosen a tourniquet** once applied unless instructed by a paramedic or physician.
- ✕ **Do NOT give food or drink** to a casualty with severe bleeding or suspected shock.
- ✕ **Do NOT use wound packing for penetrating stab wounds.**
- ✕ **Do NOT enlarge wound openings** to facilitate wound packing.
- ✕ **Do NOT assume a rapid pulse is caused only by anxiety** — suspect significant blood loss.
- ✕ **Do NOT delay calling 101** for severe or uncontrolled bleeding.
- ✕ **Do NOT apply a tourniquet directly over a joint** — place it 5-10 cm above the wound on the limb.

Equipment needed:

— Disposable gloves
— Sterile gauze or clean cloth
— Israeli Emergency Bandage (pressure dressing)
— Tourniquet (CAT or equivalent)
— Wound packing material (hemostatic gauze preferred)
— Triangular bandage (supplementary)
— Blanket or coat (for shock management)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Bleeding	דימום	di-MUM
Severe bleeding	דימום חמור	di-MUM kha-MUR
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Pressure bandage	תחבושת לחץ	takh-BO-shet LA-khats
Israeli bandage	אמצעי חבישה ישראלי	em-tsa-I kha-vi-SHA yis-re-e-LI
Direct pressure	לחץ ישיר	LA-khats ya-SHIR
Wound	פצע	PE-tsa
Shock	הלם	HE-lem
Gloves	כפפות	kfa-FOT
Ambulance	אמבולנס	am-bu-LANS
Press here hard	לחץ כאן חזק	le-KHATS KAN kha-ZAK
Where is the bleeding?	איפה הדימום?	EI-fo ha-di-MUM?

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — How to Stop Bleeding (MDA 101 Public Guidance)
URL: https://www.mdais.org/101/bleeding
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Continued Care



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT give food or drink** – moisten lips only if thirsty.
- ✕ **Do NOT elevate legs** if head, neck, or spinal injury is suspected.
- ✕ **Do NOT elevate legs** if the patient has a broken hip or leg.
- ✕ **Do NOT elevate legs** if it causes pain or further injury.
- ✕ **Do NOT elevate legs** if the patient has difficulty breathing – keep semi-upright.
- ✕ **Do NOT overheat** the patient when maintaining body temperature.
- ✕ **Do NOT move** a patient with suspected spinal injury unnecessarily.
- ✕ **Do NOT delay calling 101** – shock is a life-threatening emergency.
- ✕ **Do NOT remove a tourniquet** once applied – leave removal to MDA paramedics.
- ✕ **Do NOT leave** the patient unattended.

Equipment needed:

— Tourniquet -- widely available in Israeli first aid kits
— Israeli bandage (emergency bandage / amtza'sh) -- for wound packing and pressure
— Blanket or coat -- to maintain body temperature
— Ground insulation -- to protect from cold ground
— EpiPen (epinephrine auto-injector) -- for anaphylactic shock if prescribed

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Shock	הלם	HE-lem
Bleeding	דימום	di-MUM
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Israeli bandage	אמצ"ש	am-TZA-sh
Pulse	דופק	DO-fek
Blood pressure	לחץ דם	LA-khats DAM
Blanket	שמיכה	smi-KHA
Conscious	בהכרה	be-ha-ka-RA
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Raise the legs	להרים את הרגליים	le-ha-RIM et ha-rag-LA-yim

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — MDA-Approved First Aid Curriculum and BLS Guide
URL: https://www.mdais.org/101/first-aid
Publication date: Standing public guidance
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


🔥 ENSURE scene is safe. Stay low if indoors (smoke rises).
Smoke inhalation (fire/fumes)?
N
Tear gas exposure. See Page 2.
Y
Can you safely move the person to fresh air?
N
📱 WAIT for Fire & Rescue (102) and MDA (101). Do not risk your safety.
Y
MOVE to fresh air immediately.
📱 CALL 101. Also call 102 if fire ongoing. Report location and casualties.
👁️ Is the person conscious?
N
🌬️ Breathing?
Y
PLACE in recovery position (on side).
❤️ BEGIN CPR immediately (30:2).
Y
🧑 SIT upright for easier breathing.
👁️ MONITOR for respiratory distress. CHECK for burns (face, neck, hands). Cool burns with clean water.
👐 COVER with blanket. Keep warm until MDA arrives.

Tear Gas Exposure



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT re-enter** a smoke-filled or burning structure without proper firefighting equipment and training.
✕ **Do NOT rub eyes** after tear gas exposure – this worsens irritation and spreads the chemical.
✕ **Do NOT use milk** or other substances to flush eyes – use only clean water or saline due to infection risk.
✕ **Do NOT remove clothing** stuck to burns.
✕ **Do NOT assume stability** – symptoms may worsen over hours as airway swelling develops.
✕ **Do NOT leave** a smoke inhalation patient alone – continuous monitoring is required.
✕ **Do NOT ignore** soot around the mouth/nose or singed facial hair – these indicate potentially serious airway injury.

Equipment needed:

— Phone to call 101 (MDA) and 102 (Fire and Rescue)
— Clean water for eye flushing (tear gas) and burn cooling
— Clean cloth or non-adherent dressings for burns
— Blanket for warmth
— Soap for skin decontamination (tear gas)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Smoke	עשן	a-SHAN
Fire	אש / שריפה	ESH / sre-FA
Tear gas	גז מדמיע	GAZ mad-MI-a
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
Burns	כוויות	kvi-YOT
Fresh air	אוויר צח	a-VIR TSAKH
Water	מים	MA-yim
Blanket	שמיכה	smi-KHA
Ambulance	אמבולנס	am-bu-LANS
Fire department	כיבוי אש	ki-BUY ESH
Emergency	חירום	khe-RUM
Can you breathe?	אתה יכול לנשום?	a-TA ya-KHOL lin-SHOM?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — MDA First Aid Guidelines
URL: https://www.mdais.org/media/4954/mda-first-aid-guidelines-en.pdf
Publication date: Standing public guidance
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT apply a tourniquet** – it traps venom locally and causes severe tissue damage.
- ✕ **Do NOT suck the venom** out of the wound – this is ineffective and dangerous.
- ✕ **Do NOT cut or incise** the bite area – increases infection risk and does not help remove venom.
- ✕ **Do NOT cool or apply ice** to the bite area – can worsen tissue damage.
- ✕ **Do NOT give food or drink** – may complicate hospital treatment.
- ✕ **Do NOT induce vomiting.**
- ✕ **Do NOT give aspirin, ibuprofen, or naproxen** – viper venom is often anticoagulant.
- ✕ **Do NOT use alcohol-based disinfectants** on the wound.
- ✕ **Do NOT attempt to catch or kill the snake** – protected under Israeli law; risks additional bites.
- ✕ **Do NOT allow the victim to walk** or move the bitten limb – movement accelerates venom circulation.

Equipment needed:

— Splint or improvised immobilisation material
— Soap and clean water for wound cleaning
— Phone or camera to photograph the snake

Medically Dangerous Snakes in Israel

Species	Hebrew Name	Antivenom
Daboia palaestinae (Palestine viper)	צפע ארצישראלי	Available (Kamada Ltd.)
Echis coloratus (Painted saw-scaled viper)	אפעה צבעונית	Available (Kamada Ltd.)
Atractaspis engaddensis (Israeli mole viper)	שחור עין גדי	Not available

Peak season: spring/summer. Israel Poison Information Center: 04-7771900 (24/7, Rambam Hospital, Haifa)


Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Snake	נחש	na-KHASH
Snakebite	הכשת נחש	ha-ka-SHAT na-KHASH
Venom / poison	ארס	E-res
Swelling	נפיחות	ne-fi-KHUT
Splint	סד	SAD
Lie down	לשכב	lish-KAV
Calm down	להירגע	le-hi-ra-GE-a
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
I was bitten by a snake	ננשכתי על ידי נחש	ni-SHAKH-ti al ye-DEI na-KHASH

Source: Magen David Adom (MDA) — How to Treat a Snake Bite
URL: https://www.mdais.org/101/snake-bite
Publication date: Standing public guidance
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution


👐 REASSURE the victim. Move away from spider.
CLEAN bite with mild soap and water.
APPLY cool compress (wrapped cloth or ice pack). 15 min per hour.
ELEVATE bitten area above heart level. Minimise movement.
📷 Can the spider be safely photographed?
Y
PHOTOGRAPH and note features. Share with medical responders.
N
Report any observed details to paramedics.
Severe pain, muscle cramps, difficulty breathing, nausea, spreading redness, or systemic symptoms?
Y
📱 CALL 101 immediately. Also call Poison Center: 04-7771900. Proceed to Page 2.
N
👁️ MONITOR the bite. Call Poison Center (04-7771900) for guidance. Seek medical attention if symptoms worsen.

Species Assessment & Monitoring



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply a tourniquet** – spider venom is not stopped by tourniquets; causes tissue damage.
✕ **Do NOT cut, suck, or apply suction** to the bite wound – ineffective and increases infection risk.
✕ **Do NOT apply heat** to the bite – use only cool compresses.
✕ **Do NOT attempt to catch the spider** at risk of additional bites – photograph from a safe distance.
✕ **Do NOT excise a recluse bite** – early surgical excision is contraindicated and worsens outcomes.
✕ **Do NOT apply ice directly** to skin – wrap in cloth first.
✕ **Do NOT ignore mild symptoms** – recluse bites are often painless at first but progress to serious necrosis over hours to days.
✕ **Do NOT assume it is a spider bite** without confirmation – 83.5% of cases presenting as "spider bites" are actually other conditions.

Equipment needed:

— Phone to call 101 (MDA) and 04-7771900 (Israel Poison Center)
— Phone or camera to photograph the spider if safe
— Mild soap and clean water for wound cleaning
— Clean cloth or wrapped ice pack for cool compress

Medically Significant Spiders in Israel

Species	Venom Type	Antivenom
Mediterranean recluse (Loxosceles rufescens) "violin spider"	Necrotoxic – painless bite progressing to tissue necrosis	Not available
Mediterranean black widow (Latrodectus tredecimguttatus)	Neurotoxic – pain, muscle cramps, abdominal spasms	Available for severe cases

Israel Poison Information Center: 04-7771900 (24/7, Rambam Hospital, Haifa)


Daniel Rosehill · share with attribution


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Spider	עכביש	ak-a-VISH
Spider bite	עקיצת עכביש	a-ki-TSAT ak-a-VISH
Swelling	נפיחות	ne-fi-KHUT
Pain	כאב	ke-EV
Muscle cramps	התכווצויות שרירים	hit-kav-tsu-YOT shri-RIM
Poison center	מרכז רעלים	mer-KAZ re-a-LIM
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
I was bitten	ננשכתי	ni-SHAKH-ti

Source: Magen David Adom (MDA) — First Aid for Spider Bites
URL: https://www.mdais.org/media/4954/mda-first-aid-guidelines-en.pdf
Publication date: Standing public guidance
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Personal reference only — not medical advice
Daniel Rosehill · share with attribution


ASSESS using FAST:
F – Face: ask to smile, check for one-sided drooping.
A – Arms: extend both, one drifting down?
S – Speech: repeat a sentence, slurred or unable?
T – Time: r xact onset.
CALL 101 immediately. Even one FAST sign = emergency evaluation.
RECORD exact time of symptom onset. If woke with symptoms, record time sleep began.
Is the person conscious?
N
Breathing?
Y
BEGIN CPR immediately.
PLACE in recovery position (on side).
Y
LAY down with head slightly raised and supported.
LOOSEN restrictive clothing: scarves, ties, tight collars.



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Monitoring & Handover



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT give food or drink** – swallowing reflex may be impaired (dysphagia); aspiration risk.
✕ **Do NOT give any medications**, especially aspirin or blood thinners – approximately 15% of strokes are hemorrhagic, and these medications worsen bleeding.
✕ **Do NOT delay calling 101** – do not drive the person to hospital yourself unless unavoidable. The ambulance crew pre-notifies the hospital stroke team.
✕ **Do NOT ignore transient symptoms** – even if symptoms resolve (possible TIA), the person needs emergency evaluation. Approximately 40% of TIA patients have a full stroke within three months.
✕ **Do NOT allow eating, drinking, or oral medications** until a medical professional has assessed swallowing function.

Equipment needed:

— Phone to call 101 and record symptom onset time
— Pillow or support to slightly elevate the head

FAST Reference

Letter	Assessment
F	**Face** – Ask to smile. One side drooping?
A	**Arms** – Raise both arms. One drifting down?
S	**Speech** – Repeat a sentence. Slurred or unable?
T	**Time** – Note exact onset. Treatment window: 6 hours. Target: hospital within 60 min.

Even a single positive FAST sign = call 101 immediately.


Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Stroke	שבץ מוחי / אירוע מוחי	SHVATZ mo-KHI / ei-RU-a mo-KHI
Face	פנים	pa-NIM
Arm	זרוע	zro-A
Speech	דיבור	di-BUR
Smile	לחייך	le-kha-YEKH
Headache	כאב ראש	ke-EV ROSH
Dizziness	סחרחורת	skhar-KHO-ret
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Can you raise your arms?	אתה יכול להרים את הידיים?	a-TA ya-KHOL le-ha-RIM et ha-ya-DA-yim?
Can you speak?	אתה יכול לדבר?	a-TA ya-KHOL le-da-BER?


Source: Magen David Adom (MDA) — Stroke Recognition and First Aid
URL: https://www.mdais.org/101/stroke
Publication date: Standing public guidance
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Monitoring & Handover



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT remove the tourniquet** once applied – only medical professionals should decide on release or conversion.
✕ **Do NOT apply to junctional zones** (groin, armpit, neck), head, or abdomen – use direct pressure, wound packing, or pressure points for these areas.
✕ **Do NOT apply over a joint** – place on the limb above the wound, preferably over a single bone.
✕ **Do NOT loosen or adjust** to "check if bleeding has stopped" – this can restart hemorrhage.
✕ **Do NOT delay application** in life-threatening hemorrhage due to fear of limb loss – loss of a limb is preferable to loss of life.
✕ **Do NOT allow the victim to walk** or move the affected limb after tourniquet application.
✕ **Do NOT attempt tourniquet conversion** (replacing tourniquet with direct pressure) – this is a professional-level decision under medical direction only.

Equipment needed:

— CAT tourniquet (Combat Application Tourniquet) -- preferred
— Improvised tourniquet material (shirt, towel, belt) as last resort
— Scissors or Leatherman Raptor for cutting clothing
— Pen or marker to document tourniquet application time
— Phone to call 101 (MDA)

MDA Progressive Hemorrhage Control

Step	Method
1	Direct pressure with hand or gauze
2	Israeli bandage (amtza'sh) for sustained pressure
3	Wound packing
4	Pressure points
5	Tourniquet (last resort or immediate in mass casualty)

In multi-casualty or active-threat situations, skip directly to tourniquet. 500,000 Israeli civilians trained by MDA in 2024.

Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Bleeding	דימום	di-MUM
Israeli bandage	אמצ"ש	am-TZA-sh
Wound	פצע	PE-tsa
Pressure	לחץ	LA-khats
Tighten	להדק	le-ha-DEK
Time	זמן / שעה	ZMAN / sha-A
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
There is heavy bleeding	יש דימום חזק	YESH di-MUM kha-ZAK
I applied a tourniquet	הנחתי חוסם עורקים	hi-NAKH-ti kho-SEM or-KIM

Source: Magen David Adom (MDA) — MDA First Aid Guidelines and "First Seven Minutes" initiative
URL: https://www.mdais.org/media/4954/mda-first-aid-guidelines-en.pdf
Publication date: Standing public guidance
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution