



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Continued Monitoring



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT lay the person down** — the upright position is essential for breathing.
✕ **Do NOT leave them alone** during an attack.
✕ **Do NOT give controller medications** (inhaled corticosteroids) for acute relief — no immediate effect.
✕ **Do NOT delay calling 101** if no inhaler is available.
✕ **Do NOT ignore a 'silent chest'** — absence of wheezing during obvious distress = life-threatening.

Equipment needed:


— Rescue inhaler (Ventolin / Salbutamol)
— Spacer device (merkhav) if available

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Inhaler	משאף	mash-AF
Spacer	מרחב	mer-KHAV
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
Asthma	אסתמה	AST-ma
Asthma attack	התקף אסתמה	het-KEF AST-ma
Wheezing	צפצופים	tsif-tsu-FIM
Chest tightness	לחץ בחזה	LA-khats ba-kha-ZE
Puffs (doses)	שאיפות	she-i-FOT
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Can you breathe?	אתה יכול לנשום?	a-TA ya-KHOL lin-SHOM?
Where is the inhaler?	איפה המשאף?	EI-fo ha-mash-AF?

Source: Magen David Adom (MDA) — MDA ALS Protocol Manual 2022
URL: https://www.mdais.org/101/first-aid
Publication date: 2022-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Daniel Rosehill · share with attribution


CHECK responsiveness and breathing (max 10 s).
Gasping is NOT normal breathing.
Unresponsive and not breathing normally?
N
Recovery position if unconscious but breathing.
Provide appropriate first aid.
CALL 101 immediately. Use speakerphone.
Dispatcher can unlock nearest smart stand AED.
BEGIN CPR immediately.
100–120 compressions/min 5–6 cm depth. Full recoil.
Another person available to retrieve AED?
N
If AED is nearby, retrieve quickly. Otherwise continue CPR.
Y
SEND them for AED. Continue CPR.
LOCATE AED: Eifo-Defi app, MDA smart stands, public buildings.

AED Pad Placement & Shock Delivery



Continued AED / CPR Cycles



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT use AED on a conscious, breathing person** – AEDs are only for cardiac arrest.
- ✕ **Do NOT apply pads in water** – move to dry ground and dry the chest first.
- ✕ **Do NOT use near flammable gases** – risk of fire or explosion.
- ✕ **Do NOT place pads directly over a pacemaker** – place at least 2.5 cm away.
- ✕ **Do NOT place pads over medication patches** – remove and wipe skin first.
- ✕ **Do NOT touch patient during analysis or shock delivery** – electrocution risk.
- ✕ **Do NOT turn off AED or disconnect pads** – even if person regains consciousness.
- ✕ **Do NOT delay defibrillation** – survival drops 7–10% per minute of delay.
- ✕ **Do NOT interrupt compressions for more than 10 seconds**.
- ✕ **Do NOT let pads touch or overlap** on the chest.
- ✕ **Do NOT use AED on a breathing, pulsed unconscious person** – use recovery position instead.

Equipment needed:

— AED (from MDA smart stands, public locations, or Eifo-Defi app)
— Razor (for shaving chest hair -- often in AED kits)
— Towel or cloth (for drying the chest)
— Scissors (for cutting clothing -- often in AED kits)
— Phone (to call 101 and use speakerphone)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
Stand clear!	התרחק!	hit-rakh-KU!
Cardiac arrest	דום לב	dom lev
Chest compressions	עיסויי חזה	i-su-YEI kha-ZE
Shock	הלם / שוק	HE-lem / shok
Not breathing	לא נושם	lo no-SHEM
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
AED	דפיברילטור חיצוני	de-fib-ri-LA-tor khi-tso-NI
Smart stand	עמדה חכמה	am-DA kha-kha-MA


Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — AED Operation Protocol / AHA 2025 Guidelines
URL: https://www.mdais.org/kursim/defibrillator
Publication date: 2025-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution


👁️ RECOGNISE signs: rash, swelling, breathing difficulty, weak pulse, confusion.
Signs of anaphylaxis? (breathing, circulation, or consciousness affected)
N
👁️ Mild reaction only. Monitor for worsening. Call 101 if deteriorates.
📱 CALL 101 immediately. Speakerphone on. Identify allergen if possible.
REMOVE offending agent. Scrape bee stinger sideways. Stop any infusion.
🌬️ MANAGE airway. Clear mouth. Loosen clothing around neck and chest.
Is the patient conscious?
Y
🧑 Breathing difficulty: SIT upright. Shock/hypotension: LIE down, elevate legs.
🌬️ Breathing?
N
Y
N
PLACE in recovery position. Proceed to EpiPen.
❤️ BEGIN CPR immediately.

EpiPen & Monitoring



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT delay calling 101** – anaphylaxis is life-threatening.
✕ **Do NOT delay epinephrine** – it is the only effective first-line treatment. Over 50% of deaths occur within 60 minutes.
✕ **Do NOT substitute antihistamines for epinephrine** – antihistamines are not effective against anaphylaxis.
✕ **Do NOT make a hypotensive patient stand or sit up suddenly** – this can cause cardiac arrest.
✕ **Do NOT leave the patient unattended** after epinephrine – symptoms may recur (biphasic reaction).
✕ **Do NOT assume safety after symptom resolution** – hospital evaluation is mandatory.
✕ **Do NOT waste time removing clothing** – EpiPen can be administered through trousers.
✕ **Do NOT inject EpiPen into a vein** – intramuscular outer thigh only.
✕ **Do NOT give food or drink** to a patient experiencing anaphylaxis.

Equipment needed:

— Epinephrine auto-injector (EpiPen) -- adult 0.3 mg
— Phone to call MDA at 101

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Allergic reaction	תגובה אלרגית	tgu-VA a-LER-git
Anaphylaxis	אנפילקסיס	a-na-fi-LAK-sis
EpiPen	אפיפן	E-pi-pen
Swelling	נפיחות	ne-fi-KHUT
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
Rash / hives	פריחה / סרפדת	pri-KHA / sar-PE-det
Injection	זריקה	zri-KA
Thigh	ירך	ya-REKH
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you allergic?	יש לך אלרגיה?	yesh le-KHA a-ler-GI-ya?
Where is the EpiPen?	איפה האפיפן?	EI-fo ha-E-pi-pen?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — ALS Protocol Manual April 2024
URL: https://www.mdais.org/101/allergy
Publication date: 2024-04-01
Imported: 2026-03-15 · **Last verified:** 2026-03-15


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Secondary Assessment & Monitoring



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT approach** until security forces confirm the scene is safe – secondary devices are a known tactic.
✕ **Do NOT remove embedded shrapnel or debris** – may cause uncontrollable bleeding.
✕ **Do NOT remove a tourniquet** once applied – only medical professionals should decide.
✕ **Do NOT remove clothing stuck to burns** – cut around it.
✕ **Do NOT let the victim move or walk** until MDA arrives.
✕ **Do NOT use head-tilt for airway management** in blast/trauma – use jaw-thrust to protect cervical spine.
✕ **Do NOT administer epinephrine in hypovolemic trauma** – blood vessels are already constricted from shock.
✕ **Do NOT assume only one injury site** – blast victims have injuries from multiple mechanisms.

Equipment needed:

— Phone to call 101 (MDA)
— Sterile gauze or clean cloth for direct pressure
— Israeli bandage (emergency bandage)
— Gauze for wound packing
— Tourniquet (CAT preferred; improvised as last resort)
— Non-adherent dressings for burns
— Clean water for cooling burns
— Blanket for warmth


Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Explosion	פיצוץ	pi-TSUTS
Shrapnel	רסיסים	rsi-SIM
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Bandage	תחבושת	takh-BO-shet
Bleeding	דימום	di-MUM
Pressure	לחץ	LA-khats
Wound	פצע	PE-tsa
Burns	כוויות	kvi-YOT
Shock	הלם	HE-lem
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you injured?	אתה פצוע?	a-TA pa-TSU-a?
Where does it hurt?	איפה כואב?	EI-fo ko-EV?

Source: Magen David Adom (MDA) — First Aid Guidelines: Blast and Shrapnel Injuries
URL: https://www.mdais.org/media/4954/mda-first-aid-guidelines-en.pdf
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Personal reference only — not medical advice
Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Chemical & Electrical Burns / Monitoring



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply ice or ice-cold water** – extreme cold causes additional tissue and blood vessel damage.
✕ **Do NOT apply toothpaste, coffee, egg, yogurt, butter, or home remedies** – these trap heat and increase infection risk.
✕ **Do NOT rupture or pop blisters** – the fluid protects underlying tissue during healing.
✕ **Do NOT use fat-based creams or ointments** on fresh burns.
✕ **Do NOT remove clothing stuck to the burn** – cut around it instead.
✕ **Do NOT attempt chemical neutralisation** – do not apply acid to base burn or vice versa.
✕ **Do NOT approach electrical burn casualty** until power source is confirmed disconnected.

Equipment needed:

— Cool running water source
— Cling film (food wrap) or clean plastic bag
— Clean non-fluffy dressings or bandages
— Scissors (to cut clothing around stuck areas)
— Blanket (to keep patient warm)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Burn	כוויה	kvi-YA
Water	מים	MA-yim
Cold water	מים קרים	MA-yim ka-RIM
Blister	שלפוחית	shal-pu-KHIT
Bandage	תחבושת	takh-BO-shet
Cling film	ניילון נצמד	NAI-lon nits-MAD
Chemical	כימיקל	khi-mi-KAL
Electricity	חשמל	KHASH-mal
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Does it hurt?	כואב לך?	ko-EV le-KHA?
Where is the burn?	איפה הכוויה?	EI-fo ha-kvi-YA?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Burns First Aid Guidelines
URL: https://www.mdais.org/101/burns
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Personal reference only — not medical advice


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Unconscious Choking & Self-Treatment



Self-Treatment if Choking Alone

LEAN over a firm object (chair back, countertop edge).
PRESS upper abdomen forcefully against the object.
Repeat until the object is expelled.
📱 **CALL** 101 as soon as you can speak.

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT perform abdominal thrusts on a person who can cough and speak** – you may worsen the obstruction.
✕ **Do NOT perform back blows on an adult** – MDA/Israeli protocol explicitly describes this as incorrect and potentially harmful for adults.
✕ **Do NOT perform blind finger sweeps** – only remove an object if you can clearly see it.
✕ **Do NOT give water or liquid** to a choking person.
✕ **Do NOT continue abdominal thrusts on an unconscious person** – transition to CPR.
✕ **Do NOT use abdominal thrusts on pregnant or very obese individuals** – use chest thrusts instead.

Note: Israeli Protocol Divergence


MDA/Israeli protocol **does not recommend back blows** for adults. This differs from AHA (2025) which recommends alternating 5 back blows and 5 abdominal thrusts, and from ERC (2025) which recommends 5 back blows first. Follow MDA guidance when in Israel.

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Choking	חנק	KHE-nek
Foreign body	גוף זר	guf zar
Can you breathe?	אתה יכול לנשום?	a-TA ya-KHOL lin-SHOM?
Can you speak?	אתה יכול לדבר?	a-TA ya-KHOL le-da-BER?
Cough	שיעול	shi-UL
Stomach / abdomen	בטן	BE-ten
Chest	חזה	kha-ZE
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you choking?	אתה נחנק?	a-TA nekh-NAK?


Source: Magen David Adom (MDA) — How to Treat Choking from a Foreign Body
URL: https://www.mdais.org/101/asphyxia
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Compressions, Breaths & AED



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT delay calling 101** – early activation is critical for survival.
- ✕ **Do NOT spend more than 10 seconds checking breathing** – delays reduce survival.
- ✕ **Do NOT interrupt compressions for more than 10 seconds** for any reason.
- ✕ **Do NOT lean on the chest between compressions** – allow full recoil.
- ✕ **Do NOT compress deeper than 6 cm** to avoid internal injury.
- ✕ **Do NOT over-ventilate** – 1 second per breath, just enough for visible chest rise.
- ✕ **Do NOT touch the person during AED analysis or shock delivery**.
- ✕ **Do NOT delay AED use** – apply as soon as it arrives.
- ✕ **Do NOT wait for AED to re-analyse** before resuming compressions after shock.
- ✕ **Do NOT move the person unnecessarily** – perform CPR where they are unless unsafe.
- ✕ **Do NOT assume gasping is normal breathing** – it is a sign of cardiac arrest.

Equipment needed:

— AED (Automated External Defibrillator)
— Barrier device or pocket mask (for rescue breaths, if available)
— Phone (to call 101 and use speakerphone for dispatcher-guided CPR)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
CPR	החייאה	ha-kha-ya-A
Cardiac arrest	דום לב	dom lev
Chest compressions	עיסויי חזה	i-su-YEI kha-ZE
Rescue breaths	הנשמות	han-sha-MOT
Are you okay?	אתה בסדר?	a-TA be-SE-der?
Not breathing	לא נושם	lo no-SHEM
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
Stand clear!	התרחקו!	hit-rakh-KU!
Recovery position	תנוחת התאוששות	tnu-KHAT hit-o-she-SHUT
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Call an ambulance!	הזמינו אמבולנס!	haz-MI-nu am-bu-LANS!

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — CPR Protocol / AHA 2025 Guidelines
URL: https://www.mdais.org/101/cpr-hands-only
Publication date: 2025-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


ASSESS severity.
Signs of heat stroke? (Temp above 40 C, hot/dry skin, NO sweating, confusion, unconscious)
Y
HEAT STROKE EMERGENCY. Go to page 2.
N
STOP all physical activity. Ensure complete rest.
MOVE to shaded, cool place. Air-conditioned room if available.
REMOVE excess clothing. Loosen tight garments.
REHYDRATE gradually with small sips of water or ORS.
ORS: 1L water + 6 tsp sugar 1. 0.5 tsp salt.
COOL with wet cloths, fan, or cool water spray.
Symptoms improved after 30 minutes of rest and rehydration?
N
CALL 101. May need IV fluids.
Y
Continue gradual rehydration and rest. Monitor for relapse.

Heat Stroke Emergency



Reference Notes

DO NOT:

✕ **Do NOT give large quantities of water quickly** – rapid fluid replacement disrupts electrolyte balance.
✕ **Do NOT give sugary drinks, coffee, or alcohol**.
✕ **Do NOT wet the head of a heat stroke patient** – choking risk if consciousness is altered.
✕ **Do NOT ignore the key distinction**: dehydration = continued sweating; heat stroke = ABSENT sweating with high body temperature.
✕ **Do NOT delay calling 101 for heat stroke** – it is a medical emergency requiring hospital treatment.
✕ **Do NOT give fluids to an unconscious person**.

Equipment needed:


— Water or ORS (Oral Rehydration Solution)
— Fan or method of creating air flow
— Wet cloths or towels for cooling
— Ice packs (for heat stroke)
— Straw (for controlled sipping)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Dehydration	התייבשות	hit-yab-SHUT
Heat stroke	מכת חום	ma-KAT khom
Water	מים	MA-yim
Shade	צל	tsel
Dizzy	סחרחורת	skhar-kho-RET
Sweating	הזעה	haz-a-A
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Drink water	שתה מים	shte MA-yim
Are you thirsty?	אתה צמא?	a-TA tsa-ME?
Come to the shade	בוא לצל	bo la-TSEL

Source: Magen David Adom (MDA) — Dehydration and Heat-Related Illness First Aid
URL: https://www.mdais.org/101/first-aid
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution


🔍 ASSESS scene. Check for medical ID, insulin pump, glucometer.
Glucometer available?
N
ASSESS by signs. If unsure, treat as hypoglycemia (safer).
Y
CHECK blood glucose. Below 70 = hypo. Above 250 = hyper.
Signs of hypoglycemia? Sweating, pallor, trembling, confusion, rapid pulse?
N
Hyperglycemia pathway. See Page 3.
Y
Hypoglycemia pathway. Continue below.
Patient conscious and able to swallow?
N
Reduced or no consciousness. See Page 2.
Y
💊 GIVE sugar immediately. Juice, glucose tablets, glucogel, sugar cubes.
Improved within 10–15 minutes?
N
📱 CALL 101 immediately. Patient needs IV dextrose.
Y
👁 MONITOR. Give complex carbs (bread, crackers).


Unconscious Hypoglycemia



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Hyperglycemia Pathway



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT administer insulin** — insulin dosing requires medical expertise. Leave to MDA paramedics.
✕ **Do NOT give food or drink to an unconscious patient** — risk of choking and aspiration.
✕ **Do NOT give diet drinks or sugar-free foods** for hypoglycemia — they will not raise blood sugar.
✕ **Do NOT give sugary foods high in fat** (chocolate bars, ice cream) as first-line treatment — fat slows absorption.
✕ **Do NOT withhold sugar when in doubt** — giving sugar to a hypo patient is life-saving; giving sugar to a hyper patient will not cause significant immediate harm.
✕ **Do NOT leave the patient unattended** — condition can deteriorate rapidly.
✕ **Do NOT delay calling 101** if unconscious, seizing, or not improving after sugar.
✕ **Do NOT disconnect an insulin pump roughly** — disconnect gently to avoid damage.

Equipment needed:

— Phone to call 101 (MDA) or 1221 (United Hatzalah)
— Glucometer and test strips (if available)
— Rapidly absorbed sugar (glucose tablets, glucogel, juice, sugar cubes, honey)
— Complex carbohydrate for follow-up (bread, crackers)
— Glucagon injection kit (if available and trained)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Diabetes	סוכרת	su-KE-ret
Low blood sugar	סוכר נמוך	SU-kar na-MUKH
High blood sugar	סוכר גבוה	SU-kar ga-VO-ah
Sugar	סוכר	SU-kar
Insulin	אינסולין	in-su-LIN
Glucometer	מד סוכר	mad SU-kar
Juice	מיץ	mitz
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you diabetic?	אתה סוכרתי?	a-TA su-kar-TI?
Did you eat today?	אכלת היום?	a-KHAL-ta ha-YOM?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — How to Treat a Diabetes Patient
URL: https://www.mdais.org/101/diabetes
Publication date: 2026-03-15
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT enter the water** unless you are a trained swimmer — untrained rescuers frequently become victims.
✕ **Do NOT attempt to remove water from the lungs** — MDA states this is impossible, dangerous, and wastes critical time.
✕ **Do NOT perform the Heimlich manoeuvre** or turn the person upside down to "drain" water.
✕ **Do NOT use an AED on a wet patient** or wet surface — dry the chest and move to dry area first.
✕ **Do NOT allow a near-drowning victim to refuse hospital transport** — secondary drowning and electrolyte imbalances can be fatal hours later.
✕ **Do NOT stop CPR** until the person breathes on their own or MDA takes over.
✕ **Do NOT delay calling 101** — call immediately, even before attempting rescue if alone.

Equipment needed:

— AED (defibrillator) — commonly at lifeguard stations and public facilities
— Reaching aids (pole, rope, flotation device) for water rescue
— Dry blankets or clothing to keep victim warm after extraction

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Drowning	טביעה	tvi-A
Water	מים	MA-yim
Swimming pool	בריכה	bre-KHA
Beach	חוף	khof
Lifeguard	מציל	ma-TSIL
He is not breathing	הוא לא נושם	hu lo no-SHEM
CPR	החייאה	ha-kha-ya-A
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
Ambulance	אמבולנס	am-bu-LANS
Help!	הצילו!	ha-tsi-LU!
Call an ambulance	תזמין אמבולנס	taz-MIN am-bu-LANS
Recovery position	תנוחת התאוששות	tnu-KHAT hit-o-she-SHUT

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Drowning First Aid Guidance
URL: https://www.mdais.org/101/drowning
Publication date: 2026-03-15
Imported: 2026-03-15 · **Last verified:** 2026-03-16


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Burns, Spine, and Shock Assessment



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT touch the victim** while they may still be in contact with the electrical source — chain electrocution can kill the rescuer.
✕ **Do NOT use wet or damp objects** to separate the victim from the source.
✕ **Do NOT approach high-voltage incidents** — maintain at least 10 metres distance. Call 101, 102, and 103 (IEC).
✕ **Do NOT apply water to burns** while the victim is still connected to the electrical source.
✕ **Do NOT apply creams, ointments, or home remedies** to electrical burns.
✕ **Do NOT assume the injury is minor** based on small visible burns — internal damage along the current pathway may be severe.
✕ **Do NOT give food or drink** to an electrocution victim.
✕ **Do NOT perform CPR on a conscious victim.**

Equipment needed:

— Dry non-conducting object (wooden broom handle, dry rope, dry rubber mat)
— Dry insulating surface to stand on (dry wood, dry rubber mat, newspapers)
— AED (automated external defibrillator)
— Cool running water source (for burn cooling after scene is safe)
— Cling film or clean non-fluffy dressings (for burn covering)
— Blanket (for shock prevention)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Electrocution	התחשמלות	hit-khash-me-LUT
Electricity	חשמל	KHASH-mal
Electric shock	מכת חשמל	ma-KAT KHASH-mal
Power line	קו מתח	kav ME-takh
High voltage	מתח גבוה	ME-takh ga-VO-ah
Circuit breaker	מפסק חשמל	maf-SEK KHASH-mal
Burns	כוויות	kvi-YOT
He is unconscious	הוא מחוסר הכרה	hu me-khu-SAR ha-ka-RA
CPR	החייאה	ha-kha-ya-A
Ambulance	אמבולנס	am-bu-LANS
Fire department	כבאות	ka-ba-UT
Electric company	חברת חשמל	khev-RAT KHASH-mal

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Electrocution First Aid Guidelines
URL: https://www.mdais.org/101/electrocution
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Blunt Trauma / Laceration



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT rub the eye** when a foreign body is present.
✕ **Do NOT attempt to remove an embedded or penetrating object.**
✕ **Do NOT apply pressure** to an injured eye or embedded object.
✕ **Do NOT wash the eye** if there is a laceration to the eyeball.
✕ **Do NOT delay flushing for chemical splashes** — start immediately.
✕ **Do NOT press a cold compress onto the eyeball** for blunt trauma.

Equipment needed:


— Clean running water source for flushing
— Sterile eye pads or clean cloth
— Cold compress (for blunt trauma)
— Cup or cone (to protect embedded objects)
— Bandages to loosely cover both eyes if needed

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Eye	עין	A-yin
Eye injury	פציעת עין	ptsi-AT A-yin
Chemical burn	כוויה כימית	kvi-YA khi-MIT
Foreign body	גוף זר	guf ZAR
I cannot see	אני לא רואה	a-NI lo ro-E
It hurts	כואב לי	ko-EV li
Water	מים	MA-yim
Flush / rinse	שטיפה	shti-FA
Bandage	תחבושת	takh-BO-shet
Emergency room	חדר מיון	KHE-der mi-YUN
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM

Source: Magen David Adom (MDA) — Eye Injury First Aid
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-03-16
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution


📱 CALL 101 (MDA). Keep on speaker for guidance.
Signs of fracture? (deformity, severe pain, inability to move limb, crepitus)
N
Likely sprain. See Page 2: RICE.
Y
Fracture pathway. Continue below.
Bone visible or protruding through the skin? (open fracture)
Y
Open fracture. CONTROL bleeding: press around (not on) bone. COVER wound with sterile dressing. Monitor for shock.
N
IMMOBILISE the limb in position found. Minimum movement, maximum stability.
Arm or upper limb?
N
PAD both sides of leg with pillows or rolled blankets to prevent movement.
Y
SUPPORT with improvised sling (shirt, scarf). Attach to chest.
APPLY ice wrapped in cloth (15-min intervals). REMOVE rings, watches, jewellery from affected limb.
👁 MONITOR for shock (pale skin, rapid breathing). Keep warm until MDA arrives.

Sprain — RICE Protocol



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT attempt to straighten or realign a broken limb** — this may tear arteries and nerves.
✕ **Do NOT push a protruding bone back in** during an open fracture.
✕ **Do NOT apply ice directly to skin** — always wrap in cloth first.
✕ **Do NOT apply heat** to a fresh fracture or sprain — heat increases swelling.
✕ **Do NOT allow the patient to walk** on a suspected leg fracture.
✕ **Do NOT remove embedded objects** from a wound associated with a fracture.

Equipment needed:

— Triangular bandage or improvised sling (shirt, scarf)
— Ice pack or cooling spray (wrapped in cloth)
— Elastic bandage (for compression of sprains)
— Pillows, blankets, or rolled towels (for immobilisation and elevation)
— Sterile dressings or clean cloth (for open fracture wounds)
— Tourniquet (for severe bleeding in open fractures)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Fracture	שבר	SHE-ver
Broken bone	עצם שבורה	E-tsem shvu-RA
Sprain	נקע	NE-ka
Arm	יד / זרוע	yad / zro-A
Leg	רגל	RE-gel
It hurts	כואב לי	ko-EV li
I cannot move it	אני לא יכול להזיז	a-NI lo ya-KHOL le-ha-ZIZ
Sling	מתלה	mit-LE
Ice	קרח	KE-rakh
Bandage	תחבושת	takh-BO-shet
Ambulance	אמבולנס	am-bu-LANS
Where does it hurt?	איפה כואב לך?	EI-fo ko-EV le-KHA?

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — Fractures and Sprains First Aid Guidelines
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


⚠️ Immediate life-threatening danger? (fire, active shooting, structural collapse)
Y
MOVE only as last resort — log-roll keeping spine aligned.
📱 CALL 101 immediately. Report location, mechanism, consciousness, bleeding.
Isolated penetrating trauma (stab/gunshot) without blunt component?
Y
Spinal immobilisation not required. MANAGE wounds and bleeding.
N
👐 STABILISE head manually. Hands on both sides, cup the ears. Hold head neutral, aligned with body. Maintain until EMS arrives.
🌬️ Patient conscious and talking?
Y
Airway is open. Maintain stabilisation. Proceed to bleeding control (Page 2).
N
OPEN airway with jaw thrust only. Push jaw forward from behind angles. Do NOT tilt head.
❤️ BEGIN CPR with spinal precautions. Use jaw thrust for ventilations (2 responders).
N
🌬️ Patient breathing and has pulse?
Y
👁️ Maintain stabilisation and jaw thrust. Proceed to Page 2.

Bleeding, Assessment, and Monitoring



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT move the patient** unless there is immediate danger to life (fire, active threat, collapse).
- ✕ **Do NOT twist, flex, or extend the neck.**
- ✕ **Do NOT attempt to sit the patient up.**
- ✕ **Do NOT use head-tilt chin-lift** — use jaw thrust only.
- ✕ **Do NOT remove a helmet** without proper training and a second person.
- ✕ **Do NOT give food or drink.**
- ✕ **Do NOT leave the patient alone.**
- ✕ **Do NOT apply direct pressure to a suspected skull fracture.**
- ✕ **Do NOT plug ears or nose** if blood or clear fluid is draining.
- ✕ **Do NOT pull the patient by arms or legs.**
- ✕ **Do NOT extract from a vehicle** unless the vehicle is on fire or there is another immediate threat.
- ✕ **Do NOT give oral fluids** even if dehydrated — risk of vomiting and aspiration.
- ✕ **Do NOT attempt to straighten the head** if the patient resists or reports pain — stabilise in position found.

Equipment needed:

— Disposable gloves
— Clean dressings or bandages for wound care
— Sterile dressing for skull fracture coverage
— Cervical collar (if available and properly sized)
— Blanket or thermal protection
— Tourniquet (CAT) for associated limb haemorrhage
— High-visibility vest (if responding at a road accident)


Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Head injury	פגיעת ראש	pgi-AT rosh
Spinal injury	פגיעת עמוד שדרה	pgi-AT a-MUD shid-RA
Concussion	זעזוע מוח	ziz-U-a MO-akh
Neck	צוואר	tsa-VAR
Spine	עמוד שדרה	a-MUD shid-RA
Do not move	לא לזוז	lo la-ZUZ
Can you feel your legs?	אתה מרגיש את הרגליים?	a-TA mar-GISH et ha-rag-LA-yim?
Can you move your fingers?	אתה יכול להזיז אצבעות?	a-TA ya-KHOL le-ha-ZIZ ets-ba-OT?
Helmet	קסדה	kas-DA
Recovery position	תנוחת התאוששות	tnu-KHAT hit-o-she-SHUT
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM

Source: Magen David Adom (MDA) — BLS Guide 2016 / ALS Professional Manual 2024
URL: https://www.mdais.org/101/first-aid
Publication date: 2024-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice
Daniel Rosehill · share with attribution


📱 CALL 101 immediately. State suspected heart attack. Stay on line for guidance.
🧑 STOP all physical activity. POSITION half-sitting with back supported, knees slightly bent. LOOSEN tight clothing.
💊 Patient allergic to aspirin?
Y
Skip aspirin. Inform MDA team about allergy.
N
💊 GIVE 100–300 mg aspirin. Patient must CHEW (not swallow whole). Standard Israeli tablet = 100 mg.
Patient has their own prescribed nitroglycerin?
Y
HELP patient take their nitroglycerin as prescribed.
👁️ MONITOR continuously. Stay with patient at all times. Send bystander to fetch AED.
👁️ Continue monitoring. Keep half-sitting. Keep calm.
N
🌬️ Patient loses consciousness and stops breathing normally?
Y
Cardiac arrest. See Page 2: CPR / AED.

CPR and AED — Cardiac Arrest



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT allow the patient to walk, exert, or dismiss symptoms.**
✕ **Do NOT give food or beverages.**
✕ **Do NOT leave the patient alone** (except briefly to call for help).
✕ **Do NOT give aspirin if the patient is allergic.**
✕ **Do NOT give nitroglycerin** not prescribed to this specific patient.
✕ **Do NOT wait to see if symptoms resolve** — every minute of delay increases irreversible heart damage.
✕ **Do NOT drive the patient to hospital** — call 101 and wait for ambulance. MDA has cardiac monitoring equipment.
✕ **Do NOT substitute paracetamol (Acamol) for aspirin** — only aspirin provides the antiplatelet effect.
✕ **Do NOT give aspirin as a whole tablet** — it must be chewed for rapid absorption.

Equipment needed:

— Aspirin tablets (100 mg, 1–3 tablets for 100–300 mg total)
— AED (Automated External Defibrillator) — if available
— Phone (to call 101 MDA and optionally 1221 United Hatzalah)
— Nitroglycerin (only if prescribed to the patient)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Heart attack	התקף לב	het-KEF lev
Chest pain	כאב בחזה	ke-EV ba-kha-ZE
Heart	לב	lev
Aspirin	אספירין	as-pi-RIN
Chew	ללעוס	lil-OS
Nitroglycerin	ניטרוגליצרין	nit-ro-gli-tse-RIN
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
CPR	החייאה	ha-kha-ya-A
He is not breathing	הוא לא נושם	hu lo no-SHEM
Are you allergic to aspirin?	יש לך אלרגיה לאספירין?	yesh le-KHA a-ler-GI-ya le-as-pi-RIN?
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Heart Attack Protocol / ALS Protocol 2024 / BLS Guide 2016
URL: https://www.mdais.org/101/mi-signs
Publication date: 2024-04-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16


Personal reference only — not medical advice

Daniel Rosehill · share with attribution




Daniel Rosehill · share with attribution

Heat Stroke — Continued




Heat Exhaustion — Continued



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Continued Monitoring — Heat Stroke



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT administer fever-reducing medications** (paracetamol, ibuprofen) — antipyretics do not work for environmental hyperthermia and may cause harm.
✕ **Do NOT give fluids** to a person who is not fully conscious or who cannot swallow safely.
✕ **Do NOT delay calling 101** for suspected heat stroke.
✕ **Do NOT ignore early signs of heat exhaustion** — it can progress to life-threatening heat stroke.
✕ **Do NOT leave the person unattended** while waiting for emergency services.

Equipment needed:


— Wet towels or cloths
— Cool/cold water (for spraying, pouring, and drinking)
— Cold packs or ice packs (if available)
— Fan or other means of air circulation (if available)

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Heatstroke	מכת חום	ma-KAT KHOM
Heat exhaustion	תשישות חום	tshi-SHUT KHOM
Shade	צל	TSEL
Water	מים	MA-yim
Confused	מבולבל	me-vul-BAL
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Cool down	לקרר	le-ka-RER
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Can you drink?	אתה יכול לשתות?	a-TA ya-KHOL lish-TOT?
Move to shade	בוא לצל	BO la-TSEL

Source: Magen David Adom (MDA) — MDA First Aid Guidelines, Heat-Related Illness
URL: https://www.mdais.org/101/first-aid
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution


📱 CALL 101 immediately. Hypothermia can deteriorate rapidly.
🧑 MOVE to warm, dry shelter. Shield from wind, rain, cold.
REMOVE wet clothing. Replace with dry layers.
COVER with blankets including the head. Warm room to 22 C if indoors.
🌬️ Conscious and able to swallow?
N
No fluids.
👁️ MONITOR breathing closely. Prepare for CPR.
Y
💦 PROVIDE warm (not hot) beverages. No alcohol. No caffeine.
🔥 APPLY heat packs to chest, abdomen, back ONLY.
Use cloth barrier on skin.
🌬️ Person stopped breathing?
N
👁️ CONTINUE gradual rewarming until MDA arrives.
Y
BEGIN CPR immediately.

Reference Notes

DO NOT:

✕ **Do NOT rub extremities** — friction can cause tissue damage and cardiac complications.
✕ **Do NOT use direct intense heat** (hot water baths, heating pads directly on skin) — rapid rewarming can cause dangerous cardiac arrhythmias.
✕ **Do NOT give alcohol** — it dilates blood vessels and accelerates heat loss.
✕ **Do NOT give caffeine** — it worsens hypothermia.
✕ **Do NOT warm extremities first** — cold blood rushing to the heart can trigger cardiac arrest (afterdrop).
✕ **Do NOT apply vigorous external rewarming** — gradual rewarming is essential.

Equipment needed:


— Blankets or thermal blankets (space blankets)
— Dry clothing
— Heat packs with cloth barrier
— Warm (not hot) beverages (for conscious patients only)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Hypothermia	היפותרמיה	hi-po-TER-mi-ya
Cold exposure	חשיפה לקור	kha-shi-FA la-KOR
Shivering	רעד	RA-ad
Blanket	שמיכה	smi-KHA
Warm drink	משקה חם	mash-KE KHAM
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נושם	no-SHEM
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you cold?	קר לך?	KAR le-KHA?
Cover him/her	תכסו אותו/אותה	tekh-SU o-TO / o-TA

Source: Magen David Adom (MDA) — MDA Emergency Medicine Manual 2024
URL: https://www.mdais.org/101/first-aid
Publication date: 2024-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Personal reference only — not medical advice
Daniel Rosehill · share with attribution


⚠️ ENSURE scene safety. Protect yourself from contamination.
📱 CALL 101. CALL Poison Center: 04-7771900.
PRESERVE substance container, packaging, remaining material.
Was the poison ingested (swallowed)?
Y
Ingestion pathway. See Page 2.
N
Was the poison inhaled (fumes, gas)?
Y
MOVE to fresh air. See Page 2.
N
Skin or eye contact?
Y
💦 FLUSH with running water 15–20 min. Remove contaminated clothing.
N
Other route (injection, bite). MONITOR and await MDA.

Ingestion and Inhalation Pathways



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT induce vomiting** unless specifically directed by the Poison Information Center (04-7771900) or a physician.
✕ **Do NOT give food or drink** unless directed by medical professionals (exception: water/milk for conscious patients who ingested caustic substances).
✕ **Do NOT attempt to neutralise the poison** (e.g., giving acid for base ingestion or vice versa).
✕ **Do NOT use home remedies** (salt water, mustard, raw eggs, etc.).
✕ **Do NOT give activated charcoal** without medical direction.
✕ **Do NOT enter a confined space** with potential toxic fumes without proper protection.
✕ **Do NOT delay calling for help** — even mild symptoms can deteriorate rapidly.
✕ **Do NOT administer antidotes** — this is a hospital-level intervention only.
✕ **Do NOT leave the patient alone.**

Equipment needed:


— Gloves (to protect against contamination)
— Running water source (for skin or eye decontamination)

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Poisoning	הרעלה	har-a-LA
Poison	רעל	RA-al
Swallowed	בלע	ba-LA
Vomiting	הקאה	ha-ka-A
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נושם	no-SHEM
Cleaning product	חומר ניקוי	KHO-mer ni-KUI
Poison Center	מרכז הרעלות	mer-KAZ har-a-LOT
Ambulance	אמבולנס	am-bu-LANS
What did they swallow?	מה הוא בלע?	MA HU ba-LA?
How much?	כמה?	KA-ma?


Source: Magen David Adom (MDA) — Poisoning First Aid Guidance
URL: https://www.mdais.org/101/poisoningn
Poison Center: 04-7771900 (Rambam Health Care Campus, Haifa — 24/7)
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Placing in Recovery Position



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT place in recovery position if the person is not breathing** — begin CPR instead.
✕ **Do NOT move a person with suspected spinal injury** unless they are in immediate danger.
✕ **Do NOT leave the person unattended** — monitor breathing continuously.
✕ **Do NOT place on their back if unconscious and breathing** — risk of airway obstruction and aspiration.
✕ **Do NOT forget to call 101** — an unconscious person always requires professional medical evaluation.

Equipment needed:


— Blanket (to keep the person warm)
— Phone to call MDA at 101

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Recovery position	תנוחת התאוששות	ta-nu-KHAT hit-o-she-SHUT
Lateral position	תנוחת צד	ta-nu-KHAT TSAD
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נושם	no-SHEM
Airway	נתיב אוויר	na-TIV a-VIR
Are you OK?	אתה בסדר?	a-TA be-SE-der?
He/she is not responding	הוא/היא לא מגיב/ה	HU/HI LO ma-GIV/ma-gi-VA
Roll onto side	הפוך על הצד	ha-FOKH al ha-TSAD
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM

Source: Magen David Adom (MDA) — MDA Emergency Medicine Manual 2024; BLS Guide 2016
URL: https://www.mdais.org/101/first-aid
Publication date: 2024-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply a tourniquet** — it traps venom locally and causes tissue damage.
✕ **Do NOT suck the venom out** — this is ineffective and potentially harmful.
✕ **Do NOT cut the sting area** — increases infection risk and does not remove venom.
✕ **Do NOT apply ice or cool the sting area** — Israeli Ministry of Health (2026) explicitly advises against cooling.
✕ **Do NOT give the victim food or drink** — risks venom dispersal and may complicate hospital treatment.
✕ **Do NOT allow the victim to walk or move the stung limb** — movement accelerates venom circulation.

Equipment needed:


— Phone to call 101 (MDA) and optionally 04-7771900 (Poison Center)
— Phone or camera to photograph the scorpion if safe to do so

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Scorpion	עקרב	ak-RAV
Scorpion sting	עקיצת עקרב	a-ki-TSAT ak-RAV
Deathstalker (yellow scorpion)	עקרב צהוב	ak-RAV tsa-HOV
Venom	ארס	E-res
Antivenom	נוגדן	nog-DAN
Swelling	נפיחות	ne-fi-KHUT
Pain	כאב	KE-ev
Poison Center	מרכז הרעלות	mer-KAZ har-a-LOT
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Where were you stung?	איפה נעקצת?	EI-fo ne-ek-TSATS-ta?
Stay still	תישאר במקום	ti-sha-ER ba-ma-KOM


Source: Magen David Adom (MDA) — First Aid for Scorpion Stings; Israeli Ministry of Health
URL: https://www.mdais.org/101/first-aid
Poison Center: 04-7771900 (Rambam Health Care Campus, Haifa — 24/7)
Note: All scorpion sting victims must be observed in ED for minimum 6 hours.
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Post-Seizure Assessment



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT restrain the person** or try to hold them down — you cannot stop a seizure by force, and restraining can cause fractures or dislocations.
✕ **Do NOT put anything in the person's mouth** — no spoons, wallets, sticks, or fingers. The person cannot swallow their tongue. Objects risk breaking teeth.
✕ **Do NOT attempt mouth-to-mouth during the seizure** — people usually resume breathing once the seizure ends.
✕ **Do NOT pour water on the person** or attempt to "wake them up".
✕ **Do NOT offer food, drinks, or medication by mouth** until the person is fully alert and able to swallow safely.
✕ **Do NOT crowd around the person** — give them space and privacy. Ask bystanders to step back.
✕ **Do NOT move the person** unless they are in immediate danger (road, ledge, fire, water).

Equipment needed:

— Soft padding for under the head (jacket, towel, cushion, blanket)
— Watch or phone to time the seizure duration
— Prescribed rescue medication if available (Buccolam, Stesolid, or Rivotril — prescription only)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Seizure	התקף	het-KEF
Epilepsy	אפילפסיה	e-pi-LEP-si-ya
Convulsions	פרכוסים	pir-ku-SIM
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נושם	no-SHEM
Recovery position	תנוחת התאוששות	ta-nu-KHAT hit-o-she-SHUT
How long?	כמה זמן?	KA-ma ZMAN?
First time?	פעם ראשונה?	PA-am ri-sho-NA?
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Epilepsy Association	האגודה לאפילפסיה	ha-a-gu-DA le-e-pi-LEP-si-ya

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — First Aid Provider Course 2026; ALS Manual 2024
URL: https://www.mdais.org/101/first-aid
Israeli Epilepsy Association: 02-5000283
Note: Status epilepticus = seizure > 5 minutes = medical emergency.
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


⚠️ ENSURE scene safety. Wear gloves if available.
📱 CALL 101 immediately.
Embedded or impaled object in the wound?
Y
STABILISE object in place with padding. Apply pressure around it, not on it. Skip to shock monitoring.
N
👐 APPLY strong, direct pressure to the wound with gloved hand or sterile gauze.
If dressing soaks through: ADD layers on top. Never remove original.
Direct pressure controlled bleeding?
Y
MAINTAIN pressure. ELEVATE limb above heart level if possible. Monitor for shock.
N
APPLY Israeli Emergency Bandage (pressure dressing). Pad over wound, wrap through pressure bar, secure.
Israeli bandage controlled bleeding?
Y
MAINTAIN dressing. Monitor for shock.
N
Escalate: see Page 2 (tourniquet / wound packing).

Escalation — Tourniquet / Wound Packing



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT remove an embedded or impaled object** from the wound.
- ✕ **Do NOT lift or remove a blood-soaked dressing** to check the wound — add layers on top.
- ✕ **Do NOT remove or loosen a tourniquet** once applied unless instructed by a paramedic or physician.
- ✕ **Do NOT give food or drink** to a casualty with severe bleeding or suspected shock.
- ✕ **Do NOT use wound packing for penetrating stab wounds.**
- ✕ **Do NOT enlarge wound openings** to facilitate wound packing.
- ✕ **Do NOT assume a rapid pulse is caused only by anxiety** — suspect significant blood loss.
- ✕ **Do NOT delay calling 101** for severe or uncontrolled bleeding.
- ✕ **Do NOT apply a tourniquet directly over a joint** — place it 5-10 cm above the wound on the limb.

Equipment needed:

— Disposable gloves
— Sterile gauze or clean cloth
— Israeli Emergency Bandage (pressure dressing)
— Tourniquet (CAT or equivalent)
— Wound packing material (hemostatic gauze preferred)
— Triangular bandage (supplementary)
— Blanket or coat (for shock management)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Bleeding	דימום	di-MUM
Severe bleeding	דימום חמור	di-MUM kha-MUR
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Pressure bandage	תחבושת לחץ	takh-BO-shet LA-khats
Israeli bandage	אמצעי חבישה ישראלי	em-tsa-I kha-vi-SHA yis-re-e-LI
Direct pressure	לחץ ישיר	LA-khats ya-SHIR
Wound	פצע	PE-tsa
Shock	הלם	HE-lem
Gloves	כפפות	kfa-FOT
Ambulance	אמבולנס	am-bu-LANS
Press here hard	לחץ כאן חזק	le-KHATS KAN kha-ZAK
Where is the bleeding?	איפה הדימום?	EI-fo ha-di-MUM?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — How to Stop Bleeding (MDA 101 Public Guidance)
URL: https://www.mdais.org/101/bleeding
Imported: 2026-03-15 · **Last verified:** 2026-03-15


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Continued Care



Reference Notes

DO NOT:

- ✕ **Do NOT give food or drink** – moisten lips only if thirsty.
- ✕ **Do NOT elevate legs** if head, neck, or spinal injury is suspected.
- ✕ **Do NOT elevate legs** if the patient has a broken hip or leg.
- ✕ **Do NOT elevate legs** if it causes pain or further injury.
- ✕ **Do NOT elevate legs** if the patient has difficulty breathing – keep semi-upright.
- ✕ **Do NOT overheat** the patient when maintaining body temperature.
- ✕ **Do NOT move** a patient with suspected spinal injury unnecessarily.
- ✕ **Do NOT delay calling 101** – shock is a life-threatening emergency.
- ✕ **Do NOT remove a tourniquet** once applied – leave removal to MDA paramedics.
- ✕ **Do NOT leave** the patient unattended.

Equipment needed:

— Tourniquet -- widely available in Israeli first aid kits
— Israeli bandage (emergency bandage / amtza'sh) -- for wound packing and pressure
— Blanket or coat -- to maintain body temperature
— Ground insulation -- to protect from cold ground
— EpiPen (epinephrine auto-injector) -- for anaphylactic shock if prescribed

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Shock	הלם	HE-lem
Bleeding	דימום	di-MUM
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Israeli bandage	אמצ"ש	am-TZA-sh
Pulse	דופק	DO-fek
Blood pressure	לחץ דם	LA-khats DAM
Blanket	שמיכה	smi-KHA
Conscious	בהכרה	be-ha-ka-RA
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Raise the legs	להרים את הרגליים	le-ha-RIM et ha-rag-LA-yim

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — MDA-Approved First Aid Curriculum and BLS Guide
URL: https://www.mdais.org/101/first-aid
Publication date: Standing public guidance
Imported: 2026-03-15 · **Last verified:** 2026-03-15


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Tear Gas Exposure



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT re-enter** a smoke-filled or burning structure without proper firefighting equipment and training.
✕ **Do NOT rub eyes** after tear gas exposure – this worsens irritation and spreads the chemical.
✕ **Do NOT use milk** or other substances to flush eyes – use only clean water or saline due to infection risk.
✕ **Do NOT remove clothing** stuck to burns.
✕ **Do NOT assume stability** – symptoms may worsen over hours as airway swelling develops.
✕ **Do NOT leave** a smoke inhalation patient alone – continuous monitoring is required.
✕ **Do NOT ignore** soot around the mouth/nose or singed facial hair – these indicate potentially serious airway injury.

Equipment needed:

— Phone to call 101 (MDA) and 102 (Fire and Rescue)
— Clean water for eye flushing (tear gas) and burn cooling
— Clean cloth or non-adherent dressings for burns
— Blanket for warmth
— Soap for skin decontamination (tear gas)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Smoke	עשן	a-SHAN
Fire	אש / שריפה	ESH / sre-FA
Tear gas	גז מדמיע	GAZ mad-MI-a
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
Burns	כוויות	kvi-YOT
Fresh air	אוויר צח	a-VIR TSAKH
Water	מים	MA-yim
Blanket	שמיכה	smi-KHA
Ambulance	אמבולנס	am-bu-LANS
Fire department	כיבוי אש	ki-BUY ESH
Emergency	חירום	khe-RUM
Can you breathe?	אתה יכול לנשום?	a-TA ya-KHOL lin-SHOM?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — MDA First Aid Guidelines
URL: https://www.mdais.org/media/4954/mda-first-aid-guidelines-en.pdf
Publication date: Standing public guidance
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT apply a tourniquet** – it traps venom locally and causes severe tissue damage.
- ✕ **Do NOT suck the venom** out of the wound – this is ineffective and dangerous.
- ✕ **Do NOT cut or incise** the bite area – increases infection risk and does not help remove venom.
- ✕ **Do NOT cool or apply ice** to the bite area – can worsen tissue damage.
- ✕ **Do NOT give food or drink** – may complicate hospital treatment.
- ✕ **Do NOT induce vomiting.**
- ✕ **Do NOT give aspirin, ibuprofen, or naproxen** – viper venom is often anticoagulant.
- ✕ **Do NOT use alcohol-based disinfectants** on the wound.
- ✕ **Do NOT attempt to catch or kill the snake** – protected under Israeli law; risks additional bites.
- ✕ **Do NOT allow the victim to walk** or move the bitten limb – movement accelerates venom circulation.

Equipment needed:

— Splint or improvised immobilisation material
— Soap and clean water for wound cleaning
— Phone or camera to photograph the snake

Medically Dangerous Snakes in Israel

Species	Hebrew Name	Antivenom
Daboia palaestinae (Palestine viper)	צפע ארצישראלי	Available (Kamada Ltd.)
Echis coloratus (Painted saw-scaled viper)	אפעה צבעונית	Available (Kamada Ltd.)
Atractaspis engaddensis (Israeli mole viper)	שחור עין גדי	Not available

Peak season: spring/summer. Israel Poison Information Center: 04-7771900 (24/7, Rambam Hospital, Haifa)


Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Snake	נחש	na-KHASH
Snakebite	הכשת נחש	ha-ka-SHAT na-KHASH
Venom / poison	ארס	E-res
Swelling	נפיחות	ne-fi-KHUT
Splint	סד	SAD
Lie down	לשכב	lish-KAV
Calm down	להירגע	le-hi-ra-GE-a
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
I was bitten by a snake	ננשכתי על ידי נחש	ni-SHAKH-ti al ye-DEI na-KHASH

Source: Magen David Adom (MDA) — How to Treat a Snake Bite
URL: https://www.mdais.org/101/snake-bite
Publication date: Standing public guidance
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution


👐 REASSURE the victim. Move away from spider.
CLEAN bite with mild soap and water.
APPLY cool compress (wrapped cloth or ice pack). 15 min per hour.
ELEVATE bitten area above heart level. Minimise movement.
📷 Can the spider be safely photographed?
Y
PHOTOGRAPH and note features. Share with medical responders.
N
Report any observed details to paramedics.
Severe pain, muscle cramps, difficulty breathing, nausea, spreading redness, or systemic symptoms?
Y
📱 CALL 101 immediately. Also call Poison Center: 04-7771900. Proceed to Page 2.
N
👁️ MONITOR the bite. Call Poison Center (04-7771900) for guidance. Seek medical attention if symptoms worsen.

Species Assessment & Monitoring



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply a tourniquet** – spider venom is not stopped by tourniquets; causes tissue damage.
✕ **Do NOT cut, suck, or apply suction** to the bite wound – ineffective and increases infection risk.
✕ **Do NOT apply heat** to the bite – use only cool compresses.
✕ **Do NOT attempt to catch the spider** at risk of additional bites – photograph from a safe distance.
✕ **Do NOT excise a recluse bite** – early surgical excision is contraindicated and worsens outcomes.
✕ **Do NOT apply ice directly** to skin – wrap in cloth first.
✕ **Do NOT ignore mild symptoms** – recluse bites are often painless at first but progress to serious necrosis over hours to days.
✕ **Do NOT assume it is a spider bite** without confirmation – 83.5% of cases presenting as "spider bites" are actually other conditions.

Equipment needed:

— Phone to call 101 (MDA) and 04-7771900 (Israel Poison Center)
— Phone or camera to photograph the spider if safe
— Mild soap and clean water for wound cleaning
— Clean cloth or wrapped ice pack for cool compress

Medically Significant Spiders in Israel

Species	Venom Type	Antivenom
Mediterranean recluse (Loxosceles rufescens) "violin spider"	Necrotoxic – painless bite progressing to tissue necrosis	Not available
Mediterranean black widow (Latrodectus tredecimguttatus)	Neurotoxic – pain, muscle cramps, abdominal spasms	Available for severe cases

Israel Poison Information Center: 04-7771900 (24/7, Rambam Hospital, Haifa)


Daniel Rosehill · share with attribution


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Spider	עכביש	ak-a-VISH
Spider bite	עקיצת עכביש	a-ki-TSAT ak-a-VISH
Swelling	נפיחות	ne-fi-KHUT
Pain	כאב	ke-EV
Muscle cramps	התכווצויות שרירים	hit-kav-tsu-YOT shri-RIM
Poison center	מרכז רעלים	mer-KAZ re-a-LIM
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
I was bitten	ננשכתי	ni-SHAKH-ti

Source: Magen David Adom (MDA) — First Aid for Spider Bites
URL: https://www.mdais.org/media/4954/mda-first-aid-guidelines-en.pdf
Publication date: Standing public guidance
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Personal reference only — not medical advice
Daniel Rosehill · share with attribution


ASSESS using FAST:
F – Face: ask to smile, check for one-sided drooping.
A – Arms: extend both, one drifting down?
S – Speech: repeat a sentence, slurred or unable?
T – Time: r xact onset.
CALL 101 immediately. Even one FAST sign = emergency evaluation.
RECORD exact time of symptom onset. If woke with symptoms, record time sleep began.
Is the person conscious?
N
Breathing?
Y
BEGIN CPR immediately.
PLACE in recovery position (on side).
Y
LAY down with head slightly raised and supported.
LOOSEN restrictive clothing: scarves, ties, tight collars.



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Monitoring & Handover



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT give food or drink** – swallowing reflex may be impaired (dysphagia); aspiration risk.
✕ **Do NOT give any medications**, especially aspirin or blood thinners – approximately 15% of strokes are hemorrhagic, and these medications worsen bleeding.
✕ **Do NOT delay calling 101** – do not drive the person to hospital yourself unless unavoidable. The ambulance crew pre-notifies the hospital stroke team.
✕ **Do NOT ignore transient symptoms** – even if symptoms resolve (possible TIA), the person needs emergency evaluation. Approximately 40% of TIA patients have a full stroke within three months.
✕ **Do NOT allow eating, drinking, or oral medications** until a medical professional has assessed swallowing function.

Equipment needed:

— Phone to call 101 and record symptom onset time
— Pillow or support to slightly elevate the head

FAST Reference

Letter	Assessment
F	**Face** – Ask to smile. One side drooping?
A	**Arms** – Raise both arms. One drifting down?
S	**Speech** – Repeat a sentence. Slurred or unable?
T	**Time** – Note exact onset. Treatment window: 6 hours. Target: hospital within 60 min.

Even a single positive FAST sign = call 101 immediately.


Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Stroke	שבץ מוחי / אירוע מוחי	SHVATZ mo-KHI / ei-RU-a mo-KHI
Face	פנים	pa-NIM
Arm	זרוע	zro-A
Speech	דיבור	di-BUR
Smile	לחייך	le-kha-YEKH
Headache	כאב ראש	ke-EV ROSH
Dizziness	סחרחורת	skhar-KHO-ret
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Can you raise your arms?	אתה יכול להרים את הידיים?	a-TA ya-KHOL le-ha-RIM et ha-ya-DA-yim?
Can you speak?	אתה יכול לדבר?	a-TA ya-KHOL le-da-BER?


Source: Magen David Adom (MDA) — Stroke Recognition and First Aid
URL: https://www.mdais.org/101/stroke
Publication date: Standing public guidance
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Monitoring & Handover



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT remove the tourniquet** once applied – only medical professionals should decide on release or conversion.
✕ **Do NOT apply to junctional zones** (groin, armpit, neck), head, or abdomen – use direct pressure, wound packing, or pressure points for these areas.
✕ **Do NOT apply over a joint** – place on the limb above the wound, preferably over a single bone.
✕ **Do NOT loosen or adjust** to "check if bleeding has stopped" – this can restart hemorrhage.
✕ **Do NOT delay application** in life-threatening hemorrhage due to fear of limb loss – loss of a limb is preferable to loss of life.
✕ **Do NOT allow the victim to walk** or move the affected limb after tourniquet application.
✕ **Do NOT attempt tourniquet conversion** (replacing tourniquet with direct pressure) – this is a professional-level decision under medical direction only.

Equipment needed:

— CAT tourniquet (Combat Application Tourniquet) -- preferred
— Improvised tourniquet material (shirt, towel, belt) as last resort
— Scissors or Leatherman Raptor for cutting clothing
— Pen or marker to document tourniquet application time
— Phone to call 101 (MDA)

MDA Progressive Hemorrhage Control

Step	Method
1	Direct pressure with hand or gauze
2	Israeli bandage (amtza'sh) for sustained pressure
3	Wound packing
4	Pressure points
5	Tourniquet (last resort or immediate in mass casualty)

In multi-casualty or active-threat situations, skip directly to tourniquet. 500,000 Israeli civilians trained by MDA in 2024.


Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Bleeding	דימום	di-MUM
Israeli bandage	אמצ"ש	am-TZA-sh
Wound	פצע	PE-tsa
Pressure	לחץ	LA-khats
Tighten	להדק	le-ha-DEK
Time	זמן / שעה	ZMAN / sha-A
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
There is heavy bleeding	יש דימום חזק	YESH di-MUM kha-ZAK
I applied a tourniquet	הנחתי חוסם עורקים	hi-NAKH-ti kho-SEM or-KIM

Source: Magen David Adom (MDA) — MDA First Aid Guidelines and "First Seven Minutes" initiative
URL: https://www.mdais.org/media/4954/mda-first-aid-guidelines-en.pdf
Publication date: Standing public guidance
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

AED Analysis & Shock Cycle



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT use an AED on a conscious, normally-breathing child** – AEDs are only for cardiac arrest.
✕ **Do NOT use an AED in water** – move the child to dry ground first before applying pads.
✕ **Do NOT delay defibrillation** because paediatric pads are unavailable – use adult pads with anteroposterior placement.
✕ **Do NOT allow pads to touch or overlap** – use anteroposterior placement if the chest is too small.
✕ **Do NOT touch the child** during AED rhythm analysis or shock delivery.
✕ **Do NOT turn off the AED or remove pads** once applied, even if the child regains consciousness.
✕ **Do NOT perform adult-depth compressions** on a child or infant – compress to one-third of chest depth only.
✕ **Do NOT use an AED near flammable gases** or oxygen-enriched atmospheres.

Equipment needed:

— AED (Automated External Defibrillator)
— Paediatric AED pads (preferred for children under 8 / under 25 kg)
— Adult AED pads (if paediatric pads unavailable)
— Scissors or shears (to cut away clothing)
— Towel or cloth (to dry the chest)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
AED	מכשיר החייאה	makh-SHIR ha-kha-ya-A
Electrode pads	אלקטרודות	e-lek-TRO-dot
Shock	הלם / שוק	HE-lem / shok
Cardiac arrest	דום לב	dom lev
CPR	החייאה	ha-kha-ya-A
Stand clear!	התרחק!	hit-ra-KHU!
Child	ילד / ילדה	YE-led / yal-DA
Infant	תינוק / תינוקת	ti-NOK / ti-NO-ket
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Not breathing	לא נושם	lo no-SHEM

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — AED Operation for Children and Infants (AHA 2025 Guidelines as adopted by MDA)
URL: https://www.mdais.org/kursim/defibrillator
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


👁️ RECOGNISE
anaphylaxis.
Breathing difficulty, swelling, cardiovascular compromise?
Signs of anaphylaxis present? (breathing, circulation, or consciousness affected)
N
👁️ MONITOR
closely.
Be ready to escalate.
Y
PROCEED
immediately.
Do not wait for worsening.
📱 CALL 101 immediately.
State age, weight, allergy, EpiPen status.
REMOVE allergen if possible.
Food from mouth. Scrape bee stinger.
🌬️ MANAGE airway.
Loosen clothing. Check tongue.
Is the child conscious?
N
Breathing: recovery position.
Not breathing:
BEGIN CPR immediately.
Y
🧑 Breathing difficulty:
ALLOW upright position.
No difficulty: lie flat, elevate legs.

EpiPen Administration & Monitoring



EpiPen Dosing by Weight

15-30 kg: EpiPen Junior 0.15 mg (green label)
Over 30 kg: EpiPen 0.3 mg (yellow label)
Under 15 kg: No auto-injector for this range, but 0.15 mg Junior is still recommended over withholding treatment.
Note: Some Israeli allergists prescribe 0.3 mg from 20 kg – follow individual prescription.

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT delay epinephrine administration** – it is the only effective treatment for anaphylaxis.
✕ **Do NOT substitute antihistamines** (Benadryl/diphenhydramine) for epinephrine in anaphylaxis.
✕ **Do NOT force the child to lie down** if having breathing difficulty – allow upright position of comfort.
✕ **Do NOT over-extend an infant's neck** when managing the airway – use neutral position only.
✕ **Do NOT inject epinephrine into fingers, hands, feet, or buttocks** – outer thigh only.
✕ **Do NOT touch the orange tip** of the EpiPen.
✕ **Do NOT induce vomiting** if the allergen was ingested.
✕ **Do NOT allow a child with anaphylaxis to leave** without hospital evaluation – biphasic reactions can be fatal.
✕ **Do NOT store EpiPen in refrigerator, car, or direct sunlight** – Israeli heat makes degradation a particular risk.
✕ **Do NOT give oral medications** to a child with throat swelling or difficulty swallowing.

Equipment needed:

— EpiPen Junior 0.15 mg (green label) -- for children 15-30 kg
— EpiPen 0.3 mg (yellow label) -- for children over 30 kg
— Phone to call MDA at 101

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Allergic reaction	תגובה אלרגית	tgu-VA a-LER-git
Anaphylaxis	אנפילקסיס	a-na-fi-LAK-sis
EpiPen	אפיפן	E-pi-pen
Swelling	נפיחות	ne-fi-KHUT
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
Thigh	ירך	ya-REKH
Allergy	אלרגיה	a-ler-GI-ya
Bee sting	עקיצת דבורה	a-ki-TSAT dvo-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Does the child have an EpiPen?	יש לילד אפיפן?	yesh la-YE-led E-pi-pen?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — ALS Manual 2024 / BLS Guide 2016
URL: https://www.mdais.org/media/5033/als-2024.pdf
Publication date: 2024-04-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT lay the child down** – upright position is essential for breathing.
✕ **Do NOT give cold drinks** during an attack – cold can worsen bronchospasm.
✕ **Do NOT leave the child alone** during an attack.
✕ **Do NOT administer someone else's prescription inhaler** without medical direction.
✕ **Do NOT delay calling 101** if the child is not improving.
✕ **Do NOT ignore drowsiness** during an asthma attack – dangerous sign of CO2 buildup requiring immediate intervention.
✕ **Do NOT assume a quiet chest means improvement** – a silent chest during distress is a life-threatening sign.

Equipment needed:

— Rescue inhaler (Ventolin/salbutamol) -- child's own prescribed inhaler
— Spacer device (for children under 6)
— Supplemental oxygen and face mask (if available)
— Pulse oximeter (if available)
— Phone (to call 101)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Inhaler	משאף	mash-AF
Spacer	מרחב	mer-KHAV
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
Asthma	אסתמה	AST-ma
Asthma attack	התקף אסתמה	het-KEF AST-ma
Wheezing	צפצופים	tsif-tsu-FIM
Chest tightness	לחץ בחזה	LA-khats ba-kha-ZE
Puffs (doses)	שאיפות	she-i-FOT
Child	ילד / ילדה	YE-led / yal-DA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Can you breathe?	אתה יכול לנשום?	a-TA ya-KHOL lin-SHOM?
Where is the inhaler?	איפה המשאף?	EI-fo ha-mash-AF?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) – ALS Professional Manual (April 2024), supplemented by Ichud Hatzalah guidelines
URL: https://www.mdais.org/101/first-aid
Publication date: 2024-04-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Disability, Exposure & Ongoing Care

D — ASSESS neurological status (AVPU scale).
Check pupil size and reactivity.

↓

E — EXPOSE and examine for additional wounds, burns, embedded objects, fractures.

Then **COVER** immediately to prevent hypothermia.

↓

COVER open wounds with sterile dressings.
STABILISE embedded objects with padding around them.

↓

👐 **KEEP** child with parent or caregiver.
Speak calmly and honestly.

↓

👁️ **MONITOR** continuously.
Keep warm and covered.
Be prepared for CPR.

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT enter an unsafe blast scene** – risk of secondary devices and structural collapse.
- ✕ **Do NOT remove embedded objects** (shrapnel, fragments) – stabilise them in place.
- ✕ **Do NOT assume a quiet child is uninjured** – children may be in psychological shock.
- ✕ **Do NOT remove a tourniquet once applied** – hospital personnel will manage removal.
- ✕ **Do NOT forget to prevent hypothermia** after examination – children lose heat rapidly.
- ✕ **Do NOT separate the child from their parent** or caregiver unnecessarily.
- ✕ **Do NOT ignore asymmetric chest rise** – indicates thoracic trauma requiring urgent attention.
- ✕ **Do NOT give food or drink** if significant injuries present – may need anaesthesia at hospital.

Equipment needed:

— Tourniquet (CAT or equivalent -- see tourniquet protocol for child-specific sizing)
— Sterile gauze and bandages
— Blankets (for hypothermia prevention)
— Rigid splints (for fracture immobilisation)
— Phone (to call 101)
— Supplemental oxygen and mask (if available)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Explosion	פיצוץ	pi-TSUTS
Shrapnel	רסיסים	re-si-SIM
Rocket / missile	רקטה / טיל	ra-KE-ta / til
Shelter (safe room)	מקלט / ממ"ד	mik-LAT / ma-MAD
Bleeding	דימום	di-MUM
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Wound	פצע	PE-tsa
Breathing	נשימה	ne-shi-MA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you hurt?	אתה פצוע?	a-TA pa-TSU-a?
Where does it hurt?	איפה כואב?	EI-fo ko-EV?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) – PHTLS-aligned trauma protocols, supplemented by Ichud Hatzalah missile strike response guidelines
URL: https://www.mdais.org/101/first-aid
Publication date: 2024-04-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Personal reference only — not medical advice


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Chemical & Electrical Burns



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply ice or ice-cold water** – extreme cold causes additional tissue and blood vessel damage.
✕ **Do NOT apply toothpaste, coffee grounds, raw eggs, yogurt, or butter** – these create a heat-trapping layer and increase infection risk.
✕ **Do NOT puncture or pop blisters** – the fluid protects tissue during healing.
✕ **Do NOT apply ointments, oils, or powders** to serious burns before medical evaluation.
✕ **Do NOT use gauze pads or cotton wool directly on burns** – fibres adhere to damaged skin.
✕ **Do NOT remove clothing stuck to the burn** – cut around it instead.
✕ **Do NOT immerse the whole child in cold water** – cool the burn, not the child. High hypothermia risk.
✕ **Do NOT attempt chemical neutralisation** on chemical burns.
✕ **Do NOT leave the child unattended if unconscious** – call 101 immediately.

Equipment needed:

— Cool (lukewarm) running water source
— Sterile non-adherent dressings or clean dry sheet
— Scissors (to cut clothing around stuck areas)
— Blanket or towel (warmth during burn cooling)
— Age-appropriate pain relief (ibuprofen or paracetamol, dosed by weight)
— Drinking water (for hydration)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Burn	כוויה	kvi-YA
Scald	כוויה מנוזל חם	kvi-YA mi-no-ZEL kham
Cool water	מים פושרים	MA-yim posh-RIM
Blister	שלפוחית	shal-pu-KHIT
Bandage / dressing	תחבושת	takh-BO-shet
Pain	כאב	ke-EV
Child	ילד / ילדה	YE-led / yal-DA
Infant	תינוק / תינוקת	ti-NOK / ti-NO-ket
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
It hurts	כואב	ko-EV

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) – Burns First Aid Guidelines
URL: https://www.mdais.org/101/burns
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16
Key child difference: Minimum 20 minutes cooling (vs 10 for adults). Cool the burn, not the whole child. All burns in infants under 1 year require immediate medical attention.


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

CPR Transition (Unconscious Victim)



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT perform abdominal thrusts on infants under 1 year** – use only back blows and chest thrusts. Risk of organ damage.
✕ **Do NOT insert fingers blindly into the mouth** (blind finger sweeps) – only remove objects you can clearly see.
✕ **Do NOT force anything into the mouth** – risk of pushing the object deeper.
✕ **Do NOT give food or fluids** to a choking victim.
✕ **Do NOT intervene if the child is coughing effectively** – allow the body to clear the obstruction.
✕ **Do NOT use this protocol for choking caused by asthma, infection, or head injury** – seek immediate medical care.
✕ **Do NOT use excessive force** during abdominal thrusts that would lift the child off the ground.
✕ **Do NOT fully extend an infant's head** when opening the airway – use slight tilt position only.

Equipment needed:

— AED with paediatric pads (if available, for CPR transition)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Choking	חנק	KHE-nek
Foreign body	גוף זר	guf zar
Coughing	שיעול	shi-UL
Back blows	מכות גב	ma-KOT gav
Chest thrusts	לחיצות חזה	le-khi-TSOT kha-ZE
Abdominal thrusts	לחיצות בטן	le-khi-TSOT BE-ten
Breathing	נשימה	ne-shi-MA
CPR	החייאה	ha-kha-ya-A
Child	ילד / ילדה	YE-led / yal-DA
Infant	תינוק / תינוקת	ti-NOK / ti-NO-ket
The child is choking!	הילד נחנק!	ha-YE-led nekh-NAK!
Ambulance	אמבולנס	am-bu-LANS

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) – MDA 101 Public First Aid Guidance Series
URL: https://www.mdais.org/101/asphyxia
Imported: 2026-03-15 · **Last verified:** 2026-03-16
Notes: Infant (0-1 year), child (1-8 years). Physical size should guide technique selection, not age alone.


Personal reference only — not medical advice

Daniel Rosehill · share with attribution




Personal reference only — not medical advice
Daniel Rosehill · share with attribution

CPR Technique (C-A-B)



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

AED & Continuation



Key Paediatric CPR Parameters

Compression rate: 100-120/min
Infant depth: 4 cm (1/3 chest)
Two-rescuer ratio: 15:2
Lone rescuer: 2 min CPR, then call 101
Child depth: 5 cm (1/3 chest)
Solo ratio: 30:2
Breath duration: 1 second each
Emergency number: 101 (MDA)

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT hyperextend an infant's neck** when opening the airway – use neutral/sniffing position only.
✕ **Do NOT insert fingers into the mouth** (risk of pushing objects deeper).
✕ **Do NOT deliver deep lung breaths to infants** – use gentle puffs from your cheeks.
✕ **Do NOT use excessive volume or pressure** when delivering rescue breaths (risk of gastric distension).
✕ **Do NOT skip rescue breaths** in paediatric CPR – compression-only CPR is NOT adequate for children and infants.
✕ **Do NOT delay CPR to call for help when alone** – perform 2 minutes of CPR first, then call 101.
✕ **Do NOT allow AED pads to touch or overlap** on a small child or infant.
✕ **Do NOT stop CPR to check for pulse** unless there are clear signs of recovery (breathing, movement).
✕ **Do NOT compress over the xiphoid process** (the very bottom tip of the sternum).

Equipment needed:

— AED with paediatric pads or paediatric dose attenuator (if available)
— Barrier device or pocket mask (optional, for rescue breaths)
— Bag-valve-mask -- infant size (250-350 cc) or paediatric size (if available)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
CPR	החייאה	ha-kha-ya-A
Cardiac arrest	דום לב	dom lev
Chest compressions	לחיצות חזה	le-khi-TSOT kha-ZE
Rescue breaths	הנשמות	han-sha-MOT
Not breathing	לא נושם	lo no-SHEM
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
Recovery position	תנוחת התאוששות	tnu-KHAT hit-o-she-SHUT
Child	ילד / ילדה	YE-led / yal-DA
Infant	תינוק / תינוקת	ti-NOK / ti-NO-ket
Ambulance	אמבולנס	am-bu-LANS
The child is not breathing!	הילד לא נושם!	ha-YE-led lo no-SHEM!

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — BLS Guide 2016 / ALS Professional Handbook 2024
URL: https://www.mdais.org/kursim/cpr-children
Publication date: 2024-04-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16
Notes: Unified from MDA and United Hatzalah. In children, cardiac arrest is almost always secondary to respiratory failure – rescue breaths are critically important.

Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT give cola, fruit juice, plain water alone, or tea** — these do not replace electrolytes and can worsen dehydration.
✕ **Do NOT use homemade salt/sugar solutions** unless ORS is completely unavailable.
✕ **Do NOT force large volumes of fluid at once** — small frequent sips are more effective.
✕ **Do NOT delay calling 101 for an infant under 3 months** with any signs of dehydration.
✕ **Do NOT ignore a sunken fontanelle** in an infant — this is a key dehydration indicator.
✕ **Do NOT give anti-diarrhoea medications** to children without medical direction.
✕ **Do NOT stop breastfeeding** — continue alongside ORS.

Equipment needed:

— ORS (Oral Rehydration Solution) packets
— Spoon or oral syringe (for infant fluid administration)
— Phone (to call 101)
— Thermometer (to check for fever)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Dehydration	התייבשות	hit-yab-SHUT
Oral rehydration	שתייה מחדש	shti-YA me-kha-DASH
ORS / electrolytes	תמיסת מלחים	tmi-SAT mla-KHIM
Fontanelle	מרפס	mar-PES
Vomiting	הקאות	ha-ka-OT
Diarrhoea	שלשולים	shil-shu-LIM
Thirsty	צמא	tsa-ME
Pharmacy	בית מרקחת	beit mir-KA-khat
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
The child is dehydrated	הילד מיובש	ha-YE-led me-yu-BASH
He/she is vomiting	הוא/היא מקיא/ה	hu/hi ma-KI/ma-ki-A

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — MDA Dehydration First Aid Guidelines — Pediatric
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Personal reference only — not medical advice

Daniel Rosehill · share with attribution




Daniel Rosehill · share with attribution

Severe Hypoglycemia — Unconscious / Seizing



Daniel Rosehill · share with attribution

DKA Assessment & Uncertainty Rule



When in Doubt — Give Sugar

If uncertain whether hypoglycemia or hyperglycemia: **give sugar**.

Sugar for a hypoglycemic child is life-saving.
Sugar for a hyperglycemic child will not cause immediate harm.
Withholding sugar from a hypoglycemic child can be fatal.

This applies only when the child is conscious and can swallow safely.

Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT administer insulin** — this is strictly a medical intervention and can be fatal if given inappropriately.
- ✕ **Do NOT give food, drink, or oral medication to an unconscious or seizing child** — choking hazard.
- ✕ **Do NOT use chocolate as the initial sugar source** — fat and protein slow absorption and delay correction.
- ✕ **Do NOT administer glucagon unless trained and authorised** to do so.
- ✕ **Do NOT delay calling 101** for severe hypoglycemia, seizures, or loss of consciousness.
- ✕ **Do NOT attempt to manage hyperglycemia or DKA with first aid** — requires medical intervention.
- ✕ **Do NOT use diet drinks or sugar-free products** as a sugar source.

Equipment needed:


— Glucose gel syringe (15 g glucose)
— Fast-acting sugar source (juice, glucose tablets, sugar cubes, honey)
— Complex carbohydrate snack (bread, crackers, fruit)
— Glucagon emergency kit (Hypo-Kit) if available and authorised
— Blood glucose meter (glucometer) if available

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Diabetes	סוכרת	su-KE-ret
Low blood sugar	סוכר נמוך	SU-kar na-MUKH
High blood sugar	סוכר גבוה	SU-kar ga-VO-ah
Sugar	סוכר	SU-kar
Glucagon	גלוקגון	glu-ka-GON
Insulin pump	משאבת אינסולין	mash-E-vet in-su-LIN
Seizure	פרכוס	pir-KUS
Unconscious	חסר הכרה	kha-SAR ha-ka-RA
Sweet drink	שתייה מתוקה	shti-YA me-tu-KA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Does the child have diabetes?	יש לילד סוכרת?	yesh la-YE-led su-KE-ret?

Source: Magen David Adom (MDA) — How to Treat a Diabetes Patient
URL: https://www.mdais.org/101/diabetes
Imported: 2026-03-15 · **Last verified:** 2026-03-16


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

CPR Continuation & Recovery



All drowning victims MUST go to hospital

Secondary drowning (delayed pulmonary oedema) can develop hours after the event, even if the child initially appears to recover fully. Electrolyte imbalances can also occur.

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT attempt to remove water from the lungs** — this is impossible and can endanger the victim.
✕ **Do NOT perform abdominal thrusts (Heimlich)** to expel water — not effective for drowning and risks aspiration.
✕ **Do NOT delay CPR** to attempt water removal.
✕ **Do NOT use compression-only CPR for drowning** — rescue breaths are essential (hypoxic emergency).
✕ **Do NOT assume the child is fine** if they regain consciousness — always transport to hospital.
✕ **Do NOT enter the water** unless you are a trained water rescuer with a flotation device.
✕ **Do NOT use excessive ventilation force** — can cause gastric distention in children.

Equipment needed:


— Reaching aid (pole, rope, towel) or throwing aid (flotation device)
— Firm, flat surface for CPR
— Blanket or covering for warmth (hypothermia prevention)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Drowning	טביעה	tvi-A
Water	מים	MA-yim
Swimming pool	בריכת שחייה	bre-KHAT skhi-YA
CPR	החייאה	ha-kha-ya-A
Rescue breaths	הנשמות	han-sha-MOT
Chest compressions	לחיצות חזה	lkhi-TSOT kha-ZE
Breathing	נשימה	nshi-MA
Recovery position	תנוחת התאוששות	tnu-KHAT hit-o-she-SHUT
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
The child is not breathing	הילד לא נושם	ha-YE-led lo no-SHEM
Call an ambulance	תזמין אמבולנס	taz-MIN am-bu-LANS

Source: Magen David Adom (MDA) — Drowning First Aid Protocol for Children
URL: https://www.mdais.org/101/first-aid
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice
Daniel Rosehill · share with attribution


⚠️ STOP — do NOT touch the child while in contact with source.
⚡ High-voltage source? (power lines, transformer)
Y
📱 CALL 101, 102, and 103. Keep distance. Wait for emergency services.
N
DISCONNECT power at circuit breaker or unplug. If unreachable, use dry non-conductive object.
📱 CALL 101 immediately.
👁️ Child responsive?
Y
KEEP still and calm. Proceed to burn assessment (Page 2).
N
👁️ Breathing and pulse present?
Y
PLACE in recovery position. Monitor breathing.
N
❤️ BEGIN CPR. Use AED if available. Continue until MDA arrives.

Burn Assessment & Shock Management



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT touch the child while in contact with the electrical source** — chain electrocution risk.
✕ **Do NOT use metal objects or wet/damp materials** to separate the child from the source.
✕ **Do NOT apply ice to electrical burns.**
✕ **Do NOT apply creams, ointments, toothpaste, or home remedies** to electrical burns.
✕ **Do NOT assume the child is fine** because external injuries appear minor — internal damage may be hidden.
✕ **Do NOT delay calling 101** — every pediatric electrocution requires hospital assessment.
✕ **Do NOT move the child unnecessarily** — spinal/traumatic injuries may be present.
✕ **Do NOT approach high-voltage sources** — call 101, 102, and 103; maintain safe distance.
✕ **Do NOT stop CPR** based on apparent lack of vital signs — electrocution victims can be resuscitated after prolonged efforts.
✕ **Do NOT give food or drink** to a child in shock after electrocution.

Equipment needed:

— Non-conductive object (wooden broom handle, dry stick, plastic chair)
— AED / defibrillator (with pediatric pads if available)
— Cool running water source (for burn cooling)
— Sterile non-adherent dressings
— Blanket (for warmth and shock treatment)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Electrocution	התחשמלות	hit-khash-me-LUT
Electric shock	מכת חשמל	ma-KAT khash-MAL
Circuit breaker	מפסק ראשי	maf-SEK ra-shi
Power line	קו חשמל	kav khash-MAL
Burn	כוויה	kvi-YA
Shock (medical)	הלם	HE-lem
CPR	החייאה	ha-kha-ya-A
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Turn off the electricity	תכבה את החשמל	te-kha-BE et ha-khash-MAL
The child was electrocuted	הילד התחשמל	ha-YE-led hit-khash-MEL

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Electrocution First Aid Guidelines — Pediatric
URL: https://www.mdais.org/101/electrocution
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


👁️ Chemical burn to the eye?
Y
💦 IRRIGATE immediately with clean flowing water for 15–30 minutes. Tilt head: inner to outer.
TRANSPORT urgently to hospital. Bring chemical container.
N
👁️ Foreign object in the eye?
Y
💦 FLUSH gently with clean lukewarm water for 15 minutes. Seek eye evaluation.
N
👁️ Object embedded or penetrating the eye surface?
Y
COVER both eyes with sterile dressings. No pressure.
📱 Call 101.
👁️ Penetrating injury?
Y
SHIELD eye with rigid cover (paper cup). Cover other eye too. Position supine.
📱 Call 101.
N
APPLY cold compress gently over closed eyelid for 10–15 minutes.
SEEK eye evaluation within 24 hours.
N
👁️ Vision changes, double vision, or severe pain?
Y
📱 CALL 101. Urgent hospital evaluation.

Reference Notes

DO NOT:

- ✕ **Do NOT rub the injured eye** — worsens damage from foreign objects and chemical burns.
- ✕ **Do NOT apply pressure or bandages to a penetrating eye injury.**
- ✕ **Do NOT attempt to remove an embedded object** from the eye.
- ✕ **Do NOT delay irrigation for chemical burns** — begin immediately.
- ✕ **Do NOT use any eye drops or ointments** without medical direction.
- ✕ **Do NOT allow the child to rub or touch the injured eye.**
- ✕ **Do NOT apply antiseptic or alcohol** to the eye.

Equipment needed:

— Clean water source (tap water or sterile water for irrigation)
— Rigid eye shield or paper cup
— Sterile gauze or clean cloth
— Cold compress or ice wrapped in cloth
— Phone (to call 101)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Eye	עין	A-yin
Eye injury	פגיעה בעין	pgi-A ba-A-yin
Chemical burn	כוויה כימית	kvi-YA khi-MIT
Foreign object	גוף זר	guf ZAR
Irrigate / rinse	לשטוף	lish-TOF
Water	מים	MA-yim
Bandage	תחבושת	takh-BO-shet
Vision	ראייה	re-i-YA
Pain	כאב	ke-EV
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Something got in the eye	נכנס משהו לעין	nikh-NAS MA-she-hu la-A-yin
Rinse with water	תשטוף עם מים	tish-TOF im MA-yim

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Eye Injury First Aid Guidelines — Pediatric
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Shock Prevention & Transport



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT attempt to straighten, pull, or realign a broken bone** — can damage arteries, nerves, and growth plates.
✕ **Do NOT move the child unnecessarily**, especially if spinal injury is suspected.
✕ **Do NOT apply heat** to the injury (no hot compresses, no warm water).
✕ **Do NOT apply ice directly to skin** — always wrap in cloth first.
✕ **Do NOT tie splints or bandages too tightly** — monitor circulation below the injury.
✕ **Do NOT allow the child to bear weight** on an injured leg.
✕ **Do NOT remove clothing by pulling over the injury** — cut it away if necessary.
✕ **Do NOT give aspirin to children** (Reye's syndrome risk); use paracetamol (Acamol) or ibuprofen (Nurofen for Children).
✕ **Do NOT push a protruding bone back** into the wound in an open fracture.
✕ **Do NOT give the child food or drink** — surgery under anaesthesia may be required.

Equipment needed:

— Improvised sling material (shirt, scarf, sweater, triangular bandage)
— Improvised splint material (cardboard, rolled newspapers, rulers)
— Padding material (clothing, towels, blankets)
— Ice pack or cold compress (or bag of frozen vegetables)
— Cloth or towel for wrapping ice pack
— Sterile dressings or clean cloth (for open fractures)
— Bandages or fabric strips for securing splints
— Blanket for warmth and shock prevention
— Tourniquet (for severe uncontrollable bleeding, if available)


Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Fracture / break	שבר	SHE-ver
Sprain	נקע	NE-ka
Splint	סד	SAD
Sling	מתלה	mit-LE
Bone	עצם	E-tsem
Swelling	נפיחות	ne-fi-KHUT
Pain	כאב	ke-EV
Growth plate	לוחית צמיחה	lu-KHIT tsmi-KHA
Open fracture	שבר פתוח	SHE-ver pa-TU-akh
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
It hurts here	כואב לי פה	ko-EV li PO
Can you move it?	אתה יכול לזוז?	a-TA ya-KHOL la-ZUZ?


Source: Magen David Adom (MDA) — Fractures and Sprains First Aid Protocol — Children
URL: https://www.mdais.org/101/first-aid
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice
Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Continued Monitoring



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT move the child** unless in immediate life-threatening danger (fire, traffic, structural collapse).
- ✕ **Do NOT remove a helmet** — leave for MDA paramedics (requires two trained rescuers).
- ✕ **Do NOT allow the child to stand or walk** — standing/walking does NOT exclude spinal injury.
- ✕ **Do NOT give food or drink** — imaging or surgery under anaesthesia may be needed.
- ✕ **Do NOT apply pressure to a depressed skull fracture** or push back any protruding matter.
- ✕ **Do NOT insert anything into a wound.**
- ✕ **Do NOT attempt to straighten the neck** if in an abnormal position — hold as found.
- ✕ **Do NOT use head-tilt chin-lift** to open the airway — use jaw thrust only.
- ✕ **Do NOT shake the child** to assess responsiveness.
- ✕ **Do NOT leave the child alone** — continuous monitoring is essential.
- ✕ **Do NOT stop the flow of clear fluid** from ears or nose (may be cerebrospinal fluid).

Equipment needed:

— Disposable gloves
— Clean cloth or sterile bandages for wound control
— Blanket or coat for warmth


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Head injury	פגיעת ראש	pgi-AT ROSH
Spinal injury	פגיעת עמוד שדרה	pgi-AT a-MUD shid-RA
Spine	עמוד שדרה	a-MUD shid-RA
Neck	צוואר	tsa-VAR
Helmet	קסדה	kas-DA
Consciousness	הכרה	ha-ka-RA
Unconscious	חסר הכרה	kha-SAR ha-ka-RA
Vomiting	הקאות	ha-ka-OT
Seizure	פרכוס	pir-KUS
Fontanelle	מרפס	mar-PES
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Do not move the child	אל תזיזו את הילד	al ta-ZI-zu et ha-YE-led
The child fell	הילד נפל	ha-YE-led na-FAL

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — MDA BLS Guide (2016), MDA ALS Professional Manual (2024)
URL: https://www.mdais.org/101/first-aid
Imported: 2026-03-15 · **Last verified:** 2026-03-16


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

CPR and AED Use



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT give aspirin to a child** — not indicated for pediatric chest pain; risk of Reye's syndrome.
✕ **Do NOT give nitroglycerin or cardiac medication** unless prescribed and directed by a physician.
✕ **Do NOT dismiss chest pain** in a child who has been ill, has a known heart condition, collapsed during activity, or has family history of sudden cardiac death.
✕ **Do NOT drive the child to hospital yourself** if cardiac arrest is suspected — call 101 for an equipped ambulance.
✕ **Do NOT leave the child unattended** while waiting for the ambulance.
✕ **Do NOT give the child food or drink** (in case hospital procedures are needed).
✕ **Do NOT give any medications** unless instructed by the 101 dispatcher or a physician.

Equipment needed:

— AED with pediatric pads or dose attenuator (if available)
— Phone to call 101 (MDA) or 1221 (United Hatzalah)
— MDA app (for AED locator and emergency calling)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Heart attack	התקף לב	het-KEF LEV
Chest pain	כאב בחזה	ke-EV ba-kha-ZE
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
CPR	החייאה	ha-kha-ya-A
Defibrillator	דפיברילטור	de-fib-ri-LA-tor
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Pulse	דופק	DO-fek
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Child	ילד / ילדה	YE-led / yal-DA
Does the chest hurt?	כואב לך בחזה?	ko-EV le-KHA ba-kha-ZE?
Can you breathe?	אתה יכול לנשום?	a-TA ya-KHOL lin-SHOM?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Heart Attack and Chest Pain in Children
URL: https://www.mdais.org/101/first-aid
Publication date: 2022-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Airway and Continued Monitoring



Child found in vehicle

Temperatures inside a locked car in Israeli summer can reach 70 C.
REMOVE the child from the vehicle immediately – break the window if necessary.
Report to MDA that the child was found in a vehicle, including estimated time of exposure.

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT give fever-reducing medications** (Acamol, Nurofen, naproxen) — these do not reduce heatstroke temperature and may cause harm.
✕ **Do NOT use ice-cold water immersion** for young children without medical supervision.
✕ **Do NOT force fluids** if the child is vomiting or has altered consciousness.
✕ **Do NOT leave the child unattended** at any point.
✕ **Do NOT give fluids to an unconscious child.**
✕ **Do NOT delay calling 101** if heat stroke is suspected.
✕ **Do NOT leave a child unattended in a vehicle** — even for a moment, even with windows open or A/C running.

Equipment needed:

— Cool water (for pouring or spraying over the body)
— Fan or means of fanning
— Ice packs or cold wet cloths
— Oral rehydration solution (ORS) or cool drinking water
— Thermometer (if available)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Heatstroke	מכת חום	ma-KAT KHOM
Heat exhaustion	תשישות חום	tshi-SHUT KHOM
Water	מים	MA-yim
Shade	צל	TSEL
Hot	חם	KHAM
Cold water	מים קרים	MA-yim ka-RIM
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Child	ילד / ילדה	YE-led / yal-DA
Are you thirsty?	אתה צמא?	a-TA tsa-ME?
Drink water	תשתה מים	tish-TE MA-yim

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Heat Stroke Guidelines
URL: https://www.mdais.org/news/300820
Publication date: 2024-04-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Transport and Monitoring

📱 **CALL** 101 and transport urgently to hospital. Keep child horizontal.

↓

👁️ **MONITOR** breathing, pulse, and consciousness continuously until handover.

Be prepared to begin CPR if child deteriorates.

Hypothermia Severity Stages

Stage	Temperature	Signs
Stage 1	37–35 C	Shivering, normal judgement
Stage 2	35–32 C	Severe shivering, confusion, slurred speech, pallor
Stage 3	32–28 C	Deteriorating consciousness, muscle rigidity
Stage 4	Below 28 C	Unconsciousness, absent reflexes, absent breathing

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT give alcohol, caffeine, or allow smoking** — these worsen hypothermia.
✕ **Do NOT make excessive or rough movements** — risk of cardiac arrhythmias in a cold heart.
✕ **Do NOT apply direct heat to skin** without a cloth barrier — risk of burns.
✕ **Do NOT rewarm extremities before the core** — can cause dangerous drop in core temperature as cold blood returns to the heart.
✕ **Do NOT give fluids by mouth** to an unconscious or semi-conscious child — aspiration risk.
✕ **Do NOT assume a severely hypothermic child is dead** — "no one is dead until they are warm and dead."
✕ **Do NOT use hot water immersion** for rewarming in the field — risk of burns and cardiac arrhythmias.
✕ **Do NOT jostle or move the child unnecessarily** during transport.

Equipment needed:

— Warm blankets or sleeping bags
— Dry clothing
— Heat packs or warm water bottles (wrapped in cloth)
— Phone (to call 101)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Hypothermia	היפותרמיה	hi-po-ter-MI-ya
Cold	קר	KAR
Blanket	שמיכה	smi-KHA
Shivering	רעד	RA-ad
Pulse	דופק	DO-fek
Breathing	נשימה	nshi-MA
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Child	ילד / ילדה	YE-led / yal-DA
Are you cold?	קר לך?	KAR le-KHA?
Warm drink	משקה חם	mash-KE KHAM

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — Hypothermia First Aid Guidelines
URL: https://www.mdais.org/101/first-aid
Publication date: 2024-04-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Non-Ingestion Routes and Calling for Help



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT induce vomiting** — regardless of the substance ingested. Causes additional oesophageal burns (caustic agents), aspiration risk, and ineffective removal.
✕ **Do NOT give activated charcoal at home** unless specifically instructed by the Poison Center or a physician.
✕ **Do NOT give food to the poisoned child.** Water or milk for caustic dilution only, when conscious.
✕ **Do NOT attempt home remedies** (raw eggs, olive oil, etc.).
✕ **Do NOT wait for symptoms** before calling for guidance — some toxic substances cause delayed organ damage (liver, kidneys).
✕ **Do NOT discard the container or packaging** — always bring it to the hospital.
✕ **Do NOT mix cleaning products** — especially bleach with acids (releases toxic chlorine gas).
✕ **Do NOT enter a toxic fume environment** without proper protection.

Equipment needed:

— Cloth (to wipe substance from child's mouth if present)
— Water or milk (for dilution of caustic substances only, age-appropriate amounts)
— Running water source (for skin and eye decontamination, 15--20 minutes flushing)
— Container or bag (to preserve substance packaging and any vomit for medical teams)

Key Phone Numbers

Service	Number
MDA (emergency)	**101**
Israel Poison Information Center (Rambam)	**04-7771900** (24/7)
United Hatzalah	**1221**
Fire Department	**102**


Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Poison / poisoning	הרעלה	har-a-LA
Poison center	מרכז הרעלות	mer-KAZ har-a-LOT
Swallowed	בלע	ba-LA
Cleaning product	חומר ניקוי	KHO-mer ni-KUY
Bleach (economica)	אקונומיקה	e-ko-NO-mi-ka
Vomiting	הקאה	ha-ka-A
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Child	ילד / ילדה	YE-led / yal-DA
What did they swallow?	מה הוא בלע?	MA HU ba-LA?
How much?	כמה?	KA-ma?


Source: Magen David Adom (MDA) — Poisoning First Aid Guidelines; Rambam National Poison Information Center
URL: https://www.mdais.org/101/poisoningn
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice
Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Monitoring



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT use the recovery position if the child is not breathing** — begin CPR instead.
✕ **Do NOT use the recovery position if spinal injury is suspected** — stabilise the spine.
✕ **Do NOT place infants under 1 year in the standard lateral recovery position** — use the forearm cradle position.
✕ **Do NOT leave the child unattended** in the recovery position — monitor breathing continuously.
✕ **Do NOT allow the child's head to flex forward** (especially in children with larger heads) — this can obstruct the airway.
✕ **Do NOT forget to cover the child** to prevent hypothermia.

Equipment needed:


— Blanket or coat (to prevent hypothermia)
— Phone (to call 101)

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Recovery position	תנוחת החלמה	tnu-KHAT ha-khla-MA
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נשימה	nshi-MA
Airway	דרכי אוויר	dar-KHEY a-VIR
On their side	על הצד	al ha-TSAD
Infant	תינוק	ti-NOK
Spinal injury	פגיעה בעמוד השדרה	pgi-A be-a-MUD ha-shid-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Child	ילד / ילדה	YE-led / yal-DA
Is the child breathing?	הילד נושם?	ha-YE-led no-SHEM?
Roll them on their side	תגלגל אותו על הצד	te-gal-GEL o-TO al ha-TSAD

Source: Magen David Adom (MDA) — Recovery Position Guidelines
URL: https://www.mdais.org/101/first-aid
Publication date: 2016-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply a tourniquet** — concentrates venom in the limb and causes tissue damage.
✕ **Do NOT cut or incise the sting site** — causes additional injury and does not remove venom.
✕ **Do NOT attempt to suck out the venom** — ineffective and risks infection.
✕ **Do NOT apply alcohol or antiseptic** to the wound — interferes with medical assessment.
✕ **Do NOT give the child medication** without medical advice — may interact with venom effects or mask symptoms.
✕ **Do NOT underestimate a sting based on scorpion colour** — black scorpions can be as dangerous as yellow ones.
✕ **Do NOT delay calling 101 for children** — children deteriorate faster due to higher venom-to-body-weight ratio.
✕ **Do NOT apply ice directly to skin** — always wrap in cloth.
✕ **Do NOT allow the child to walk or move** — movement accelerates venom distribution.
✕ **Do NOT "wait and see" with children** — all scorpion stings in children require emergency medical evaluation.

Equipment needed:

— Ice pack or cold compress (wrapped in cloth)
— Splint or rigid object for limb immobilisation
— Phone (to call 101 / 04-7771900 and photograph the scorpion)
— Cloth or towel (to wrap ice pack)
— Blanket or clothing (to keep child comfortable while lying still)


Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Scorpion	עקרב	ak-RAV
Sting	עקיצה	a-ki-TSA
Venom / poison	ארס	E-res
Antivenom	נוגדן	no-ge-DAN
Swelling	נפיחות	ne-fi-KHUT
Pain	כאב	ke-EV
Ice	קרח	KE-rakh
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Child	ילד / ילדה	YE-led / yal-DA
A scorpion stung the child	עקרב עקץ את הילד	ak-RAV a-KATS et ha-YE-led
Where is the sting?	איפה העקיצה?	EI-fo ha-a-ki-TSA?


Source: Magen David Adom (MDA) — Scorpion Sting First Aid Guidelines
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Personal reference only — not medical advice
Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Post-Seizure and Febrile Management



Post-Seizure Fever Management (Febrile Seizures)

Once the child has recovered and is no longer actively seizing:

- **ADMINISTER** age-appropriate fever-reducing medication (paracetamol or ibuprofen) as directed by the family doctor.
- Dress child in light clothing. Ventilate the room.
- Give plenty of fluids.
- A lukewarm bath (20 min) may be given only when NOT actively seizing.

Note: Antipyretics improve comfort but have NOT been proven to prevent febrile seizures (Israeli Ministry of Health).

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT force the child's mouth open** — the child will not swallow their tongue.
✕ **Do NOT insert anything into the mouth** — this can injure teeth or jaw.
✕ **Do NOT attempt CPR during the seizure** — this can only cause harm.
✕ **Do NOT put the child in a bath or shower during the seizure.**
✕ **Do NOT splash water on the child** during the seizure.
✕ **Do NOT try to wake the child** or stop the seizure by restraining or holding them down.
✕ **Do NOT give medications by mouth** during the seizure.
✕ **Do NOT attempt to cool the child rapidly** with cold water or ice.
✕ **Do NOT insert an airway device** during an active seizure.
✕ **Do NOT attempt forced airway opening** during an active seizure.

Equipment needed:

— Lukewarm damp towel (for febrile seizures)
— Pillows or blankets (to pad surroundings)
— Prescribed seizure-stopping medication if available (e.g., rectal diazepam)
— Fever-reducing medication for post-seizure fever management (paracetamol or ibuprofen, as directed by doctor)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Seizure / convulsion	פרכוס	pir-KUS
Febrile seizure	פרכוס חום	pir-KUS KHOM
Fever	חום	KHOM
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Breathing	נשימה	nshi-MA
Medication	תרופה	tru-FA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Child	ילד / ילדה	YE-led / yal-DA
The child is having a seizure	הילד מפרכס	ha-YE-led me-far-KES
How long has the seizure lasted?	כמה זמן הפרכוס?	KA-ma ZMAN ha-pir-KUS?
Does the child have a fever?	יש לילד חום?	YESH la-YE-led KHOM?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — How to Treat Febrile Seizures
URL: https://www.mdais.org/101/seizures
Publication date: 2022-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-16


Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Escalation — Wound Packing / Tourniquet



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

- ✕ **Do NOT remove an embedded or impaled object** from the wound.
- ✕ **Do NOT lift or remove a blood-soaked dressing** — add layers on top.
- ✕ **Do NOT remove or loosen a tourniquet** once applied unless instructed by a paramedic or physician.
- ✕ **Do NOT loosen a tourniquet because the child is screaming** — pain is expected.
- ✕ **Do NOT give food or drink** to a child with severe bleeding or suspected shock.
- ✕ **Do NOT use wound packing for penetrating stab wounds.**
- ✕ **Do NOT enlarge wound openings** to facilitate wound packing.
- ✕ **Do NOT assume tachycardia is only from anxiety or pain** — suspect blood loss.
- ✕ **Do NOT wait for hypotension** to suspect shock in a child — it is a late, pre-arrest sign.
- ✕ **Do NOT delay calling 101** for severe or uncontrolled bleeding in a child.
- ✕ **Do NOT apply a tourniquet over a joint** — place 5–10 cm above the wound.
- ✕ **Do NOT use a manufactured tourniquet on a child under 2** — use sustained direct pressure.

Equipment needed:


— Disposable gloves
— Sterile gauze or clean cloth
— Israeli Emergency Bandage (pressure dressing)
— Tourniquet (CAT Gen 7 — for children aged 2+ with limb circumference 13 cm+)
— Wound packing material (hemostatic gauze preferred)
— Blanket or coat (for shock management and warmth)
— Comfort item for the child (toy, stuffed animal) if available

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Bleeding	דימום	di-MUM
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Bandage	תחבושת	takh-BO-shet
Pressure	לחץ	LA-khats
Wound	פצע	PE-tsa
Shock	הלם	HE-lem
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Where is the bleeding?	איפה הדימום?	EI-fo ha-di-MUM?
The child is bleeding	הילד מדמם	ha-YE-led me-da-MEM
Apply pressure	תלחץ	til-KHATZ
Gloves	כפפות	kfa-FOT


Source: Magen David Adom (MDA) — MDA 101 First Aid Public Guidance (Bleeding Control) — Pediatric Adaptation
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-15



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Continued Monitoring



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT sit the child upright** — this reduces blood flow to the brain and worsens shock.
✕ **Do NOT give the child anything to eat or drink** — moisten lips only.
✕ **Do NOT leave the child unattended** — continuous monitoring is essential due to risk of sudden deterioration.
✕ **Do NOT delay calling 101** — early hospital treatment with IV fluids is critical for survival.
✕ **Do NOT assume normal blood pressure means no shock** — children maintain BP through strong compensatory mechanisms until sudden decompensation.
✕ **Do NOT attribute tachycardia solely to anxiety, fear, or pain** — suspect shock until proven otherwise.
✕ **Do NOT move a child with suspected spinal injury** — stabilise and wait for EMS.
✕ **Do NOT remove blood-soaked dressings** — add additional layers on top.

Equipment needed:

— Blanket or coat (for warmth and insulation from ground)
— Sterile gauze or clean cloth (for bleeding control)
— Disposable gloves
— Pillow or folded blanket (for leg elevation)


Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Shock	הלם	HE-lem
Pulse / heart rate	דופק	DO-fek
Blood pressure	לחץ דם	LA-khats DAM
Pale	חיוור	khi-VER
Cold	קר	KAR
Breathing	נשימה	ne-shi-MA
Consciousness	הכרה	ha-ka-RA
Recovery position	תנוחת התאוששות	tnu-KHAT hit-o-she-SHUT
Blanket	שמיכה	smi-KHA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
The child is in shock	הילד בהלם	ha-YE-led be-HE-lem

Personal reference only — not medical advice

Daniel Rosehill · share with attribution


Source: Magen David Adom (MDA) — MDA BLS Guide (2016) and MDA 101 First Aid Public Guidance — Pediatric Shock
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-15



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Tear Gas Exposure & Poisoning Signs



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT re-enter a burning building or hazardous area** — wait for Fire Department rescue teams.
✕ **Do NOT ignore light smoke exposure in a child** — children should be assessed even with mild exposure.
✕ **Do NOT assume a well-appearing child is uninjured** — respiratory deterioration can be delayed by hours.
✕ **Do NOT rub the eyes after tear gas exposure** — this worsens irritation.
✕ **Do NOT give food or drink to a child with stridor or breathing difficulty** — aspiration risk.
✕ **Do NOT examine the throat of a child with stridor** — this can worsen airway swelling.
✕ **Do NOT leave the child unattended** — monitor breathing continuously.
✕ **Do NOT position a child with breathing difficulty lying flat** — keep upright.

Equipment needed:

— Supplemental oxygen and face mask (if available)
— Clean water (for eye irrigation after tear gas)
— Blanket (to prevent hypothermia)
— Cool mist humidifier (if available, for stridor management)
— Phone (to call 101 and 102)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Smoke	עשן	a-SHAN
Fire	שריפה	sre-FA
Tear gas	גז מדמיע	GAZ mad-MI-a
Hard to breathe	קשה לנשום	ka-SHE lin-SHOM
Oxygen	חמצן	kham-TSAN
Cough	שיעול	shi-UL
Stridor	סטרידור	STRI-dor
Fire department	כיבוי אש	ki-BUY ESH
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
The child inhaled smoke	הילד שאף עשן	ha-YE-led sha-AF a-SHAN
Can you breathe?	אתה יכול לנשום?	a-TA ya-KHOL lin-SHOM?

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — MDA Smoke and Inhalation Injury First Aid Guidelines — Pediatric
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Personal reference only — not medical advice


Daniel Rosehill · share with attribution



Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Monitoring & Transport



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply a tourniquet** — concentrates venom and causes tissue damage.
✕ **Do NOT cut the bite site** — causes additional injury and does not remove venom.
✕ **Do NOT attempt to suck out the venom** — ineffective and risks infection.
✕ **Do NOT apply ice or cold compresses** — can cause tissue damage and does not slow absorption.
✕ **Do NOT apply alcohol, antiseptic, or any substance** to the wound.
✕ **Do NOT give the child food or drink** — risk of vomiting; may need anaesthesia.
✕ **Do NOT chase, catch, kill, or handle the snake** — risk of additional bites.
✕ **Do NOT use a pressure immobilisation bandage** — not recommended for viper bites.
✕ **Do NOT allow the child to walk or move** — movement accelerates venom spread.
✕ **Do NOT wait for symptoms** before seeking emergency care.

Equipment needed:

— Splint or rigid object for limb immobilisation
— Water and soap (for wound cleaning)
— Phone (to call 101 and photograph the snake)
— Blanket or clothing (to keep child warm while lying still)

Hebrew Terminology — מונחים בעברית


English	עברית	Pronunciation
Snake	נחש	na-KHASH
Snakebite	הכשת נחש	ha-ka-SHAT na-KHASH
Viper	צפע	TSE-fa
Palestine viper	צפע ארצישראלי	TSE-fa er-ets yis-re-e-LI
Venom	ארס	E-res
Antivenom	נוגדן נגד ארס	nog-DAN NE-ged E-res
Swelling	נפיחות	ne-fi-KHUT
Splint	סד	SAD
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
The child was bitten by a snake	הילד הוכש על ידי נחש	ha-YE-led hu-KASH al ye-DEI na-KHASH
Keep still	תישאר במקום	ti-sha-ER ba-ma-KOM

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — MDA Snakebite First Aid Guidelines — Pediatric
URL: https://www.mdais.org/101/snake-bite
Publication date: 2026-01-01
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Personal reference only — not medical advice

Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT apply a tourniquet** — tourniquets worsen envenomation outcomes by concentrating venom.
✕ **Do NOT cut the bite site** — causes additional injury and does not remove venom.
✕ **Do NOT attempt to suck out the venom** — ineffective and risks infection.
✕ **Do NOT apply antiseptic, alcohol, or any substance** other than water and soap.
✕ **Do NOT bandage tightly** over the bite site.
✕ **Do NOT delay medical evaluation** for any spider bite in a child.
✕ **Do NOT ignore a painless bite** — brown recluse bites are often painless initially but develop serious necrotic wounds hours later.
✕ **Do NOT give the child medications** (painkillers, antihistamines) without medical direction.

Equipment needed:

— Water and soap (for wound cleaning)
— Ice or cold compress wrapped in cloth
— Container or glass (to safely capture the spider for identification)
— Phone (to call 101 and photograph the spider)

Venomous Spiders in Israel

Brown Recluse (shshan chum)
Small, brown, long legs
Lives indoors (wardrobes, shoes)
Bite barely felt initially
Effects appear hours later
Causes local necrotic wound

Black Widow (almana shchora)
Black with markings
Found outdoors (gardens, walls)
Immediate pain and burning
Rapid systemic symptoms
Female bite is dangerous
Antivenom available


Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Spider	עכביש	ak-a-VISH
Spider bite	עקיצת עכביש	a-ki-TSAT ak-a-VISH
Black widow	אלמנה שחורה	al-ma-NA shcho-RA
Brown recluse	שושן חום	sho-SHAN KHUM
Venom	ארס	E-res
Antivenom	נוגדן נגד ארס	nog-DAN NE-ged E-res
Swelling	נפיחות	ne-fi-KHUT
Pain	כאב	ke-EV
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
The child was bitten by a spider	הילד נעקץ על ידי עכביש	ha-YE-led ne-e-KATS al ye-DEI ak-a-VISH

Source: Magen David Adom (MDA) — MDA Spider Bite First Aid Guidelines — Pediatric
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution



Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT delay calling 101** to wait and see if symptoms resolve.
✕ **Do NOT give the child any food or drink** — stroke may impair swallowing reflex, creating aspiration and choking risk.
✕ **Do NOT give any medications, especially not aspirin** — if the stroke is haemorrhagic, aspirin will worsen it.
✕ **Do NOT drive the child to hospital yourself** unless MDA explicitly advises it. MDA ambulances begin treatment en route.
✕ **Do NOT assume a child cannot have a stroke** — children can have strokes at any age.
✕ **Do NOT restrain the child during a seizure.**
✕ **Do NOT put anything in the child's mouth** during a seizure.
✕ **Do NOT rely solely on FAST to rule out stroke in children** — adult screening tools are not validated for the pediatric population.

Pediatric Stroke — Key Differences from Adults

- Seizures at onset are more common in children than adults
- Symptoms may appear gradually rather than suddenly
- Very young children may show only irritability, feeding difficulty, or lethargy
- FAST assessment is not validated for the pediatric population
- Treatment window: alteplase effective within 4.5 hours (ages 28 days–18 years)
- Mechanical clot removal may be effective for ages 6+ within 6–24 hours


Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Stroke	שבץ מוחי	SHEV-ets mo-KHI
Face	פנים	pa-NIM
Arm	זרוע	zro-A
Speech	דיבור	di-BUR
Seizure	פרכוס	pir-KUS
Headache	כאב ראש	ke-EV ROSH
Consciousness	הכרה	ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
The child may be having a stroke	ייתכן שהילד סובל משבץ	yi-ta-KHEN she-ha-YE-led so-VEL mi-SHEV-ets
Can you smile?	אתה יכול לחייך?	a-TA ya-KHOL le-kha-YEKH?
Raise both arms	תרים את שתי הידיים	ta-RIM et shTEI ha-ya-DA-yim

Source: Magen David Adom (MDA) — MDA Stroke Recognition Protocol (FAST) and AHA 2024 First Aid Guidelines — Pediatric Stroke Considerations
URL: https://www.mdais.org/101/stroke
Publication date: 2024-11-18
Imported: 2026-03-15 · **Last verified:** 2026-03-15

Personal reference only — not medical advice
Daniel Rosehill · share with attribution



Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Monitoring & Transport

👁 **CHECK** tourniquet has not loosened.
CHECK bleeding has stopped.
MONITOR for shock: rapid weak pulse, pallor, cool mottled skin, altered consciousness.

↓

KEEP child warm to prevent hypothermia.
REASSURE child and parent.
Report time, device, and condition to arriving EMS.

Personal reference only — not medical advice
Daniel Rosehill · share with attribution

Reference Notes

DO NOT:

✕ **Do NOT remove a tourniquet once applied** — only hospital personnel should remove it.
✕ **Do NOT apply a tourniquet for venomous animal bites** (snakes, scorpions, spiders) — tourniquets worsen envenomation outcomes.
✕ **Do NOT apply a tourniquet over a joint** (elbow, knee) — place on upper arm or upper thigh.
✕ **Do NOT apply a tourniquet too loosely** — an insufficiently tight tourniquet causes venous bleeding without arterial occlusion, worsening blood loss.
✕ **Do NOT use a standard CAT tourniquet on very small infant limbs** (under 13 cm circumference) — it will not be effective.
✕ **Do NOT delay tourniquet application** to attempt prolonged direct pressure on massive haemorrhage.
✕ **Do NOT forget to record the time of application.**

Equipment needed:

— CAT (Combat Application Tourniquet) Gen 7 — for children aged ~2+ with sufficient limb size
— SWAT-T tourniquet — for all pediatric sizes including very small limbs
— RATS tourniquet — alternative for all pediatric sizes
— Improvised tourniquet materials (triangular bandage + rigid windlass object)
— Marker or pen (to record time of application)
— Sterile gauze (for wound packing if needed)
— Phone (to call 101)

Pediatric Tourniquet Sizing Guide

Age / Size	Device	Notes
Age 2+, limb 13 cm+	CAT Gen 7	Upper + lower extremity
Age 6+, limb 16 cm+	CAT Gen 7	Reliable both extremities
All pediatric sizes	SWAT-T / RATS	Effective on very small limbs
Infants < 2 years	Direct pressure	No evidence for manufactured TQ

Daniel Rosehill · share with attribution

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Tourniquet	חוסם עורקים	kho-SEM or-KIM
Bleeding	דימום	di-MUM
Tighten	הדק	ha-DEK
Windlass	מוט סיבוב	MOT si-BUV
Above the wound	מעל הפצע	me-AL ha-PE-tsa
Time of application	זמן הנחה	ZMAN ha-na-KHA
Shock	הלם	HE-lem
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
The child needs a tourniquet	הילד צריך חוסם עורקים	ha-YE-led tsa-RIKH kho-SEM or-KIM
The bleeding won’t stop	הדימום לא נעצר	ha-di-MUM LO ne-e-TSAR

Source: Magen David Adom (MDA) — MDA Tourniquet Application Guidelines — Pediatric
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16

Personal reference only — not medical advice
Daniel Rosehill · share with attribution