

Personal reference only — not medical advice

Daniel Rosehill · share with attribution

Unconscious Hypoglycemia



Hyperglycemia Pathway



Reference Notes

DO NOT:

✕ **Do NOT administer insulin** — insulin dosing requires medical expertise. Leave to MDA paramedics.
✕ **Do NOT give food or drink to an unconscious patient** — risk of choking and aspiration.
✕ **Do NOT give diet drinks or sugar-free foods** for hypoglycemia — they will not raise blood sugar.
✕ **Do NOT give sugary foods high in fat** (chocolate bars, ice cream) as first-line treatment — fat slows absorption.
✕ **Do NOT withhold sugar when in doubt** — giving sugar to a hypo patient is life-saving; giving sugar to a hyper patient will not cause significant immediate harm.
✕ **Do NOT leave the patient unattended** — condition can deteriorate rapidly.
✕ **Do NOT delay calling 101** if unconscious, seizing, or not improving after sugar.
✕ **Do NOT disconnect an insulin pump roughly** — disconnect gently to avoid damage.

Equipment needed:

— Phone to call 101 (MDA) or 1221 (United Hatzalah)
— Glucometer and test strips (if available)
— Rapidly absorbed sugar (glucose tablets, glucogel, juice, sugar cubes, honey)
— Complex carbohydrate for follow-up (bread, crackers)
— Glucagon injection kit (if available and trained)

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Diabetes	סוכרת	su-KE-ret
Low blood sugar	סוכר נמוך	SU-kar na-MUKH
High blood sugar	סוכר גבוה	SU-kar ga-VO-ah
Sugar	סוכר	SU-kar
Insulin	אינסולין	in-su-LIN
Glucometer	מד סוכר	mad SU-kar
Juice	מיץ	mitz
Unconscious	מחוסר הכרה	me-khu-SAR ha-ka-RA
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
Are you diabetic?	אתה סוכרתי?	a-TA su-kar-TI?
Did you eat today?	אכלת היום?	a-KHAL-ta ha-YOM?

Daniel Rosehill · share with attribution

Source: Magen David Adom (MDA) — How to Treat a Diabetes Patient
URL: https://www.mdais.org/101/diabetes
Publication date: 2026-03-15
Imported: 2026-03-15 · **Last verified:** 2026-03-16

Daniel Rosehill · share with attribution