

Personal reference only — not medical advice

Daniel Rosehill · share with attribution

1 / 3

Reference Notes

DO NOT:

✕ **Do NOT give cola, fruit juice, plain water alone, or tea** — these do not replace electrolytes and can worsen dehydration.
✕ **Do NOT use homemade salt/sugar solutions** unless ORS is completely unavailable.
✕ **Do NOT force large volumes of fluid at once** — small frequent sips are more effective.
✕ **Do NOT delay calling 101 for an infant under 3 months** with any signs of dehydration.
✕ **Do NOT ignore a sunken fontanelle** in an infant — this is a key dehydration indicator.
✕ **Do NOT give anti-diarrhoea medications** to children without medical direction.
✕ **Do NOT stop breastfeeding** — continue alongside ORS.

Equipment needed:

— ORS (Oral Rehydration Solution) packets
— Spoon or oral syringe (for infant fluid administration)
— Phone (to call 101)
— Thermometer (to check for fever)

Hebrew Terminology — מונחים בעברית

English	עברית	Pronunciation
Dehydration	התייבשות	hit-yab-SHUT
Oral rehydration	שתייה מחדש	shti-YA me-kha-DASH
ORS / electrolytes	תמיסת מלחים	tmi-SAT mla-KHIM
Fontanelle	מרפס	mar-PES
Vomiting	הקאות	ha-ka-OT
Diarrhoea	שלשולים	shil-shu-LIM
Thirsty	צמא	tsa-ME
Pharmacy	בית מרקחת	beit mir-KA-khat
Ambulance	אמבולנס	am-bu-LANS
Emergency	חירום	khe-RUM
The child is dehydrated	הילד מיובש	ha-YE-led me-yu-BASH
He/she is vomiting	הוא/היא מקיא/ה	hu/hi ma-KI/ma-ki-A

Source: Magen David Adom (MDA) — MDA Dehydration First Aid Guidelines — Pediatric
URL: https://www.mdais.org/101/first-aid
Publication date: 2026-01-01
Imported: 2026-03-16 · **Last verified:** 2026-03-16