Fludarabine and clofarabine induce caspase 3 cleavage and a heat shock response in AML cells

DEMO MANUSCRIPT, NOT A REAL PAPER. Built by the Below the Watermark team for Gator Hack V from openly licensed figures, with two duplications we made on purpose. The last page lists the sources and our edits.

Abstract. Combination chemotherapy with nucleoside analogs is used in acute myeloid leukemia (AML). We examined progranulin, heat shock proteins and markers of apoptosis (cleaved caspase 3, PARP1 cleavage and gamma-H2AX) by western blot in cells exposed to fludarabine and clofarabine.




Figure 1. Progranulin and vinculin over time and after treatment (western blot).




Figure 2. Heat shock proteins and Nrf 2 / HO 1 after oxidative stress (western blot).




Figure 3. Cleaved caspase 3, PARP1 and gamma-H2AX after combination treatment (western blot).

Sources and our edits (all figures CC BY):
- First figure: Fig. 2 of PMC10755782 (doi:10.1016/j.jbc.2023.105475). We replaced the M10 Vinculin strip in panel A with the HO 1 blot from the second figure, resized. This is the within-paper duplication.
- Second figure: Fig. 4 of PMC10352194 (doi:10.1007/s12192-023-01349-6), unaltered.
- Third figure: part of Fig. 1 of PMC10701888 (doi:10.3389/fonc.2023.1287444), cropped and resized by us. This is the across-paper duplication.

Nothing here reflects anything the original authors did; their published figures are unaltered.