

Table of contents

1 Mapping Oscillatory and 1/f Broadband Spectral Features Across Cortical Regions and Recording Modalities	1
1.1 Introduction	1
1.2 Datasets	4
1.3 Methods	6
1.4 Results	8
1.5 Discussion	11
1.6 Conclusion	12
1.7 Ideas	12
1.8 References	13

1 Mapping Oscillatory and 1/f Broadband Spectral Features Across Cortical Regions and Recording Modalities

1.1 Introduction

Cortical organisation of the brain is often defined by cytoarchitecture and connectivity. However, cortical regions also differ in their intrinsic activity, including their electrophysiological dynamics in the resting state. Intracranial recordings show that different brain areas have distinct spectral profiles: characteristic patterns of oscillatory peaks and broadband spectral shape that together shape regional electrophysiological signatures (Frauscher et al. 2018; Groppe et al. 2013). The importance of understanding resting-state intrinsic activity also help in understanding/predicting task response (Im-mink et al. 2021) and the is often used and basic for comparison when looking for biomarkers to detect early disease onset, of as the help in diagnosis.

An important reference point for understanding these signatures is a normative atlas of intracranial EEG published by Frauscher et al. (2018). This atlas initially included 1785 channels from 106 epilepsy patients, distributed across 38 cortical regions during resting wakefulness with eyes closed. Using k-means clustering and comparison against a no-peak reference set, the authors identified region-specific patterns of oscillatory peaks across canonical frequency bands. They also characterised the non-oscillatory component of the signal using a scale-free model. This atlas remains one of the most comprehensive open-access resources for normative iEEG spectral features, and it provides a useful external reference for evaluating spectral maps obtained with non-invasive methods. Comparing these features across recording modalities is methodologically difficult. iEEG provides a relatively direct measure of local field potentials with limited volume conduction, making it a useful reference for regional spectral organisation. Its signal quality advantage is substantial: Ball et al. (2009) reported that the signal-to-noise ratio

of invasive EEG is 20 to more than 100 times better than simultaneously obtained scalp EEG. However, iEEG is clinically constrained, spatially sparse, and biased by implantation coverage and patient populations.

In source-reconstructed MEG, similar regional structure has been documented through complementary descriptors. Keitel and Gross (2016) showed that resting-state spectral profiles are distinctive enough to fingerprint individual cortical areas. Mellem et al. (2017) found that within most regions multiple frequency bands dominate at once, so regional identity is distributed across the spectrum. Mahjoory et al. (2020) reported, across 187 participants, a posterior-to-anterior decrease in dominant peak frequency that anticorrelates with cortical thickness as a proxy for hierarchical level. Niso et al. (2019) demonstrated that source-space MEG power maps of this kind are reproducible with a standardised pipeline. Shafiei et al. (2023) brought this picture even sharper linking intrinsic electrophysiological activity in MEG with cortical micro-architecture. This approach have however still some limitation,as the orginal papers discussed and Tait et al. (2021) s that source reconstruction can introduce systematic distortions and variability that propagate into spectral parameterisation.

Source-reconstructed HD-EEG, by contrast, like MEG is non-invasive method and can provide whole-head coverage, but source reconstruction can introduce spatial blurring and method-specific biases that affect the estimated spectral profiles (Liu et al. 2018; Lai et al. 2018; Piastra et al. 2020). While MEG will provide a lot of excellent qualities most labs while have not access to it due to its costs, and there the question about spatial organisation of oscialltions is less established. The present chapter uses @Frauscher et al. (2018) as a benchmark for source-reconstructed high-density EEG. The aim is not only simply to ask whether non-invasive recordings can reproduce intracranial spectral maps. Rather, this chapter is one part of the broader thesis question of how oscillatory and 1/f broadband features of the power spectrum should be quantified, interpreted, and compared across datasets.

Source-reconstructed HD EEG is useful for this purpose because it provides a non-invasive estimate of cortical activity that is realively easy and cheaply available and used, but reconstructed sources are not always reliable. Their spatial and spectral properties depend on the inverse method, electrode density, head model, signal-to-noise ratio, and spatial leakage. Therefore, the chapter treats cross-modal comparison not as a simple validation exercise, but as a way to examine which spectral features are robust to changes in recording modality and which are more sensitive to the assumptions of spectral parameterisation and source reconstruction. the initial work in this chapter was done at the beginning of the phd project, with the update in 2026.

Related work provides useful points of comparison for the present chapter. Afnan et al. (2023) validated MEG source imaging of resting-state oscillatory patterns against the Frauscher iEEG atlas. Using 45 healthy participants and the same 38 regions of interest, they found that MEG-estimated spectra were more comparable to iEEG after aperiodic components were removed, and that lateral regions were recovered more accurately than deep or medial structures. Janiukstyte et al. (2023) presented source-localised resting-state scalp EEG maps and showed that systematic spatial variation in spectral features can be detected with scalp EEG, although with reduced spatial specificity. These studies show that regional spectral organisation can be partly preserved across modalities. In the present chapter, they are treated as converging benchmarks rather than as the original motivation: the focus here is specifically on the HD-EEG source-reconstruction pipeline used in this thesis and on the separation of periodic and aperiodic spectral features.

Given certain assumption about the signal, the power spectrum may be operationally decomposed into two component types: narrowband rhythmic oscillations and a broadband (power decreasing with frequency $1/f$ -like) background. Classical analyses focused predominantly on oscillations, but the broadband component also carries meaningful neurophysiological information (Donoghue et al. 2020). As we discussed in Chapter 1, this separation is operationally useful but rests on modelling assumptions that do not guarantee the two components types reflect independent physiological processes. Still, failing to model the broadband background can produce apparent difference in oscillatory peaks that actually reflect spectral variation in the background rather than true difference to narrowband rhythms. Conversely, temporally varying oscillatory activity can contribute to what looks like broadband structure.

The $1/f$ like broadband component also shows regional structure. Frauscher et al. (2018) found regional variation in the scale-free component of intracranial EEG, including stronger scale-free values in visual regions (also observing the knee-like structure in low frequencies of the spectrum). Armonaitė et al. (2026) extended this line of work by showing that scale-free spectral properties across brain regions can serve as regional functional fingerprints, with the high-frequency exponent distinguishing cortical from subcortical areas. These findings suggest that regional electrophysiological identity may be expressed not only in oscillatory peaks, but also in broadband parameters such as spectral slope and offset.

This chapter addresses three related questions. First, which aspects of regional spectral organisation reported in intracranial recordings are also visible in source-reconstructed HD-EEG? Second, do oscillatory and aperiodic features show comparable spatial organisation, or do they reveal different aspects of cortical spectral structure? Third, how do different ways of quantifying oscillatory activity relate to one another once broadband $1/f$ structure is modelled?

To answer these questions, I compare spectral features across the Frauscher iEEG atlas and source-reconstructed HD-EEG data. I examine both oscillatory and broadband components of the power spectrum, using the original Frauscher-style peak detection procedure alongside spectral parameterisation. To assess robustness against methodological choices, I use two complementary decomposition approaches: `specparam` and IRASA. As discussed in Chapter 1 ([?@sec-specparam-geometry](#)), the parameters produced by these methods are model- and range-dependent descriptors of spectral shape. The exponent depends on the frequency range over which it is estimated ([Gerster et al. 2022](#); [Boncompte et al. 2026](#)), and the offset and exponent are structurally coupled by the geometry of the fit. These caveats apply throughout the chapter. Thus, the goal is not to treat any single parameter as a direct physiological quantity, but to determine which spectral features are regionally organised, which are preserved across modalities, and which are sensitive to the assumptions of the analysis pipeline.

There are some limitation in the realisation of current approach. The Frauscher iEEG atlas was recorded with eyes closed, whereas the HD-EEG dataset used here was recorded with eyes open. Eye state affects oscillatory activity, most strongly posterior alpha power, and may also affect aperiodic exponent and offset ([Geller et al. 2014](#)). Where cross-modal agreement is observed, it is therefore observed despite differences in recording condition. Where cross-modal agreement is weak, modality and condition cannot be fully separated with the data used here. This limitation is considered again in the Discussion.

1.2 Datasets

1.2.a Intracranial data

The intracranial EEG data were obtained from the MNI Open iEEG Atlas ([Frauscher et al. 2018](#)), a normative resource compiled from 106 patients with refractory epilepsy recorded at three centres. Only channels implanted in brain regions that were determined to be non-epileptogenic throughout the monitoring period were retained, yielding 1772 channels (13 channels were removed from initial release as were later judged erroneous on visual inspection) distributed across 38 cortical regions defined by the MICCAI atlas. Figure 1 provides a spatial overview of channel coverage in the atlas, distinguishing depth/stereo-EEG contacts from cortical grid and strip contacts across both hemispheres.




Figure 1: Spatial distribution of the MNI Open iEEG Atlas channels and regional reference centroids in MNI space. **(A)** Glass-brain visualization of the 1785 intracranial EEG channels, with depth and stereo-EEG contacts shown in blue and cortical grid/strip contacts shown in orange. For visualization, left-hemisphere channel coordinates were assigned negative x-values based on the hemisphere label. **(B)** Glass-brain visualization of the regional reference centroids used in the grouped regional analyses, shown as enlarged magenta markers.

These 38 regions were derived from the consolidation of 20 smaller MICCAI labels into 8 functional groups to ensure sufficient channel density (min. 5 per region), while 16 labels were omitted due to lack of coverage. For each channel, 60 seconds of resting wakefulness data with eyes closed were available.

Signals were band-pass filtered at 0.5-80 Hz and resampled to 200 Hz. Power spectral density (PSD) was estimated using Welch’s method with 2-second Hamming windows and 50% overlap, resulting in a frequency resolution of 0.5 Hz. Spectral densities were normalised to unit total power, making them independent of signal amplitude. Full preprocessing details are given in Frauscher et al. (2018).

1.2.b Reconstructed sources

The HD-EEG dataset ($n = 19$) was harmonized with the MNI Open iEEG Atlas to allow for direct cross-modal comparison. High-density EEG dataset was described in Samogin et al. (2019) and Samogin et al. (2020). While the HD-EEG data were recorded with eyes open, the processing pipeline was strictly matched to the iEEG atlas: source activity was extracted from using average coordinates from the electrodes within 76 MICCAI-defined ROIs in MNI space, resulting in 1444 virtual sensors, with the grouped regional reference coordinates illustrated in Figure Figure 1, panel B.

For every virtual sensor 60s activity was extracted according to the parameters defined in Frauscher’s paper (sampling frequency 200Hz). Spectral densities were computed using identical Welch parameters (2s windows, 50% overlap, 0.5 Hz resolution).

All spectral data were normalized to unit total power to ensure that the comparison focused on the distribution of frequency peaks (e.g., the occipital alpha or hippocampal delta) rather than absolute amplitude differences between scalp and intracranial sensors.

1.3 Methods

1.3.a Clustering

1.3.a.a Oscillatory peak detection: k-means clustering

To enable direct comparison with the original atlas results, the peak detection procedure of Frauscher et al. (2018) was reproduced. The normalised power spectra were classified into k groups using the k-means algorithm, with 160 features corresponding to the frequency steps from 0.5 to 80 Hz in 0.5 Hz increments, using Euclidean distance and 100 repetitions. The classification was repeated with increasing k until a group without peaks was identified, defined as the group whose mean normalised spectrum was lower than the maximum among the other groups. The 50% of channels closest to the group mean were retained as the no-peak set.

The 160 frequency bins were grouped into 22 frequency intervals based on the condition that each bin contains at least 4% of total mean power, yielding boundaries at 0.5, 0.75, 1.25, 1.75, 2.25, 3.25, 3.75, 4.25, 5.25, 6.25, 6.75, 7.75, 8.25, 9.25, 10.25, 11.75, 13.25, 15.25, 17.25, 20.25, 24.25, 31.75, and 80 Hz. The presence of a peak in a given frequency interval for a particular brain region was determined by comparing the distribution of spectral densities of all channels in that region against the distribution of the no-peak set at the same interval, using a one-sided two-sample Kolmogorov-Smirnov test at the 5% significance level with Dunnett’s correction for multiple comparisons across 42 regions (38 cortical regions plus 4 lobes) and 22 frequency intervals.

For the HD-EEG source-reconstructed data, the same procedure was applied. However, the 22 frequency intervals proposed by Frauscher et al. (2018) did not satisfy the 4% power condition for the reconstructed source spectra. New frequency bins satisfying this condition were therefore defined separately for HD-EEG dataset. This deviation from the original procedure is noted as a methodological caveat.

1.3.b Spectral parameterisation: specparam

PSDs were parameterised using the specparam algorithm (formerly FOOOF; Donoghue et al. (2020)) in fixed aperiodic mode. ! We should use knee here as even frauscher describe the there was knee here.

The algorithm fits an aperiodic component of the form

$$\log_{10} P(f) = b - \chi \cdot \log_{10}(f)$$

where b is the offset and χ is the exponent, and then identifies periodic peaks above this background as Gaussian functions. The following settings were used:

- Fitting frequency range: 1-40 Hz
- Peak width limits: 1.0-8.0 Hz
- Maximum number of peaks: 6
- Minimum peak height: 0.1 (log power)
- Peak threshold: 2.0 standard deviations

For each region in each dataset, the primary aperiodic parameters of interest were the exponent (χ) and offset (b). Oscillatory peaks were characterised by their centre frequency, power, and bandwidth. Model fits were evaluated using R^2 and mean absolute error; fits with R^2 below 0.90 were flagged for inspection.

1.3.c Aperiodic estimation: IRASA

As a complementary approach to specparam, the aperiodic component of each regional time course was estimated using Irregular Resampling Auto-Spectral Analysis (IRASA; Wen and Liu (2016)). IRASA exploits the self-similar properties of fractal processes: the power spectrum is computed at multiple resampling factors ($h = 1.1$ to 1.9 in steps of 0.05), and for each factor, the geometric mean of the up-sampled and down-sampled spectra is taken. The median across resampling factors yields the fractal (aperiodic) component, and the oscillatory residual is obtained by subtraction from the original spectrum.

The fractal exponent was estimated by fitting a line in log-log space to the IRASA-derived aperiodic spectrum over the 1-40 Hz range, matching the specparam fitting range. The slope (with sign reversed) provides an IRASA-based estimate of the aperiodic exponent.

1.3.d Computing correspondence

Regional aggregation and atlas-level correspondence

For each modality, spectral parameters were aggregated within atlas regions. For the iEEG atlas, parameters were averaged (mean) across all channels assigned to each of the 38 MICCAI regions. For the HD-EEG datasets, parameters were first computed per subject per label and then averaged across subjects to produce group-level regional estimates.

Cross-modal correspondence was quantified using Spearman rank correlations across regions. For each parameter (exponent, offset, peak prevalence per frequency band, dominant peak frequency), the vector of regional values from one modality was correlated with the corresponding vector from another modality, yielding a cross-modal correlation

coefficient. Significance was assessed using permutation testing (10,000 permutations, shuffling region labels) to account for the small number of spatial units.

1.3.e Oscillatory re-analysis after aperiodic correction

To evaluate whether regional oscillatory patterns are robust to the presence of aperiodic structure, the oscillatory analyses were repeated on aperiodic-corrected spectra using two approaches:

1. **specparam correction.** The fitted aperiodic component was subtracted from each PSD, producing a flattened spectrum. The Frauscher peak detection procedure (k-means clustering, no-peak set, KS test) was then re-applied to these flattened spectra.
2. **IRASA correction.** The IRASA-derived fractal component was subtracted from each PSD, yielding the oscillatory residual. The same peak detection procedure was applied to this residual.

Stability of oscillatory findings was operationalised as follows: for each region $\times$ frequency bin combination, the presence of a significant peak before correction was compared with the presence of a significant peak after correction. Peaks that survived both corrections were considered robust; peaks that disappeared after correction were considered potentially confounded by aperiodic variation. This comparison was performed separately for the specparam and IRASA corrections, and the agreement between the two methods was assessed.

1.4 Results

1.4.a Reproducing the Frauscher oscillatory atlas

The k-means clustering procedure applied to the iEEG atlas yielded eight spectral groups, closely matching the original classification reported by Frauscher et al. (2018). Seven groups were characterised by prominent peaks in one or more canonical frequency bands (delta, theta, alpha, low beta, high beta, gamma, and mixed), while one group - the no-peak set - exhibited a smooth, monotonically decreasing spectrum with no visible narrowband peaks. The overall structure of these clusters was highly similar to the published result, although the absolute scaling of normalised spectral density differed slightly. The no-peak set served as the reference distribution for all subsequent peak detection tests.

Applying the same procedure to the HD-EEG source-reconstructed data (Mantini dataset) yielded six spectral groups. The reduced number of clusters indicates a narrower range of spectral shapes than in iEEG, consistent with the spatial smoothing introduced by source reconstruction. The HD-EEG no-peak cluster had a flatter profile than its iEEG counterpart, with stronger concentration of power in the alpha range across most clusters,

likely reflecting both the eyes-open recording condition and the dominance of posterior sources in the source model. Figure 2 presents these clustering results side by side.






Figure 2: Figure 2. Spectral clustering in iEEG and HD-EEG source-reconstructed data. Panel A shows the eight iEEG clusters and Panel B shows the six HD-EEG clusters. In both modalities, one no-peak cluster is visible, while HD-EEG shows fewer distinct spectral classes, consistent with stronger spatial smoothing.

A methodological issue arose when applying the frequency binning procedure to the HD-EEG data. The 22 frequency intervals proposed by Frauscher et al. (2018) were defined so that each bin contained at least 4% of total mean power in the iEEG data. This condition was not satisfied for the source-reconstructed spectra, where power was more concentrated in the alpha range and the higher-frequency bins fell below the 4% threshold. Dataset-specific frequency bins were therefore computed for the HD-EEG analyses, as described in the Methods. This deviation is noted as a caveat when comparing peak detection results across modalities.

1.4.b Regional oscillatory profiles

Peak detection was performed per region and frequency bin using the Frauscher procedure, separately for each dataset.





1.4.c Regional broadband profiles

Regional broadband parameters were estimated using both specparam (fixed mode, 1-40 Hz) and IRASA, and aggregated at the atlas-region level for each dataset.

1.4.d Cross-modal correspondence

1.4.e Effect of aperiodic correction on oscillatory findings

1.5 Discussion

1.5.a Regional spectral organisation reflects cortical hierarchy

The finding that oscillatory profiles are regionally organised is consistent with alpha being tied to laminar and feedback/feedforward circuit architecture (Jensen and Bonnefond 2026), implying that at least some of the regional oscillatory structure recovered here reflects broader circuit organisation rather than isolated narrowband generators.

1.5.b Partial cross-modal preservation and its limits

1.5.c Aperiodic correction changes the interpretation of oscillatory findings

1.5.d Methodological caveats and limitations

Spectral parameterisation depends on several modelling choices, including the assumed aperiodic functional form, the fitting frequency range, and peak detection thresholds. Results should therefore be interpreted with caution ([Donoghue et al. 2021](#)).

A subset of source-reconstructed spectra exhibited near-flat spectral slopes with exponents close to zero and low model fit quality ($R^2 < 0.70$). These spectra typically showed no clear $1/f$ structure and, where peaks were present, these were confined to the beta range. Such profiles are unlikely to reflect genuine neural dynamics with an unusually flat broadband component. More plausibly, they arise from source labels that receive little coherent neural signal after inverse modelling, so that the reconstructed time course is dominated by spatially smoothed noise or leakage from neighbouring regions ([Tait et al. 2021](#)). Recent biophysical modelling suggests that high-frequency broadband power in scalp EEG may not be neural in origin ([Brake and Khadra 2025](#); [Brake et al. 2024](#)), and this concern extends to source-reconstructed data, where the inverse solution cannot recover signal that was not present at the sensor level. Fitting `specparam`'s fixed aperiodic model to such spectra produces nominally valid parameter estimates, but these should not be interpreted as reflecting cortical broadband dynamics ([Wilson et al. 2024](#); [Gerster et al. 2022](#)). In the present analyses, sources with R^2 below 0.90 were flagged, and the sensitivity of regional results to the inclusion or exclusion of these low-quality fits was assessed.

1.6 Conclusion

This chapter demonstrates the value of jointly examining oscillatory and aperiodic spectral features when mapping regional brain activity. Regional spectral profiles show partial preservation across iEEG and HD-EEG source-reconstructed recordings, suggesting that large-scale spatial organisation is partly recoverable across modalities. At the same time, accounting for the aperiodic background alters the interpretation of oscillatory findings, highlighting the risk of conflating broadband spectral variation with narrowband oscillations. These results underscore the need for cautious, model-aware interpretation of parameterised spectral measures in human EEG research.

1.7 Ideas

The posterior-to-anterior frequency gradient recovered here is consistent with reports from MEG using both parameterised ([Mahjoory et al. 2020](#)) and non-parametric data-driven ([Capilla et al. 2022](#)) approaches, and from iEEG using spectral parameterisation ([Kalamangalam et al. 2020](#)). Our finding that this gradient is partially preserved in HD-

EEG extends the pattern across an additional modality and a coarser spatial resolution regime.

1.8 References

- Afnan, Jawata, Nicolás von Ellenrieder, Jean-Marc Lina, et al. 2023. “Validating MEG Source Imaging of Resting State Oscillatory Patterns with an Intracranial EEG Atlas.” *NeuroImage* 274 (July): 120158. <https://doi.org/10.1016/j.neuroimage.2023.120158>.
- Armonaite, Karolina, Franca Tecchio, Baingio Pinna, Camillo Porcaro, and Livio Conti. 2026. “Scale-Free Neurodynamics as Functional Fingerprint of Brain Regions.” *Bioengineering* 13 (3): 323. <https://doi.org/10.3390/bioengineering13030323>.
- Ball, Tonio, Markus Kern, Isabella Mutschler, Ad Aertsen, and Andreas Schulze-Bonhage. 2009. “Signal Quality of Simultaneously Recorded Invasive and Non-Invasive EEG.” *NeuroImage* 46 (3): 708–16. <https://doi.org/10.1016/j.neuroimage.2009.02.028>.
- Boncompte, Gonzalo, Martin Irani, Jean-Philippe Lachaux, Vicente Medel, and Tomas Ossandon. 2026. “Aperiodic Exponent of Brain Field Potentials Is Dependent on the Frequency Range It Is Estimated.” *IEEE Transactions on Neural Systems and Rehabilitation Engineering* 34: 500–506. <https://doi.org/10.1109/TNSRE.2025.3649641>.
- Brake, Niklas, Flavie Duc, Alexander Rokos, et al. 2024. “A Neurophysiological Basis for Aperiodic EEG and the Background Spectral Trend.” *Nature Communications* 15 (1): 1514. <https://doi.org/10.1038/s41467-024-45922-8>.
- Brake, Niklas, and Anmar Khadra. 2025. “Contributions of Action Potentials to Scalp EEG: Theory and Biophysical Simulations.” *PLOS Computational Biology* 21 (2): e1012794. <https://doi.org/10.1371/journal.pcbi.1012794>.
- Donoghue, Thomas, Matar Haller, Erik J. Peterson, et al. 2020. “Parameterizing Neural Power Spectra into Periodic and Aperiodic Components.” *Nature Neuroscience* 23 (12): 1655–65. <https://doi.org/10.1038/s41593-020-00744-x>.
- Donoghue, Thomas, Natalie Schaworonkow, and Bradley Voytek. 2021. “Methodological Considerations for Studying Neural Oscillations.” *European Journal of Neuroscience* n/a (n/a). <https://doi.org/10.1111/ejn.15361>.
- Frauscher, Birgit, Nicolas von Ellenrieder, Rina Zelman, et al. 2018. “Atlas of the Normal Intracranial Electroencephalogram: Neurophysiological Awake Activity in Different Cortical Areas.” *Brain* 141 (4): 1130–44. <https://doi.org/10.1093/brain/awy035>.
- Geller, Aaron S., John F. Burke, Michael R. Sperling, et al. 2014. “Eye Closure Causes Widespread Low-Frequency Power Increase and Focal Gamma Attenuation in the Human

Electrocorticogram.” *Clinical Neurophysiology* 125 (9): 1764–73. <https://doi.org/10.1016/j.clinph.2014.01.021>.

Gerster, Moritz, Gunnar Waterstraat, Vladimir Litvak, et al. 2022. “Separating Neural Oscillations from Aperiodic 1/f Activity: Challenges and Recommendations.” *Neuroinformatics*, ahead of print, April. <https://doi.org/10.1007/s12021-022-09581-8>.

Groppe, David M., Stephan Bickel, Corey J. Keller, et al. 2013. “Dominant Frequencies of Resting Human Brain Activity as Measured by the Electrocorticogram.” *NeuroImage* 79 (October): 223–33. <https://doi.org/10.1016/j.neuroimage.2013.04.044>.

Immink, Maarten A., Zachariah R. Cross, Alex Chatburn, James Baumeister, Matthias Schlesewsky, and Ina Bornkessel-Schlesewsky. 2021. “Resting-State Aperiodic Neural Dynamics Predict Individual Differences in Visuomotor Performance and Learning.” *Human Movement Science* 78 (August): 102829. <https://doi.org/10.1016/j.humov.2021.102829>.

Janiukstyte, Vytene, Thomas W. Owen, Umair J. Chaudhary, et al. 2023. “Normative Brain Mapping Using Scalp EEG and Potential Clinical Application.” *Scientific Reports* 13 (1): 13442. <https://doi.org/10.1038/s41598-023-39700-7>.

Jensen, Ole, and Mathilde Bonnefond. 2026. “The Alpha Rhythm: From Physiology to Behaviour.” *Physiological Reviews*, ahead of print, February. <https://doi.org/10.1152/physrev.00001.2025>.

Keitel, Anne, and Joachim Gross. 2016. “Individual Human Brain Areas Can Be Identified from Their Characteristic Spectral Activation Fingerprints.” *PLOS Biology* 14 (6): e1002498. <https://doi.org/10.1371/journal.pbio.1002498>.

Lai, Margherita, Matteo Demuru, Arjan Hillebrand, and Matteo Fraschini. 2018. “A Comparison Between Scalp- and Source-Reconstructed EEG Networks Scientific Reports.” *Scientific Reports* 8 (1): 12269. <https://doi.org/10.1038/s41598-018-30869-w>.

Liu, Quanying, Marco Ganzetti, Nicole Wenderoth, and Dante Mantini. 2018. “Detecting Large-Scale Brain Networks Using EEG: Impact of Electrode Density, Head Modeling and Source Localization.” *Frontiers in Neuroinformatics* 12 (March): 4. <https://doi.org/10.3389/fninf.2018.00004>.

Mahjoory, Keyvan, Jan-Mathijs Schoffelen, Anne Keitel, and Joachim Gross. 2020. “The Frequency Gradient of Human Resting-State Brain Oscillations Follows Cortical Hierarchies.” *eLife* 9 (August): e53715. <https://doi.org/10.7554/eLife.53715>.

Mellem, Monika S., Sophie Wohltjen, Stephen J. Gotts, Avniel Singh Ghuman, and Alex Martin. 2017. “Intrinsic Frequency Biases and Profiles Across Human Cortex.” *Journal of Neurophysiology* 118 (5): 2853–64. <https://doi.org/10.1152/jn.00061.2017>.

- Niso, Guiomar, Francois Tadel, Elizabeth Bock, Martin Cousineau, Andrés Santos, and Sylvain Baillet. 2019. “Brainstorm Pipeline Analysis of Resting-State Data From the Open MEG Archive.” *Frontiers in Neuroscience* 13 (April). <https://doi.org/10.3389/fnins.2019.00284>.
- Piastra, Maria Carla, Andreas Nüßing, Johannes Vorwerk, Maureen Clerc, Christian Engwer, and Carsten H. Wolters. 2020. “A Comprehensive Study on Electroencephalography and Magnetoencephalography Sensitivity to Cortical and Subcortical Sources.” *Human Brain Mapping* 42 (4): 978–92. <https://doi.org/10.1002/hbm.25272>.
- Samogin, Jessica, Quanying Liu, Marco Marino, Nicole Wenderoth, and Dante Mantini. 2019. “Shared and Connection-Specific Intrinsic Interactions in the Default Mode Network.” *NeuroImage* 200 (October): 474–81. <https://doi.org/10.1016/j.neuroimage.2019.07.007>.
- Samogin, Jessica, Marco Marino, Camillo Porcaro, et al. 2020. “Frequency-Dependent Functional Connectivity in Resting State Networks.” *Human Brain Mapping* 41 (18): 5187–98. <https://doi.org/10.1002/hbm.25184>.
- Shafiei, Golia, Ben D. Fulcher, Bradley Voytek, Theodore D. Satterthwaite, Sylvain Baillet, and Bratislav Misic. 2023. “Neurophysiological Signatures of Cortical Micro-Architecture.” *Nature Communications* 14 (1): 6000. <https://doi.org/10.1038/s41467-023-41689-6>.
- Tait, Luke, Ayşegül Özkan, Maciej J. Szul, and Jiaxiang Zhang. 2021. “A Systematic Evaluation of Source Reconstruction of Resting MEG of the Human Brain with a New High-Resolution Atlas: Performance, Precision, and Parcellation.” *Human Brain Mapping* 42 (14): 4685–707. <https://doi.org/10.1002/hbm.25578>.
- Wen, Haiguang, and Zhongming Liu. 2016. “Separating Fractal and Oscillatory Components in the Power Spectrum of Neurophysiological Signal.” *Brain Topography* 29 (1): 13–26. <https://doi.org/10.1007/s10548-015-0448-0>.
- Wilson, Luc E., Jason da Silva Castanheira, Benjamin Lévesque Kinder, and Sylvain Baillet. 2024. *Model Selection for Spectral Parameterization*. bioRxiv. <https://doi.org/10.1101/2024.08.01.606216>.