

PSYCHOLOGY – DEPT. OF DATA ANALYSIS

PhD-student: Daniel Borek – Promotor: Prof. dr. Daniele Marinazzo
Doctoral advisory committee: Prof. dr. Dante Mantini, Prof dr. Tom Verguts

PARAMETRIZATION OF POWER SPECTRA TO BETTER UNDERSTAND COGNITIVE PROCESSES

0 My PhD research is part of the GOA-BOF project "Brain Interactions and Neural Dynamics (BIND)", which investigates memory and cognitive control using neuroimaging data.

Oscillations

- 1 Neural oscillatory activity play an important role during maintaining and processing information on different scales
- a proposed mechanism for human flexibility (memory, attention) and cognitive control (Verguts, 2017)
- "when oscillations do exist, they often manifest as "bumps" on top of the 1/f slope in the power spectrum" (He, 2014)
- Non-oscillatory, 1/f-like activity in the background carries physiological information too (Voytek, 2015)
- oscillatory content of the signal may change during a cognitive task or resting state

Possible pitfalls


- 2 standard: comparison of mean alpha-band power between conditions
- nonzero power in a narrow frequency band is not necessarily a sign of oscillating activity there
- a priori definition of canonical bands not consistent across studies
- information about 1/f component is put aside

Parametrizing spectra


- 3 data-driven approach built on standard Fourier analysis
- model of the power spectrum as a combination of additive distinct functional processes (aperiodic and periodic)
- an aperiodic 'background' component, reflecting 1/f like characteristics, modeled with an exponential fit, oscillatory components as Gaussians

foof toolbox algorithm in Python in Haller, 2018


1. Input Power Spectra




2. Fit 1/f Background




3. Remove Background Process



4. Fit Gaussians to Oscillatory Peaks



Haller, 2018



PROJECT

- Investigate the oscillatory dynamics during processes underlying cognitive flexibility
- Estimation of different spectral components in the signal. combine existing methods & improve them
- Learn more about the regional distribution of neurophysiological activity

Idea 1: compare localisations of oscillatory components in reconstructed sources and intracranial recordings during resting state

- datasets: Frauscher, 2018 (106 patients, 1772 channels, 60 s closed eyes activity from healthy brain regions) and Liu, 2017 (19 participants, sources extracted to match those regions)
- quantify the effect of solving the inverse problem
- validate a parametrical approach to a signal by reproducing results about topographical distributions of oscillations.
- code as a toolbox in Python

Idea 2: modeling alpha power changes and frequency shifts in attentional and cognitive control processes

- alpha oscillations in the EEG as a reliable marker of attention allocation (Janssens, 2017)
- collaborate between BIND group members
- inter- and intra-individual variability in alpha peak frequency
- take into account non-stationarity of the signal (single bursts)

Contact

Beginjnhoflaan 464, 9000 Ghent DanielBorek@UGent.be danieltomasz

Acknowledgements

This research is funded by Special Research Fund of Ghent University.

REFERENCES

- Frauscher, B., von Ellenrieder, N., Zemann, R., Doležalová, I., Minotti, L., Olivier, A., ... Gotman, J. (2018). Atlas of the normal intracranial electroencephalogram: neurophysiological awake activity in different cortical areas. *Brain*, 141(4), 1130–1144. <https://doi.org/10/gc5ct7>
- Haller, M., Donoghue, T., Peterson, E., Varma, P., Sebastian, P., Gao, R., ... Voytek, B. (2018). Parameterizing neural power spectra. Retrieved from *BioRxiv*, 299859. <https://doi.org/10.1101/299859>
- He, B. J. (2014). Scale-free brain activity: past, present and future. *Trends in Cognitive Sciences*, 18(9), 480–487. <https://doi.org/10/gctmp2>
- Janssens, C., De Loof, E., Boehler, C. N., Pourtois, G., & Verguts, T. (2018). Occipital alpha power reveals fast attentional inhibition of incongruent distractors. *Psychophysiology*, 55(3), e13011. <https://doi.org/10.1111/psyp.13011>
- Liu, Q., Farahibozorg, S., Porcaro, C., Wenderoth, N., & Mantini, D. (2017). Detecting large-scale networks in the human brain using high-density electroencephalography: Imaging Brain Networks with High Density EEG. *Human Brain Mapping*, 38(9), 4631–4643. <https://doi.org/10/gbsmn3>
- Verguts, T. (2017). Binding by Random Bursts: A Computational Model of Cognitive Control. *Journal of Cognitive Neuroscience*, 29(6), 1103–1118. https://doi.org/10.1162/jocn_a_01117
- Voytek, B., Kramer, M. A., Case, J., Lepage, K. Q., Tempesta, Z. R., Knight, R. T., & Gazzaley, A. (2015). Age-Related Changes in 1/f Neural Electrophysiological Noise. *Journal of Neuroscience*, 35(38), 13257–13265. <https://doi.org/10.1523/JNEUROSCI.2332-14.2015>