

On spaces with small dense sets

Dániel T. Soukup

<http://www.logic.univie.ac.at/~soukupd73/>



universität
wien

Introduction

Goal: study topological spaces X which have a **dense set** $\bigcup\{D_n : n \in \omega\}$ so that D_n is *small*.

- separable spaces: D_n can be chosen singleton/finite/countable.
- **d-separable spaces**: D_n can be chosen discrete;
- **e-separable spaces**: D_n can be chosen closed and discrete;
- how do **products/powers** behave?
- study related **cardinal functions**.

Joint work with Rodrigo R. Dias.

- disclaimer: space = regular, topological space.

Introduction

Goal: study topological spaces X which have a **dense set** $\bigcup\{D_n : n \in \omega\}$ so that D_n is *small*.

- separable spaces: D_n can be chosen singleton/finite/countable.
- **d-separable spaces**: D_n can be chosen discrete;
- **e-separable spaces**: D_n can be chosen closed and discrete;
- how do **products/powers** behave?
- study related **cardinal functions**.

Joint work with Rodrigo R. Dias.

- disclaimer: space = regular, topological space.

Introduction

Goal: study topological spaces X which have a **dense set** $\bigcup\{D_n : n \in \omega\}$ so that D_n is *small*.

- separable spaces: D_n can be chosen singleton/finite/countable.
- **d-separable spaces**: D_n can be chosen discrete;
- **e-separable spaces**: D_n can be chosen closed and discrete;
- how do **products/powers** behave?
- study related **cardinal functions**.

Joint work with Rodrigo R. Dias.

- disclaimer: space = regular, topological space.

Introduction

Goal: study topological spaces X which have a **dense set** $\bigcup\{D_n : n \in \omega\}$ so that D_n is *small*.

- separable spaces: D_n can be chosen singleton/finite/countable.
- **d-separable spaces**: D_n can be chosen discrete;
- **e-separable spaces**: D_n can be chosen closed and discrete;
- how do **products/powers** behave?
- study related **cardinal functions**.

Joint work with Rodrigo R. Dias.

- disclaimer: space = regular, topological space.

Introduction

Goal: study topological spaces X which have a **dense set** $\bigcup\{D_n : n \in \omega\}$ so that D_n is *small*.

- separable spaces: D_n can be chosen singleton/finite/countable.
- **d-separable spaces**: D_n can be chosen discrete;
- **e-separable spaces**: D_n can be chosen closed and discrete;
- how do **products/powers** behave?
- study related **cardinal functions**.

Joint work with Rodrigo R. Dias.

- disclaimer: space = regular, topological space.

Introduction

Goal: study topological spaces X which have a **dense set** $\bigcup\{D_n : n \in \omega\}$ so that D_n is *small*.

- separable spaces: D_n can be chosen singleton/finite/countable.
- **d-separable spaces**: D_n can be chosen discrete;
- **e-separable spaces**: D_n can be chosen closed and discrete;
- how do **products/powers** behave?
- study related **cardinal functions**.

Joint work with Rodrigo R. Dias.

- disclaimer: space = regular, topological space.

Introduction

Goal: study topological spaces X which have a **dense set** $\bigcup\{D_n : n \in \omega\}$ so that D_n is *small*.

- separable spaces: D_n can be chosen singleton/finite/countable.
- **d-separable spaces**: D_n can be chosen discrete;
- **e-separable spaces**: D_n can be chosen closed and discrete;
- how do **products/powers** behave?
- study related **cardinal functions**.

Joint work with Rodrigo R. Dias.

- disclaimer: space = regular, topological space.

Introduction

Goal: study topological spaces X which have a **dense set** $\bigcup\{D_n : n \in \omega\}$ so that D_n is *small*.

- separable spaces: D_n can be chosen singleton/finite/countable.
- **d-separable spaces**: D_n can be chosen discrete;
- **e-separable spaces**: D_n can be chosen closed and discrete;
- how do **products/powers** behave?
- study related **cardinal functions**.

Joint work with Rodrigo R. Dias.

- disclaimer: space = regular, topological space.



Introduction

Goal: study topological spaces X which have a **dense set** $\bigcup\{D_n : n \in \omega\}$ so that D_n is *small*.

- separable spaces: D_n can be chosen singleton/finite/countable.
- **d-separable spaces**: D_n can be chosen discrete;
- **e-separable spaces**: D_n can be chosen closed and discrete;
- how do **products/powers** behave?
- study related **cardinal functions**.

Joint work with Rodrigo R. Dias.

- disclaimer: space = regular, topological space.



Separable spaces and products

Recall: X is separable iff there is a countable dense subset of X i.e. $d(X) = \aleph_0$.

If X is separable then X has a basis of size $\leq \mathfrak{c} = 2^{\aleph_0}$ and $|X| \leq 2^{\mathfrak{c}}$.

[Pondiczery 1944] If X is separable then $X^{\mathfrak{c}}$ is separable but $X^{\mathfrak{c}^+}$ is not.

- fix a countable dense D in X ,
- $\mathfrak{c} = |\mathbb{R}|$, let $\mathcal{Q} \subseteq \mathcal{P}(\mathfrak{c})$ correspond to rational intervals in \mathfrak{c} ,
- let $f \in E$ iff $f \in X^{\mathfrak{c}}$ and there are $\{I_k : k < m\}$ from \mathcal{Q} and $d, d_k \in D$ so that $f \upharpoonright I_k = d_k$ and $f \upharpoonright \mathfrak{c} \setminus \bigcup_{k < m} I_k = d$;
- E is still countable and dense in $X^{\mathfrak{c}}$.

Separable spaces and products

Recall: X is separable iff there is a countable dense subset of X i.e. $d(X) = \aleph_0$.

If X is separable then X has a basis of size $\leq \mathfrak{c} = 2^{\aleph_0}$ and $|X| \leq 2^{\mathfrak{c}}$.

[Pondiczery 1944] If X is separable then $X^{\mathfrak{c}}$ is separable but $X^{\mathfrak{c}^+}$ is not.

- fix a countable dense D in X ,
- $\mathfrak{c} = |\mathbb{R}|$, let $\mathcal{Q} \subseteq \mathcal{P}(\mathfrak{c})$ correspond to rational intervals in \mathfrak{c} ,
- let $f \in E$ iff $f \in X^{\mathfrak{c}}$ and there are $\{I_k : k < m\}$ from \mathcal{Q} and $d, d_k \in D$ so that $f \upharpoonright I_k = d_k$ and $f \upharpoonright \mathfrak{c} \setminus \bigcup_{k < m} I_k = d$;
- E is still countable and dense in $X^{\mathfrak{c}}$.

Separable spaces and products

Recall: X is separable iff there is a countable dense subset of X i.e. $d(X) = \aleph_0$.

If X is separable then X has a basis of size $\leq \mathfrak{c} = 2^{\aleph_0}$ and $|X| \leq 2^{\mathfrak{c}}$.

[Pondiczery 1944] If X is separable then $X^{\mathfrak{c}}$ is separable but $X^{\mathfrak{c}^+}$ is not.

- fix a countable dense D in X ,
- $\mathfrak{c} = |\mathbb{R}|$, let $\mathcal{Q} \subseteq \mathcal{P}(\mathfrak{c})$ correspond to rational intervals in \mathfrak{c} ,
- let $f \in E$ iff $f \in X^{\mathfrak{c}}$ and there are $\{I_k : k < m\}$ from \mathcal{Q} and $d, d_k \in D$ so that $f \upharpoonright I_k = d_k$ and $f \upharpoonright \mathfrak{c} \setminus \bigcup_{k < m} I_k = d$;
- E is still countable and dense in $X^{\mathfrak{c}}$.

Separable spaces and products

Recall: X is separable iff there is a countable dense subset of X i.e. $d(X) = \aleph_0$.

If X is separable then X has a basis of size $\leq \mathfrak{c} = 2^{\aleph_0}$ and $|X| \leq 2^{\mathfrak{c}}$.

[Pondiczery 1944] If X is separable then $X^{\mathfrak{c}}$ is separable but $X^{\mathfrak{c}^+}$ is not.

- fix a countable dense D in X ,
- $\mathfrak{c} = |\mathbb{R}|$, let $\mathcal{Q} \subseteq \mathcal{P}(\mathfrak{c})$ correspond to rational intervals in \mathfrak{c} ,
- let $f \in E$ iff $f \in X^{\mathfrak{c}}$ and there are $\{I_k : k < m\}$ from \mathcal{Q} and $d, d_k \in D$ so that $f \upharpoonright I_k = d_k$ and $f \upharpoonright \mathfrak{c} \setminus \bigcup_{k < m} I_k = d$;
- E is still countable and dense in $X^{\mathfrak{c}}$.

Separable spaces and products

Recall: X is separable iff there is a countable dense subset of X i.e. $d(X) = \aleph_0$.

If X is separable then X has a basis of size $\leq \mathfrak{c} = 2^{\aleph_0}$ and $|X| \leq 2^{\mathfrak{c}}$.

[Pondiczery 1944] If X is separable then $X^{\mathfrak{c}}$ is separable but $X^{\mathfrak{c}^+}$ is not.

- fix a countable dense D in X ,
- $\mathfrak{c} = |\mathbb{R}|$, let $\mathcal{Q} \subseteq \mathcal{P}(\mathfrak{c})$ correspond to rational intervals in \mathfrak{c} ,
- let $f \in E$ iff $f \in X^{\mathfrak{c}}$ and there are $\{I_k : k < m\}$ from \mathcal{Q} and $d, d_k \in D$ so that $f \upharpoonright I_k = d_k$ and $f \upharpoonright \mathfrak{c} \setminus \bigcup_{k < m} I_k = d$;
- E is still countable and dense in $X^{\mathfrak{c}}$.

Separable spaces and products

Recall: X is separable iff there is a countable dense subset of X i.e. $d(X) = \aleph_0$.

If X is separable then X has a basis of size $\leq \mathfrak{c} = 2^{\aleph_0}$ and $|X| \leq 2^{\mathfrak{c}}$.

[Pondiczery 1944] If X is separable then $X^{\mathfrak{c}}$ is separable but $X^{\mathfrak{c}^+}$ is not.

- fix a countable dense D in X ,
- $\mathfrak{c} = |\mathbb{R}|$, let $\mathcal{Q} \subseteq \mathcal{P}(\mathfrak{c})$ correspond to rational intervals in \mathfrak{c} ,
- let $f \in E$ iff $f \in X^{\mathfrak{c}}$ and there are $\{I_k : k < m\}$ from \mathcal{Q} and $d, d_k \in D$ so that $f \upharpoonright I_k = d_k$ and $f \upharpoonright \mathfrak{c} \setminus \bigcup_{k < m} I_k = d$;
- E is still countable and dense in $X^{\mathfrak{c}}$.

Separable spaces and products

Recall: X is separable iff there is a countable dense subset of X i.e. $d(X) = \aleph_0$.

If X is separable then X has a basis of size $\leq \mathfrak{c} = 2^{\aleph_0}$ and $|X| \leq 2^{\mathfrak{c}}$.

[Pondiczery 1944] If X is separable then $X^{\mathfrak{c}}$ is separable but $X^{\mathfrak{c}^+}$ is not.

- fix a countable dense D in X ,
- $\mathfrak{c} = |\mathbb{R}|$, let $\mathcal{Q} \subseteq \mathcal{P}(\mathfrak{c})$ correspond to rational intervals in \mathfrak{c} ,
- let $f \in E$ iff $f \in X^{\mathfrak{c}}$ and there are $\{I_k : k < m\}$ from \mathcal{Q} and $d, d_k \in D$ so that $f \upharpoonright I_k = d_k$ and $f \upharpoonright \mathfrak{c} \setminus \bigcup_{k < m} I_k = d$;
- E is still countable and dense in $X^{\mathfrak{c}}$.

[Kurepa 1936] X is d-separable iff there is a σ -discrete dense subset of X .

- introduced as property K_0 in his study of Suslin's problem .

If X is separable or metrizable then X is d-separable.

- [K 1936] a Suslin-continuum is not d-separable.
 - size of discrete sets $\leq \aleph_0 < \aleph_1 \leq$ size of dense sets.
- [Todorcevic 1981]
 - In ZFC, there is a non d-separable continuum.
 - Consistently, any LOTS of density \aleph_1 is d-separable.

[Kurepa 1936] X is d-separable iff there is a σ -discrete dense subset of X .

- introduced as property K_0 in his study of Suslin's problem .

If X is separable or metrizable then X is d-separable.

- [K 1936] a Suslin-continuum is not d-separable.
 - size of discrete sets $\leq \aleph_0 < \aleph_1 \leq$ size of dense sets.
- [Todorcevic 1981]
 - In ZFC, there is a non d-separable continuum.
 - Consistently, any LOTS of density \aleph_1 is d-separable.

[Kurepa 1936] X is d-separable iff there is a σ -discrete dense subset of X .

- introduced as property K_0 in his study of Suslin's problem .

If X is separable or metrizable then X is d-separable.

- [K 1936] a Suslin-continuum is not d-separable.
 - size of discrete sets $\leq \aleph_0 < \aleph_1 \leq$ size of dense sets.
- [Todorcevic 1981]
 - In ZFC, there is a non d-separable continuum.
 - Consistently, any LOTS of density \aleph_1 is d-separable.

[Kurepa 1936] X is d-separable iff there is a σ -discrete dense subset of X .

- introduced as property K_0 in his study of Suslin's problem .

If X is separable or metrizable then X is d-separable.

- [K 1936] a Suslin-continuum is not d-separable.
 - size of discrete sets $\leq \aleph_0 < \aleph_1 \leq$ size of dense sets.
- [Todorcevic 1981]
 - In ZFC, there is a non d-separable continuum.
 - Consistently, any LOTS of density \aleph_1 is d-separable.

[Kurepa 1936] X is d-separable iff there is a σ -discrete dense subset of X .

- introduced as property K_0 in his study of Suslin's problem .

If X is separable or metrizable then X is d-separable.

- [K 1936] a Suslin-continuum is not d-separable.
 - size of discrete sets $\leq \aleph_0 < \aleph_1 \leq$ size of dense sets.
- [Todorcevic 1981]
 - In ZFC, there is a non d-separable continuum.
 - Consistently, any LOTS of density \aleph_1 is d-separable.

[Kurepa 1936] X is d-separable iff there is a σ -discrete dense subset of X .

- introduced as property K_0 in his study of Suslin's problem .

If X is separable or metrizable then X is d-separable.

- [K 1936] a Suslin-continuum is not d-separable.
 - size of discrete sets $\leq \aleph_0 < \aleph_1 \leq$ size of dense sets.
- [Todorcevic 1981]
 - In ZFC, there is a non d-separable continuum.
 - Consistently, any LOTS of density \aleph_1 is d-separable.

[Kurepa 1936] X is d-separable iff there is a σ -discrete dense subset of X .

- introduced as property K_0 in his study of Suslin's problem .

If X is separable or metrizable then X is d-separable.

- [K 1936] a Suslin-continuum is not d-separable.
 - size of discrete sets $\leq \aleph_0 < \aleph_1 \leq$ size of dense sets.
- [Todorcevic 1981]
 - In ZFC, there is a non d-separable continuum.
 - Consistently, any LOTS of density \aleph_1 is d-separable.

[Kurepa 1936] X is d-separable iff there is a σ -discrete dense subset of X .

- introduced as property K_0 in his study of Suslin's problem .

If X is separable or metrizable then X is d-separable.

- [K 1936] a Suslin-continuum is not d-separable.
 - size of discrete sets $\leq \aleph_0 < \aleph_1 \leq$ size of dense sets.
- [Todorcevic 1981]
 - In ZFC, there is a non d-separable continuum.
 - Consistently, any LOTS of density \aleph_1 is d-separable.

[Kurepa 1936] X is d-separable iff there is a σ -discrete dense subset of X .

- introduced as property K_0 in his study of Suslin's problem .

If X is separable or metrizable then X is d-separable.

- [K 1936] a Suslin-continuum is not d-separable.
 - size of discrete sets $\leq \aleph_0 < \aleph_1 \leq$ size of dense sets.
- [Todorcevic 1981]
 - In ZFC, there is a non d-separable continuum.
 - Consistently, any LOTS of density \aleph_1 is d-separable.

Products and d-separable spaces

[Arhangel'skii 1981] Products of d-separable spaces are d-separable.

[Juhász, Szentmiklóssy 2008] X^κ is d-separable if there is a discrete subset of X^κ of size $d(X)$.

Note: $\{x \in 2^\kappa : |x^{-1}(1)| = n\} \subseteq D(2)^\kappa$ is discrete for any $n \in \omega$.

$\Rightarrow D(\kappa) \hookrightarrow D(2)^\kappa \hookrightarrow X^\kappa$.

[Amirdzanov 1977, JS 2008] $X^{d(X)}$ is d-separable for any X .

Products and d-separable spaces

[Arhangel'skii 1981] Products of d-separable spaces are d-separable.

[Juhász, Szentmiklóssy 2008] X^κ is d-separable if there is a discrete subset of X^κ of size $d(X)$.

Note: $\{x \in 2^\kappa : |x^{-1}(1)| = n\} \subseteq D(2)^\kappa$ is discrete for any $n \in \omega$.

$\Rightarrow D(\kappa) \hookrightarrow D(2)^\kappa \hookrightarrow X^\kappa$.

[Amirdzanov 1977, JS 2008] $X^{d(X)}$ is d-separable for any X .

Products and d-separable spaces

[Arhangel'skii 1981] Products of d-separable spaces are d-separable.

[Juhász, Szentmiklóssy 2008] X^κ is d-separable if there is a discrete subset of X^κ of size $d(X)$.

Note: $\{x \in 2^\kappa : |x^{-1}(1)| = n\} \subseteq D(2)^\kappa$ is discrete for any $n \in \omega$.

$\Rightarrow D(\kappa) \hookrightarrow D(2)^\kappa \hookrightarrow X^\kappa$.

[Amirdzanov 1977, JS 2008] $X^{d(X)}$ is d-separable for any X .

Products and d-separable spaces

[Arhangel'skii 1981] Products of d-separable spaces are d-separable.

[Juhász, Szentmiklóssy 2008] X^κ is d-separable if there is a discrete subset of X^κ of size $d(X)$.

Note: $\{x \in 2^\kappa : |x^{-1}(1)| = n\} \subseteq D(2)^\kappa$ is discrete for any $n \in \omega$.

$\Rightarrow D(\kappa) \hookrightarrow D(2)^\kappa \hookrightarrow X^\kappa$.

[Amirdzanov 1977, JS 2008] $X^{d(X)}$ is d-separable for any X .

Products and d-separable spaces

[Arhangel'skii 1981] Products of d-separable spaces are d-separable.

[Juhász, Szentmiklóssy 2008] X^κ is d-separable if there is a discrete subset of X^κ of size $d(X)$.

Note: $\{x \in 2^\kappa : |x^{-1}(1)| = n\} \subseteq D(2)^\kappa$ is discrete for any $n \in \omega$.

$\Rightarrow D(\kappa) \hookrightarrow D(2)^\kappa \hookrightarrow X^\kappa$.

[Amirdzanov 1977, JS 2008] $X^{d(X)}$ is d-separable for any X .

Products and d-separable spaces

[Arhangel'skii 1981] Products of d-separable spaces are d-separable.

[Juhász, Szentmiklóssy 2008] X^κ is d-separable if there is a discrete subset of X^κ of size $d(X)$.

Note: $\{x \in 2^\kappa : |x^{-1}(1)| = n\} \subseteq D(2)^\kappa$ is discrete for any $n \in \omega$.

$\Rightarrow D(\kappa) \hookrightarrow D(2)^\kappa \hookrightarrow X^\kappa$.

[Amirdzanov 1977, JS 2008] $X^{d(X)}$ is d-separable for any X .

Compact spaces and powers

[JS 2008] X^ω is d-separable for any compact X .

- $(\omega^*)^n$ is not d-separable for any $n \in \omega$.

[JS 2008] X^2 has a discrete subset of size $d(X)$ for any compact X .

- Wlog: any non empty open subset has weight $w(X)$ (not trivial).
- find $(x_\alpha, y_\alpha) \in U_\alpha \times V_\alpha \subseteq X^2$ so that $U_\alpha \cap V_\alpha = \emptyset$ for $\alpha < d(X)$,
- there is a open $H \neq \emptyset$ so that $K = \overline{H} \subseteq X \setminus \overline{\{x_\alpha, y_\alpha : \alpha < \beta\}}$,
- $\{U_\alpha, V_\alpha : \alpha < \beta\}$ generate a coarser topology on K than the original compact so cannot be Hausdorff.
- Let $x_\beta, y_\beta \in K$ witness this; then $(x_\beta, y_\beta) \notin U_\alpha \times V_\alpha$ for $\alpha < \beta$.
- Now take any disjoint open U_β, V_β with $(x_\beta, y_\beta) \in U_\beta \times V_\beta \subseteq H^2$.

Compact spaces and powers

[JS 2008] X^ω is d-separable for any compact X .

- $(\omega^*)^n$ is not d-separable for any $n \in \omega$.

[JS 2008] X^2 has a discrete subset of size $d(X)$ for any compact X .

- Wlog: any non empty open subset has weight $w(X)$ (not trivial).
- find $(x_\alpha, y_\alpha) \in U_\alpha \times V_\alpha \subseteq X^2$ so that $U_\alpha \cap V_\alpha = \emptyset$ for $\alpha < d(X)$,
- there is a open $H \neq \emptyset$ so that $K = \overline{H} \subseteq X \setminus \overline{\{x_\alpha, y_\alpha : \alpha < \beta\}}$,
- $\{U_\alpha, V_\alpha : \alpha < \beta\}$ generate a coarser topology on K than the original compact so cannot be Hausdorff.
- Let $x_\beta, y_\beta \in K$ witness this; then $(x_\beta, y_\beta) \notin U_\alpha \times V_\alpha$ for $\alpha < \beta$.
- Now take any disjoint open U_β, V_β with $(x_\beta, y_\beta) \in U_\beta \times V_\beta \subseteq H^2$.

Compact spaces and powers

[JS 2008] X^ω is d-separable for any compact X .

- $(\omega^*)^n$ is not d-separable for any $n \in \omega$.

[JS 2008] X^2 has a discrete subset of size $d(X)$ for any compact X .

- Wlog: any non empty open subset has weight $w(X)$ (not trivial).
- find $(x_\alpha, y_\alpha) \in U_\alpha \times V_\alpha \subseteq X^2$ so that $U_\alpha \cap V_\alpha = \emptyset$ for $\alpha < d(X)$,
- there is a open $H \neq \emptyset$ so that $K = \overline{H} \subseteq X \setminus \overline{\{x_\alpha, y_\alpha : \alpha < \beta\}}$,
- $\{U_\alpha, V_\alpha : \alpha < \beta\}$ generate a coarser topology on K than the original compact so cannot be Hausdorff.
- Let $x_\beta, y_\beta \in K$ witness this; then $(x_\beta, y_\beta) \notin U_\alpha \times V_\alpha$ for $\alpha < \beta$.
- Now take any disjoint open U_β, V_β with $(x_\beta, y_\beta) \in U_\beta \times V_\beta \subseteq H^2$.

Compact spaces and powers

[JS 2008] X^ω is d-separable for any compact X .

- $(\omega^*)^n$ is not d-separable for any $n \in \omega$.

[JS 2008] X^2 has a discrete subset of size $d(X)$ for any compact X .

- Wlog: any non empty open subset has weight $w(X)$ (not trivial).
- find $(x_\alpha, y_\alpha) \in U_\alpha \times V_\alpha \subseteq X^2$ so that $U_\alpha \cap V_\alpha = \emptyset$ for $\alpha < d(X)$,
- there is a open $H \neq \emptyset$ so that $K = \overline{H} \subseteq X \setminus \overline{\{x_\alpha, y_\alpha : \alpha < \beta\}}$,
- $\{U_\alpha, V_\alpha : \alpha < \beta\}$ generate a coarser topology on K than the original compact so cannot be Hausdorff.
- Let $x_\beta, y_\beta \in K$ witness this; then $(x_\beta, y_\beta) \notin U_\alpha \times V_\alpha$ for $\alpha < \beta$.
- Now take any disjoint open U_β, V_β with $(x_\beta, y_\beta) \in U_\beta \times V_\beta \subseteq H^2$.

Compact spaces and powers

[JS 2008] X^ω is d-separable for any compact X .

- $(\omega^*)^n$ is not d-separable for any $n \in \omega$.

[JS 2008] X^2 has a discrete subset of size $d(X)$ for any compact X .

- Wlog: any non empty open subset has weight $w(X)$ (not trivial).
- find $(x_\alpha, y_\alpha) \in U_\alpha \times V_\alpha \subseteq X^2$ so that $U_\alpha \cap V_\alpha = \emptyset$ for $\alpha < d(X)$,
- there is a open $H \neq \emptyset$ so that $K = \overline{H} \subseteq X \setminus \overline{\{x_\alpha, y_\alpha : \alpha < \beta\}}$,
- $\{U_\alpha, V_\alpha : \alpha < \beta\}$ generate a coarser topology on K than the original compact so cannot be Hausdorff.
- Let $x_\beta, y_\beta \in K$ witness this; then $(x_\beta, y_\beta) \notin U_\alpha \times V_\alpha$ for $\alpha < \beta$.
- Now take any disjoint open U_β, V_β with $(x_\beta, y_\beta) \in U_\beta \times V_\beta \subseteq H^2$.

Compact spaces and powers

[JS 2008] X^ω is d-separable for any compact X .

- $(\omega^*)^n$ is not d-separable for any $n \in \omega$.

[JS 2008] X^2 has a discrete subset of size $d(X)$ for any compact X .

- Wlog: any non empty open subset has weight $w(X)$ (not trivial).
- find $(x_\alpha, y_\alpha) \in U_\alpha \times V_\alpha \subseteq X^2$ so that $U_\alpha \cap V_\alpha = \emptyset$ for $\alpha < d(X)$,
- there is a open $H \neq \emptyset$ so that $K = \overline{H} \subseteq X \setminus \overline{\{x_\alpha, y_\alpha : \alpha < \beta\}}$,
- $\{U_\alpha, V_\alpha : \alpha < \beta\}$ generate a coarser topology on K than the original compact so cannot be Hausdorff.
- Let $x_\beta, y_\beta \in K$ witness this; then $(x_\beta, y_\beta) \notin U_\alpha \times V_\alpha$ for $\alpha < \beta$.
- Now take any disjoint open U_β, V_β with $(x_\beta, y_\beta) \in U_\beta \times V_\beta \subseteq H^2$.

Compact spaces and powers

[JS 2008] X^ω is d-separable for any compact X .

- $(\omega^*)^n$ is not d-separable for any $n \in \omega$.

[JS 2008] X^2 has a discrete subset of size $d(X)$ for any compact X .

- Wlog: any non empty open subset has weight $w(X)$ (not trivial).
- find $(x_\alpha, y_\alpha) \in U_\alpha \times V_\alpha \subseteq X^2$ so that $U_\alpha \cap V_\alpha = \emptyset$ for $\alpha < d(X)$,
- there is a open $H \neq \emptyset$ so that $K = \overline{H} \subseteq X \setminus \overline{\{x_\alpha, y_\alpha : \alpha < \beta\}}$,
- $\{U_\alpha, V_\alpha : \alpha < \beta\}$ generate a coarser topology on K than the original compact so cannot be Hausdorff.
- Let $x_\beta, y_\beta \in K$ witness this; then $(x_\beta, y_\beta) \notin U_\alpha \times V_\alpha$ for $\alpha < \beta$.
- Now take any disjoint open U_β, V_β with $(x_\beta, y_\beta) \in U_\beta \times V_\beta \subseteq H^2$.

Compact spaces and powers

[JS 2008] X^ω is d-separable for any compact X .

- $(\omega^*)^n$ is not d-separable for any $n \in \omega$.

[JS 2008] X^2 has a discrete subset of size $d(X)$ for any compact X .

- Wlog: any non empty open subset has weight $w(X)$ (not trivial).
- find $(x_\alpha, y_\alpha) \in U_\alpha \times V_\alpha \subseteq X^2$ so that $U_\alpha \cap V_\alpha = \emptyset$ for $\alpha < d(X)$,
- there is a open $H \neq \emptyset$ so that $K = \overline{H} \subseteq X \setminus \overline{\{x_\alpha, y_\alpha : \alpha < \beta\}}$,
- $\{U_\alpha, V_\alpha : \alpha < \beta\}$ generate a coarser topology on K than the original compact so cannot be Hausdorff.
- Let $x_\beta, y_\beta \in K$ witness this; then $(x_\beta, y_\beta) \notin U_\alpha \times V_\alpha$ for $\alpha < \beta$.
- Now take any disjoint open U_β, V_β with $(x_\beta, y_\beta) \in U_\beta \times V_\beta \subseteq H^2$.

[JS 2008] X^ω is d-separable for any compact X .

- $(\omega^*)^n$ is not d-separable for any $n \in \omega$.

[JS 2008] X^2 has a discrete subset of size $d(X)$ for any compact X .

- Wlog: any non empty open subset has weight $w(X)$ (not trivial).
- find $(x_\alpha, y_\alpha) \in U_\alpha \times V_\alpha \subseteq X^2$ so that $U_\alpha \cap V_\alpha = \emptyset$ for $\alpha < d(X)$,
- there is a open $H \neq \emptyset$ so that $K = \overline{H} \subseteq X \setminus \overline{\{x_\alpha, y_\alpha : \alpha < \beta\}}$,
- $\{U_\alpha, V_\alpha : \alpha < \beta\}$ generate a coarser topology on K than the original compact so cannot be Hausdorff.
- Let $x_\beta, y_\beta \in K$ witness this; then $(x_\beta, y_\beta) \notin U_\alpha \times V_\alpha$ for $\alpha < \beta$.
- Now take any disjoint open U_β, V_β with $(x_\beta, y_\beta) \in U_\beta \times V_\beta \subseteq H^2$.

Some open problems

[JS 2008] Is there a compact space X in ZFC so that X has no discrete subsets of size $d(X)$?

- a compact L -space (e.g. a Suslin-continuum) is a consistent example.

[Burke, Tkachuk 2013] $\text{CH} + \diamond(S_{\omega_1}^{\omega_2})$ implies that X^ω is not d -separable for some countably compact X .

- $d(X) = \aleph_2$ but every discrete subset of X^ω has size $\leq \aleph_1$.

[BT 2013] Is there a countably compact space X in ZFC so that X^ω is not d -separable?

Some open problems

[JS 2008] Is there a compact space X in ZFC so that X has no discrete subsets of size $d(X)$?

- a compact L -space (e.g. a Suslin-continuum) is a consistent example.

[Burke, Tkachuk 2013] $\text{CH} + \diamond(S_{\omega_1}^{\omega_2})$ implies that X^ω is not d -separable for some countably compact X .

- $d(X) = \aleph_2$ but every discrete subset of X^ω has size $\leq \aleph_1$.

[BT 2013] Is there a countably compact space X in ZFC so that X^ω is not d -separable?

Some open problems

[JS 2008] Is there a compact space X in ZFC so that X has no discrete subsets of size $d(X)$?

- a compact L -space (e.g. a Suslin-continuum) is a consistent example.

[Burke, Tkachuk 2013] $\text{CH} + \diamond(S_{\omega_1}^{\omega_2})$ implies that X^ω is not d -separable for some countably compact X .

- $d(X) = \aleph_2$ but every discrete subset of X^ω has size $\leq \aleph_1$.

[BT 2013] Is there a countably compact space X in ZFC so that X^ω is not d -separable?

Some open problems

[JS 2008] Is there a compact space X in ZFC so that X has no discrete subsets of size $d(X)$?

- a compact L -space (e.g. a Suslin-continuum) is a consistent example.

[Burke, Tkachuk 2013] $\text{CH} + \diamond(S_{\omega_1}^{\omega_2})$ implies that X^ω is not d -separable for some countably compact X .

- $d(X) = \aleph_2$ but every discrete subset of X^ω has size $\leq \aleph_1$.

[BT 2013] Is there a countably compact space X in ZFC so that X^ω is not d -separable?

Some open problems

[JS 2008] Is there a compact space X in ZFC so that X has no discrete subsets of size $d(X)$?

- a compact L -space (e.g. a Suslin-continuum) is a consistent example.

[Burke, Tkachuk 2013] $\text{CH} + \diamond(S_{\omega_1}^{\omega_2})$ implies that X^ω is not d -separable for some countably compact X .

- $d(X) = \aleph_2$ but every discrete subset of X^ω has size $\leq \aleph_1$.

[BT 2013] Is there a countably compact space X in ZFC so that X^ω is not d -separable?

Some open problems

[JS 2008] Is there a compact space X in ZFC so that X has no discrete subsets of size $d(X)$?

- a compact L -space (e.g. a Suslin-continuum) is a consistent example.

[Burke, Tkachuk 2013] $\text{CH} + \diamond(S_{\omega_1}^{\omega_2})$ implies that X^ω is not d -separable for some countably compact X .

- $d(X) = \aleph_2$ but every discrete subset of X^ω has size $\leq \aleph_1$.

[BT 2013] Is there a countably compact space X in ZFC so that X^ω is not d -separable?

[Kurepa 1936] X is e-separable iff there is a σ -closed discrete dense subset of X .

- Kurepa named these K'_0 -spaces.

If X is separable or metrizable then X is e-separable.

If X is (countably) compact and e-separable then X is separable.

- $D(2)^{\mathfrak{c}^+}$ is d-separable but not e-separable.

[Kurepa 1936] X is e-separable iff there is a σ -closed discrete dense subset of X .

- Kurepa named these K'_0 -spaces.

If X is separable or metrizable then X is e-separable.

If X is (countably) compact and e-separable then X is separable.

- $D(2)^{\mathfrak{c}^+}$ is d-separable but not e-separable.

[Kurepa 1936] X is e-separable iff there is a σ -closed discrete dense subset of X .

- Kurepa named these K'_0 -spaces.

If X is separable or metrizable then X is e-separable.

If X is (countably) compact and e-separable then X is separable.

- $D(2)^{\mathfrak{c}^+}$ is d-separable but not e-separable.

[Kurepa 1936] X is e-separable iff there is a σ -closed discrete dense subset of X .

- Kurepa named these K'_0 -spaces.

If X is separable or metrizable then X is e-separable.

If X is (countably) compact and e-separable then X is separable.

- $D(2)^{\mathfrak{c}^+}$ is d-separable but not e-separable.

[Kurepa 1936] X is e-separable iff there is a σ -closed discrete dense subset of X .

- Kurepa named these K'_0 -spaces.

If X is separable or metrizable then X is e-separable.

If X is (countably) compact and e-separable then X is separable.

- $D(2)^{\mathfrak{c}^+}$ is d-separable but not e-separable.

[Kurepa 1936] X is e-separable iff there is a σ -closed discrete dense subset of X .

- Kurepa named these K'_0 -spaces.

If X is separable or metrizable then X is e-separable.

If X is (countably) compact and e-separable then X is separable.

- $D(2)^{\mathfrak{c}^+}$ is d-separable but not e-separable.

Prior focus in research

[Faber 1974] Any e-separable GO-space is perfect (closed sets are G_δ).

- for perfect spaces: e-separable \leftrightarrow d-separable.

[Maurice 1970s] Is there, in ZFC, a perfect GO-space which is not e-separable?

[Reed 1970s] Is there, in ZFC, a perfect, first countable X which is not e-separable?

- [Qiao, Tall 2003] connecting to non-Archimedean spaces.
- [Benett, Lutzer 2006] Answer is “No” if density is \aleph_1 .

Prior focus in research

[Faber 1974] Any e-separable GO-space is perfect (closed sets are G_δ).

- for perfect spaces: e-separable \leftrightarrow d-separable.

[Maurice 1970s] Is there, in ZFC, a perfect GO-space which is not e-separable?

[Reed 1970s] Is there, in ZFC, a perfect, first countable X which is not e-separable?

- **[Qiao, Tall 2003]** connecting to non-Archimedean spaces.
- **[Benett, Lutzer 2006]** Answer is “No” if density is \aleph_1 .

[Faber 1974] Any e-separable GO-space is perfect (closed sets are G_δ).

- for perfect spaces: e-separable \leftrightarrow d-separable.

[Maurice 1970s] Is there, in ZFC, a perfect GO-space which is not e-separable?

[Reed 1970s] Is there, in ZFC, a perfect, first countable X which is not e-separable?

- **[Qiao, Tall 2003]** connecting to non-Archimedean spaces.
- **[Benett, Lutzer 2006]** Answer is “No” if density is \aleph_1 .

[Faber 1974] Any e-separable GO-space is perfect (closed sets are G_δ).

- for perfect spaces: e-separable \leftrightarrow d-separable.

[Maurice 1970s] Is there, in ZFC, a perfect GO-space which is not e-separable?

[Reed 1970s] Is there, in ZFC, a perfect, first countable X which is not e-separable?

- [Qiao, Tall 2003] connecting to non-Archimedean spaces.
- [Benett, Lutzer 2006] Answer is “No” if density is \aleph_1 .

[Faber 1974] Any e-separable GO-space is perfect (closed sets are G_δ).

- for perfect spaces: e-separable \leftrightarrow d-separable.

[Maurice 1970s] Is there, in ZFC, a perfect GO-space which is not e-separable?

[Reed 1970s] Is there, in ZFC, a perfect, first countable X which is not e-separable?

- **[Qiao, Tall 2003]** connecting to non-Archimedean spaces.
- **[Benett, Lutzer 2006]** Answer is “No” if density is \aleph_1 .

[Faber 1974] Any e-separable GO-space is perfect (closed sets are G_δ).

- for perfect spaces: e-separable \leftrightarrow d-separable.

[Maurice 1970s] Is there, in ZFC, a perfect GO-space which is not e-separable?

[Reed 1970s] Is there, in ZFC, a perfect, first countable X which is not e-separable?

- **[Qiao, Tall 2003]** connecting to non-Archimedean spaces.
- **[Benett, Lutzer 2006]** Answer is “No” if density is \aleph_1 .

[Faber 1974] Any e-separable GO-space is perfect (closed sets are G_δ).

- for perfect spaces: e-separable \leftrightarrow d-separable.

[Maurice 1970s] Is there, in ZFC, a perfect GO-space which is not e-separable?

[Reed 1970s] Is there, in ZFC, a perfect, first countable X which is not e-separable?

- **[Qiao, Tall 2003]** connecting to non-Archimedean spaces.
- **[Benett, Lutzer 2006]** Answer is “No” if density is \aleph_1 .

Powers and large closed discrete sets

- Products of d -separable are d -separable.
- X^κ is d -separable if it has a discrete subset of size $d(X)$.

[Alas] If X is e -separable then X^κ is e -separable for all $\kappa \leq \mathfrak{c}$.

X^κ is e -separable if there is a closed discrete subset of X^κ of size $d(X^\kappa)$.

- Suppose that $\{d_\xi\}_{\xi < \delta} \subseteq X'$ is dense in X' and $\{e_\xi\}_{\xi < \delta} \subseteq X^{\kappa \setminus I'}$ is closed discrete in $X^{\kappa \setminus I'}$.
- $\{d_\xi \cup e_\xi\}_{\xi < \delta} \subseteq X^\kappa$ is closed discrete in X^κ .
- Write κ as increasing union $\bigcup \{I_n : n \in \omega\}$ with $|I_n| = |\kappa \setminus I_n| = \kappa$ and repeat the above.

Powers and large closed discrete sets

- Products of d -separable are d -separable.
- X^κ is d -separable if it has a discrete subset of size $d(X)$.

[Alas] If X is e -separable then X^κ is e -separable for all $\kappa \leq \mathfrak{c}$.

X^κ is e -separable if there is a closed discrete subset of X^κ of size $d(X^\kappa)$.

- Suppose that $\{d_\xi\}_{\xi < \delta} \subseteq X'$ is dense in X' and $\{e_\xi\}_{\xi < \delta} \subseteq X^{\kappa \setminus I}$ is closed discrete in $X^{\kappa \setminus I}$.
- $\{d_\xi \cup e_\xi\}_{\xi < \delta} \subseteq X^\kappa$ is closed discrete in X^κ .
- Write κ as increasing union $\bigcup \{I_n : n \in \omega\}$ with $|I_n| = |\kappa \setminus I_n| = \kappa$ and repeat the above.

Powers and large closed discrete sets

- Products of d -separable are d -separable.
- X^κ is d -separable if it has a discrete subset of size $d(X)$.

[Alas] If X is e -separable then X^κ is e -separable for all $\kappa \leq \mathfrak{c}$.

X^κ is e -separable if there is a closed discrete subset of X^κ of size $d(X^\kappa)$.

- Suppose that $\{d_\xi\}_{\xi < \delta} \subseteq X'$ is dense in X' and $\{e_\xi\}_{\xi < \delta} \subseteq X^{\kappa \setminus I}$ is closed discrete in $X^{\kappa \setminus I}$.
- $\{d_\xi \cup e_\xi\}_{\xi < \delta} \subseteq X^\kappa$ is closed discrete in X^κ .
- Write κ as increasing union $\bigcup \{I_n : n \in \omega\}$ with $|I_n| = |\kappa \setminus I_n| = \kappa$ and repeat the above.

Powers and large closed discrete sets

- Products of d -separable are d -separable.
- X^κ is d -separable if it has a discrete subset of size $d(X)$.

[Alas] If X is e -separable then X^κ is e -separable for all $\kappa \leq \mathfrak{c}$.

X^κ is e -separable if there is a closed discrete subset of X^κ of size $d(X^\kappa)$.

- Suppose that $\{d_\xi\}_{\xi < \delta} \subseteq X'$ is dense in X' and $\{e_\xi\}_{\xi < \delta} \subseteq X^{\kappa \setminus I}$ is closed discrete in $X^{\kappa \setminus I}$.
- $\{d_\xi \cup e_\xi\}_{\xi < \delta} \subseteq X^\kappa$ is closed discrete in X^κ .
- Write κ as increasing union $\bigcup \{I_n : n \in \omega\}$ with $|I_n| = |\kappa \setminus I_n| = \kappa$ and repeat the above.

Powers and large closed discrete sets

- Products of d -separable are d -separable.
- X^κ is d -separable if it has a discrete subset of size $d(X)$.

[Alas] If X is e -separable then X^κ is e -separable for all $\kappa \leq \mathfrak{c}$.

X^κ is e -separable if there is a closed discrete subset of X^κ of size $d(X^\kappa)$.

- Suppose that $\{d_\xi\}_{\xi < \delta} \subseteq X'$ is dense in X' and $\{e_\xi\}_{\xi < \delta} \subseteq X^{\kappa \setminus I'}$ is closed discrete in $X^{\kappa \setminus I'}$.
- $\{d_\xi \cup e_\xi\}_{\xi < \delta} \subseteq X^\kappa$ is closed discrete in X^κ .
- Write κ as increasing union $\bigcup \{I_n : n \in \omega\}$ with $|I_n| = |\kappa \setminus I_n| = \kappa$ and repeat the above.

Powers and large closed discrete sets

- Products of d -separable are d -separable.
- X^κ is d -separable if it has a discrete subset of size $d(X)$.

[Alas] If X is e -separable then X^κ is e -separable for all $\kappa \leq \mathfrak{c}$.

X^κ is e -separable if there is a closed discrete subset of X^κ of size $d(X^\kappa)$.

- Suppose that $\{d_\xi\}_{\xi < \delta} \subseteq X'$ is dense in X' and $\{e_\xi\}_{\xi < \delta} \subseteq X^{\kappa \setminus I}$ is closed discrete in $X^{\kappa \setminus I}$.
- $\{d_\xi \cup e_\xi\}_{\xi < \delta} \subseteq X^\kappa$ is closed discrete in X^κ .
- Write κ as increasing union $\bigcup \{I_n : n \in \omega\}$ with $|I_n| = |\kappa \setminus I_n| = \kappa$ and repeat the above.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Mycielski 1964] $D(\omega)^\kappa$ contains a closed discrete set of size κ for every κ less than the 1st weakly inaccessible cardinal.

This helps if X has an infinite closed discrete subset i.e. not countably compact.

Suppose that X is not countably compact. Then $X^{d(X)}$ is e-separable if $d(X) < \text{the 1st weakly inaccessible cardinal}$.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Mycielski 1964] $D(\omega)^\kappa$ contains a closed discrete set of size κ for every κ less than the 1st weakly inaccessible cardinal.

This helps if X has an infinite closed discrete subset i.e. not countably compact.

Suppose that X is not countably compact. Then $X^{d(X)}$ is e-separable if $d(X) < \text{the 1st weakly inaccessible cardinal}$.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Mycielski 1964] $D(\omega)^\kappa$ contains a closed discrete set of size κ for every κ less than the 1st weakly inaccessible cardinal.

This helps if X has an infinite closed discrete subset i.e. not countably compact.

Suppose that X is not countably compact. Then $X^{d(X)}$ is e-separable if $d(X) < \text{the 1}^{st} \text{ weakly inaccessible cardinal}$.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Mycielski 1964] $D(\omega)^\kappa$ contains a closed discrete set of size κ for every κ less than the 1st weakly inaccessible cardinal.

This helps if X has an infinite closed discrete subset i.e. not countably compact.

Suppose that X is not countably compact. Then $X^{d(X)}$ is e-separable if $d(X) < \text{the 1st weakly inaccessible cardinal}$.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Lós 1959] $D(\omega)^{2^\kappa}$ contains a closed discrete set of size κ for every κ less than the 1^{st} measurable cardinal.

Suppose that X is not countably compact. Then $X^{2^{d(X)}}$ is e-separable if $d(X) < \text{the } 1^{\text{st}} \text{ measurable cardinal}$.

What happens at a measurable?

If $\kappa > \omega$ is measurable then $D(\omega)^\kappa$ has no closed discrete subsets of size κ so is not e-separable.

Note: $D(\omega)^\kappa$ has closed discrete sets of all sizes $< \kappa$.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Lós 1959] $D(\omega)^{2^\kappa}$ contains a closed discrete set of size κ for every κ less than the 1^{st} measurable cardinal.

Suppose that X is not countably compact. Then $X^{2^{d(X)}}$ is e-separable if $d(X) < \text{the } 1^{\text{st}} \text{ measurable cardinal}$.

What happens at a measurable?

If $\kappa > \omega$ is measurable then $D(\omega)^\kappa$ has no closed discrete subsets of size κ so is not e-separable.

Note: $D(\omega)^\kappa$ has closed discrete sets of all sizes $< \kappa$.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Lós 1959] $D(\omega)^{2^\kappa}$ contains a closed discrete set of size κ for every κ less than the 1^{st} measurable cardinal.

Suppose that X is not countably compact. Then $X^{2^{d(X)}}$ is e-separable if $d(X) < \text{the } 1^{\text{st}} \text{ measurable cardinal}$.

What happens at a measurable?

If $\kappa > \omega$ is measurable then $D(\omega)^\kappa$ has no closed discrete subsets of size κ so is not e-separable.

Note: $D(\omega)^\kappa$ has closed discrete sets of all sizes $< \kappa$.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Lós 1959] $D(\omega)^{2^\kappa}$ contains a closed discrete set of size κ for every κ less than the 1^{st} measurable cardinal.

Suppose that X is not countably compact. Then $X^{2^{d(X)}}$ is e-separable if $d(X) < \text{the } 1^{\text{st}} \text{ measurable cardinal}$.

What happens at a measurable?

If $\kappa > \omega$ is measurable then $D(\omega)^\kappa$ has no closed discrete subsets of size κ so is not e-separable.

Note: $D(\omega)^\kappa$ has closed discrete sets of all sizes $< \kappa$.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Lós 1959] $D(\omega)^{2^\kappa}$ contains a closed discrete set of size κ for every κ less than the 1^{st} measurable cardinal.

Suppose that X is not countably compact. Then $X^{2^{d(X)}}$ is e-separable if $d(X) < \text{the } 1^{\text{st}} \text{ measurable cardinal}$.

What happens at a measurable?

If $\kappa > \omega$ is measurable then $D(\omega)^\kappa$ has no closed discrete subsets of size κ so is not e-separable.

Note: $D(\omega)^\kappa$ has closed discrete sets of all sizes $< \kappa$.

Are there large closed discrete sets?

Recall: X^κ is e-separable if there is a closed discrete subset of size $d(X^\kappa)$.

[Lós 1959] $D(\omega)^{2^\kappa}$ contains a closed discrete set of size κ for every κ less than the 1^{st} measurable cardinal.

Suppose that X is not countably compact. Then $X^{2^{d(X)}}$ is e-separable if $d(X) < \text{the } 1^{\text{st}} \text{ measurable cardinal}$.

What happens at a measurable?

If $\kappa > \omega$ is measurable then $D(\omega)^\kappa$ has no closed discrete subsets of size κ so is not e-separable.

Note: $D(\omega)^\kappa$ has closed discrete sets of all sizes $< \kappa$.

Products - two reduction steps

Recall: $X^{\mathfrak{c}}$ is e -separable if X is e -separable.

Focus: products of $\kappa \leq \mathfrak{c}$ terms b.c. $D(2)^{\mathfrak{c}^+}$ is not e -separable.

Suppose that $\kappa \leq \mathfrak{c}$. Then the following are equivalent:

- every product of at most κ many e -separable spaces is e -separable;
- every product of at most κ many discrete spaces is e -separable.

\Rightarrow finite products preserve e -separability.

The product of at most κ many discrete spaces of cardinality at least κ is e -separable.

\Rightarrow a product of κ spaces is **non e -separable because of terms of size $< \kappa$.**

Products - two reduction steps

Recall: $X^{\mathfrak{c}}$ is e -separable if X is e -separable.

Focus: products of $\kappa \leq \mathfrak{c}$ terms b.c. $D(2)^{\mathfrak{c}^+}$ is not e -separable.

Suppose that $\kappa \leq \mathfrak{c}$. Then the following are equivalent:

- every product of at most κ many e -separable spaces is e -separable;
- every product of at most κ many discrete spaces is e -separable.

\Rightarrow finite products preserve e -separability.

The product of at most κ many discrete spaces of cardinality at least κ is e -separable.

\Rightarrow a product of κ spaces is **non e -separable because of terms of size $< \kappa$.**

Products - two reduction steps

Recall: $X^{\mathfrak{c}}$ is e -separable if X is e -separable.

Focus: products of $\kappa \leq \mathfrak{c}$ terms b.c. $D(2)^{\mathfrak{c}^+}$ is not e -separable.

Suppose that $\kappa \leq \mathfrak{c}$. Then the following are equivalent:

- every product of at most κ many e -separable spaces is e -separable;
- every product of at most κ many discrete spaces is e -separable.

\Rightarrow finite products preserve e -separability.

The product of at most κ many discrete spaces of cardinality at least κ is e -separable.

\Rightarrow a product of κ spaces is **non e -separable because of terms of size $< \kappa$.**

Products - two reduction steps

Recall: $X^{\mathfrak{c}}$ is e -separable if X is e -separable.

Focus: products of $\kappa \leq \mathfrak{c}$ terms b.c. $D(2)^{\mathfrak{c}^+}$ is not e -separable.

Suppose that $\kappa \leq \mathfrak{c}$. Then the following are equivalent:

- every product of at most κ many e -separable spaces is e -separable;
- every product of at most κ many discrete spaces is e -separable.

\Rightarrow finite products preserve e -separability.

The product of at most κ many discrete spaces of cardinality at least κ is e -separable.

\Rightarrow a product of κ spaces is **non e -separable because of terms of size $< \kappa$.**

Products - two reduction steps

Recall: $X^{\mathfrak{c}}$ is e -separable if X is e -separable.

Focus: products of $\kappa \leq \mathfrak{c}$ terms b.c. $D(2)^{\mathfrak{c}^+}$ is not e -separable.

Suppose that $\kappa \leq \mathfrak{c}$. Then the following are equivalent:

- every product of at most κ many e -separable spaces is e -separable;
- every product of at most κ many discrete spaces is e -separable.

\Rightarrow finite products preserve e -separability.

The product of at most κ many discrete spaces of cardinality at least κ is e -separable.

\Rightarrow a product of κ spaces is **non e -separable because of terms of size $< \kappa$.**

Products - two reduction steps

Recall: $X^{\mathfrak{c}}$ is e -separable if X is e -separable.

Focus: products of $\kappa \leq \mathfrak{c}$ terms b.c. $D(2)^{\mathfrak{c}^+}$ is not e -separable.

Suppose that $\kappa \leq \mathfrak{c}$. Then the following are equivalent:

- every product of at most κ many e -separable spaces is e -separable;
- every product of at most κ many discrete spaces is e -separable.

\Rightarrow finite products preserve e -separability.

The product of at most κ many discrete spaces of cardinality at least κ is e -separable.

\Rightarrow a product of κ spaces is **non e -separable because of terms of size $< \kappa$.**

Closed discrete sets in products

[Mrowka 1970] Define $\kappa \notin \mathcal{M}^*$ iff any product of κ many discrete spaces $X = \prod \{X_\alpha : \alpha < \kappa\}$ with each of size $< \kappa$ has no closed discrete set of size κ .

[M 1970, Jech] If $\kappa \notin \mathcal{M}^*$ then κ is weakly inaccessible.

Suppose that $\kappa \leq \mathfrak{c}$ is minimal so that there is a family of κ e-separable spaces with non e-separable product. Then $\kappa \notin \mathcal{M}^*$.

The preservation theorem:

Suppose there are no weakly inaccessible cardinals $\leq \mathfrak{c}$. Then the product of at most \mathfrak{c} many e-separable spaces is e-separable.

Closed discrete sets in products

[Mrowka 1970] Define $\kappa \notin \mathcal{M}^*$ iff any product of κ many discrete spaces $X = \prod \{X_\alpha : \alpha < \kappa\}$ with each of size $< \kappa$ has no closed discrete set of size κ .

[M 1970, Jech] If $\kappa \notin \mathcal{M}^*$ then κ is weakly inaccessible.

Suppose that $\kappa \leq \mathfrak{c}$ is minimal so that there is a family of κ e-separable spaces with non e-separable product. Then $\kappa \notin \mathcal{M}^*$.

The preservation theorem:

Suppose there are no weakly inaccessible cardinals $\leq \mathfrak{c}$. Then the product of at most \mathfrak{c} many e-separable spaces is e-separable.

Closed discrete sets in products

[Mrowka 1970] Define $\kappa \notin \mathcal{M}^*$ iff any product of κ many discrete spaces $X = \prod \{X_\alpha : \alpha < \kappa\}$ with each of size $< \kappa$ has no closed discrete set of size κ .

[M 1970, Jech] If $\kappa \notin \mathcal{M}^*$ then κ is weakly inaccessible.

Suppose that $\kappa \leq \mathfrak{c}$ is minimal so that there is a family of κ e-separable spaces with non e-separable product. Then $\kappa \notin \mathcal{M}^*$.

The preservation theorem:

Suppose there are no weakly inaccessible cardinals $\leq \mathfrak{c}$. Then the product of at most \mathfrak{c} many e-separable spaces is e-separable.

Closed discrete sets in products

[Mrowka 1970] Define $\kappa \notin \mathcal{M}^*$ iff any product of κ many discrete spaces $X = \prod \{X_\alpha : \alpha < \kappa\}$ with each of size $< \kappa$ has no closed discrete set of size κ .

[M 1970, Jech] If $\kappa \notin \mathcal{M}^*$ then κ is weakly inaccessible.

Suppose that $\kappa \leq \mathfrak{c}$ is minimal so that there is a family of κ e-separable spaces with non e-separable product. Then $\kappa \notin \mathcal{M}^*$.

The preservation theorem:

Suppose there are no weakly inaccessible cardinals $\leq \mathfrak{c}$. Then the product of at most \mathfrak{c} many e-separable spaces is e-separable.

Closed discrete sets in products

[Mrowka 1970] Define $\kappa \notin \mathcal{M}^*$ iff any product of κ many discrete spaces $X = \prod \{X_\alpha : \alpha < \kappa\}$ with each of size $< \kappa$ has no closed discrete set of size κ .

[M 1970, Jech] If $\kappa \notin \mathcal{M}^*$ then κ is weakly inaccessible.

Suppose that $\kappa \leq \mathfrak{c}$ is minimal so that there is a family of κ e-separable spaces with non e-separable product. Then $\kappa \notin \mathcal{M}^*$.

The preservation theorem:

Suppose there are no weakly inaccessible cardinals $\leq \mathfrak{c}$. Then the product of at most \mathfrak{c} many e-separable spaces is e-separable.

\mathcal{M}^* and weak compactness

$\mathcal{L}_{\kappa,\omega}$ is the infinitary language which allows conjunctions and disjunctions of $< \kappa$ formulas and universal or existential quantification over finitely many variables.

$\mathcal{L}_{\kappa,\omega}$ is weakly compact iff every set of at most κ sentences Σ from $\mathcal{L}_{\kappa,\omega}$ has a model provided that every $S \in [\Sigma]^{<\kappa}$ has a model.

[M 1970] $\mathcal{L}_{\kappa,\omega}$ is weakly compact iff $\kappa \notin \mathcal{M}^*$.

\mathcal{M}^* and weak compactness

$\mathcal{L}_{\kappa,\omega}$ is the infinitary language which allows conjunctions and disjunctions of $< \kappa$ formulas and universal or existential quantification over finitely many variables.

$\mathcal{L}_{\kappa,\omega}$ is weakly compact iff every set of at most κ sentences Σ from $\mathcal{L}_{\kappa,\omega}$ has a model provided that every $S \in [\Sigma]^{<\kappa}$ has a model.

[M 1970] $\mathcal{L}_{\kappa,\omega}$ is weakly compact iff $\kappa \notin \mathcal{M}^*$.

\mathcal{M}^* and weak compactness

$\mathcal{L}_{\kappa,\omega}$ is the infinitary language which allows conjunctions and disjunctions of $< \kappa$ formulas and universal or existential quantification over finitely many variables.

$\mathcal{L}_{\kappa,\omega}$ is weakly compact iff every set of at most κ sentences Σ from $\mathcal{L}_{\kappa,\omega}$ has a model provided that every $S \in [\Sigma]^{<\kappa}$ has a model.

[M 1970] $\mathcal{L}_{\kappa,\omega}$ is weakly compact iff $\kappa \notin \mathcal{M}^*$.

\mathcal{M}^* and weak compactness

$\mathcal{L}_{\kappa,\omega}$ is the infinitary language which allows conjunctions and disjunctions of $< \kappa$ formulas and universal or existential quantification over finitely many variables.

$\mathcal{L}_{\kappa,\omega}$ is *weakly compact* iff every set of at most κ sentences Σ from $\mathcal{L}_{\kappa,\omega}$ has a model provided that every $S \in [\Sigma]^{<\kappa}$ has a model.

[M 1970] $\mathcal{L}_{\kappa,\omega}$ is weakly compact iff $\kappa \notin \mathcal{M}^*$.

Weak compactness below \mathfrak{c}

[Erdős, Tarski 1961] κ is a weakly compact cardinal iff $\kappa \rightarrow (\kappa)_2^2$.

κ is a weakly compact cardinal iff it is strongly inaccessible and $\mathcal{L}_{\kappa,\omega}$ is weakly compact.

[Cudnovskii, Boos] If $V \models \kappa$ is a weakly compact cardinal then $V^{\mathbb{C}_{\kappa^+}} \models \kappa < \mathfrak{c} = \kappa^+$ and $\mathcal{L}_{\kappa,\omega}$ is weakly compact.

Weak compactness below \mathfrak{c}

[Erdős, Tarski 1961] κ is a weakly compact cardinal iff $\kappa \rightarrow (\kappa)_2^2$.

κ is a weakly compact cardinal iff it is strongly inaccessible and $\mathcal{L}_{\kappa,\omega}$ is weakly compact.

[Cudnovskii, Boos] If $V \models \kappa$ is a weakly compact cardinal then $V^{\mathbb{C}_{\kappa^+}} \models \kappa < \mathfrak{c} = \kappa^+$ and $\mathcal{L}_{\kappa,\omega}$ is weakly compact.

Weak compactness below \mathfrak{c}

[Erdős, Tarski 1961] κ is a weakly compact cardinal iff $\kappa \rightarrow (\kappa)_2^2$.

κ is a weakly compact cardinal iff it is strongly inaccessible and $\mathcal{L}_{\kappa,\omega}$ is weakly compact.

[Cudnovskii, Boos] If $V \models \kappa$ is a weakly compact cardinal then $V^{\mathbb{C}_{\kappa^+}} \models \kappa < \mathfrak{c} = \kappa^+$ and $\mathcal{L}_{\kappa,\omega}$ is weakly compact.

Weak compactness below \mathfrak{c}

[Erdős, Tarski 1961] κ is a weakly compact cardinal iff $\kappa \rightarrow (\kappa)_2^2$.

κ is a weakly compact cardinal iff it is strongly inaccessible and $\mathcal{L}_{\kappa,\omega}$ is weakly compact.

[Cudnovskii, Boos] If $V \models \kappa$ is a weakly compact cardinal then $V^{\mathbb{C}_{\kappa^+}} \models \kappa < \mathfrak{c} = \kappa^+$ and $\mathcal{L}_{\kappa,\omega}$ is weakly compact.

Non e-separable products of size $\leq \mathfrak{c}$

Consistently, there are $< \mathfrak{c}$ -many discrete spaces (each of size $< \mathfrak{c}$) with non e-separable product.

- Use a weakly compact cardinal to find a model with $\kappa < \mathfrak{c}$ and $\mathcal{L}_{\kappa, \omega}$ weakly compact i.e. $\kappa \notin \mathcal{M}^*$.
- $X = \prod \{D(\alpha) : \alpha < \kappa\}$ has no closed discrete sets of size κ .
- κ is regular (and limit) so σ -closed discrete sets have size $< \kappa$.
- $d(X) = \kappa$ so X is not e-separable.

Non e-separable products of size $\leq \mathfrak{c}$

Consistently, there are $< \mathfrak{c}$ -many discrete spaces (each of size $< \mathfrak{c}$) with non e-separable product.

- Use a weakly compact cardinal to find a model with $\kappa < \mathfrak{c}$ and $\mathcal{L}_{\kappa, \omega}$ weakly compact i.e. $\kappa \notin \mathcal{M}^*$.
- $X = \prod \{D(\alpha) : \alpha < \kappa\}$ has no closed discrete sets of size κ .
- κ is regular (and limit) so σ -closed discrete sets have size $< \kappa$.
- $d(X) = \kappa$ so X is not e-separable.

Non e-separable products of size $\leq \mathfrak{c}$

Consistently, there are $< \mathfrak{c}$ -many discrete spaces (each of size $< \mathfrak{c}$) with non e-separable product.

- Use a weakly compact cardinal to find a model with $\kappa < \mathfrak{c}$ and $\mathcal{L}_{\kappa, \omega}$ weakly compact i.e. $\kappa \notin \mathcal{M}^*$.
- $X = \prod \{D(\alpha) : \alpha < \kappa\}$ has no closed discrete sets of size κ .
- κ is regular (and limit) so σ -closed discrete sets have size $< \kappa$.
- $d(X) = \kappa$ so X is not e-separable.

Non e-separable products of size $\leq \mathfrak{c}$

Consistently, there are $< \mathfrak{c}$ -many discrete spaces (each of size $< \mathfrak{c}$) with non e-separable product.

- Use a weakly compact cardinal to find a model with $\kappa < \mathfrak{c}$ and $\mathcal{L}_{\kappa, \omega}$ weakly compact i.e. $\kappa \notin \mathcal{M}^*$.
- $X = \prod \{D(\alpha) : \alpha < \kappa\}$ has no closed discrete sets of size κ .
- κ is regular (and limit) so σ -closed discrete sets have size $< \kappa$.
- $d(X) = \kappa$ so X is not e-separable.

Non e-separable products of size $\leq \mathfrak{c}$

Consistently, there are $< \mathfrak{c}$ -many discrete spaces (each of size $< \mathfrak{c}$) with non e-separable product.

- Use a weakly compact cardinal to find a model with $\kappa < \mathfrak{c}$ and $\mathcal{L}_{\kappa, \omega}$ weakly compact i.e. $\kappa \notin \mathcal{M}^*$.
- $X = \prod \{D(\alpha) : \alpha < \kappa\}$ has no closed discrete sets of size κ .
- κ is regular (and limit) so σ -closed discrete sets have size $< \kappa$.
- $d(X) = \kappa$ so X is not e-separable.

Non e-separable products of size $\leq \mathfrak{c}$

Consistently, there are $< \mathfrak{c}$ -many discrete spaces (each of size $< \mathfrak{c}$) with non e-separable product.

- Use a weakly compact cardinal to find a model with $\kappa < \mathfrak{c}$ and $\mathcal{L}_{\kappa, \omega}$ weakly compact i.e. $\kappa \notin \mathcal{M}^*$.
- $X = \prod \{D(\alpha) : \alpha < \kappa\}$ has no closed discrete sets of size κ .
- κ is regular (and limit) so σ -closed discrete sets have size $< \kappa$.
- $d(X) = \kappa$ so X is not e-separable.

Some cardinal functions

Recall some classical cardinal functions:

- **density**: $d(X) = \min\{|D| : D \subseteq X \text{ is dense in } X\}$;
- **spread**: $s(X) = \sup\{|S| : S \subseteq X \text{ is discrete in } X\}$.

If X is d-separable:

$$d_s(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-discrete in } X\}$$

- **extent**: $e(X) = \sup\{|E| : E \subseteq X \text{ is closed discrete in } X\}$.

If X is e-separable:

$$d_e(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-closed discrete in } X\}$$

Some cardinal functions

Recall some classical cardinal functions:

- **density**: $d(X) = \min\{|D| : D \subseteq X \text{ is dense in } X\}$;
- **spread**: $s(X) = \sup\{|S| : S \subseteq X \text{ is discrete in } X\}$.

If X is d-separable:

$$d_s(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-discrete in } X\}$$

- **extent**: $e(X) = \sup\{|E| : E \subseteq X \text{ is closed discrete in } X\}$.

If X is e-separable:

$$d_e(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-closed discrete in } X\}$$

Some cardinal functions

Recall some classical cardinal functions:

- **density**: $d(X) = \min\{|D| : D \subseteq X \text{ is dense in } X\}$;
- **spread**: $s(X) = \sup\{|S| : S \subseteq X \text{ is discrete in } X\}$.

If X is d-separable:

$$d_s(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-discrete in } X\}$$

- **extent**: $e(X) = \sup\{|E| : E \subseteq X \text{ is closed discrete in } X\}$.

If X is e-separable:

$$d_e(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-closed discrete in } X\}$$

Some cardinal functions

Recall some classical cardinal functions:

- **density**: $d(X) = \min\{|D| : D \subseteq X \text{ is dense in } X\}$;
- **spread**: $s(X) = \sup\{|S| : S \subseteq X \text{ is discrete in } X\}$.

If X is d-separable:

$$d_s(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-discrete in } X\}$$

- **extent**: $e(X) = \sup\{|E| : E \subseteq X \text{ is closed discrete in } X\}$.

If X is e-separable:

$$d_e(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-closed discrete in } X\}$$

Some cardinal functions

Recall some classical cardinal functions:

- **density**: $d(X) = \min\{|D| : D \subseteq X \text{ is dense in } X\}$;
- **spread**: $s(X) = \sup\{|S| : S \subseteq X \text{ is discrete in } X\}$.

If X is d-separable:

$$d_s(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-discrete in } X\}$$

- **extent**: $e(X) = \sup\{|E| : E \subseteq X \text{ is closed discrete in } X\}$.

If X is e-separable:

$$d_e(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-closed discrete in } X\}$$

Some cardinal functions

Recall some classical cardinal functions:

- **density**: $d(X) = \min\{|D| : D \subseteq X \text{ is dense in } X\}$;
- **spread**: $s(X) = \sup\{|S| : S \subseteq X \text{ is discrete in } X\}$.

If X is d-separable:

$$d_s(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-discrete in } X\}$$

- **extent**: $e(X) = \sup\{|E| : E \subseteq X \text{ is closed discrete in } X\}$.

If X is e-separable:

$$d_e(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-closed discrete in } X\}$$

Some cardinal functions

Recall some classical cardinal functions:

- **density**: $d(X) = \min\{|D| : D \subseteq X \text{ is dense in } X\}$;
- **spread**: $s(X) = \sup\{|S| : S \subseteq X \text{ is discrete in } X\}$.

If X is d-separable:

$$d_s(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-discrete in } X\}$$

- **extent**: $e(X) = \sup\{|E| : E \subseteq X \text{ is closed discrete in } X\}$.

If X is e-separable:

$$d_e(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-closed discrete in } X\}$$

Some cardinal functions

Recall some classical cardinal functions:

- **density**: $d(X) = \min\{|D| : D \subseteq X \text{ is dense in } X\}$;
- **spread**: $s(X) = \sup\{|S| : S \subseteq X \text{ is discrete in } X\}$.

If X is d-separable:

$$d_s(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-discrete in } X\}$$

- **extent**: $e(X) = \sup\{|E| : E \subseteq X \text{ is closed discrete in } X\}$.

If X is e-separable:

$$d_e(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-closed discrete in } X\}$$

Some cardinal functions

Recall some classical cardinal functions:

- **density**: $d(X) = \min\{|D| : D \subseteq X \text{ is dense in } X\}$;
- **spread**: $s(X) = \sup\{|S| : S \subseteq X \text{ is discrete in } X\}$.

If X is d-separable:

$$d_s(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-discrete in } X\}$$

- **extent**: $e(X) = \sup\{|E| : E \subseteq X \text{ is closed discrete in } X\}$.

If X is e-separable:

$$d_e(X) = \min\{|D| : D \subseteq X \text{ is dense and } \sigma\text{-closed discrete in } X\}$$

Inequalities

- $d(X) \leq d_s(X) \leq s(X)$ for any d -separable X ;
- $d(X) \leq d_e(X) \leq e(X)$ for any e -separable X .

There is a 0-dimensional e -separable space X such that

$$\mathfrak{c} = d(X) < d_e(X) = e(X) = w(X) = 2^{\mathfrak{c}}.$$

The plan is that $X = Y \cup E \subseteq 2^{2^{\mathfrak{c}}}$, both Y, E are dense and

- $|Y| = \mathfrak{c}$ and σ -closed discrete sets are nowhere dense,
 - let $|Y_0| = \mathfrak{c}$ dense and find $Y_0 \subseteq Y$ countably compact, $|Y| = \mathfrak{c}$.
- E is σ -closed discrete with size and density $2^{\mathfrak{c}}$,
 - use the trick for glueing dense and closed discrete sets with $\sigma(2^{2^{\mathfrak{c}}})$.

- $d(X) \leq d_s(X) \leq s(X)$ for any d -separable X ;
- $d(X) \leq d_e(X) \leq e(X)$ for any e -separable X .

There is a 0-dimensional e -separable space X such that

$$\mathfrak{c} = d(X) < d_e(X) = e(X) = w(X) = 2^{\mathfrak{c}}.$$

The plan is that $X = Y \cup E \subseteq 2^{2^{\mathfrak{c}}}$, both Y, E are dense and

- $|Y| = \mathfrak{c}$ and σ -closed discrete sets are nowhere dense,
 - let $|Y_0| = \mathfrak{c}$ dense and find $Y_0 \subseteq Y$ countably compact, $|Y| = \mathfrak{c}$.
- E is σ -closed discrete with size and density $2^{\mathfrak{c}}$,
 - use the trick for glueing dense and closed discrete sets with $\sigma(2^{2^{\mathfrak{c}}})$.

- $d(X) \leq d_s(X) \leq s(X)$ for any d -separable X ;
- $d(X) \leq d_e(X) \leq e(X)$ for any e -separable X .

There is a 0-dimensional e -separable space X such that

$$\mathfrak{c} = d(X) < d_e(X) = e(X) = w(X) = 2^{\mathfrak{c}}.$$

The plan is that $X = Y \cup E \subseteq 2^{2^{\mathfrak{c}}}$, both Y, E are dense and

- $|Y| = \mathfrak{c}$ and σ -closed discrete sets are nowhere dense,
 - let $|Y_0| = \mathfrak{c}$ dense and find $Y_0 \subseteq Y$ countably compact, $|Y| = \mathfrak{c}$.
- E is σ -closed discrete with size and density $2^{\mathfrak{c}}$,
 - use the trick for glueing dense and closed discrete sets with $\sigma(2^{2^{\mathfrak{c}}})$.

- $d(X) \leq d_s(X) \leq s(X)$ for any d -separable X ;
- $d(X) \leq d_e(X) \leq e(X)$ for any e -separable X .

There is a 0-dimensional e -separable space X such that

$$\mathfrak{c} = d(X) < d_e(X) = e(X) = w(X) = 2^{\mathfrak{c}}.$$

The plan is that $X = Y \cup E \subseteq 2^{2^{\mathfrak{c}}}$, both Y, E are dense and

- $|Y| = \mathfrak{c}$ and σ -closed discrete sets are nowhere dense,
 - let $|Y_0| = \mathfrak{c}$ dense and find $Y_0 \subseteq Y$ countably compact, $|Y| = \mathfrak{c}$.
- E is σ -closed discrete with size and density $2^{\mathfrak{c}}$,
 - use the trick for glueing dense and closed discrete sets with $\sigma(2^{2^{\mathfrak{c}}})$.

- $d(X) \leq d_s(X) \leq s(X)$ for any d -separable X ;
- $d(X) \leq d_e(X) \leq e(X)$ for any e -separable X .

There is a 0-dimensional e -separable space X such that

$$\mathfrak{c} = d(X) < d_e(X) = e(X) = w(X) = 2^{\mathfrak{c}}.$$

The plan is that $X = Y \cup E \subseteq 2^{2^{\mathfrak{c}}}$, both Y, E are dense and

- $|Y| = \mathfrak{c}$ and σ -closed discrete sets are nowhere dense,
 - let $|Y_0| = \mathfrak{c}$ dense and find $Y_0 \subseteq Y$ countably compact, $|Y| = \mathfrak{c}$.
- E is σ -closed discrete with size and density $2^{\mathfrak{c}}$,
 - use the trick for glueing dense and closed discrete sets with $\sigma(2^{2^{\mathfrak{c}}})$.

- $d(X) \leq d_s(X) \leq s(X)$ for any d -separable X ;
- $d(X) \leq d_e(X) \leq e(X)$ for any e -separable X .

There is a 0-dimensional e -separable space X such that

$$\mathfrak{c} = d(X) < d_e(X) = e(X) = w(X) = 2^{\mathfrak{c}}.$$

The plan is that $X = Y \cup E \subseteq 2^{2^{\mathfrak{c}}}$, both Y, E are dense and

- $|Y| = \mathfrak{c}$ and σ -closed discrete sets are nowhere dense,
 - let $|Y_0| = \mathfrak{c}$ dense and find $Y_0 \subseteq Y$ countably compact, $|Y| = \mathfrak{c}$.
- E is σ -closed discrete with size and density $2^{\mathfrak{c}}$,
 - use the trick for glueing dense and closed discrete sets with $\sigma(2^{2^{\mathfrak{c}}})$.

- $d(X) \leq d_s(X) \leq s(X)$ for any d -separable X ;
- $d(X) \leq d_e(X) \leq e(X)$ for any e -separable X .

There is a 0-dimensional e -separable space X such that

$$\mathfrak{c} = d(X) < d_e(X) = e(X) = w(X) = 2^{\mathfrak{c}}.$$

The plan is that $X = Y \cup E \subseteq 2^{2^{\mathfrak{c}}}$, both Y, E are dense and

- $|Y| = \mathfrak{c}$ and σ -closed discrete sets are nowhere dense,
 - let $|Y_0| = \mathfrak{c}$ dense and find $Y_0 \subseteq Y$ countably compact, $|Y| = \mathfrak{c}$.
- E is σ -closed discrete with size and density $2^{\mathfrak{c}}$,
 - use the trick for glueing dense and closed discrete sets with $\sigma(2^{2^{\mathfrak{c}}})$.

- $d(X) \leq d_s(X) \leq s(X)$ for any d -separable X ;
- $d(X) \leq d_e(X) \leq e(X)$ for any e -separable X .

There is a 0-dimensional e -separable space X such that

$$\mathfrak{c} = d(X) < d_e(X) = e(X) = w(X) = 2^{\mathfrak{c}}.$$

The plan is that $X = Y \cup E \subseteq 2^{2^{\mathfrak{c}}}$, both Y, E are dense and

- $|Y| = \mathfrak{c}$ and σ -closed discrete sets are nowhere dense,
 - let $|Y_0| = \mathfrak{c}$ dense and find $Y_0 \subseteq Y$ countably compact, $|Y| = \mathfrak{c}$.
- E is σ -closed discrete with size and density $2^{\mathfrak{c}}$,
 - use the trick for glueing dense and closed discrete sets with $\sigma(2^{2^{\mathfrak{c}}})$.

- $d(X) \leq d_s(X) \leq s(X)$ for any d -separable X ;
- $d(X) \leq d_e(X) \leq e(X)$ for any e -separable X .

There is a 0-dimensional e -separable space X such that

$$\mathfrak{c} = d(X) < d_e(X) = e(X) = w(X) = 2^{\mathfrak{c}}.$$

The plan is that $X = Y \cup E \subseteq 2^{2^{\mathfrak{c}}}$, both Y, E are dense and

- $|Y| = \mathfrak{c}$ and σ -closed discrete sets are nowhere dense,
 - let $|Y_0| = \mathfrak{c}$ dense and find $Y_0 \subseteq Y$ countably compact, $|Y| = \mathfrak{c}$.
- E is σ -closed discrete with size and density $2^{\mathfrak{c}}$,
 - use the trick for glueing dense and closed discrete sets with $\sigma(2^{2^{\mathfrak{c}}})$.

How to get $d(X) < d_s(X)$ for a d-separable X ?

- [Moore 2006] there is a dense $Y \subseteq \omega^{\omega_1}$ such that any σ -discrete is nowhere dense.
- if $\omega_2 \leq \mathfrak{c}$ then there is a countable dense $D \subseteq 2^{\omega_2}$,
- so $Y \hookrightarrow D^{\omega_1} \subseteq (2^{\omega_2})^{\omega_1} \simeq 2^{\omega_2}$ densely.
- $X = Y \cup \sigma(2^{\omega_2})$ is d-separable but there are no σ -discrete dense sets of size \aleph_1 .

Is there, in ZFC, a dense $Y \subseteq 2^{\omega_2}$ of size \aleph_1 such that any σ -discrete subset is nowhere dense?

How to get $d(X) < d_s(X)$ for a d-separable X ?

- [Moore 2006] there is a dense $Y \subseteq \omega^{\omega_1}$ such that any σ -discrete is nowhere dense.
- if $\omega_2 \leq \mathfrak{c}$ then there is a countable dense $D \subseteq 2^{\omega_2}$,
- so $Y \hookrightarrow D^{\omega_1} \subseteq (2^{\omega_2})^{\omega_1} \simeq 2^{\omega_2}$ densely.
- $X = Y \cup \sigma(2^{\omega_2})$ is d-separable but there are no σ -discrete dense sets of size \aleph_1 .

Is there, in ZFC, a dense $Y \subseteq 2^{\omega_2}$ of size \aleph_1 such that any σ -discrete subset is nowhere dense?

How to get $d(X) < d_s(X)$ for a d-separable X ?

- **[Moore 2006]** there is a dense $Y \subseteq \omega^{\omega_1}$ such that any σ -discrete is nowhere dense.
- if $\omega_2 \leq \mathfrak{c}$ then there is a countable dense $D \subseteq 2^{\omega_2}$,
- so $Y \hookrightarrow D^{\omega_1} \subseteq (2^{\omega_2})^{\omega_1} \simeq 2^{\omega_2}$ densely.
- $X = Y \cup \sigma(2^{\omega_2})$ is d-separable but there are no σ -discrete dense sets of size \aleph_1 .

Is there, in ZFC, a dense $Y \subseteq 2^{\omega_2}$ of size \aleph_1 such that any σ -discrete subset is nowhere dense?

How to get $d(X) < d_s(X)$ for a d-separable X ?

- [Moore 2006] there is a dense $Y \subseteq \omega^{\omega_1}$ such that any σ -discrete is nowhere dense.
- if $\omega_2 \leq \mathfrak{c}$ then there is a countable dense $D \subseteq 2^{\omega_2}$,
- so $Y \hookrightarrow D^{\omega_1} \subseteq (2^{\omega_2})^{\omega_1} \simeq 2^{\omega_2}$ densely.
- $X = Y \cup \sigma(2^{\omega_2})$ is d-separable but there are no σ -discrete dense sets of size \aleph_1 .

Is there, in ZFC, a dense $Y \subseteq 2^{\omega_2}$ of size \aleph_1 such that any σ -discrete subset is nowhere dense?

How to get $d(X) < d_s(X)$ for a d-separable X ?

- [Moore 2006] there is a dense $Y \subseteq \omega^{\omega_1}$ such that any σ -discrete is nowhere dense.
- if $\omega_2 \leq \mathfrak{c}$ then there is a countable dense $D \subseteq 2^{\omega_2}$,
- so $Y \hookrightarrow D^{\omega_1} \subseteq (2^{\omega_2})^{\omega_1} \simeq 2^{\omega_2}$ densely.
- $X = Y \cup \sigma(2^{\omega_2})$ is d-separable but there are no σ -discrete dense sets of size \aleph_1 .

Is there, in ZFC, a dense $Y \subseteq 2^{\omega_2}$ of size \aleph_1 such that any σ -discrete subset is nowhere dense?

How to get $d(X) < d_s(X)$ for a d-separable X ?

- [Moore 2006] there is a dense $Y \subseteq \omega^{\omega_1}$ such that any σ -discrete is nowhere dense.
- if $\omega_2 \leq \mathfrak{c}$ then there is a countable dense $D \subseteq 2^{\omega_2}$,
- so $Y \hookrightarrow D^{\omega_1} \subseteq (2^{\omega_2})^{\omega_1} \simeq 2^{\omega_2}$ densely.
- $X = Y \cup \sigma(2^{\omega_2})$ is d-separable but there are no σ -discrete dense sets of size \aleph_1 .

Is there, in ZFC, a dense $Y \subseteq 2^{\omega_2}$ of size \aleph_1 such that any σ -discrete subset is nowhere dense?

How to get $d(X) < d_s(X)$ for a d-separable X ?

- [Moore 2006] there is a dense $Y \subseteq \omega^{\omega_1}$ such that any σ -discrete is nowhere dense.
- if $\omega_2 \leq \mathfrak{c}$ then there is a countable dense $D \subseteq 2^{\omega_2}$,
- so $Y \hookrightarrow D^{\omega_1} \subseteq (2^{\omega_2})^{\omega_1} \simeq 2^{\omega_2}$ densely.
- $X = Y \cup \sigma(2^{\omega_2})$ is d-separable but there are no σ -discrete dense sets of size \aleph_1 .

Is there, in ZFC, a dense $Y \subseteq 2^{\omega_2}$ of size \aleph_1 such that any σ -discrete subset is nowhere dense?

Last questions

Does $d(X) = d_s(X)$ for compact, d-separable X ?

[Moore 2008] There is an L -space X such that X^2 is d-separable.

[Peng 2015] There is an L -space X such that X^2 is e-separable.

Note: $s(X) = e(X) = \aleph_0 < d(X) = \aleph_1$ so X is not d-separable.

Is there a non-separable, countably compact X so that X^2 is e-separable?

Last questions

Does $d(X) = d_s(X)$ for compact, d-separable X ?

[Moore 2008] There is an L -space X such that X^2 is d-separable.

[Peng 2015] There is an L -space X such that X^2 is e-separable.

Note: $s(X) = e(X) = \aleph_0 < d(X) = \aleph_1$ so X is not d-separable.

Is there a non-separable, countably compact X so that X^2 is e-separable?

Last questions

Does $d(X) = d_s(X)$ for compact, d-separable X ?

[Moore 2008] There is an L -space X such that X^2 is d-separable.

[Peng 2015] There is an L -space X such that X^2 is e-separable.

Note: $s(X) = e(X) = \aleph_0 < d(X) = \aleph_1$ so X is not d-separable.

Is there a non-separable, countably compact X so that X^2 is e-separable?

Last questions

Does $d(X) = d_s(X)$ for compact, d-separable X ?

[Moore 2008] There is an L -space X such that X^2 is d-separable.

[Peng 2015] There is an L -space X such that X^2 is e-separable.

Note: $s(X) = e(X) = \aleph_0 < d(X) = \aleph_1$ so X is not d-separable.

Is there a non-separable, countably compact X so that X^2 is e-separable?

Last questions

Does $d(X) = d_s(X)$ for compact, d-separable X ?

[Moore 2008] There is an L -space X such that X^2 is d-separable.

[Peng 2015] There is an L -space X such that X^2 is e-separable.

Note: $s(X) = e(X) = \aleph_0 < d(X) = \aleph_1$ so X is not d-separable.

Is there a non-separable, countably compact X so that X^2 is e-separable?

Last questions

Does $d(X) = d_s(X)$ for compact, d-separable X ?

[Moore 2008] There is an L -space X such that X^2 is d-separable.

[Peng 2015] There is an L -space X such that X^2 is e-separable.

Note: $s(X) = e(X) = \aleph_0 < d(X) = \aleph_1$ so X is not d-separable.

Is there a non-separable, countably compact X so that X^2 is e-separable?

Thank you for your attention!

