

Constructive Algebra – Exercises

Autumn School Proof & Computation
Fischbachau, September 16–22, 2018

Throughout, let \mathbf{R} be a commutative ring with 1. The subset of invertible elements (*units*) of \mathbf{R} will be denoted \mathbf{R}^\times .

Exercise 1. (Constructive prime spectrum) The (single-conclusion) entailment relation \triangleright of *radical ideal* of \mathbf{R} and the (multi-conclusion) entailment relation \vdash of *proper prime ideal* [1] are inductively generated by all instances of the following axioms, respectively:

| | |
|-----------------------------|---------------------|
| $\triangleright 0$ | $\vdash 0$ |
| $a \triangleright ra$ | $a \vdash ra$ |
| $a, b \triangleright a + b$ | $a, b \vdash a + b$ |
| $a^2 \triangleright a$ | $ab \vdash a, b$ |
| | $1 \vdash$ |

Find direct, non-inductive descriptions for \triangleright and \vdash . Show that \vdash is a *conservative extension* of \triangleright , i.e., for all $a \in R$ and $U \in \text{Fin}(R)$,

$$U \triangleright a \quad \text{if and only if} \quad U \vdash a.$$

What does this say from a classical point of view? (There is a versatile criterion on the interplay of single- and multi-conclusion entailment relations due to Dana Scott [8], which has turned out widely applicable in algebra and order theory [7].)

Exercise 2. (Nil- and Jacobson radicals) Recall that the *nilradical* of \mathbf{R} is

$$\text{Nil}(\mathbf{R}) = \{ a \in \mathbf{R} : \exists n \in \mathbb{N} (a^n = 0) \}.$$

The *Jacobson radical* of \mathbf{R} is defined as

$$\text{Jac}(\mathbf{R}) = \{ a \in \mathbf{R} : \forall b \in \mathbf{R} (1 - ab \in \mathbf{R}^\times) \}.$$

- (a) Use the entailment relation \triangleright of radical ideal (cf. Exercise 1) to show that the sum of a unit and a nilpotent element is a unit.
- (b) Show that the polynomial ring $\mathbf{R}[X]$ is *rad-nil*, i.e., $\text{Jac}(\mathbf{R}[X]) = \text{Nil}(\mathbf{R}[X])$. (Hint: what do we know about the coefficients of an invertible polynomial?)

Exercise 3. (Local rings, constructively [2, 4, 6, 9]) A ring \mathbf{R} is *local* if, for all $a \in \mathbf{R}$, either a is a unit or $1 - a$ is a unit.

- (a) Show that \mathbf{R} is local if and only if, for all $a, b \in \mathbf{R}$, if $a + b$ is a unit, then so is a or b .
- (b) Let \mathbf{R} be local. Show that $\text{Jac}(\mathbf{R}) = \{ a \in \mathbf{R} : a \in \mathbf{R}^\times \rightarrow \mathbf{R} = 0 \}$ [6, Theorem III.6.5].

Exercise 4. (Minimal prime ideals) Suppose that \mathbf{R} is *reduced*, i.e., for all $a \in R$, if $a^2 = 0$, then $a = 0$. Matlis [5, Proposition 1.2(2)] has shown that a finitely generated ideal \mathfrak{a} of \mathbf{R} is contained in a minimal prime ideal of \mathbf{R} if and only if $\text{Ann}(\mathfrak{a}) \neq 0$, where

$$\text{Ann}(\mathfrak{a}) = \{ x \in \mathbf{R} : x\mathfrak{a} = 0 \}.$$

Use the entailment relation $\vdash_{\mathbf{m}}$ of maximal filter of \mathbf{R} [3] to give a constructive interpretation of Matlis' result. (Hint: write $\mathfrak{a} = \langle a_1, \dots, a_k \rangle$ and describe $\vdash_{\mathbf{m}} a_1, \dots, a_k$.)

References

- [1] Jan Cederquist and Thierry Coquand. “Entailment relations and distributive lattices”. In: *Logic Colloquium '98. Proceedings of the Annual European Summer Meeting of the Association for Symbolic Logic, Prague, Czech Republic, August 9–15, 1998*. Ed. by Samuel R. Buss, Petr Hájek, and Pavel Pudlák. Vol. 13. Lect. Notes Logic. Natick, MA: A. K. Peters, 2000, pp. 127–139.
- [2] Thierry Coquand. *A remark about the theory of local rings*. 2008. URL: <http://www.cse.chalmers.se/~coquand/local.pdf>.
- [3] Thierry Coquand and Henri Lombardi. “A logical approach to abstract algebra”. In: *Mathematical Structures in Computer Science* 16.5 (2006), pp. 885–900.
- [4] Henri Lombardi and Claude Quitté. *Commutative Algebra: Constructive Methods. Finite Projective Modules*. Dordrecht: Springer Netherlands, 2015.
- [5] Eben Matlis. “The minimal prime spectrum of a reduced ring”. In: *Illinois Journal of Mathematics* 27.3 (1983), pp. 353–391.
- [6] Ray Mines, Fred Richman, and Wim Ruitenburg. *A Course in Constructive Algebra*. Universitext. New York: Springer-Verlag, 1988.
- [7] Davide Rinaldi, Peter Schuster, and Daniel Wessel. “Eliminating disjunctions by disjunction elimination”. In: *Indag. Math. (N.S.)* 29.1 (2018). Virtual Special Issue – L.E.J. Brouwer, fifty years later, pp. 226–259.
- [8] Dana Scott. “Completeness and axiomatizability in many-valued logic”. In: *Proceedings of the Tarski Symposium (Proc. Sympos. Pure Math., Vol. XXV, Univ. California, Berkeley, Calif., 1971)*. Ed. by Leon Henkin et al. Providence, RI: Amer. Math. Soc., 1974, pp. 411–435.
- [9] Ihsen Yengui. *Constructive commutative algebra. Projective modules over polynomial rings and dynamical Gröbner bases*. Vol. 2138. Lecture Notes in Mathematics. Cham: Springer, 2015.