

Cross-temporal Probabilistic Forecast Reconciliation: online appendix

Daniele Girolimetto

Department of Statistical Sciences, University of Padova

Email: daniele.girolimetto@phd.unipd.it

George Athanasopoulos

Department of Econometrics and Business Statistics, Monash University

Email: george.athanasopoulos@monash.edu

Tommaso Di Fonzo

Department of Statistical Sciences, University of Padova

Email: tommaso.difonzo@unipd.it

Rob J Hyndman

Department of Econometrics and Business Statistics, Monash University

Email: rob.hyndman@monash.edu

30 March 2023

A Alternative forms of the cross-temporal covariance matrix

In this appendix, some derivations of the solutions proposed in Section 4 to obtain an estimator of the cross-temporal covariance matrix are reported. Starting from the the definition of cross-temporal covariance matrix we obtain the first equivalence in (10). Therefore, we have that

$$\begin{aligned} & \lambda \widehat{\mathbf{\Omega}}_{hf-bts,D} + (1 - \lambda) \widehat{\mathbf{\Omega}}_{hf-bts} \\ & \Downarrow \\ & \widehat{\mathbf{\Omega}}_{HB} = \mathbf{S}_{ct} \left[\lambda \widehat{\mathbf{\Omega}}_{hf-bts,D} + (1 - \lambda) \widehat{\mathbf{\Omega}}_{hf-bts} \right] \mathbf{S}_{ct}' \\ & = \lambda \mathbf{S}_{ct} \widehat{\mathbf{\Omega}}_{hf-bts,D} \mathbf{S}_{ct}' + (1 - \lambda) \mathbf{S}_{ct} \widehat{\mathbf{\Omega}}_{hf-bts} \mathbf{S}_{ct}'. \end{aligned}$$

The high-frequency time series representation (the second equivalence) can be derived in the following manner:

$$\begin{aligned} \mathbf{\Omega} &= \mathbf{S}_{ct} \mathbf{\Omega}_{hf-bts} \mathbf{S}_{ct}' \\ &= (\mathbf{S}_{cs} \otimes \mathbf{S}_{te}) \mathbf{\Omega}_{hf-bts} (\mathbf{S}_{cs} \otimes \mathbf{S}_{te})' \\ &= (\mathbf{I}_n \otimes \mathbf{S}_{te}) (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*}) \mathbf{\Omega}_{hf-bts} (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*})' (\mathbf{I}_n \otimes \mathbf{S}_{te})' \\ &= (\mathbf{I}_n \otimes \mathbf{S}_{te}) \mathbf{\Omega}_{hf} (\mathbf{I}_n \otimes \mathbf{S}_{te})' \end{aligned}$$

where $\mathbf{\Omega}_{hf} = (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*}) \mathbf{\Omega}_{hf-bts} (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*})'$ and $\mathbf{S}_{ct} = \mathbf{S}_{cs} \otimes \mathbf{S}_{te} = (\mathbf{I}_n \otimes \mathbf{S}_{te}) (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*})'$. We can apply the shrinkage estimator as

$$\begin{aligned} & \lambda \widehat{\mathbf{\Omega}}_{hf,D} + (1 - \lambda) \widehat{\mathbf{\Omega}}_{hf} \\ & \Downarrow \\ & \widehat{\mathbf{\Omega}}_H = (\mathbf{I}_n \otimes \mathbf{S}_{te}) \left[\lambda \widehat{\mathbf{\Omega}}_{hf,D} + (1 - \lambda) \widehat{\mathbf{\Omega}}_{hf} \right] (\mathbf{I}_n \otimes \mathbf{S}_{te})' \\ & = \lambda (\mathbf{I}_n \otimes \mathbf{S}_{te}) \widehat{\mathbf{\Omega}}_{hf,D} (\mathbf{I}_n \otimes \mathbf{S}_{te})' + (1 - \lambda) (\mathbf{I}_n \otimes \mathbf{S}_{te}) \widehat{\mathbf{\Omega}}_{hf} (\mathbf{I}_n \otimes \mathbf{S}_{te})'. \end{aligned}$$

The bottom time series representation (the third equivalence) follows by

$$\begin{aligned} \mathbf{\Omega} &= \mathbf{S}_{ct} \mathbf{\Omega}_{hf-bts} \mathbf{S}_{ct}' \\ &= (\mathbf{S}_{cs} \otimes \mathbf{S}_{te}) \mathbf{\Omega}_{hf-bts} (\mathbf{S}_{cs} \otimes \mathbf{S}_{te})' \\ &= (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*}) (\mathbf{I}_n \otimes \mathbf{S}_{te}) \mathbf{\Omega}_{hf-bts} (\mathbf{I}_n \otimes \mathbf{S}_{te})' (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*})' \\ &= (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*}) \mathbf{\Omega}_{bts} (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*})', \end{aligned}$$

where $\mathbf{\Omega}_{bts} = (\mathbf{I}_n \otimes \mathbf{S}_{te}) \mathbf{\Omega}_{hf-bts} (\mathbf{I}_n \otimes \mathbf{S}_{te})'$ and $\mathbf{S}_{ct} = \mathbf{S}_{cs} \otimes \mathbf{S}_{te} = (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*}) (\mathbf{I}_n \otimes \mathbf{S}_{te})'$. Finally we have that

$$\begin{aligned} & \lambda \widehat{\mathbf{\Omega}}_{bts,D} + (1 - \lambda) \widehat{\mathbf{\Omega}}_{bts} \\ & \Downarrow \\ & \widehat{\mathbf{\Omega}}_B = (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*}) \left[\lambda \widehat{\mathbf{\Omega}}_{bts,D} + (1 - \lambda) \widehat{\mathbf{\Omega}}_{bts} \right] (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*})' \\ & = \lambda (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*}) \widehat{\mathbf{\Omega}}_{bts,D} (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*})' + \end{aligned}$$




Figure A.1: The four graphs on the left represent the number of different parameters in the covariance matrix for the various approaches presented for different values of m and n (n_b , the number of bottom time series, is about 60% of the total). On the right, we have the boxplot of the percentage reduction in the number of parameters compared to the global approach.

$$(1 - \lambda) (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*}) \hat{\mathbf{\Omega}}_{bts} (\mathbf{S}_{cs} \otimes \mathbf{I}_{m+k^*})'.$$

In general, the covariance matrix of the reconciled forecasts is equal to $\mathbf{M} \hat{\mathbf{\Omega}} \mathbf{M}'$ where $\mathbf{M} = \mathbf{S}_{ct} \mathbf{G}$ is the projection matrix. When using the HB approach, the covariance matrix of the reconciliation and the base forecasts will be identical. Indeed, it can be shown (see Panagiotelis et al. 2021 for more details) that if \mathbf{M} is a projection matrix (6) then $\mathbf{M} \mathbf{S}_{ct} = \mathbf{S}_{ct} \mathbf{G} \mathbf{S}_{ct} = \mathbf{S}_{ct}$, and we obtain that

$$\begin{aligned} \mathbf{M} \hat{\mathbf{\Omega}}_{HB} \mathbf{M}' &= \mathbf{M} \mathbf{S}_{ct} \hat{\mathbf{\Omega}}_{hf-bts, HB} \mathbf{S}_{ct}' \mathbf{M}' \\ &= \mathbf{S}_{ct} \mathbf{G} \mathbf{S}_{ct} \hat{\mathbf{\Omega}}_{hf-bts, HB} \mathbf{S}_{ct}' \mathbf{G}' \mathbf{S}_{ct}' \\ &= \mathbf{S}_{ct} \hat{\mathbf{\Omega}}_{hf-bts, HB} \mathbf{S}_{ct}' = \hat{\mathbf{\Omega}}_{HB}. \end{aligned}$$

Figure A.1 shows the number of parameters for different values of m and n , with n_b fixed to approximately 60% of n . The right panel reports the boxplot of the percentage reductions in the number of parameters compared to the global approach.

B Cross-temporal covariance matrix for the Monte Carlo simulation

We assume two AR(2) processes with correlated errors such that

$$y_{i,t} = \phi_{i,1}y_{i,t-1} + \phi_{i,2}y_{i,t-2} + \varepsilon_{i,t}$$

where $\varepsilon_t \sim \mathcal{N}_2(\mathbf{0}_{(2 \times 1)}, \mathbf{\Omega}_{cs})$ and $i \in \{B, C\}$. Let $y_{i,T+h}$ be the true observation for the i^{th} series and $\tilde{y}_{i,T+h}$ the corresponding forecasts such that

$$\begin{aligned} y_{i,T+1} &= \phi_{i,1}y_{i,T} + \phi_{i,2}y_{i,T-1} + \varepsilon_{i,T+1} & \text{and} & & \tilde{y}_{i,T+1} &= \phi_{i,1}y_{i,T} + \phi_{i,2}y_{i,T-1} \\ y_{i,T+2} &= \phi_{i,1}y_{i,T+1} + \phi_{i,2}y_{i,T} + \varepsilon_{i,T+2} & & & \tilde{y}_{i,T+2} &= \phi_{i,1}\tilde{y}_{i,T+1} + \phi_{i,2}y_{i,T} \end{aligned} ,$$

then

$$\begin{aligned} y_{i,T+1} - \tilde{y}_{i,T+1} &= \varepsilon_{i,T+1} \\ y_{i,T+2} - \tilde{y}_{i,T+2} &= \varepsilon_{i,T+2} + \phi_{i,1}\varepsilon_{i,T+1}. \end{aligned}$$

Finally, we can compute each element of the high frequency bottom time series covariance matrix

$$\begin{aligned} Var(y_{i,T+1} - \tilde{y}_{i,T+1}) &= \sigma_i^2, \quad \forall i \in \{B, C\} \\ Var(y_{i,T+2} - \tilde{y}_{i,T+2}) &= \sigma_i^2(1 + \phi_{i,1}^2), \quad \forall i \in \{B, C\} \\ Cov[(y_{i,T+2} - \tilde{y}_{i,T+2}), (y_{i,T+1} - \tilde{y}_{i,T+1})] &= Cov[(y_{i,T+1} - \tilde{y}_{i,T+1}), (y_{i,T+2} - \tilde{y}_{i,T+2})] \\ &= \phi_{i,1}\sigma_i^2, \quad \forall i \in \{B, C\} \\ Cov[(y_{i,T+1} - \tilde{y}_{i,T+1}), (y_{j,T+1} - \tilde{y}_{j,T+1})] &= Cov[(y_{j,T+1} - \tilde{y}_{j,T+1}), (y_{i,T+1} - \tilde{y}_{i,T+1})] \\ &= \sigma_{i,j}, \quad \forall i, j \in \{B, C\}, \quad i \neq j \\ Cov[(y_{i,T+2} - \tilde{y}_{i,T+2}), (y_{j,T+1} - \tilde{y}_{j,T+1})] &= Cov[(y_{j,T+1} - \tilde{y}_{j,T+1}), (y_{i,T+2} - \tilde{y}_{i,T+2})] \\ &= \phi_{i,1}\sigma_{i,j}, \quad \forall i, j \in \{B, C\}, \quad i \neq j \\ Cov[(y_{i,T+2} - \tilde{y}_{i,T+2}), (y_{j,T+2} - \tilde{y}_{j,T+2})] &= Cov[(y_{j,T+2} - \tilde{y}_{j,T+2}), (y_{i,T+2} - \tilde{y}_{i,T+2})] \\ &= \sigma_{i,j}(1 + \phi_{i,1}\phi_{j,1}), \quad \forall i, j \in \{B, C\}, \quad i \neq j. \end{aligned}$$

In conclusion,

$$\mathbf{\Omega}_{hf-bts} = \begin{bmatrix} \sigma_B^2 & \phi_{B,1}\sigma_B^2 & \sigma_B^2(1 + \phi_{B,1}^2) & \sigma_C^2 \\ \sigma_{BC} & \phi_{B,1}\sigma_{BC} & \phi_{C,1}\sigma_{BC} & \sigma_C^2(1 + \phi_{C,1}^2) \\ \phi_{C,1}\sigma_{BC} & \sigma_{BC}(1 + \phi_{B,1}\phi_{C,1}) & \phi_{C,1}\sigma_C^2 & \sigma_C^2(1 + \phi_{C,1}^2) \end{bmatrix}$$

and

$$\mathbf{\Omega}_{ct} = \mathbf{S}_{ct}\mathbf{\Omega}_{hf-bts}\mathbf{S}_{ct}'.$$

C Monte Carlo Simulation: one-step residuals and shrinkage covariance matrix

In Section 4.1, we discussed the use of one-step residuals in estimating the covariance matrix. In particular we point out that one-step residuals lead to a biased estimate of the covariance matrix where some correlation are zeros by definition (see Figure C.2). In addition, Tables C.1, C.2 and C.3 show the Frobenius norm, CRPS, and ES skill scores as explained in the paper to investigate the effectiveness of one-step residuals. Moreover, in Tables C.4 and C.5, we have utilized a shrinkage matrix rather than the sample covariance matrix to assess the performance of our approach.

Reconciliation approach	Generation of the base forecasts paths								
	Gaussian approach: sample covariance matrix								
	ctjb	In-sample residuals				Multi-step residuals			
		G	B	H	HB	G	B	H	HB
base	8.260	17.638	16.733	22.178	21.789	7.748	6.549	3.409	2.215
ct(bu)	3.195	21.789	21.789	21.789	21.789	2.215	2.215	2.215	2.215
ct(shr _{cs} , bu _{te})	3.202	21.942	21.789	21.942	21.789	2.224	2.215	2.224	2.215
ct(wlsv _{te} , bu _{cs})	3.183	18.237	18.237	21.789	21.789	2.188	2.188	2.215	2.215
oct(wlsv)	3.766	19.174	18.611	22.304	21.789	3.082	2.191	2.910	2.215
oct(bdshr)	3.203	18.559	18.416	21.937	21.789	2.195	2.184	2.224	2.215
oct(shr)	5.217	25.015	23.457	23.413	21.789	2.260	2.202	2.226	2.215
oct(bshr)	5.282	23.772	23.997	22.146	21.789	2.720	2.220	2.756	2.215
oct(hshr)	6.161	11.336	10.940	23.598	21.789	4.138	4.167	2.225	2.215
oct(hbshr)	5.731	11.379	10.940	22.146	21.789	5.085	4.167	2.756	2.215
oct _h (shr)	3.251	20.965	19.992	22.079	21.789	2.260	2.202	2.226	2.215
oct _h (bshr)	3.602	21.306	21.022	22.146	21.789	2.720	2.220	2.756	2.215
oct _h (hshr)	4.869	11.405	10.940	22.037	21.789	4.138	4.167	2.225	2.215
oct _h (hbshr)	5.731	11.379	10.940	22.146	21.789	5.085	4.167	2.756	2.215

Table C.1: Frobenius norm between the true and the estimated covariance matrix for different reconciliation approaches and different techniques for simulating the base forecasts. Entries in bold represent the lowest value for each column, while the blue entry represent the global minimum. The reconciliation approaches are described in Table 2.




Figure C.2: Comparison of estimated covariance and correlation matrices (first simulation) for base forecasts using a parametric Gaussian (with one-step residuals) approach. The true covariance and correlation matrices are shown in Figure 7.

Reconciliation approach	Generation of the base forecasts paths								
	Gaussian approach: sample covariance matrix								
	ctjb	In-sample residuals				Multi-step residuals			
		G	B	H	HB	G	B	H	HB
$\forall k \in \{2, 1\}$									
base	1.000	1.008	1.009	1.044	1.047	0.998	0.999	1.002	1.004
ct(bu)	0.901	0.930	0.929	0.929	0.929	0.900	0.900	0.900	0.900
ct(shr _{cs} , bu _{te})	0.901	0.929	0.928	0.929	0.928	0.900	0.899	0.900	0.900
ct(wlsv _{te} , bu _{cs})	0.910	0.930	0.929	0.939	0.939	0.916	0.916	0.916	0.917
oct(wlsv)	0.922	0.942	0.944	0.951	0.953	0.930	0.930	0.930	0.931
oct(bdshr)	0.910	0.930	0.930	0.939	0.938	0.916	0.915	0.916	0.916
oct(shr)	0.941	0.999	0.985	0.983	0.973	0.903	0.902	0.902	0.903
oct(bshr)	0.951	0.995	1.000	0.983	0.986	0.922	0.922	0.921	0.922
oct(hshr)	0.987	0.995	0.993	1.039	1.026	0.972	0.972	0.974	0.975
oct(hbshr)	0.987	0.995	0.996	1.024	1.028	0.985	0.985	0.987	0.989
oct _h (shr)	0.904	0.929	0.928	0.932	0.932	0.903	0.902	0.902	0.903
oct _h (bshr)	0.923	0.948	0.952	0.951	0.954	0.922	0.922	0.921	0.922
oct _h (hshr)	0.974	0.982	0.982	1.012	1.012	0.972	0.972	0.974	0.975
oct _h (hbshr)	0.987	0.995	0.996	1.024	1.028	0.985	0.985	0.987	0.989
$k = 1$									
base	1.000	1.017	1.019	1.017	1.019	0.998	0.999	0.999	1.000
ct(bu)	0.978	0.994	0.994	0.994	0.994	0.976	0.976	0.977	0.977
ct(shr _{cs} , bu _{te})	0.977	0.993	0.993	0.994	0.993	0.976	0.976	0.976	0.976
ct(wlsv _{te} , bu _{cs})	0.986	1.002	1.002	1.003	1.003	0.993	0.993	0.993	0.993
oct(wlsv)	0.998	1.014	1.015	1.015	1.016	1.006	1.006	1.007	1.007
oct(bdshr)	0.986	1.002	1.002	1.003	1.003	0.992	0.992	0.993	0.993
oct(shr)	1.037	1.082	1.067	1.064	1.056	0.979	0.978	0.979	0.979
oct(bshr)	1.041	1.071	1.074	1.060	1.062	0.998	0.998	0.998	0.998
oct(hshr)	1.080	1.090	1.091	1.119	1.105	1.050	1.050	1.053	1.053
oct(hbshr)	1.065	1.080	1.081	1.088	1.090	1.063	1.064	1.066	1.068
oct _h (shr)	0.980	0.996	0.995	0.996	0.996	0.979	0.978	0.979	0.979
oct _h (bshr)	0.999	1.016	1.018	1.016	1.018	0.998	0.998	0.998	0.998
oct _h (hshr)	1.052	1.067	1.066	1.074	1.075	1.050	1.050	1.053	1.053
oct _h (hbshr)	1.065	1.080	1.081	1.088	1.090	1.063	1.064	1.066	1.068
$k = 2$									
base	1.000	0.998	0.999	1.071	1.075	0.998	0.999	1.005	1.008
ct(bu)	0.831	0.869	0.869	0.869	0.869	0.830	0.829	0.829	0.830
ct(shr _{cs} , bu _{te})	0.830	0.869	0.868	0.868	0.868	0.830	0.829	0.829	0.830
ct(wlsv _{te} , bu _{cs})	0.840	0.863	0.862	0.879	0.878	0.846	0.844	0.845	0.846
oct(wlsv)	0.851	0.875	0.877	0.891	0.893	0.859	0.859	0.859	0.861
oct(bdshr)	0.839	0.863	0.863	0.879	0.878	0.845	0.844	0.845	0.846
oct(shr)	0.854	0.922	0.909	0.908	0.897	0.833	0.831	0.832	0.832
oct(bshr)	0.869	0.925	0.931	0.911	0.915	0.851	0.851	0.851	0.852
oct(hshr)	0.901	0.908	0.904	0.966	0.952	0.900	0.899	0.901	0.902
oct(hbshr)	0.915	0.917	0.919	0.964	0.969	0.913	0.913	0.914	0.917
oct _h (shr)	0.834	0.868	0.865	0.872	0.872	0.833	0.831	0.832	0.832
oct _h (bshr)	0.852	0.886	0.890	0.890	0.894	0.851	0.851	0.851	0.852
oct _h (hshr)	0.902	0.904	0.904	0.953	0.952	0.900	0.899	0.901	0.902
oct _h (hbshr)	0.915	0.917	0.919	0.964	0.969	0.913	0.913	0.914	0.917

Table C.2: Simulation experiment. AvgRelCRPS defined in Section 5.1. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

Reconciliation approach	Generation of the base forecasts paths								
	Gaussian approach: sample covariance matrix								
	ctjb	In-sample residuals				Multi-step residuals			
		G	B	H	HB	G	B	H	HB
$\forall k \in \{2, 1\}$									
base	1.000	1.005	1.009	1.039	1.046	0.996	0.999	1.000	1.004
ct(bu)	0.897	0.924	0.923	0.924	0.923	0.895	0.896	0.897	0.895
ct(shr _{cs} , bu _{te})	0.896	0.924	0.923	0.923	0.922	0.895	0.895	0.896	0.896
ct(wlsv _{te} , bu _{cs})	0.906	0.924	0.923	0.933	0.932	0.912	0.911	0.910	0.912
oct(wlsv)	0.916	0.935	0.937	0.944	0.945	0.923	0.923	0.923	0.924
oct(bdshr)	0.906	0.923	0.923	0.932	0.932	0.910	0.910	0.911	0.912
oct(shr)	0.938	0.993	0.980	0.977	0.969	0.898	0.898	0.898	0.897
oct(bshr)	0.947	0.990	0.995	0.979	0.981	0.915	0.915	0.915	0.915
oct(hshr)	0.978	0.987	0.985	1.027	1.016	0.963	0.964	0.966	0.967
oct(hbshr)	0.977	0.986	0.985	1.012	1.016	0.974	0.976	0.977	0.978
oct _h (shr)	0.900	0.923	0.922	0.926	0.925	0.898	0.898	0.897	0.898
oct _h (bshr)	0.916	0.940	0.943	0.942	0.945	0.914	0.916	0.915	0.916
oct _h (hshr)	0.967	0.974	0.974	1.002	1.002	0.964	0.964	0.966	0.967
oct _h (hbshr)	0.978	0.984	0.986	1.012	1.015	0.975	0.976	0.977	0.980
$k = 1$									
base	1.000	1.014	1.020	1.015	1.019	0.997	1.000	0.997	1.000
ct(bu)	0.969	0.985	0.983	0.985	0.984	0.967	0.967	0.968	0.968
ct(shr _{cs} , bu _{te})	0.968	0.984	0.983	0.984	0.983	0.968	0.967	0.968	0.968
ct(wlsv _{te} , bu _{cs})	0.977	0.991	0.991	0.992	0.992	0.984	0.983	0.981	0.984
oct(wlsv)	0.989	1.002	1.004	1.003	1.004	0.994	0.995	0.995	0.997
oct(bdshr)	0.977	0.989	0.991	0.992	0.992	0.981	0.982	0.983	0.985
oct(shr)	1.028	1.070	1.056	1.053	1.046	0.969	0.969	0.970	0.969
oct(bshr)	1.034	1.061	1.065	1.051	1.053	0.985	0.987	0.986	0.987
oct(hshr)	1.066	1.075	1.076	1.099	1.090	1.037	1.037	1.039	1.039
oct(hbshr)	1.050	1.065	1.065	1.070	1.073	1.048	1.049	1.049	1.052
oct _h (shr)	0.971	0.985	0.985	0.986	0.986	0.969	0.969	0.969	0.969
oct _h (bshr)	0.987	1.002	1.005	1.002	1.005	0.986	0.987	0.987	0.988
oct _h (hshr)	1.040	1.053	1.053	1.059	1.058	1.036	1.036	1.040	1.040
oct _h (hbshr)	1.051	1.064	1.063	1.071	1.073	1.047	1.049	1.051	1.052
$k = 2$									
base	1.000	0.997	0.999	1.063	1.073	0.996	0.998	1.003	1.008
ct(bu)	0.831	0.867	0.867	0.867	0.867	0.829	0.829	0.830	0.828
ct(shr _{cs} , bu _{te})	0.829	0.867	0.866	0.866	0.865	0.828	0.829	0.829	0.829
ct(wlsv _{te} , bu _{cs})	0.839	0.860	0.860	0.877	0.876	0.844	0.844	0.844	0.845
oct(wlsv)	0.849	0.872	0.875	0.887	0.890	0.858	0.856	0.856	0.857
oct(bdshr)	0.839	0.861	0.861	0.876	0.875	0.845	0.843	0.845	0.844
oct(shr)	0.856	0.921	0.909	0.907	0.898	0.832	0.831	0.832	0.831
oct(bshr)	0.868	0.924	0.930	0.911	0.915	0.849	0.848	0.849	0.848
oct(hshr)	0.897	0.905	0.901	0.959	0.947	0.895	0.896	0.898	0.899
oct(hbshr)	0.910	0.912	0.912	0.957	0.961	0.906	0.909	0.909	0.910
oct _h (shr)	0.835	0.865	0.862	0.870	0.868	0.833	0.833	0.831	0.832
oct _h (bshr)	0.850	0.881	0.885	0.886	0.889	0.847	0.849	0.849	0.850
oct _h (hshr)	0.900	0.902	0.901	0.947	0.948	0.897	0.896	0.897	0.899
oct _h (hbshr)	0.910	0.910	0.914	0.957	0.961	0.907	0.908	0.909	0.912

Table C.3: Simulation experiment. ES ratio indices defined in Section 5.1. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

Reconciliation approach	Generation of the base forecasts paths								
	Gaussian approach: shrinkage covariance matrix								
	ctjb	In-sample residuals				Multi-step residuals			
		G	B	H	HB	G	B	H	HB
	$\forall k \in \{2, 1\}$								
base	1.007	1.009	1.044	1.046	0.997	0.999	1.002	1.003	1.000
ct(bu)	0.929	0.929	0.929	0.929	0.899	0.900	0.900	0.900	0.901
ct(shr _{cs} , bu _{te})	0.929	0.928	0.929	0.928	0.899	0.899	0.900	0.900	0.901
ct(wlsv _{te} , bu _{cs})	0.930	0.930	0.939	0.938	0.915	0.916	0.917	0.916	0.910
oct(wlsv)	0.943	0.944	0.951	0.952	0.929	0.930	0.931	0.930	0.922
oct(bdshr)	0.930	0.930	0.938	0.938	0.915	0.916	0.916	0.916	0.910
oct(shr)	0.994	0.982	0.980	0.973	0.902	0.902	0.903	0.902	0.941
oct(bshr)	0.995	0.998	0.983	0.986	0.921	0.922	0.922	0.922	0.951
oct(hshr)	0.994	0.994	1.035	1.025	0.971	0.972	0.974	0.974	0.987
oct(hbshr)	0.995	0.997	1.025	1.027	0.984	0.986	0.988	0.988	0.987
oct _h (shr)	0.929	0.928	0.932	0.932	0.902	0.902	0.903	0.902	0.904
oct _h (bshr)	0.948	0.951	0.951	0.953	0.921	0.922	0.922	0.922	0.923
oct _h (hshr)	0.982	0.982	1.011	1.011	0.971	0.972	0.974	0.974	0.974
oct _h (hbshr)	0.995	0.997	1.025	1.027	0.984	0.986	0.988	0.988	0.987
	$k = 1$								
base	1.017	1.019	1.017	1.019	0.998	0.999	0.999	0.999	1.000
ct(bu)	0.994	0.994	0.994	0.994	0.976	0.976	0.977	0.976	0.978
ct(shr _{cs} , bu _{te})	0.993	0.993	0.993	0.993	0.975	0.976	0.976	0.976	0.977
ct(wlsv _{te} , bu _{cs})	1.002	1.002	1.003	1.003	0.992	0.993	0.993	0.993	0.986
oct(wlsv)	1.015	1.015	1.015	1.016	1.005	1.007	1.007	1.007	0.998
oct(bdshr)	1.002	1.002	1.003	1.002	0.992	0.992	0.993	0.992	0.986
oct(shr)	1.076	1.065	1.061	1.056	0.978	0.978	0.979	0.978	1.037
oct(bshr)	1.070	1.072	1.060	1.062	0.997	0.998	0.998	0.998	1.041
oct(hshr)	1.090	1.092	1.114	1.105	1.049	1.050	1.053	1.052	1.080
oct(hbshr)	1.080	1.081	1.089	1.090	1.062	1.064	1.066	1.066	1.065
oct _h (shr)	0.996	0.995	0.996	0.996	0.978	0.978	0.979	0.978	0.980
oct _h (bshr)	1.016	1.018	1.016	1.018	0.997	0.998	0.998	0.998	0.999
oct _h (hshr)	1.066	1.067	1.075	1.075	1.049	1.050	1.053	1.052	1.052
oct _h (hbshr)	1.080	1.081	1.089	1.090	1.062	1.064	1.066	1.066	1.065
	$k = 2$								
base	0.997	0.999	1.071	1.074	0.997	0.999	1.005	1.008	1.000
ct(bu)	0.869	0.868	0.868	0.868	0.829	0.829	0.830	0.830	0.831
ct(shr _{cs} , bu _{te})	0.868	0.867	0.868	0.867	0.829	0.829	0.830	0.829	0.830
ct(wlsv _{te} , bu _{cs})	0.863	0.862	0.878	0.878	0.845	0.845	0.846	0.846	0.840
oct(wlsv)	0.876	0.877	0.891	0.892	0.859	0.860	0.860	0.860	0.851
oct(bdshr)	0.863	0.863	0.878	0.877	0.844	0.845	0.846	0.845	0.839
oct(shr)	0.918	0.906	0.906	0.897	0.832	0.832	0.833	0.832	0.854
oct(bshr)	0.924	0.928	0.911	0.915	0.850	0.851	0.852	0.851	0.869
oct(hshr)	0.907	0.905	0.962	0.951	0.898	0.899	0.902	0.902	0.901
oct(hbshr)	0.917	0.919	0.964	0.968	0.912	0.913	0.915	0.916	0.915
oct _h (shr)	0.867	0.864	0.872	0.871	0.832	0.832	0.833	0.832	0.834
oct _h (bshr)	0.886	0.890	0.890	0.893	0.850	0.851	0.852	0.851	0.852
oct _h (hshr)	0.904	0.905	0.952	0.952	0.898	0.899	0.902	0.902	0.902
oct _h (hbshr)	0.917	0.919	0.964	0.968	0.912	0.913	0.915	0.916	0.915

Table C.4: Simulation experiment. AvgRelCRPS defined in Section 5.1. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

Reconciliation approach	Generation of the base forecasts paths								
	Gaussian approach: shrinkage covariance matrix								
	ctjb	In-sample residuals				Multi-step residuals			
		G	B	H	HB	G	B	H	HB
	$\forall k \in \{2, 1\}$								
base	1.005	1.008	1.039	1.045	0.996	0.999	1.000	1.003	1.000
ct(bu)	0.923	0.923	0.923	0.923	0.895	0.896	0.897	0.897	0.897
ct(shr _{cs} , bu _{te})	0.923	0.922	0.922	0.922	0.896	0.895	0.895	0.895	0.896
ct(wlsv _{te} , bu _{cs})	0.924	0.924	0.932	0.932	0.910	0.911	0.911	0.911	0.906
oct(wlsv)	0.935	0.937	0.944	0.945	0.922	0.924	0.923	0.923	0.916
oct(bdshr)	0.924	0.924	0.932	0.931	0.909	0.911	0.911	0.910	0.906
oct(shr)	0.989	0.978	0.975	0.968	0.897	0.898	0.898	0.898	0.938
oct(bshr)	0.990	0.993	0.978	0.981	0.915	0.915	0.915	0.915	0.947
oct(hshr)	0.986	0.985	1.024	1.015	0.963	0.964	0.966	0.967	0.978
oct(hbshr)	0.985	0.986	1.012	1.015	0.973	0.976	0.977	0.978	0.977
oct _h (shr)	0.923	0.922	0.925	0.925	0.897	0.898	0.898	0.898	0.900
oct _h (bshr)	0.941	0.943	0.942	0.945	0.913	0.915	0.915	0.915	0.916
oct _h (hshr)	0.974	0.975	1.001	1.001	0.964	0.964	0.966	0.966	0.967
oct _h (hbshr)	0.985	0.986	1.013	1.016	0.973	0.976	0.977	0.978	0.978
	$k = 1$								
base	1.014	1.018	1.015	1.019	0.997	0.999	0.997	0.998	1.000
ct(bu)	0.983	0.984	0.984	0.984	0.967	0.967	0.969	0.969	0.969
ct(shr _{cs} , bu _{te})	0.983	0.982	0.982	0.983	0.966	0.967	0.966	0.966	0.968
ct(wlsv _{te} , bu _{cs})	0.991	0.992	0.993	0.992	0.983	0.983	0.983	0.983	0.977
oct(wlsv)	1.002	1.004	1.004	1.004	0.994	0.995	0.994	0.996	0.989
oct(bdshr)	0.990	0.991	0.992	0.991	0.981	0.983	0.984	0.982	0.977
oct(shr)	1.065	1.054	1.051	1.045	0.969	0.970	0.970	0.969	1.028
oct(bshr)	1.061	1.063	1.050	1.052	0.986	0.986	0.987	0.985	1.034
oct(hshr)	1.076	1.077	1.095	1.088	1.036	1.036	1.040	1.038	1.066
oct(hbshr)	1.064	1.065	1.071	1.073	1.047	1.048	1.050	1.050	1.050
oct _h (shr)	0.984	0.985	0.986	0.986	0.969	0.969	0.969	0.968	0.971
oct _h (bshr)	1.003	1.005	1.003	1.005	0.985	0.987	0.987	0.986	0.987
oct _h (hshr)	1.054	1.054	1.059	1.059	1.036	1.037	1.038	1.039	1.040
oct _h (hbshr)	1.063	1.065	1.071	1.074	1.046	1.048	1.049	1.051	1.051
	$k = 2$								
base	0.996	0.998	1.064	1.073	0.995	0.999	1.003	1.007	1.000
ct(bu)	0.867	0.866	0.867	0.866	0.829	0.829	0.830	0.830	0.831
ct(shr _{cs} , bu _{te})	0.867	0.866	0.866	0.866	0.830	0.829	0.830	0.830	0.829
ct(wlsv _{te} , bu _{cs})	0.861	0.861	0.875	0.875	0.843	0.845	0.845	0.845	0.839
oct(wlsv)	0.873	0.874	0.888	0.889	0.856	0.857	0.857	0.856	0.849
oct(bdshr)	0.862	0.861	0.876	0.874	0.843	0.844	0.844	0.844	0.839
oct(shr)	0.918	0.907	0.905	0.898	0.831	0.832	0.832	0.832	0.856
oct(bshr)	0.924	0.928	0.911	0.915	0.849	0.849	0.849	0.849	0.868
oct(hshr)	0.904	0.901	0.957	0.946	0.895	0.896	0.898	0.900	0.897
oct(hbshr)	0.912	0.913	0.956	0.961	0.905	0.909	0.909	0.911	0.910
oct _h (shr)	0.866	0.863	0.869	0.869	0.830	0.831	0.832	0.832	0.835
oct _h (bshr)	0.882	0.886	0.886	0.889	0.846	0.848	0.849	0.848	0.850
oct _h (hshr)	0.901	0.902	0.947	0.946	0.896	0.896	0.898	0.899	0.900
oct _h (hbshr)	0.912	0.914	0.958	0.961	0.905	0.908	0.910	0.909	0.910

Table C.5: Simulation experiment. ES ratio indices defined in Section 5.1. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

D Forecast reconciliation of the Australian GDP dataset

Athanasopoulos et al. (2020) proposed using state-of-the-art forecast reconciliation methods to improve the accuracy of macroeconomic forecasts and facilitate aligned decision-making. In their empirical analysis, they applied cross-sectional forecast reconciliation to 95 Australian QNA time series that represent the Gross Domestic Product (GDP) calculated using both the income and expenditure approaches. These two approaches correspond to two distinct hierarchical structures, with GDP at the top and 15 lower-level aggregates in the income approach, and GDP as the top-level aggregate in a hierarchy of 79 time series in the expenditure approach (for more information, see Athanasopoulos et al. 2020, pp. 702–705 and figures 21.4–21.7). Bisaglia et al. (2020) showed how to obtain a “one-number” forecast where the GDP reconciled forecasts are coherent for both the expenditure and income sides. Di Fonzo & Girolimetto (2022c,d) extended the one number forecasts idea to obtain fully reconciled probabilistic forecasts, and Di Fonzo & Girolimetto (2023a) computed cross-temporally reconciled point forecasts.

D.1 One-step residuals and shrinkage covariance matrix

Reconciliation approach	Generation of the base forecasts paths									
	ctjb	Gaussian approach*				ctjb	Gaussian approach*			
		G_h	H_h	G_{oh}	H_{oh}		G_h	H_h	G_{oh}	H_{oh}
		$\forall k \in \{4, 2, 1\}$					$k = 1$			
base	1.000	0.979	0.995	0.968	0.976	1.000	0.988	0.988	0.971	0.971
ct(shr_{cs}, bu_{te})	0.937	0.956	0.956	0.976	0.976	0.992	1.008	1.008	1.029	1.029
ct(wls_{cs}, bu_{te})	0.930	0.917	0.917	0.898	0.898	0.986	0.974	0.975	0.956	0.956
oct($wlsv$)	0.926	0.919	0.920	0.900	0.900	0.984	0.981	0.979	0.959	0.959
oct($bdshr$)	0.940	0.965	0.945	0.992	0.957	0.997	1.019	1.003	1.044	1.018
oct(shr)	0.944	1.020	0.940	1.094	0.988	1.015	1.095	1.010	1.160	1.059
oct($hshr$)	0.988	0.972	1.002	0.974	1.001	1.048	1.037	1.060	1.034	1.061
oct _o ($wlsv$)	0.926	0.911	0.912	0.896	0.895	0.984	0.971	0.970	0.954	0.954
oct _o ($bdshr$)	0.978	0.964	0.946	0.952	0.930	1.034	1.016	1.003	1.005	0.989
oct _o (shr)	0.950	0.946	0.922	0.925	0.903	1.014	1.003	0.985	0.987	0.968
oct _o ($hshr$)	0.989	0.966	0.984	0.954	0.965	1.047	1.028	1.038	1.012	1.023
oct _{oh} (shr)	1.102	1.059	1.001	1.094	0.988	1.172	1.109	1.066	1.160	1.059
oct _{oh} ($hshr$)	1.006	0.983	1.009	0.974	1.001	1.068	1.046	1.059	1.034	1.061
		$k = 2$					$k = 4$			
base	1.000	0.984	0.993	0.968	0.976	1.000	0.966	1.004	0.964	0.981
ct(shr_{cs}, bu_{te})	0.949	0.966	0.966	0.987	0.987	0.874	0.896	0.896	0.914	0.914
ct(wls_{cs}, bu_{te})	0.942	0.928	0.928	0.909	0.909	0.866	0.853	0.853	0.834	0.834
oct($wlsv$)	0.938	0.929	0.931	0.911	0.911	0.860	0.853	0.855	0.835	0.834
oct($bdshr$)	0.953	0.976	0.956	1.003	0.969	0.874	0.904	0.880	0.931	0.889
oct(shr)	0.955	1.031	0.951	1.113	1.002	0.866	0.940	0.864	1.015	0.909
oct($hshr$)	1.001	0.985	1.014	0.987	1.016	0.919	0.900	0.935	0.904	0.931
oct _o ($wlsv$)	0.938	0.921	0.923	0.907	0.906	0.860	0.847	0.848	0.832	0.830
oct _o ($bdshr$)	0.991	0.974	0.957	0.964	0.942	0.914	0.905	0.883	0.892	0.865
oct _o (shr)	0.965	0.958	0.934	0.938	0.916	0.877	0.882	0.852	0.854	0.831
oct _o ($hshr$)	1.002	0.979	0.996	0.967	0.978	0.922	0.898	0.923	0.888	0.898
oct _{oh} (shr)	1.120	1.069	1.013	1.113	1.002	1.020	1.002	0.928	1.015	0.909
oct _{oh} ($hshr$)	1.021	0.996	1.021	0.987	1.016	0.934	0.912	0.951	0.904	0.931

*The Gaussian method employs a sample covariance matrix:

G_h and H_h use multi-step residuals and G_{oh} and H_{oh} use overlapping and multi-step residuals.

Table D.6: AvgRelCRPS indices defined in Section 5.1 for the Australian QNA dataset. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

Reconciliation approach	Generation of the base forecasts paths									
	ctjb	Gaussian approach*				ctjb	Gaussian approach*			
		G_h	H_h	G_{oh}	H_{oh}		G_h	H_h	G_{oh}	H_{oh}
		$\forall k \in \{4, 2, 1\}$					$k = 1$			
base	1.000	0.970	0.988	0.960	0.970	1.000	0.977	0.977	0.965	0.965
ct(shr_{cs}, bu_{te})	0.897	0.944	0.944	0.973	0.973	0.964	1.001	1.001	1.033	1.033
ct(wls_{cs}, bu_{te})	0.886	0.880	0.880	0.860	0.860	0.954	0.944	0.945	0.928	0.928
oct($wlsv$)	0.890	0.890	0.894	0.872	0.872	0.958	0.957	0.957	0.938	0.939
oct($bdshr$)	0.905	0.956	0.934	0.992	0.954	0.972	1.014	0.994	1.048	1.018
oct(shr)	0.895	0.979	0.895	1.053	0.944	0.973	1.060	0.969	1.121	1.015
oct($hshr$)	0.951	0.940	0.973	0.959	0.992	1.017	1.010	1.034	1.023	1.055
oct _o ($wlsv$)	0.891	0.879	0.881	0.864	0.864	0.958	0.945	0.945	0.931	0.931
oct _o ($bdshr$)	0.940	0.928	0.910	0.918	0.895	1.004	0.986	0.971	0.980	0.961
oct _o (shr)	0.900	0.899	0.876	0.878	0.858	0.973	0.963	0.944	0.949	0.930
oct _o ($hshr$)	0.956	0.936	0.955	0.922	0.936	1.021	1.004	1.012	0.987	1.000
oct _{oh} (shr)	1.059	1.015	0.956	1.053	0.945	1.130	1.063	1.019	1.121	1.016
oct _{oh} ($hshr$)	0.986	0.968	0.999	0.959	0.992	1.053	1.034	1.049	1.024	1.055
		$k = 2$					$k = 4$			
base	1.000	0.972	0.985	0.959	0.969	1.000	0.959	1.000	0.957	0.976
ct(shr_{cs}, bu_{te})	0.915	0.961	0.960	0.991	0.991	0.818	0.874	0.874	0.899	0.900
ct(wls_{cs}, bu_{te})	0.904	0.896	0.896	0.877	0.877	0.807	0.805	0.805	0.782	0.783
oct($wlsv$)	0.909	0.907	0.912	0.889	0.889	0.811	0.813	0.819	0.794	0.794
oct($bdshr$)	0.925	0.976	0.953	1.013	0.974	0.825	0.883	0.860	0.920	0.876
oct(shr)	0.913	1.000	0.914	1.076	0.963	0.807	0.885	0.808	0.967	0.861
oct($hshr$)	0.973	0.960	0.993	0.978	1.014	0.871	0.856	0.897	0.881	0.913
oct _o ($wlsv$)	0.908	0.895	0.898	0.881	0.882	0.812	0.802	0.806	0.786	0.786
oct _o ($bdshr$)	0.960	0.947	0.929	0.938	0.915	0.860	0.856	0.836	0.841	0.816
oct _o (shr)	0.921	0.919	0.896	0.898	0.878	0.814	0.821	0.796	0.794	0.775
oct _o ($hshr$)	0.977	0.956	0.976	0.942	0.957	0.876	0.854	0.882	0.844	0.856
oct _{oh} (shr)	1.082	1.029	0.973	1.076	0.963	0.971	0.954	0.882	0.967	0.861
oct _{oh} ($hshr$)	1.007	0.988	1.017	0.979	1.014	0.904	0.888	0.934	0.881	0.913

*The Gaussian method employs a sample covariance matrix:

G_h and H_h use multi-step residuals and G_{oh} and H_{oh} use overlapping and multi-step residuals.

Table D.7: ES ratio indices defined in Section 5.1 for the Australian QNA dataset. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

Reconciliation approach	Generation of the base forecasts paths									
	ctjb	Gaussian approach*				ctjb	Gaussian approach*			
		G _h	H _h	G _{oh}	H _{oh}		G _h	H _h	G _{oh}	H _{oh}
		∀k ∈ {4, 2, 1}					k = 1			
base	1.000	0.979	1.011	0.968	0.987	1.000	0.988	0.988	0.971	0.971
ct(shr _{cs} , bu _{te})	0.937	0.960	0.961	0.962	0.960	0.992	1.001	1.001	1.004	1.000
ct(wls _{cs} , bu _{te})	0.930	0.951	0.953	0.911	0.915	0.986	0.997	0.998	0.964	0.967
oct(wlsv)	0.926	0.972	0.957	0.918	0.917	0.984	1.010	1.003	0.971	0.970
oct(bdshr)	0.940	0.986	0.966	0.981	0.956	0.997	1.015	1.006	1.016	1.000
oct(shr)	0.944	0.999	0.962	1.051	0.995	1.015	1.047	1.021	1.105	1.058
oct(hshr)	0.988	1.000	1.021	0.979	1.002	1.048	1.045	1.066	1.034	1.053
oct _o (wlsv)	0.926	0.961	0.948	0.914	0.912	0.984	1.000	0.993	0.966	0.965
oct _o (bdshr)	0.978	0.956	0.949	0.949	0.934	1.034	0.984	0.983	0.988	0.977
oct _o (shr)	0.950	0.957	0.946	0.933	0.917	1.014	0.998	0.995	0.986	0.974
oct _o (hshr)	0.989	0.997	1.013	0.967	0.982	1.047	1.039	1.054	1.019	1.032
oct _{oh} (shr)	1.102	1.010	1.006	1.051	0.995	1.172	1.059	1.063	1.105	1.058
oct _{oh} (hshr)	1.006	0.989	1.004	0.979	1.002	1.068	1.037	1.050	1.034	1.053
		k = 2					k = 4			
base	1.000	0.984	1.009	0.968	0.987	1.000	0.966	1.037	0.964	1.002
ct(shr _{cs} , bu _{te})	0.949	0.972	0.972	0.974	0.971	0.874	0.910	0.911	0.910	0.910
ct(wls _{cs} , bu _{te})	0.942	0.962	0.964	0.923	0.927	0.866	0.897	0.900	0.851	0.855
oct(wlsv)	0.938	0.988	0.968	0.931	0.929	0.860	0.921	0.903	0.856	0.856
oct(bdshr)	0.953	1.004	0.979	0.996	0.970	0.874	0.942	0.914	0.932	0.900
oct(shr)	0.955	1.016	0.973	1.070	1.010	0.866	0.937	0.895	0.981	0.922
oct(hshr)	1.001	1.015	1.034	0.993	1.017	0.919	0.942	0.965	0.913	0.937
oct _o (wlsv)	0.938	0.976	0.959	0.927	0.925	0.860	0.910	0.894	0.853	0.852
oct _o (bdshr)	0.991	0.970	0.963	0.963	0.948	0.914	0.917	0.905	0.899	0.880
oct _o (shr)	0.965	0.973	0.959	0.948	0.931	0.877	0.903	0.886	0.868	0.850
oct _o (hshr)	1.002	1.013	1.026	0.980	0.996	0.922	0.943	0.962	0.905	0.921
oct _{oh} (shr)	1.120	1.026	1.019	1.070	1.010	1.020	0.947	0.939	0.981	0.922
oct _{oh} (hshr)	1.021	1.005	1.017	0.993	1.017	0.934	0.929	0.946	0.913	0.937

*The Gaussian method employs a shrinkage covariance matrix:

G_h and H_h use multi-step residuals and G_{oh} and H_{oh} use overlapping and multi-step residuals.

Table D.8: AvgRelCRPS indices defined in Section 5.1 for the Australian QNA dataset. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

Reconciliation approach	Generation of the base forecasts paths									
	ctjb	Gaussian approach*				ctjb	Gaussian approach*			
		G_h	H_h	G_{oh}	H_{oh}		G_h	H_h	G_{oh}	H_{oh}
		$\forall k \in \{4, 2, 1\}$					$k = 1$			
base	1.000	0.967	1.002	0.957	0.980	1.000	0.973	0.973	0.961	0.962
ct(shr_{cs}, bu_{te})	0.897	0.968	0.969	0.963	0.962	0.964	1.012	1.012	1.009	1.004
ct(wls_{cs}, bu_{te})	0.886	0.939	0.944	0.882	0.888	0.954	0.994	0.998	0.947	0.952
oct($wlsv$)	0.890	0.966	0.959	0.897	0.901	0.958	1.017	1.012	0.960	0.965
oct($bdshr$)	0.905	0.997	0.981	0.986	0.960	0.972	1.031	1.021	1.024	1.005
oct(shr)	0.895	0.979	0.945	1.021	0.962	0.973	1.041	1.011	1.083	1.028
oct($hshr$)	0.951	0.997	1.023	0.973	1.005	1.017	1.051	1.073	1.034	1.063
oct _o ($wlsv$)	0.891	0.950	0.945	0.889	0.892	0.958	1.002	0.997	0.953	0.956
oct _o ($bdshr$)	0.940	0.935	0.933	0.922	0.909	1.004	0.965	0.964	0.969	0.959
oct _o (shr)	0.900	0.935	0.928	0.895	0.884	0.973	0.984	0.982	0.960	0.950
oct _o ($hshr$)	0.956	0.997	1.015	0.945	0.965	1.021	1.049	1.062	1.007	1.024
oct _{oh} (shr)	1.059	0.981	0.983	1.021	0.962	1.130	1.034	1.041	1.083	1.029
oct _{oh} ($hshr$)	0.986	0.996	1.014	0.973	1.005	1.053	1.050	1.064	1.034	1.063
		$k = 2$					$k = 4$			
base	1.000	0.970	0.999	0.955	0.980	1.000	0.958	1.033	0.953	1.000
ct(shr_{cs}, bu_{te})	0.915	0.987	0.988	0.983	0.982	0.818	0.909	0.910	0.902	0.902
ct(wls_{cs}, bu_{te})	0.904	0.958	0.962	0.900	0.906	0.807	0.871	0.876	0.805	0.812
oct($wlsv$)	0.909	0.988	0.979	0.916	0.920	0.811	0.896	0.891	0.820	0.825
oct($bdshr$)	0.925	1.024	1.005	1.010	0.984	0.825	0.938	0.919	0.926	0.895
oct(shr)	0.913	1.006	0.967	1.045	0.982	0.807	0.898	0.864	0.940	0.881
oct($hshr$)	0.973	1.020	1.046	0.994	1.028	0.871	0.924	0.954	0.897	0.929
oct _o ($wlsv$)	0.908	0.972	0.964	0.908	0.911	0.812	0.882	0.876	0.812	0.816
oct _o ($bdshr$)	0.960	0.959	0.957	0.945	0.932	0.860	0.884	0.879	0.857	0.841
oct _o (shr)	0.921	0.958	0.950	0.917	0.905	0.814	0.867	0.857	0.815	0.803
oct _o ($hshr$)	0.977	1.021	1.038	0.966	0.987	0.876	0.926	0.949	0.868	0.889
oct _{oh} (shr)	1.082	1.002	1.003	1.045	0.982	0.971	0.910	0.911	0.941	0.882
oct _{oh} ($hshr$)	1.007	1.017	1.036	0.994	1.028	0.904	0.924	0.947	0.896	0.929

*The Gaussian method employs a shrinkage covariance matrix:

G_h and H_h use multi-step residuals and G_{oh} and H_{oh} use overlapping and multi-step residuals.

Table D.9: ES ratio indices defined in Section 5.1 for the Australian QNA dataset. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

E Australian Tourism Demand dataset

Table E.10: Geographic divisions of Australia in States, Zones e Regions. Zones formed by a single region are highlighted in italics and not numbered.

Series	Name	Label	Series	Name	Label
<i>Total</i>			<i>continues Regions</i>		
1	Australia	Total	49	Gippsland	BCB
<i>States</i>			50	Phillip Island	BCC
2	New South Wales (NSW)	A	51	Central Murray	BDA
3	Victoria (VIC)	B	52	Goulburn	BDB
4	Queensland (QLD)	C	53	High Country	BDC
5	South Australia (SA)	D	54	Melbourne East	BDD
6	Western Australia (WA)	E	55	Upper Yarra	BDE
7	Tasmania (TAS)	F	56	MurrayEast	BDF
8	Northern Territory (NT)	G	57	Mallee	BEA
<i>Zones</i>			58	Wimmera	BEB
9	Metro NSW	AA	59	Western Grampians	BEC
10	Nth Coast NSW	AB	60	Bendigo Loddon	BED
	<i>Sth Coast NSW</i>	AC	61	Macedon	BEE
11	Sth NSW	AD	62	Spa Country	BEF
12	Nth NSW	AE	63	Ballarat	BEG
	<i>ACT</i>	AF	64	Central Highlands	BEG
13	Metro VIC	BA	65	Gold Coast	CAA
	<i>West Coast VIC</i>	BB	66	Brisbane	CAB
14	East Coast VIC	BC	67	Sunshine Coast	CAC
15	Nth East VIC	BD	68	Central Queensland	CBA
16	Nth West VIC	BE	69	Bundaberg	CBB
17	Metro QLD	CA	70	Fraser Coast	CBC
18	Central Coast QLD	CB	71	Mackay	CBD
19	Nth Coast QLD	CC	72	Whitsundays	CCA
20	Inland QLD	CD	73	Northern	CCB
21	Metro SA	DA	74	Tropical North Queensland	CCC
22	Sth Coast SA	DB	75	Darling Downs	CDA
23	Inland SA	DC	76	Outback	CDB
24	West Coast SA	DD	77	Adelaide	DAA
25	West CoastWA	EA	78	Barossa	DAB
	<i>Nth WA</i>	EB	79	Adelaide Hills	DAC
	<i>SthWA</i>	EC	80	Limestone Coast	DBA
	<i>Sth TAS</i>	FA	81	Fleurieu Peninsula	DBB
26	Nth East TAS	FB	82	Kangaroo Island	DBC
27	Nth West TAS	FC	83	Murraylands	DCA
28	Nth Coast NT	GA	84	Riverland	DCB
29	Central NT	GB	85	Clare Valley	DCC
<i>Regions</i>			86	Flinders Range and Outback	DCD
30	Sydney	AAA	87	Eyre Peninsula	DDA
31	Central Coast	AAB	88	Yorke Peninsula	ddb
32	Hunter	ABA	89	Australia's Coral Coast	EAA
33	North Coast NSW	ABB	90	Experience Perth	EAB
34	South Coast	ACA	91	Australia's SouthWest	EAC
35	Snowy Mountains	ADA	92	Australia's North West	EBA
36	Capital Country	ADB	93	Australia's Golden Outback	ECA
37	The Murray	ADC	94	Hobart and the South	FAA
38	Riverina	ADD	95	East Coast	FBA
39	Central NSW	AEA	96	Launceston, Tamar and the North	FBB
40	New England North West	AEB	97	North West	FCA
41	Outback NSW	AEC	98	WildernessWest	FCB
42	Blue Mountains	AED	99	Darwin	GAA
43	Canberra	AFA	100	Kakadu Arnhem	GAB
44	Melbourne	BAA	101	Katherine Daly	GAC
45	Peninsula	BAB	102	Barkly	GBA
46	Geelong	BAC	103	Lasseter	GBB
47	Western	BBA	104	Alice Springs	GBC
48	Lakes	BCA	105	MacDonnell	GBD

Source: Wickramasuriya et al. (2019), Di Fonzo & Girolimetto (2022b)

E.1 Dealing with negative reconciled forecasts

One issue in working with time series data is the presence of negative values, which can cause difficulties for certain types of models or analyses. For the base forecasts, using the bootstrap approach produces forecasts naturally non negative (ETS model with the log-transformation), while this is not true for the Gaussian approach. In this case, any negative forecast is set equal to zero. For the cross-temporal reconciliation, Di Fonzo & Girolimetto (2022a, 2023b) propose two solutions: either a state-of-the-art numerical optimization procedure (osqp, Stellato et al. 2020, 2022), or a simple heuristic strategy called set-negative-to-zero (sntz). With sntz, any negative high frequency bottom time series reconciled forecasts are set to zero, and then a cross-temporal reconciliation bottom-up is used to obtain the complete set of fully coherent forecasts. Di Fonzo & Girolimetto (2023b) found that both methods produce similar quality forecasts, but the optimization method required much more time and computational effort compared to the sntz heuristic. To reduce computational demands, we used the less time-intensive heuristic approach for reconciliation.

E.2 Tables for all the temporal aggregation orders

Reconciliation approach	Generation of the base forecasts paths									
	ctjb	Gaussian approach*				ctjb	Gaussian approach*			
		G	B	H	HB		G	B	H	HB
		$\forall k \in \{12, 6, 4, 3, 2, 1\}$					$k = 1$			
base	1.000	0.971	0.971	0.973	0.973	1.000	0.972	0.972	0.972	0.972
ct(bu)	1.321	1.011	1.011	1.011	1.011	1.077	0.983	0.982	0.982	0.982
ct(shr _{cs} , bu _{te})	1.057	0.974	0.969	0.974	0.969	0.976	0.963	0.962	0.963	0.962
ct(wlsv _{te} , bu _{cs})	1.062	0.974	0.974	0.972	0.972	0.976	0.965	0.965	0.966	0.966
oct(ols)	0.989	0.989	0.989	0.987	0.987	0.982	0.986	0.988	0.986	0.989
oct(struc)	0.982	0.962	0.961	0.961	0.959	0.970	0.963	0.963	0.963	0.963
oct(wlsv)	0.987	0.959	0.959	0.958	0.957	0.952	0.957	0.957	0.957	0.957
oct(bdshr)	0.975	0.956	0.953	0.952	0.951	0.949	0.955	0.953	0.954	0.954
oct _h (hbshr)	0.989	1.018	1.020	1.016	1.018	0.982	1.004	1.007	1.004	1.009
oct _h (bshr)	0.994	1.018	1.020	1.016	1.019	0.988	1.007	1.013	1.006	1.012
oct _h (hshr)	0.969	0.993	0.993	0.990	0.991	0.953	0.977	0.977	0.979	0.979
oct _h (shr)	1.007	0.980	0.972	0.970	0.970	1.000	0.986	0.977	0.976	0.974
		$k = 2$					$k = 3$			
base	1.000	0.970	0.969	0.970	0.971	1.000	0.971	0.971	0.972	0.973
ct(bu)	1.189	0.999	0.999	0.999	0.999	1.273	1.010	1.010	1.010	1.010
ct(shr _{cs} , bu _{te})	1.015	0.972	0.970	0.972	0.970	1.041	0.977	0.974	0.977	0.974
ct(wlsv _{te} , bu _{cs})	1.016	0.971	0.971	0.970	0.970	1.046	0.976	0.976	0.974	0.974
oct(ols)	0.992	0.991	0.991	0.990	0.991	0.994	0.992	0.993	0.991	0.992
oct(struc)	0.982	0.966	0.965	0.965	0.965	0.986	0.967	0.966	0.966	0.965
oct(wlsv)	0.972	0.961	0.960	0.960	0.960	0.983	0.963	0.962	0.962	0.962
oct(bdshr)	0.964	0.958	0.957	0.956	0.956	0.972	0.960	0.958	0.957	0.957
oct _h (hbshr)	0.992	1.013	1.015	1.012	1.015	0.994	1.019	1.021	1.018	1.020
oct _h (bshr)	0.997	1.015	1.018	1.013	1.017	0.999	1.021	1.022	1.018	1.022
oct _h (hshr)	0.965	0.987	0.987	0.986	0.987	0.971	0.994	0.994	0.992	0.993
oct _h (shr)	1.005	0.986	0.978	0.976	0.975	1.009	0.986	0.978	0.976	0.976
		$k = 4$					$k = 6$			
base	1.000	0.973	0.973	0.974	0.975	1.000	0.976	0.976	0.978	0.978
ct(bu)	1.340	1.016	1.015	1.015	1.015	1.450	1.023	1.023	1.023	1.023
ct(shr _{cs} , bu _{te})	1.061	0.978	0.973	0.978	0.973	1.094	0.978	0.972	0.978	0.972
ct(wlsv _{te} , bu _{cs})	1.068	0.977	0.977	0.974	0.974	1.103	0.977	0.977	0.974	0.974
oct(ols)	0.993	0.991	0.992	0.990	0.990	0.989	0.989	0.989	0.987	0.986
oct(struc)	0.986	0.965	0.964	0.964	0.963	0.986	0.961	0.960	0.959	0.957
oct(wlsv)	0.990	0.962	0.961	0.961	0.960	1.001	0.960	0.959	0.958	0.957
oct(bdshr)	0.977	0.959	0.956	0.955	0.954	0.985	0.956	0.953	0.950	0.948
oct _h (hbshr)	0.993	1.021	1.023	1.019	1.021	0.989	1.024	1.026	1.022	1.022
oct _h (bshr)	0.997	1.022	1.022	1.019	1.022	0.994	1.022	1.022	1.020	1.022
oct _h (hshr)	0.973	0.996	0.997	0.994	0.995	0.976	1.000	1.001	0.996	0.997
oct _h (shr)	1.009	0.984	0.976	0.973	0.973	1.010	0.978	0.970	0.967	0.967
		$k = 12$								
base	1.000	0.968	0.967	0.969	0.969					
ct(bu)	1.675	1.038	1.037	1.037	1.038					
ct(shr _{cs} , bu _{te})	1.163	0.977	0.965	0.977	0.965					
ct(wlsv _{te} , bu _{cs})	1.174	0.978	0.978	0.971	0.971					
oct(ols)	0.982	0.982	0.983	0.980	0.975					
oct(struc)	0.982	0.951	0.949	0.947	0.943					
oct(wlsv)	1.025	0.954	0.953	0.949	0.947					
oct(bdshr)	1.002	0.950	0.944	0.939	0.935					
oct _h (hbshr)	0.982	1.027	1.029	1.024	1.021					
oct _h (bshr)	0.987	1.024	1.021	1.021	1.019					
oct _h (hshr)	0.978	1.003	1.005	0.996	0.997					
oct _h (shr)	1.010	0.963	0.956	0.952	0.952					

*The Gaussian method employs a sample covariance matrix and includes four techniques (G, B, H, HB) with multi-step residuals.

Table E.11: AvgRelCRPS defined in Section 5.1 for the Australian Tourism Demand dataset. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

Generation of the base forecasts paths										
Reconciliation approach	ctjb	Gaussian approach*				ctjb	Gaussian approach*			
		G	B	H	HB		G	B	H	HB
		$\forall k \in \{12, 6, 4, 3, 2, 1\}$					$k = 1$			
base	1.000	0.956	0.955	0.958	0.951	1.000	0.952	0.950	0.952	0.950
ct(bu)	2.427	0.983	0.983	0.983	0.983	1.759	0.982	0.982	0.982	0.982
ct(shr _{cs} , bu _{te})	1.243	0.886	0.879	0.886	0.879	1.098	0.929	0.928	0.930	0.927
ct(wlsv _{te} , bu _{cs})	1.499	0.977	0.977	0.971	0.972	1.241	0.975	0.975	0.973	0.977
oct(ols)	0.955	0.893	0.891	0.893	0.888	0.975	0.937	0.936	0.936	0.935
oct(struc)	1.085	0.917	0.915	0.916	0.912	1.027	0.943	0.942	0.943	0.942
oct(wlsv)	1.132	0.933	0.929	0.931	0.927	1.050	0.951	0.949	0.950	0.949
oct(bdshr)	1.047	0.904	0.897	0.897	0.891	1.009	0.936	0.933	0.934	0.931
oct _h (hbshr)	0.956	0.889	0.886	0.888	0.884	0.975	0.937	0.936	0.937	0.935
oct _h (bshr)	0.931	0.867	0.866	0.863	0.860	0.965	0.927	0.927	0.925	0.923
oct _h (hshr)	1.081	0.935	0.931	0.935	0.927	1.028	0.952	0.951	0.952	0.950
oct _h (shr)	1.068	0.899	0.878	0.875	0.864	1.023	0.935	0.923	0.921	0.916
		$k = 2$					$k = 3$			
base	1.000	0.958	0.954	0.956	0.953	1.000	0.961	0.958	0.960	0.955
ct(bu)	2.176	1.001	1.001	1.001	1.001	2.428	0.998	0.997	0.997	0.997
ct(shr _{cs} , bu _{te})	1.192	0.927	0.921	0.927	0.921	1.245	0.911	0.904	0.911	0.904
ct(wlsv _{te} , bu _{cs})	1.400	0.992	0.992	0.988	0.988	1.500	0.991	0.991	0.986	0.987
oct(ols)	0.985	0.935	0.932	0.934	0.930	0.976	0.918	0.915	0.917	0.912
oct(struc)	1.075	0.949	0.947	0.948	0.944	1.096	0.939	0.936	0.938	0.933
oct(wlsv)	1.110	0.960	0.958	0.958	0.955	1.142	0.953	0.949	0.951	0.946
oct(bdshr)	1.045	0.938	0.933	0.933	0.929	1.060	0.926	0.920	0.921	0.915
oct _h (hbshr)	0.984	0.933	0.931	0.933	0.928	0.975	0.915	0.912	0.915	0.909
oct _h (bshr)	0.967	0.917	0.916	0.913	0.908	0.954	0.895	0.895	0.892	0.887
oct _h (hshr)	1.073	0.962	0.959	0.963	0.956	1.093	0.955	0.951	0.956	0.949
oct _h (shr)	1.064	0.933	0.916	0.913	0.904	1.082	0.923	0.903	0.900	0.890
		$k = 4$					$k = 6$			
base	1.000	0.960	0.960	0.962	0.956	1.000	0.961	0.959	0.964	0.956
ct(bu)	2.585	0.996	0.996	0.995	0.996	2.849	1.004	1.003	1.003	1.004
ct(shr _{cs} , bu _{te})	1.277	0.898	0.890	0.899	0.891	1.339	0.882	0.873	0.883	0.874
ct(wlsv _{te} , bu _{cs})	1.559	0.990	0.990	0.984	0.985	1.662	0.997	0.997	0.991	0.992
oct(ols)	0.966	0.905	0.902	0.904	0.899	0.962	0.889	0.887	0.890	0.885
oct(struc)	1.106	0.930	0.927	0.928	0.924	1.132	0.923	0.919	0.922	0.916
oct(wlsv)	1.157	0.947	0.943	0.945	0.939	1.192	0.942	0.937	0.941	0.934
oct(bdshr)	1.065	0.917	0.909	0.910	0.903	1.084	0.907	0.897	0.898	0.890
oct _h (hbshr)	0.967	0.901	0.898	0.900	0.895	0.964	0.882	0.880	0.883	0.877
oct _h (bshr)	0.943	0.879	0.878	0.876	0.871	0.932	0.856	0.855	0.851	0.848
oct _h (hshr)	1.101	0.949	0.944	0.949	0.941	1.126	0.945	0.939	0.945	0.936
oct _h (shr)	1.089	0.915	0.893	0.890	0.878	1.107	0.899	0.875	0.871	0.858
		$k = 12$								
base	1.000	0.942	0.947	0.951	0.937					
ct(bu)	2.990	0.922	0.921	0.923	0.923					
ct(shr _{cs} , bu _{te})	1.326	0.779	0.767	0.777	0.766					
ct(wlsv _{te} , bu _{cs})	1.679	0.917	0.917	0.906	0.908					
oct(ols)	0.872	0.783	0.784	0.783	0.779					
oct(struc)	1.077	0.826	0.822	0.823	0.818					
oct(wlsv)	1.149	0.851	0.845	0.847	0.840					
oct(bdshr)	1.021	0.808	0.796	0.796	0.787					
oct _h (hbshr)	0.872	0.775	0.772	0.772	0.770					
oct _h (bshr)	0.833	0.741	0.741	0.737	0.735					
oct _h (hshr)	1.066	0.851	0.846	0.848	0.838					
oct _h (shr)	1.043	0.797	0.768	0.764	0.750					

*The Gaussian method employs a sample covariance matrix and includes four techniques (G, B, H, HB) with multi-step residuals.

Table E.12: ES ratio indices defined in Section 5.1 for the Australian Tourism Demand dataset. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

Reconciliation approach	Generation of the base forecasts paths									
	ctjb	Gaussian approach*				ctjb	Gaussian approach*			
		G	B	H	HB		G	B	H	HB
		$\forall k \in \{12, 6, 4, 3, 2, 1\}$					$k = 1$			
base	1.000	0.971	0.972	0.971	0.972	1.000	0.972	0.971	0.972	0.971
ct(bu)	1.321	1.017	1.018	1.017	1.017	1.077	0.983	0.983	0.983	0.983
ct(shr _{cs} , bu _{te})	1.057	1.013	0.971	1.013	0.971	0.976	0.987	0.961	0.988	0.961
ct(wlsv _{te} , bu _{cs})	1.062	1.069	1.070	0.974	0.974	0.976	0.986	0.986	0.965	0.965
oct(ols)	0.989	1.163	1.052	1.139	0.987	0.982	1.038	0.992	1.047	0.987
oct(struc)	0.982	1.099	1.039	1.037	0.960	0.970	1.007	0.971	0.999	0.962
oct(wlsv)	0.987	1.080	1.041	0.992	0.958	0.952	1.004	0.969	0.978	0.956
oct(bdshr)	0.975	1.072	1.032	0.985	0.950	0.949	0.999	0.965	0.975	0.952
oct _h (hbshr)	0.989	1.189	1.076	1.171	1.021	0.982	1.045	1.000	1.063	1.009
oct _h (bshr)	0.994	1.202	1.073	1.168	1.021	0.988	1.046	1.012	1.063	1.012
oct _h (hshr)	0.969	1.066	1.052	1.008	0.994	0.953	0.994	0.972	0.991	0.979
oct _h (shr)	1.007	1.090	1.046	1.000	0.970	1.000	1.035	0.992	0.998	0.973
		$k = 2$					$k = 3$			
base	1.000	0.969	0.969	0.968	0.968	1.000	0.971	0.970	0.969	0.970
ct(bu)	1.189	1.000	1.000	1.000	1.000	1.273	1.013	1.013	1.013	1.013
ct(shr _{cs} , bu _{te})	1.015	1.004	0.968	1.004	0.968	1.041	1.013	0.973	1.014	0.973
ct(wlsv _{te} , bu _{cs})	1.016	1.043	1.044	0.969	0.969	1.046	1.067	1.068	0.974	0.974
oct(ols)	0.992	1.118	1.037	1.092	0.989	0.994	1.153	1.053	1.124	0.990
oct(struc)	0.982	1.075	1.022	1.020	0.963	0.986	1.099	1.041	1.033	0.964
oct(wlsv)	0.972	1.064	1.021	0.987	0.958	0.983	1.083	1.041	0.993	0.960
oct(bdshr)	0.964	1.057	1.015	0.983	0.953	0.972	1.075	1.033	0.988	0.955
oct _h (hbshr)	0.992	1.136	1.055	1.116	1.014	0.994	1.178	1.075	1.153	1.020
oct _h (bshr)	0.997	1.145	1.059	1.114	1.016	0.999	1.190	1.075	1.151	1.021
oct _h (hshr)	0.965	1.050	1.029	1.001	0.986	0.971	1.067	1.051	1.009	0.994
oct _h (shr)	1.005	1.083	1.035	1.001	0.973	1.009	1.097	1.050	1.004	0.974
		$k = 4$					$k = 6$			
base	1.000	0.973	0.973	0.971	0.973	1.000	0.976	0.977	0.975	0.977
ct(bu)	1.340	1.021	1.021	1.021	1.021	1.450	1.032	1.033	1.032	1.033
ct(shr _{cs} , bu _{te})	1.061	1.018	0.974	1.018	0.974	1.094	1.023	0.974	1.024	0.974
ct(wlsv _{te} , bu _{cs})	1.068	1.087	1.089	0.976	0.976	1.103	1.108	1.110	0.978	0.978
oct(ols)	0.993	1.186	1.068	1.148	0.989	0.989	1.223	1.080	1.184	0.987
oct(struc)	0.986	1.120	1.057	1.042	0.962	0.986	1.141	1.071	1.054	0.959
oct(wlsv)	0.990	1.100	1.059	0.996	0.959	1.001	1.115	1.076	0.998	0.958
oct(bdshr)	0.977	1.091	1.049	0.989	0.952	0.985	1.103	1.064	0.989	0.949
oct _h (hbshr)	0.993	1.215	1.095	1.182	1.022	0.989	1.258	1.112	1.225	1.026
oct _h (bshr)	0.997	1.230	1.089	1.178	1.023	0.994	1.278	1.101	1.219	1.025
oct _h (hshr)	0.973	1.084	1.071	1.012	0.996	0.976	1.097	1.091	1.017	1.002
oct _h (shr)	1.009	1.108	1.062	1.003	0.972	1.010	1.113	1.070	1.000	0.968
		$k = 12$								
base	1.000	0.968	0.969	0.969	0.971					
ct(bu)	1.675	1.056	1.057	1.057	1.057					
ct(shr _{cs} , bu _{te})	1.163	1.032	0.974	1.033	0.974					
ct(wlsv _{te} , bu _{cs})	1.174	1.128	1.130	0.982	0.982					
oct(ols)	0.982	1.277	1.085	1.252	0.982					
oct(struc)	0.982	1.158	1.074	1.075	0.950					
oct(wlsv)	1.025	1.122	1.085	1.001	0.954					
oct(bdshr)	1.002	1.110	1.071	0.989	0.941					
oct _h (hbshr)	0.982	1.322	1.125	1.305	1.033					
oct _h (bshr)	0.987	1.347	1.107	1.297	1.031					
oct _h (hshr)	0.978	1.106	1.107	1.021	1.010					
oct _h (shr)	1.010	1.107	1.067	0.991	0.959					

*The Gaussian method employs a shrinkage covariance matrix and includes four techniques (G, B, H, HB) with multi-step residuals..

Table E.13: AvgRelCRPS defined in Section 5.1 for the Australian Tourism Demand dataset. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

Reconciliation approach	Generation of the base forecasts paths									
	ctjb	Gaussian approach*				ctjb	Gaussian approach*			
		G	B	H	HB		G	B	H	HB
		$\forall k \in \{12, 6, 4, 3, 2, 1\}$					$k = 1$			
base	1.000	0.958	0.984	0.972	0.992	1.000	0.954	0.958	0.954	0.958
ct(bu)	2.427	1.040	1.042	1.040	1.041	1.759	1.001	1.002	1.002	1.002
ct(shr _{CS} , bu _{te})	1.243	0.988	0.913	0.990	0.913	1.098	1.011	0.938	1.013	0.938
ct(wlsv _{te} , bu _{CS})	1.499	1.117	1.120	1.025	1.025	1.241	1.019	1.020	0.990	0.990
oct(ols)	0.955	1.000	0.984	0.985	0.922	0.975	0.983	0.961	0.987	0.945
oct(struc)	1.085	1.094	1.047	1.018	0.952	1.027	1.054	0.981	1.022	0.953
oct(wlsv)	1.132	1.137	1.065	1.059	0.969	1.050	1.078	0.989	1.043	0.960
oct(bdshr)	1.047	1.085	1.013	1.011	0.927	1.009	1.050	0.966	1.019	0.942
oct _h (hbshr)	0.956	1.018	0.981	1.016	0.919	0.975	0.991	0.961	1.002	0.947
oct _h (bshr)	0.931	1.002	1.001	0.982	0.889	0.965	0.980	0.975	0.985	0.933
oct _h (hshr)	1.081	1.109	1.039	1.076	0.973	1.028	1.061	0.978	1.052	0.963
oct _h (shr)	1.068	1.088	1.008	0.995	0.896	1.023	1.061	0.966	1.011	0.924
		$k = 2$					$k = 3$			
base	1.000	0.960	0.971	0.958	0.972	1.000	0.963	0.981	0.966	0.986
ct(bu)	2.176	1.035	1.036	1.035	1.035	2.428	1.042	1.044	1.042	1.043
ct(shr _{CS} , bu _{te})	1.192	1.020	0.942	1.021	0.942	1.245	1.009	0.931	1.011	0.931
ct(wlsv _{te} , bu _{CS})	1.400	1.104	1.106	1.018	1.019	1.500	1.127	1.130	1.029	1.029
oct(ols)	0.985	1.028	1.008	1.002	0.950	0.976	1.020	1.004	0.994	0.938
oct(struc)	1.075	1.115	1.051	1.039	0.967	1.096	1.117	1.064	1.033	0.965
oct(wlsv)	1.110	1.149	1.065	1.070	0.979	1.142	1.160	1.082	1.073	0.981
oct(bdshr)	1.045	1.105	1.024	1.033	0.949	1.060	1.109	1.032	1.029	0.943
oct _h (hbshr)	0.984	1.041	1.007	1.024	0.951	0.975	1.036	1.002	1.023	0.937
oct _h (bshr)	0.967	1.029	1.025	0.998	0.928	0.954	1.024	1.025	0.993	0.911
oct _h (hshr)	1.073	1.122	1.042	1.083	0.983	1.093	1.129	1.054	1.090	0.984
oct _h (shr)	1.064	1.110	1.019	1.018	0.922	1.082	1.116	1.030	1.015	0.915
		$k = 4$					$k = 6$			
base	1.000	0.962	0.987	0.973	0.996	1.000	0.963	0.998	0.984	1.011
ct(bu)	2.585	1.052	1.054	1.053	1.053	2.849	1.083	1.085	1.083	1.084
ct(shr _{CS} , bu _{te})	1.277	1.000	0.923	1.002	0.923	1.339	0.999	0.921	1.000	0.920
ct(wlsv _{te} , bu _{CS})	1.559	1.150	1.153	1.037	1.037	1.662	1.189	1.193	1.066	1.066
oct(ols)	0.966	1.022	1.008	0.994	0.931	0.962	1.023	1.014	1.003	0.930
oct(struc)	1.106	1.120	1.076	1.031	0.963	1.132	1.132	1.100	1.039	0.972
oct(wlsv)	1.157	1.167	1.097	1.075	0.982	1.192	1.187	1.124	1.090	0.995
oct(bdshr)	1.065	1.112	1.041	1.025	0.939	1.084	1.121	1.058	1.029	0.940
oct _h (hbshr)	0.967	1.041	1.005	1.027	0.929	0.964	1.046	1.008	1.042	0.924
oct _h (bshr)	0.943	1.028	1.028	0.994	0.900	0.932	1.029	1.032	1.000	0.887
oct _h (hshr)	1.101	1.137	1.068	1.093	0.986	1.126	1.153	1.089	1.110	0.999
oct _h (shr)	1.089	1.118	1.039	1.012	0.910	1.107	1.118	1.045	1.006	0.902
		$k = 12$								
base	1.000	0.948	1.010	1.002	1.033					
ct(bu)	2.990	1.028	1.031	1.029	1.029					
ct(shr _{CS} , bu _{te})	1.326	0.897	0.830	0.899	0.830					
ct(wlsv _{te} , bu _{CS})	1.679	1.119	1.123	1.009	1.009					
oct(ols)	0.872	0.927	0.914	0.930	0.840					
oct(struc)	1.077	1.028	1.012	0.950	0.894					
oct(wlsv)	1.149	1.089	1.041	1.006	0.922					
oct(bdshr)	1.021	1.015	0.964	0.935	0.855					
oct _h (hbshr)	0.872	0.955	0.906	0.978	0.833					
oct _h (bshr)	0.833	0.927	0.927	0.927	0.784					
oct _h (hshr)	1.066	1.056	1.005	1.026	0.926					
oct _h (shr)	1.043	1.011	0.952	0.909	0.809					

*The Gaussian method employs a shrinkage covariance matrix and includes four techniques (G, B, H, HB) with multi-step residuals.

Table E.14: ES ratio indices defined in Section 5.1 for the Australian Tourism Demand dataset. Approaches performing worse than the benchmark (bootstrap base forecasts, ctjb) are highlighted in red, the best for each column is marked in bold, and the overall lowest value is highlighted in blue. The reconciliation approaches are described in Table 2.

References

- Athanasopoulos, G., Gamakumara, P., Panagiotelis, A., Hyndman, R. J. & Affan, M. (2020), Hierarchical Forecasting, *in* P. Fuleky, ed., ‘Macroeconomic Forecasting in the Era of Big Data’, Vol. 52, Springer International Publishing, Cham, pp. 689–719.
- Bisaglia, L., Fonzo, T. D. & Girolimetto, D. (2020), Fully reconciled GDP forecasts from Income and Expenditure sides, *in* A. Pollice, N. Salvati & F. Schirripa Spagnolo, eds, ‘Book of Short Papers SIS 2020’, Pearson, pp. 951–956.
- Di Fonzo, T. & Girolimetto, D. (2022a), ‘Enhancements in cross-temporal forecast reconciliation, with an application to solar irradiance forecasts’, *arXiv:2209.07146* .
- Di Fonzo, T. & Girolimetto, D. (2022b), ‘Forecast combination-based forecast reconciliation: Insights and extensions’, *International Journal of Forecasting* . in press.
- Di Fonzo, T. & Girolimetto, D. (2022c), Fully reconciled probabilistic GDP forecasts from Income and Expenditure sides, *in* A. Balzanella, M. Bini, C. Cavicchia & R. Verde, eds, ‘Book of Short Papers SIS 2022’, Pearson, pp. 1376–1381.
- Di Fonzo, T. & Girolimetto, D. (2022d), ‘Point and probabilistic forecast reconciliation for general linearly constrained multiple time series’, *Working paper* .
- Di Fonzo, T. & Girolimetto, D. (2023a), ‘Cross-temporal forecast reconciliation: Optimal combination method and heuristic alternatives’, *International Journal of Forecasting* **39**(1), 39–57.
- Di Fonzo, T. & Girolimetto, D. (2023b), ‘Spatio-temporal reconciliation of solar forecasts’, *Solar Energy* **251**, 13–29.
- Panagiotelis, A., Athanasopoulos, G., Gamakumara, P. & Hyndman, R. J. (2021), ‘Forecast reconciliation: A geometric view with new insights on bias correction’, *International Journal of Forecasting* **37**(1), 343–359.
- Stellato, B., Banjac, G., Goulart, P., Bemporad, A. & Boyd, S. (2020), ‘OSQP: An operator splitting solver for quadratic programs’, *Mathematical Programming Computation* **12**(4), 637–672.
- Stellato, B., Banjac, G., Goulart, P., Boyd, S. & Bansal, V. (2022), *osqp: Quadratic Programming Solver using the ‘OSQP’ Library*. R package v0.6.0.7.
URL: osqp.org
- Wickramasuriya, S. L., Athanasopoulos, G. & Hyndman, R. J. (2019), ‘Optimal Forecast Reconciliation for Hierarchical and Grouped Time Series Through Trace Minimization’, *Journal of the American Statistical Association* **114**(526), 804–819.