

School of Physics & Astronomy
Queen Mary University of London

The Effects of Energy Densities on the CMB Power Spectrum

Daniil Dolgich (150066826)

Supervised by Prof. Teppei Katori

Submitted August 2019 in part fulfilment of the requirements for the degree of
Bachelor in Physics and Particle Physics at Queen Mary University of London

Declaration

I hereby certify that this Dissertation, has been written by me at the School of Physics and Astronomy, Queen Mary University of London, that all material in this dissertation which is not my own work has been properly acknowledged, and that it has not been submitted in any previous application for a higher degree.

Daniil Dolgich (150066826)

Acknowledgements

I'd like to give my appreciation to Prof. Teppei Katori for supervising my project even though it was outside of his comfort zone. There was a lot of mutual learning that made this all the more valuable. I would also like to thank him for his patience and support in regards to my health and well-being.

Consultation by Dr. Alkistis Pourtsidou was very helpful in kick-starting my understanding regarding the topics in this project. Her passion for her field inspired me to pursue this topic.

Many thanks as well to Harvey Abraham-Green and Leonie Dos Santos for offering their time and support when it was needed most.

Finally, a big thank you to my parents.

Abstract

We studied the Cosmic Microwave Background radiation (CMB) in order to better understand the lesser studied portions of the CMB Power Spectrum. To facilitate this, I created my own CMB simulator with Python 2.7, using Jupyter Notebook as the compiler and Github to commit and push my work.

I used this simulator to demonstrate the effects that different values of the energy density for Baryonic matter (Ω_b), Dark Matter (Ω_c), and Neutrinos (Ω_ν) have on the CMB power spectrum, and what observations could be made regarding the outcomes.

I found that altering the values of Ω_c caused the first peak of the CMB spectrum to greatly increase in amplitude and width, whilst smoothing out the rest of the power spectrum past that point. It also moved the maximum of the peak from $\ell \approx 200$ to $\ell \approx 300$, resulting in a huge and significant temperature fluctuation of $\sim 0.1 \mu\text{K}$.

Altering the values of Ω_c caused the overall amplitude of the spectrum to go down, but maintained well defined peaks and did not shift the relative x-axis position of the peaks in any direction.

Increasing the value of Ω_ν caused an overall smoothing of the power spectrum, as well as a large reduction in the amplitudes of the peaks, and a slight shift of the first peak maximum to the left.

We saw that having a higher ratio of baryonic matter would result in a huge increase to the CMB anisotropy, and that a high neutrino energy density creates more "empty space" (space that is occupied by extremely unreactive neutrinos instead of any other particle). These would change our perception of when important cosmological events occurred, and would also affect large cosmological bodies such as galaxy clusters.

Contents

List of Figures	vi
Listings	vii
1 Introduction	1
2 Methodology	4
2.1 Preliminary research	4
2.2 Making the simulator	4
3 Results	10
4 Conclusions	15
Bibliography	16

List of Figures

1.1	WMAP CMB Anisotropy Photo. Note the speckled appearance as a result of many hot and cold zones. Edward L. Wright (2013)	2
1.2	Cumulative image of the sky, featuring a multipole ℓ value of 260. This closely resembles the pattern in figure 1.1, and the fluctuations are also not equally distributed. You can also see the corresponding relative position on the CMB power spectrum plot in the top left corner. Clem Pryke (2000)	3
2.1	The WMAP nine-year power spectrum to which I was comparing my plot. Notice how the maximum peak at around $\ell \approx 200$ is also around 1° on the angular scale. NASA (2012)	8
2.2	My plot of the Scalar Non-lensed CMB Power Spectrum TT	9
3.1	TT Spectrum with biased Ω_b . The first peak maximum skyrockets beyond the y-axis limit while the other peaks become smeared and start to homogenise.	11
3.2	TT Spectrum with biased Ω_c . Lowered total amplitude but peaks are still well defined and the position of the peak maximum has remained the same.	12
3.3	TT Spectrum with biased Ω_ν . Huge decrease in overall amplitude, in addition to lots of smoothing of the line, suggesting increased homogeneity due to the increased amount of neutrinos in this particular simulation.	13

Listings

2.1	Setting up	5
2.2	Initialising the first plot	6
2.3	Creating the first plot	6

1 Introduction

The Cosmic Microwave Background radiation (CMB) is electromagnetic radiation (EM) from the time of the Big Bang. Over the course of around 13 billion years, the CMB has been red-shifted to longer wavelengths due to the persisting expansion of the universe. As such, it has cooled dramatically from its initial state about 300,000 years after the Big Bang. The exact nature of the origins of the CMB and the Big Bang theory are beyond the scope of this project; however, the important part of this is that the CMB is about 2.7 Kelvin above absolute zero. [Rita Tojeiro \(2006\)](#)

At cosmological scales, about 100 giga parsecs (Gpc), the universe may be taken to be homogeneous and isotropic. In other words, the universe is the same in all locations, and in all directions. The CMB is therefore also considered to be both of these things when looking at such immense scales. However, this is not quite the case for the CMB.

One can see in the image of the CMB taken by the Planck satellite (figure 1.1 that it's quite the opposite in fact. There are many zones of differing temperatures. The CMB is not wholly isotropic and this image shows that very clearly. These fluctuations are incredibly small, within the order of micro-Kelvin, and are collectively known as anisotropy. [Edward L. Wright \(2013\)](#)

The way to describe the CMB anisotropy is with spherical harmonics. [Martin White \(1997\)](#) As such, the CMB power spectrum is actually an angular power spectrum. Under the model of Inflation, these temperature fluctuations are Gaussian, and the variance is given by C_ℓ . ℓ in this instance is the multipole moment. Different values of ℓ represent a given angle in the sky, and can be thought of as angular frequency for the temperature fluctuations, with higher values of ℓ corresponding to smaller angles. Essentially, the value of ℓ can be thought of as "how many opposite zones are there", the simplest of which is a dipole, in which there are two zones, opposites of each other, just like a bar magnet. As you increase the number of zones, the number of poles will also increase. This is essentially what the CMB




Figure 1.1: WMAP CMB Anisotropy Photo. Note the speckled appearance as a result of many hot and cold zones. [Edward L. Wright \(2013\)](#)

power spectrum represents; at higher values of ℓ , there are many many more zones being observed simultaneously. This can be seen in figure 1.2.

We can see how the plots of the multipole moment against the temperature fluctuations in μK^2 results in the CMB power spectrum. The three characteristic peaks correspond to the curvature of the universe, the baryon density, and the cold dark matter density, though the exact reasons why are beyond the scope of this research.

There are three commonly considered energy densities used to construct the CMB power spectrum: Ω_b which denotes normal baryonic matter, Ω_c denotes cold dark matter, and Ω_Λ denotes dark energy. This however neglects another energy density value, specifically, Ω_ν , the energy density of neutrinos.

With this in mind, I decided to create my own CMB simulator to see exactly how different values for these energy densities affects the power spectrum, and how the overall composition of the universe would change.




Figure 1.2: Cumulative image of the sky, featuring a multipole ℓ value of 260. This closely resembles the pattern in figure 1.1, and the fluctuations are also not equally distributed. You can also see the corresponding relative position on the CMB power spectrum plot in the top left corner. [Clem Pryke \(2000\)](#)

2 Methodology

2.1 Preliminary research

Before I could begin, I had to first familiarise myself with the precise meanings of each element of the CMB power spectrum and how anisotropy was the underlying aspect.

I found online, two simulators of the CMB, one simplistic and the other incredibly in-depth. [Chris North \(2013\)](#) The simplistic one was similar to what I had planned to make myself: a simulator that allows changing of variables and instantly outputs the new graph. The second simulator did not have this feature but had a large selection of variables to tweak. [NASA \(2019\)](#)

The second simulator, upon completion of the simulation, would also provide a data file in which were contained the multipole moment values ranging from 2 to 2200, as well as the temperature fluctuations for the different versions of the CMB spectrums. The one I was concerned with was the temperature angular power spectrum (TT). I used this file as the input for all the base variables in my simulator. I also looked at the values presented for the temperature-polarisation cross-power spectrum (TE) and the polarised spectrum (EE), and decided to also simulate these graphs for the purposes of comparison; to see whether they get affected the same way as my baseline TT graph. There was no intent to delve deeper into the effects of the polarisation at this time.

2.2 Making the simulator

The language of choice was Python 2.7. Using this I was able to start work on constructing the simulator.

I chose to use Jupyter Notebook [Project Jupyter \(2019a\)](#) as my platform to code on, and a Github repository [Daniil Dolgich \(2019\)](#) to commit and push my work

to. The first step was to gather all the necessary data, as well as import all the necessary packages.

```
1 import camb
2 import matplotlib
3 import matplotlib.pyplot as plt
4 import numpy as np
5 from ipywidgets import interact
6 import ipywidgets as widgets
7
8 params = camb.read_ini('camb_interface/camb_params.ini')
```

Listing 2.1: Setting up

This imported the primary packages that I needed in order to make the graph and have it actually function. CAMB is Code for Anisotropies in the Microwave Background and is used for calculating many things within cosmology [Antony Lewis \(2013\)](#). It just so happens to also use Python as the main code. Matplot was necessary to have a way to draw the graph. NumPy was used to manipulate the data from CAMB and pass it on to Matplot to use in the formation of the graph [NumPy Developers \(2019\)](#). IPyWidgets was to create the interactive controls to vary the values of the graph and output the changes in real time [Project Jupyter \(2019b\)](#).

The next step was to create a function that would be called and "interacted" with. I defined this first run as "camb_plot_tt", and also defined the values that I would have available to change; each of the "om_h2" represent their respective energy densities, with om standing for Ω , h2 being Planck's constant squared (h^2), and with the letter corresponding to its respective object: b for baryons, c for dark matter, nu for neutrinos.

Here I set the initial values for each of the three variables as the defaults present within the CAMB data. By assigning variables in Python, I was able to save some time and avoid having to rewrite too much. 'TxT' was taken from the array and contained all the data points for plotting the TT CAMB graph. Once I assigned reasonable limits for the axes and added a handy feature to save the future plot as a .pdf file, I moved on to figuring out how to create the interactive sliders for the three variables.

```

1 %matplotlib inline
2 def camb_plot_tt(ombh2, omch2, omnuh2):
3     params.ombh2 = ombh2
4     params.omch2 = omch2
5     params.omnuh2 = omnuh2
6     results = camb.get_results(params)
7
8     unlensed = results.get_cmb_unlensed_scalar_array_dict(lmax=2200,
9         CMB_unit='muK')
10
11     tt = unlensed['TxT']
12
13     fig, ax = plt.subplots()
14     tt = np.delete(tt, [0, 1])
15     ax.plot(tt, linewidth=1)
16     ax.set_xlim([None, 2500])
17     ax.set_ylim([None, 7000])
18     ax.set_xlabel('Multipole Moment (l)', ylabel='CMB_Unit (uK^2)',
19         title='Cl TT CMB Power Spectrum')
20     fig.savefig("Cl-TT-vs-l.pdf")
21     plt.show()

```

Listing 2.2: Initialising the first plot

```

1 params = camb.read_ini('camb_interface/camb_params.ini')
2 interact(camb_plot_tt,
3     ombh2=widgets.FloatSlider(value=params.ombh2, min=0, max=1,
4         step=0.001, readout_format='.5f'),
5     omch2=widgets.FloatSlider(value=params.omch2, min=0, max=1,
6         step=0.001, readout_format='.5f'),
7     omnuh2=widgets.FloatSlider(value=params.omnuh2, min=0, max=1,
8         step=0.001, readout_format='.5f'));

```

Listing 2.3: Creating the first plot

By using the floatsliders present in iPywidgets, I was able to assign the minimum and maximum values for the variables and also assign an appropriate step value of 0.001. The readout format was chosen specifically to avoid visually truncating the value of the Ω_ν value, as it's very small at default (0.00064).

Running these three cells in sequence created the graph you see in figure 2.2, which is a near perfect match to the plot I saw during my initial research. With this now functioning program, I set about checking what happens when I change the values of the variables.

Initially however, I discovered a flaw in my programming. Because I had constrained Ω_Λ , this meant that its value was completely fixed; the total energy density Ω_{total} has to equal 1 because it is comprised of ratios, but in my simulator, you can go beyond that and end up breaking the simulator. This same error occurs if variables are changed too quickly.

Regardless, I was able to get a working version of the simulator, and decided to plot the EE and TE values to see that they also matched the ones seen on the Lambda website (see figure 2.1).




Figure 2.1: The WMAP nine-year power spectrum to which I was comparing my plot. Notice how the maximum peak at around $l \approx 200$ is also around 1° on the angular scale. [NASA \(2012\)](#)




Figure 2.2: My plot of the Scalar Non-lensed CMB Power Spectrum TT

3 Results

I chose to alter each variable individually, one at a time, in order to be sure that that specific variable was responsible for the change in the graph.

My predictions were mainly these:

Increasing Ω_b would result in the graph becoming more "smeared", causing the peaks to shrink and become less defined, as more regular matter would mean there are many more minute interactions occurring and thus muddying the spectrum.

Increasing Ω_c would result in the graph growing in intensity, as dark matter is much more massive than regular matter, and so I expect the fluctuations to also increase alongside the increase in dark matter.

Increasing Ω_ν would also result in a smearing of the graph, and a reduction in the amplitudes, but not for the same reasons as with baryonic matter; neutrinos are incredibly light and interact with very little, meaning that an increase in neutrinos would result in there being less room within each measured multipole for dark matter or baryonic matter. Essentially, neutrinos would add what could be thought of as "empty space" between particles that would otherwise interact.

The most notable thing is that past a certain point, increases in value for each variable have less and less effect on the graph. There appears to be a point at which the graph seemingly stops changing. I assume this is due to my constrained Ω_Λ not allowing any further changes to occur. However, the immediate effects are visible from even very minute changes to each variable.

Strangely, for baryonic matter, my prediction was only half wrong. In figure [3.1](#) the amplitudes of the second and third peaks became heavily smeared, as expected, but the first peak amplitude rose well past the y-limit I had assigned in my code. I cannot say as to why that is the case. Interestingly, the dipole peak seems to have




Figure 3.1: TT Spectrum with biased Ω_b . The first peak maximum skyrockets beyond the y-axis limit while the other peaks become smeared and start to homogenise.

also shifted to around $\ell \approx 300$, suggesting that the corresponding angle has also gone down below 1° .

For dark matter, this test also went against my prediction entirely. Figure 3.2 shows that increasing the dark matter actually greatly lowered the amplitudes of all the peaks. However, and quite notably, it has not smeared the graph like the baryonic change. The peaks are still quite well defined, and only their amplitudes have shrunk. This suggests that this ends up greatly reducing the anisotropy, and thus the temperature fluctuations. Additionally, the two most well-defined peaks remain at $\ell \approx 200$ and the third peak which we know from our preliminary research to be the dark matter peak; considering it is the amount of dark matter we have increased in this simulation, this is a very logical outcome.

Figure 3.3 shows the neutrino biased simulation to be quite similar in results to the the dark matter simulation initially, however, the peaks are not as well defined, indicating that the graph has indeed been smeared and smoothed, matching my initial prediction. This was the particular simulation I was interested in doing, as




Figure 3.2: TT Spectrum with biased Ω_c . Lowered total amplitude but peaks are still well defined and the position of the peak maximum has remained the same.




Figure 3.3: TT Spectrum with biased Ω_ν . Huge decrease in overall amplitude, in addition to lots of smoothing of the line, suggesting increased homogeneity due to the increased amount of neutrinos in this particular simulation.

it had been absent from the two simulators mentioned in the introduction to this document, and it shows that neutrino energy density is as impactful and important to the composition of the universe as the three most considered values for baryonic matter, dark matter and dark energy.

Comparing the three graphs shows that biasing Ω_b creates the most noticeable change, and more specifically, that it hugely increases the temperature fluctuations at the lower values of ℓ , as well as moving the maximum of the largest first peak to be at a slightly higher value of ℓ compared to the other two graphs. In figure [3.1](#), the first peak exceeds the maximum limit on the graph; I predict it reaches at least $10000\mu K^2$, meaning that the real temperature fluctuation is $\sim 0.1\mu K$, which is a very significant anisotropy.

One could infer what sort of consequences may arise on a universal scale as a result of this, such as how it would affect the rate of expansion, as well as the percentage of accompanying dark matter, dark energy, and neutrinos. Certain other phenomenon such as galaxy cluster lensing would also get affected by fluctuating rates of dark matter, and the size of the galaxy clusters would also change in accordance with the amount of dark matter present in the universe.

4 Conclusions

By matching the simulations to observed results, we can figure out precise, allowed values for specific objects and phenomena. Here, most notably, we can see how big of an impact the energy density of neutrinos has on the CMB power spectrum, and indeed, also confirms that neutrinos are not massless.

This also means that neutrinos must have a maximum allowed mass. Since energy is related to mass, if their energy density was higher, their mass would also need to be higher. As seen from the simulation, a higher amount of neutrinos would likely start to prevent further reactions from taking place.

Additionally, the dark matter result was not expected and shows that an increase in dark matter would result in much lower temperature fluctuations across the CMB, meaning that in turn, the universe could be considered homogeneous and isotropic at slightly smaller scales than what are used now.

These rudimentary simulations have shown the possible effects on CMB anisotropy, and subsequently on the distribution of different types of matter within the universe. Though these are mostly hypothetical scenarios, they can be useful to help us to predict the behaviour of certain particles, or indeed on larger scales such as galaxy clusters and possibly even black holes, though more research would need to be done on that particular case.

There is much more that can be factored in that was beyond the scope of this project; the lensing effect along with the alternate forms of the CMB spectrum in the temperature-polarisation cross-spectrum TE, and the polarised spectrum EE. Exploring this further could yield some very interesting results.

Bibliography

- Antony Lewis. Python CAMB Database, 2013. URL <https://camb.readthedocs.io/en/latest/>.
- Chris North. Planck CMB Simulator, 2013. URL <https://chrisnorth.github.io/planckapps/Simulator/#about>.
- Clem Pryke. Harmonic Multipoles and the CMB Sky, 2000. URL <http://spud.spa.umn.edu/~pryke/logbook/20000922/>.
- Daniil Dolgich. Personal Github Repository, 2019. URL <https://github.com/daniildolgich/neutrinocmbspectrum>.
- Edward L. Wright. Cosmic Microwave Background Anisotropy, 2013. URL <http://www.astro.ucla.edu/~wright/CMB-DT.html>.
- Martin White. CMB Anisotropy, 1997. URL <http://w.astro.berkeley.edu/~mwhite/rosetta/node2.html>.
- NASA. WMAP Observations, 2012. URL https://lambda.gsfc.nasa.gov/product/map/current/pub_papers/nineyear/cosmology/wmap_9yr_cosmology_images.cfm.
- NASA. CAMB Web Interface, 2019. URL https://lambda.gsfc.nasa.gov/toolbox/tb_camb_form.cfm.
- NumPy Developers. NumPy Scientific Computing, 2019. URL <https://numpy.org/>.
- Project Jupyter. Open-source web app for code and data, 2019a. URL <https://jupyter.org/>.
- Project Jupyter. Jupyter Widgets, 2019b. URL <https://ipywidgets.readthedocs.io/en/stable/examples/Widget%20List.html>.
- Rita Tojeiro. Understanding the Cosmic Microwave Background Temperature Power Spectrum, 2006. URL https://www.roe.ac.uk/ifa/postgrad/pedagogy/2006_tojeiro.pdf.