OpenGeoProver Output for conjecture "geothm_zadatak"

Wu's method used

September 30, 2016

1 Invoking the theorem prover

The used proving method is Wu's method. The input system is:

```
= -x_4 + x_1
 p_1
          -x_4 + x_2
           -x_5 + x_1
      = x_5 + x_2 -
           x_3 –
            -x_8 + x_6
      = -x_8 + x_7
      = x_9 + x_6 -
            -x_9 + x_7
      = -x_{13}x_7 + x_{11}
p_{10}
      = x_{13} + x_{12} -
            -x_{14}x_1 + x_{10}
p_{12}
p_{13}
      = -x_{14}x_2 + x_{11}
      = -x_{14}x_3 + x_{12}
p_{14}
           -x_{18}x_7 + x_{16}
p_{15}
      = x_{18} + x_{17} -
      = -x_{19}x_1 + x_{19} + x_{15} -
           -x_{19}x_2 + x_{16}
p_{18}
      = -x_{19}x_3 + x_{17}
p_{19}
      = -x_{23}x_7 + x_{23} + x_{21} -
p_{20}
      = x_{23} + x_{22} -
p_{21}
      = -x_{24}x_1 + x_{20}
p_{22}
      = -x_{24}x_2 + x_{24} + x_{21} -
p_{23}
      = -x_{24}x_3 + x_{22}
     = -x_{28}x_7 + x_{28} + x_{26} -
```

```
\begin{array}{rcl} p_{26} & = & x_{28} + x_{27} - \\ p_{27} & = & -x_{29}x_1 + x_{29} + x_{25} - \\ p_{28} & = & -x_{29}x_2 + x_{29} + x_{26} - \\ p_{29} & = & -x_{29}x_3 + x_{27} \\ p_{30} & = & x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12} \\ p_{31} & = & x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{32} & = & x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{33} & = & x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10} \end{array}
```

1.1 Triangulation, step 1

Choosing variable: Trying the variable with index 33.

Variable x_{33} selected: The number of polynomials with this variable, with indexes from 1 to 33, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{33} . No reduction needed.

The triangular system has not been changed.

1.2 Triangulation, step 2

Choosing variable: Trying the variable with index 32.

Variable x_{32} selected: The number of polynomials with this variable, with indexes from 1 to 32, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{32} . No reduction needed.

The triangular system has not been changed.

1.3 Triangulation, step 3

Choosing variable: Trying the variable with index 31.

Variable x_{31} selected: The number of polynomials with this variable, with indexes from 1 to 31, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{31} . No reduction needed.

The triangular system has not been changed.

1.4 Triangulation, step 4

Choosing variable: Trying the variable with index 30.

Variable x_{30} selected: The number of polynomials with this variable, with indexes from 1 to 30, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{30} . No reduction needed.

The triangular system has not been changed.

1.5 Triangulation, step 5

Choosing variable: Trying the variable with index 29.

Variable x_{29} selected: The number of polynomials with this variable, with indexes from 1 to 29, is 3.

Minimal degrees: 3 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{29} from all other polynomials by reducing them with polynomial p_{27} from previous step.

```
= -x_4 + x_1
p_1
     = -x_4 + x_2
p_2
 p_3
      = -x_5 + x_1
      = x_5 + x_2 -
 p_4
      = x_3 -
 p_5
      = -x_8 + x_6
 p_6
      = -x_8 + x_7
 p_7
      = x_9 + x_6 -
      = -x_9 + x_7
      = -x_{13}x_7 + x_{11}
p_{10}
      = x_{13} + x_{12} -
      = -x_{14}x_1 + x_{10}
p_{12}
      = -x_{14}x_2 + x_{11}
p_{13}
      = -x_{14}x_3 + x_{12}
      = -x_{18}x_7 + x_{16}
      = x_{18} + x_{17} -
p_{16}
      = -x_{19}x_1 + x_{19} + x_{15} -
      = -x_{19}x_2 + x_{16}
p_{18}
      = -x_{19}x_3 + x_{17}
p_{19}
      = -x_{23}x_7 + x_{23} + x_{21} -
p_{20}
      = x_{23} + x_{22} -
p_{21}
      = -x_{24}x_1 + x_{20}
p_{22}
     = -x_{24}x_2 + x_{24} + x_{21} -
p_{23}
      = -x_{24}x_3 + x_{22}
p_{25}
      = -x_{28}x_7 + x_{28} + x_{26} -
     = x_{28} + x_{27} -
p_{26}
```

```
\begin{array}{rcl} p_{27} & = & -x_{26}x_{1} + x_{26} + x_{25}x_{2} - x_{25} - x_{2} + x_{1} \\ p_{28} & = & -x_{27}x_{1} + x_{27} + x_{25}x_{3} - x_{3} \\ p_{29} & = & -x_{29}x_{1} + x_{29} + x_{25} - \\ p_{30} & = & x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12} \\ p_{31} & = & x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{32} & = & x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{33} & = & x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10} \end{array}
```

1.6 Triangulation, step 6

Choosing variable: Trying the variable with index 28.

Variable x_{28} selected: The number of polynomials with this variable, with indexes from 1 to 28, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{28} from all other polynomials by reducing them with polynomial p_{25} from previous step.

```
= -x_4 + x_1
 p_1
 p_2
     = -x_4 + x_2
     = -x_5 + x_1
 p_3
     = x_5 + x_2 -
 p_4
      = x_3 -
      = -x_8 + x_6
 p_6
     = -x_8 + x_7
 p_7
      = x_9 + x_6 -
      = -x_9 + x_7
 p_9
      = -x_{13}x_7 + x_{11}
p_{10}
      = x_{13} + x_{12} -
p_{11}
     = -x_{14}x_1 + x_{10}
p_{12}
     = -x_{14}x_2 + x_{11}
p_{13}
      = -x_{14}x_3 + x_{12}
p_{14}
      = -x_{18}x_7 + x_{16}
p_{15}
     = x_{18} + x_{17} -
p_{16}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{17}
     = -x_{19}x_2 + x_{16}
p_{18}
     = -x_{19}x_3 + x_{17}
p_{19}
      = -x_{23}x_7 + x_{23} + x_{21} -
     = x_{23} + x_{22} -
p_{21}
     = -x_{24}x_1 + x_{20}
p_{22}
```

```
-x_{24}x_2 + x_{24} + x_{21} -
p_{23}
            -x_{24}x_3 + x_{22}
p_{24}
              -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{25}
        = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{26}
        = -x_{27}x_7 + x_{27} - x_{26} + x_7
p_{27}
              -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
        = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
             x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
       = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
       = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
```

1.7 Triangulation, step 7

Choosing variable: Trying the variable with index 27.

Variable x_{27} selected: The number of polynomials with this variable, with indexes from 1 to 27, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{27} from all other polynomials by reducing them with polynomial p_{26} from previous step.

```
= -x_4 + x_1
     = -x_4 + x_2
 p_2
     = -x_5 + x_1
     = x_5 + x_2 -
      = x_3 -
 p_5
     = -x_8 + x_6
      = -x_8 + x_7
     = x_9 + x_6 -
 p_8
     = -x_9 + x_7
      = -x_{13}x_7 + x_{11}
p_{10}
     = x_{13} + x_{12} -
p_{11}
     = -x_{14}x_1 + x_{10}
p_{12}
     = -x_{14}x_2 + x_{11}
p_{13}
     = -x_{14}x_3 + x_{12}
p_{14}
     = -x_{18}x_7 + x_{16}
p_{15}
     = x_{18} + x_{17} -
p_{17}
     = -x_{19}x_1 + x_{19} + x_{15} -
p_{18} = -x_{19}x_2 + x_{16}
```

```
p_{19}
            -x_{19}x_3 + x_{17}
            -x_{23}x_7 + x_{23} + x_{21} -
p_{20}
             x_{23} + x_{22} -
p_{21}
              -x_{24}x_1 + x_{20}
             -x_{24}x_2 + x_{24} + x_{21} -
p_{23}
             -x_{24}x_3 + x_{22}
p_{24}
       = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{25}
       = x_{26}x_1 - x_{26} + x_{25}x_7x_3 - x_{25}x_3 - x_7x_3 - x_7x_1 + x_7 + x_3
p_{26}
             -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
       = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
       = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
            x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
p_{33}
```

1.8 Triangulation, step 8

Choosing variable: Trying the variable with index 26.

Variable x_{26} selected: The number of polynomials with this variable, with indexes from 1 to 26, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{26} from all other polynomials by reducing them with polynomial p_{25} from previous step.

$$\begin{array}{rcl} p_1 & = & -x_4 + x_1 \\ p_2 & = & -x_4 + x_2 \\ p_3 & = & -x_5 + x_1 \\ p_4 & = & x_5 + x_2 - \\ p_5 & = & x_3 - \\ p_6 & = & -x_8 + x_6 \\ p_7 & = & -x_8 + x_7 \\ p_8 & = & x_9 + x_6 - \\ p_9 & = & -x_9 + x_7 \\ p_{10} & = & -x_{13}x_7 + x_{11} \\ p_{11} & = & x_{13} + x_{12} - \\ p_{12} & = & -x_{14}x_1 + x_{10} \\ p_{13} & = & -x_{14}x_2 + x_{11} \\ p_{14} & = & -x_{14}x_3 + x_{12} \end{array}$$

```
p_{15}
       = -x_{18}x_7 + x_{16}
       = x_{18} + x_{17} -
p_{16}
             -x_{19}x_1 + x_{19} + x_{15} -
p_{17}
       = -x_{19}x_2 + x_{16}
       = -x_{19}x_3 + x_{17}
p_{19}
             -x_{23}x_7 + x_{23} + x_{21} -
p_{20}
       = x_{23} + x_{22} -
p_{21}
       = -x_{24}x_1 + x_{20}
p_{22}
             -x_{24}x_2 + x_{24} + x_{21} -
p_{23}
       = -x_{24}x_3 + x_{22}
p_{24}
       = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
             x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
             -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
       = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
             -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
       = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
      = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
            x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
p_{33}
```

1.9 Triangulation, step 9

Choosing variable: Trying the variable with index 25.

Variable x_{25} selected: The number of polynomials with this variable, with indexes from 1 to 25, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{25} . No reduction needed.

The triangular system has not been changed.

1.10 Triangulation, step 10

Choosing variable: Trying the variable with index 24.

Variable x_{24} selected: The number of polynomials with this variable, with indexes from 1 to 24, is 3.

Minimal degrees: 3 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{24} from all other polynomials by reducing them with polynomial p_{22} from previous step.

Finished a triangulation step, the current system is:

```
p_1
       = -x_4 + x_1
      = -x_4 + x_2
 p_2
      = -x_5 + x_1
       = x_5 + x_2 -
       = x_3 -
 p_5
      = -x_8 + x_6
 p_6
       = -x_8 + x_7
       = x_9 + x_6 -
       = -x_9 + x_7
 p_9
       = -x_{13}x_7 + x_{11}
p_{10}
      = x_{13} + x_{12} -
p_{11}
      = -x_{14}x_1 + x_{10}
p_{12}
      = -x_{14}x_2 + x_{11}
p_{13}
      = -x_{14}x_3 + x_{12}
p_{14}
      = -x_{18}x_7 + x_{16}
p_{15}
      = x_{18} + x_{17} -
p_{16}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{17}
      = -x_{19}x_2 + x_{16}
p_{18}
      = -x_{19}x_3 + x_{17}
p_{19}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{20}
      = x_{23} + x_{22} -
p_{21}
p_{22}
      = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
      = -x_{22}x_1 + x_{20}x_3
p_{23}
      = -x_{24}x_1 + x_{20}
p_{24}
       = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
            x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
            -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
      = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
       = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
p_{29}
       = -x_{29}x_1 + x_{29} + x_{25} -
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
       = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
            x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{33}
      = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
```

1.11 Triangulation, step 11

Choosing variable: Trying the variable with index 23.

Variable x_{23} selected: The number of polynomials with this variable, with indexes from 1 to 23, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{23} from all other polynomials by reducing them with polynomial p_{20} from previous step.

```
p_1 = -x_4 + x_1
      = -x_4 + x_2
 p_2
      = -x_5 + x_1
 p_3
      = x_5 + x_2 -
 p_4
      = x_3 -
      = -x_8 + x_6
 p_6
      = -x_8 + x_7
 p_7
      = x_9 + x_6 -
      = -x_9 + x_7
p_9
      = -x_{13}x_7 + x_{11}
p_{10}
      = x_{13} + x_{12} -
p_{11}
      = -x_{14}x_1 + x_{10}
p_{12}
      =
          -x_{14}x_2 + x_{11}
p_{13}
      = -x_{14}x_3 + x_{12}
p_{14}
      = -x_{18}x_7 + x_{16}
p_{15}
      = x_{18} + x_{17} -
p_{16}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{17}
      = -x_{19}x_2 + x_{16}
p_{18}
      = -x_{19}x_3 + x_{17}
p_{19}
      = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{20}
      = -x_{22}x_1 + x_{20}x_3
p_{21}
      = -x_{22}x_7 + x_{22} - x_{21} + x_7
p_{22}
      = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
      = -x_{24}x_1 + x_{20}
p_{24}
      = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
            x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
            -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
      = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
      = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
      = -x_{28}x_7 + x_{28} + x_{26} -
      = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
      = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
```

```
\begin{array}{lll} p_{31} & = & x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{32} & = & x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{33} & = & x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10} \end{array}
```

1.12 Triangulation, step 12

Choosing variable: Trying the variable with index 22.

Variable x_{22} selected: The number of polynomials with this variable, with indexes from 1 to 22, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{22} from all other polynomials by reducing them with polynomial p_{21} from previous step.

```
p_1 = -x_4 + x_1
     = -x_4 + x_2
 p_2
      = -x_5 + x_1
     = x_5 + x_2 -
 p_4
     = x_3 -
      = -x_8 + x_6
 p_6
      = -x_8 + x_7
 p_7
     = x_9 + x_6 -
      = -x_9 + x_7
      = -x_{13}x_7 + x_{11}
p_{10}
     = x_{13} + x_{12} -
p_{11}
      = -x_{14}x_1 + x_{10}
     = -x_{14}x_2 + x_{11}
p_{13}
     = -x_{14}x_3 + x_{12}
      = -x_{18}x_7 + x_{16}
     = x_{18} + x_{17} -
p_{16}
     = -x_{19}x_1 + x_{19} + x_{15} -
      = -x_{19}x_2 + x_{16}
      = -x_{19}x_3 + x_{17}
p_{19}
      = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
      = x_{21}x_1 + x_{20}x_7x_3 - x_{20}x_3 - x_7x_1
p_{21}
p_{22}
     = -x_{22}x_1 + x_{20}x_3
     = -x_{23}x_7 + x_{23} + x_{21} - 
p_{23}
      = -x_{24}x_1 + x_{20}
p_{25}
      = \quad -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 + \\
           x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
```

$$\begin{array}{rcl} & -x_3x_1+x_3+x_2x_1-x_2-x_1^2+x_1\\ p_{26} & = & -x_{26}x_1+x_{26}+x_{25}x_2-x_{25}-x_2+x_1\\ p_{27} & = & -x_{27}x_1+x_{27}+x_{25}x_3-x_3\\ p_{28} & = & -x_{28}x_7+x_{28}+x_{26}-\\ p_{29} & = & -x_{29}x_1+x_{29}+x_{25}-\\ p_{30} & = & x_{30}-x_{22}x_{16}+x_{22}x_{11}+x_{21}x_{17}-x_{21}x_{12}-x_{17}x_{11}+x_{16}x_{12}\\ p_{31} & = & x_{31}+x_{22}x_{15}-x_{22}x_{10}-x_{20}x_{17}+x_{20}x_{12}+x_{17}x_{10}-x_{15}x_{12}\\ p_{32} & = & x_{32}-x_{21}x_{15}+x_{21}x_{10}+x_{20}x_{16}-x_{20}x_{11}-x_{16}x_{10}+x_{15}x_{11}\\ p_{33} & = & x_{33}+x_{32}x_{12}+x_{31}x_{11}+x_{30}x_{10} \end{array}$$

1.13 Triangulation, step 13

Choosing variable: Trying the variable with index 21.

Variable x_{21} selected: The number of polynomials with this variable, with indexes from 1 to 21, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{21} from all other polynomials by reducing them with polynomial p_{20} from previous step.

```
p_1
     = -x_4 + x_1
     = -x_4 + x_2
     = -x_5 + x_1
     = x_5 + x_2 -
 p_{4}
     = x_3 -
     = -x_8 + x_6
     = -x_8 + x_7
 p_7
     = x_9 + x_6 -
      = -x_9 + x_7
     = -x_{13}x_7 + x_{11}
p_{10}
     = x_{13} + x_{12} -
p_{11}
      = -x_{14}x_1 + x_{10}
p_{12}
     = -x_{14}x_2 + x_{11}
p_{13}
     = -x_{14}x_3 + x_{12}
p_{14}
      = -x_{18}x_7 + x_{16}
p_{15}
     = x_{18} + x_{17} -
p_{16}
     = -x_{19}x_1 + x_{19} + x_{15} -
     = -x_{19}x_2 + x_{16}
p_{19}
     = -x_{19}x_3 + x_{17}
    = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
```

```
p_{21}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
       = -x_{22}x_1 + x_{20}x_3
p_{22}
             -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
p_{24}
       = -x_{24}x_1 + x_{20}
       = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
             x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
             -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
       = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
            -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
       =
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
       = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
      = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
            x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
p_{33}
```

1.14 Triangulation, step 14

Choosing variable: Trying the variable with index 20.

Variable x_{20} selected: The number of polynomials with this variable, with indexes from 1 to 20, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{20} . No reduction needed.

The triangular system has not been changed.

1.15 Triangulation, step 15

Choosing variable: Trying the variable with index 19.

Variable x_{19} selected: The number of polynomials with this variable, with indexes from 1 to 19, is 3.

Minimal degrees: 3 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{19} from all other polynomials by reducing them with polynomial p_{17} from previous step.

$$p_{1} = -x_{4} + x_{1}$$

$$p_{2} = -x_{4} + x_{2}$$

$$p_{3} = -x_{5} + x_{1}$$

$$p_{4} = x_{5} + x_{2} - x_{5}$$

$$p_{5} = x_{3} - x_{5}$$

```
= -x_8 + x_6
       = -x_8 + x_7
       = x_9 + x_6 -
       = -x_9 + x_7
       = -x_{13}x_7 + x_{11}
p_{10}
            x_{13} + x_{12} -
p_{11}
       = -x_{14}x_1 + x_{10}
       = -x_{14}x_2 + x_{11}
p_{13}
       = -x_{14}x_3 + x_{12}
p_{14}
       = -x_{18}x_7 + x_{16}
       = x_{18} + x_{17} -
p_{16}
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{17}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
       = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
             -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
             x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
             -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
       = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
            -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
       = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
      = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
            x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
p_{33}
```

1.16 Triangulation, step 16

Choosing variable: Trying the variable with index 18.

Variable x_{18} selected: The number of polynomials with this variable, with indexes from 1 to 18, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{18} from all other polynomials by reducing them with polynomial p_{15} from previous step.

```
= -x_4 + x_1
 p_1
      = -x_4 + x_2
           -x_5 + x_1
 p_3
      = x_5 + x_2 -
 p_4
      = x_3 -
            -x_8 + x_6
 p_6
       = -x_8 + x_7
 p_7
       = x_9 + x_6 -
           -x_9 + x_7
 p_9
       = -x_{13}x_7 + x_{11}
p_{10}
       = x_{13} + x_{12} -
p_{11}
       = -x_{14}x_1 + x_{10}
p_{12}
      = -x_{14}x_2 + x_{11}
p_{13}
      = -x_{14}x_3 + x_{12}
p_{14}
p_{15}
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{16}
      = -x_{17}x_7 - x_{16} + x_7
p_{17}
       = -x_{18}x_7 + x_{16}
p_{18}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
      = -x_{22}x_1 + x_{20}x_3
p_{22}
      = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
      = -x_{24}x_1 + x_{20}
p_{24}
      = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
            x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
            -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
      = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
       = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
      = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
      = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
      = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
     = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
     = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
p_{33}
```

1.17 Triangulation, step 17

Choosing variable: Trying the variable with index 17.

Variable x_{17} selected: The number of polynomials with this variable, with indexes from 1 to 17, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{17} from all other polynomials by reducing them with polynomial p_{16} from previous step.

```
p_1 = -x_4 + x_1
     = -x_4 + x_2
 p_2
      = -x_5 + x_1
      = x_5 + x_2 -
 p_4
      = x_3 -
      = -x_8 + x_6
 p_6
 p_7
      = -x_8 + x_7
      = x_9 + x_6 -
      = -x_9 + x_7
p_9
      = -x_{13}x_7 + x_{11}
p_{10}
      = x_{13} + x_{12} -
p_{11}
      = -x_{14}x_1 + x_{10}
p_{12}
      = -x_{14}x_2 + x_{11}
p_{13}
      = -x_{14}x_3 + x_{12}
p_{14}
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{15}
      = x_{16}x_1 - x_{16} + x_{15}x_7x_3 - x_7x_3 - x_7x_1 + x_7
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
      = -x_{18}x_7 + x_{16}
p_{18}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
      = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
      = -x_{22}x_1 + x_{20}x_3
p_{22}
      = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
      = -x_{24}x_1 + x_{20}
p_{24}
      = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
            x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
            -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
      = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
      = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
      = -x_{28}x_7 + x_{28} + x_{26} -
      = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
      = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
```

```
\begin{array}{lll} p_{31} & = & x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{32} & = & x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{33} & = & x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10} \end{array}
```

1.18 Triangulation, step 18

Choosing variable: Trying the variable with index 16.

Variable x_{16} selected: The number of polynomials with this variable, with indexes from 1 to 16, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{16} from all other polynomials by reducing them with polynomial p_{15} from previous step.

```
p_1 = -x_4 + x_1
     = -x_4 + x_2
 p_2
      = -x_5 + x_1
     = x_5 + x_2 -
 p_4
     = x_3 -
 p_5
      = -x_8 + x_6
 p_6
      = -x_8 + x_7
 p_7
     = x_9 + x_6 -
      = -x_9 + x_7
      = -x_{13}x_7 + x_{11}
p_{10}
     = x_{13} + x_{12} -
p_{11}
      = -x_{14}x_1 + x_{10}
     = -x_{14}x_2 + x_{11}
p_{13}
     = -x_{14}x_3 + x_{12}
p_{14}
      = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
           -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
      = -x_{18}x_7 + x_{16}
p_{18}
      = -x_{19}x_1 + x_{19} + x_{15} -
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
      = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
     = -x_{22}x_1 + x_{20}x_3
p_{22}
     = -x_{23}x_7 + x_{23} + x_{21} -
     = -x_{24}x_1 + x_{20}
p_{24}
     = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
```

$$\begin{array}{rcl} & x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 \\ & -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1 \\ \\ p_{26} & = & -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1 \\ p_{27} & = & -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3 \\ \\ p_{28} & = & -x_{28}x_7 + x_{28} + x_{26} - \\ \\ p_{29} & = & -x_{29}x_1 + x_{29} + x_{25} - \\ \\ p_{30} & = & x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12} \\ \\ p_{31} & = & x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ \\ p_{32} & = & x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ \\ p_{33} & = & x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10} \end{array}$$

1.19 Triangulation, step 19

Choosing variable: Trying the variable with index 15.

Variable x_{15} **selected:** The number of polynomials with this variable, with indexes from 1 to 15, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{15} . No reduction needed.

The triangular system has not been changed.

1.20 Triangulation, step 20

Choosing variable: Trying the variable with index 14.

Variable x_{14} selected: The number of polynomials with this variable, with indexes from 1 to 14, is 3.

Minimal degrees: 3 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{14} from all other polynomials by reducing them with polynomial p_{12} from previous step.

$$p_{1} = -x_{4} + x_{1}$$

$$p_{2} = -x_{4} + x_{2}$$

$$p_{3} = -x_{5} + x_{1}$$

$$p_{4} = x_{5} + x_{2} - x_{5} + x_{2} - x_{5} + x_{6}$$

$$p_{6} = -x_{8} + x_{6}$$

$$p_{7} = -x_{8} + x_{7}$$

$$p_{8} = x_{9} + x_{6} - x_{9} + x_{7}$$

$$p_{9} = -x_{9} + x_{7}$$

$$p_{10} = -x_{13}x_{7} + x_{11}$$

```
p_{11}
       = x_{13} + x_{12} -
       = -x_{11}x_1 + x_{10}x_2
p_{12}
       = -x_{12}x_1 + x_{10}x_3
p_{13}
p_{14}
       = -x_{14}x_1 + x_{10}
       = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
             -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
       = -x_{18}x_7 + x_{16}
p_{18}
       = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
      = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
            x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
             -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
       = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
       = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
      = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
       = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
      = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
      = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
```

1.21 Triangulation, step 21

Choosing variable: Trying the variable with index 13.

Variable x_{13} selected: The number of polynomials with this variable, with indexes from 1 to 13, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{13} from all other polynomials by reducing them with polynomial p_{10} from previous step.

$$p_1 = -x_4 + x_1$$

$$p_2 = -x_4 + x_2$$

$$p_3 = -x_5 + x_1$$

```
p_4
      = x_5 + x_2 -
       = x_3 -
 p_5
       = -x_8 + x_6
       = -x_8 + x_7
       = x_9 + x_6 -
 p_8
            -x_9 + x_7
 p_9
       = -x_{11}x_1 + x_{10}x_2
p_{10}
       = -x_{12}x_1 + x_{10}x_3
p_{11}
       = -x_{12}x_7 - x_{11} + x_7
p_{12}
       = -x_{13}x_7 + x_{11}
p_{13}
       = -x_{14}x_1 + x_{10}
p_{14}
       = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
             -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
       = -x_{18}x_7 + x_{16}
p_{18}
       = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
            -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
             x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
             -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
       = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
       = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
       = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
      = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
       = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
p_{33}
```

1.22 Triangulation, step 22

Choosing variable: Trying the variable with index 12.

Variable x_{12} selected: The number of polynomials with this variable, with indexes from 1 to 12, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{12} from all other polynomials by reducing them with polynomial p_{11} from previous step.

```
= -x_4 + x_1
 p_1
      = -x_4 + x_2
 p_2
      = -x_5 + x_1
      = x_5 + x_2 -
 p_4
      = x_3 -
 p_5
      = -x_8 + x_6
      = -x_8 + x_7
      = x_9 + x_6 -
 p_8
      = -x_9 + x_7
      = -x_{11}x_1 + x_{10}x_2
p_{10}
      = x_{11}x_1 + x_{10}x_7x_3 - x_7x_1
p_{11}
     = -x_{12}x_1 + x_{10}x_3
p_{12}
      = -x_{13}x_7 + x_{11}
p_{13}
      = -x_{14}x_1 + x_{10}
p_{14}
      = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
            -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
      = -x_{18}x_7 + x_{16}
p_{18}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
      = -x_{22}x_1 + x_{20}x_3
p_{22}
      = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
            x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
            -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
      = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
       = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
      = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
       = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
      = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
     = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
p_{33}
```

1.23 Triangulation, step 23

Choosing variable: Trying the variable with index 11.

Variable x_{11} selected: The number of polynomials with this variable, with indexes from 1 to 11, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{11} from all other polynomials by reducing them with polynomial p_{10} from previous step.

```
p_1 = -x_4 + x_1
     = -x_4 + x_2
     = -x_5 + x_1
      = x_5 + x_2 -
 p_4
      = x_3 -
 p_5
     = -x_8 + x_6
 p_6
      = -x_8 + x_7
      = x_9 + x_6 -
 p_8
      = -x_9 + x_7
      = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
p_{10}
      = -x_{11}x_1 + x_{10}x_2
p_{11}
     = -x_{12}x_1 + x_{10}x_3
p_{12}
      = -x_{13}x_7 + x_{11}
     = -x_{14}x_1 + x_{10}
p_{14}
      = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
           -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
      = -x_{18}x_7 + x_{16}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
      = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
      = -x_{22}x_1 + x_{20}x_3
p22
     = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
      = -x_{24}x_1 + x_{20}
p_{24}
      = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
           x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
           -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
     = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
     = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
```

```
\begin{array}{rcl} p_{28} & = & -x_{28}x_7 + x_{28} + x_{26} - \\ p_{29} & = & -x_{29}x_1 + x_{29} + x_{25} - \\ p_{30} & = & x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12} \\ p_{31} & = & x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{32} & = & x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{33} & = & x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10} \end{array}
```

1.24 Triangulation, step 24

Choosing variable: Trying the variable with index 10.

Variable x_{10} **selected:** The number of polynomials with this variable, with indexes from 1 to 10, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{10} . No reduction needed.

The triangular system has not been changed.

1.25 Triangulation, step 25

Choosing variable: Trying the variable with index 9.

Variable x_9 selected: The number of polynomials with this variable, with indexes from 1 to 9, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_9 from all other polynomials by reducing them with polynomial p_8 from previous step.

```
p_1 = -x_4 + x_1
p_2
     = -x_4 + x_2
     = -x_5 + x_1
p_3
     = x_5 + x_2 -
     = x_3 -
     = -x_8 + x_6
p_6
     = -x_8 + x_7
p_7
     = x_7 + x_6 -
     = x_9 + x_6 -
     = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
     = -x_{11}x_1 + x_{10}x_2
p_{11}
    = -x_{12}x_1 + x_{10}x_3
p_{12}
     = -x_{13}x_7 + x_{11}
     = -x_{14}x_1 + x_{10}
p_{14}
```

$$\begin{array}{rclcrcl} p_{15} & = & -x_{15}x_{7}x_{3}x_{1} + x_{15}x_{7}x_{3} - x_{15}x_{2}x_{1} + x_{15}x_{2} + x_{7}x_{3}x_{1} \\ & -x_{7}x_{3} + x_{7}x_{1}^{2} - 2x_{7}x_{1} + x_{7} + x_{2}x_{1} - x_{2} \\ \end{array} \\ p_{16} & = & -x_{16}x_{1} + x_{16} + x_{15}x_{2} - x_{2} \\ p_{17} & = & -x_{17}x_{1} + x_{17} + x_{15}x_{3} - x_{3} \\ p_{18} & = & -x_{18}x_{7} + x_{16} \\ p_{19} & = & -x_{19}x_{1} + x_{19} + x_{15} - \\ p_{20} & = & -x_{20}x_{7}x_{3}x_{1} + x_{20}x_{3}x_{1} - x_{20}x_{2}x_{1} + x_{20}x_{1} + x_{7}x_{1}^{2} - x_{1}^{2} \\ p_{21} & = & -x_{21}x_{1} + x_{20}x_{2} - x_{20} + x_{1} \\ p_{22} & = & -x_{22}x_{1} + x_{20}x_{3} \\ p_{23} & = & -x_{23}x_{7} + x_{23} + x_{21} - \\ p_{24} & = & -x_{24}x_{1} + x_{20} \\ p_{25} & = & -x_{25}x_{7}x_{3}x_{1} + x_{25}x_{7}x_{3} + x_{25}x_{3}x_{1} - x_{25}x_{3} - x_{25}x_{2}x_{1} + \\ & & x_{25}x_{2} + x_{25}x_{1} - x_{25} + x_{7}x_{3}x_{1} - x_{7}x_{3} + x_{7}x_{1}^{2} - 2x_{7}x_{1} + x_{7} \\ & & -x_{3}x_{1} + x_{3} + x_{2}x_{1} - x_{2} - x_{1}^{2} + x_{1} \\ p_{26} & = & -x_{26}x_{1} + x_{26} + x_{25}x_{2} - x_{25} - x_{2} + x_{1} \\ p_{27} & = & -x_{27}x_{1} + x_{27} + x_{25}x_{3} - x_{3} \\ p_{28} & = & -x_{28}x_{7} + x_{28} + x_{26} - \\ p_{29} & = & -x_{29}x_{1} + x_{29} + x_{25} - \\ p_{30} & = & x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12} \\ p_{31} & = & x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{11} \\ p_{32} & = & x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{33} & = & x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10} \\ \end{array}$$

1.26 Triangulation, step 26

Choosing variable: Trying the variable with index 8.

Variable x_8 selected: The number of polynomials with this variable, with indexes from 1 to 8, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_8 from all other polynomials by reducing them with polynomial p_6 from previous step.

$$p_{1} = -x_{4} + x_{1}$$

$$p_{2} = -x_{4} + x_{2}$$

$$p_{3} = -x_{5} + x_{1}$$

$$p_{4} = x_{5} + x_{2} - x_{5}$$

$$p_{5} = x_{3} - x_{6}$$

$$p_{6} = x_{7} + x_{6} - x_{7}$$

$$p_{7} = -x_{7} + x_{6}$$

```
= -x_8 + x_6
       = x_9 + x_6 -
            -x_{10}x_7x_3x_1-x_{10}x_2x_1+x_7x_1^2
p_{10}
       = -x_{11}x_1 + x_{10}x_2
       = -x_{12}x_1 + x_{10}x_3
p_{12}
            -x_{13}x_7 + x_{11}
p_{13}
       = -x_{14}x_1 + x_{10}
p_{14}
       = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
             -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
       = -x_{18}x_7 + x_{16}
p_{18}
       = -x_{19}x_1 + x_{19} + x_{15} -
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
            -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
            -x_{24}x_1 + x_{20}
p_{24}
       = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
            x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
             -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
       = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
       = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
            x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
      = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
      = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
```

1.27 Triangulation, step 27

Choosing variable: Trying the variable with index 7.

Variable x_7 selected: The number of polynomials with this variable, with indexes from 1 to 7, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_7 from all other polynomials by reducing them with polynomial p_6 from previous step.

```
p_1 = -x_4 + x_1
      = -x_4 + x_2
      = -x_5 + x_1
      = x_5 + x_2 -
 p_5
      = x_3 -
      = 2x_6 -
 p_6
      = x_7 + x_6 -
 p_7
      = -x_8 + x_6
      = x_9 + x_6 -
      = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
      = -x_{11}x_1 + x_{10}x_2
p_{11}
p_{12} = -x_{12}x_1 + x_{10}x_3
      = -x_{13}x_7 + x_{11}
p_{13}
      = -x_{14}x_1 + x_{10}
p_{14}
      = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
            -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
      = -x_{18}x_7 + x_{16}
p_{18}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
      = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
      = -x_{22}x_1 + x_{20}x_3
p_{22}
     = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
      = -x_{24}x_1 + x_{20}
p_{24}
      = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
            x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
            -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
      = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
      = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
      = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
      = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
      = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
      = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
     = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
p_{33} = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
```

1.28 Triangulation, step 28

Choosing variable: Trying the variable with index 6.

Variable x_6 **selected:** The number of polynomials with this variable, with indexes from 1 to 6, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_6 . No reduction needed.

The triangular system has not been changed.

1.29 Triangulation, step 29

Choosing variable: Trying the variable with index 5.

Variable x_5 selected: The number of polynomials with this variable, with indexes from 1 to 5, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_5 from all other polynomials by reducing them with polynomial p_3 from previous step.

```
= -x_4 + x_1
p_1
 p_2
     = -x_4 + x_2
     = x_3 -
 p_3
     = -x_2 - x_1 + 1
 p_4
      = -x_5 + x_1
      = 2x_6 -
 p_6
     = x_7 + x_6 -
 p_7
      = -x_8 + x_6
      = x_9 + x_6 -
      = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
      = -x_{11}x_1 + x_{10}x_2
     = -x_{12}x_1 + x_{10}x_3
p_{12}
     = -x_{13}x_7 + x_{11}
p_{13}
      = -x_{14}x_1 + x_{10}
      = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
           -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
      = -x_{18}x_7 + x_{16}
      = -x_{19}x_1 + x_{19} + x_{15} -
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
     = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
     = -x_{22}x_1 + x_{20}x_3
```

```
= -x_{23}x_7 + x_{23} + x_{21} -
       = -x_{24}x_1 + x_{20}
p_{24}
       = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
             x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
             -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
       = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
       = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
       = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
       = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
      = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
     = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
p_{33}
```

1.30 Triangulation, step 30

Choosing variable: Trying the variable with index 4.

Variable x_4 selected: The number of polynomials with this variable, with indexes from 1 to 4, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_4 from all other polynomials by reducing them with polynomial p_1 from previous step.

```
p_1 = x_3 -
 p_2 = -x_2 - x_1 + 1
     = -x_2 + x_1
 p_3
     = -x_4 + x_1
 p_4
     = -x_5 + x_1
 p_5
     = 2x_6 -
 p_6
     = x_7 + x_6 -
 p_7
     = -x_8 + x_6
     = x_9 + x_6 -
 p_9
     = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
p_{10}
     = -x_{11}x_1 + x_{10}x_2
p_{11}
     = -x_{12}x_1 + x_{10}x_3
p_{12}
    = -x_{13}x_7 + x_{11}
p_{13}
     = -x_{14}x_1 + x_{10}
p_{14}
p_{15}
     = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
           -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
```

```
= -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
             -x_{18}x_7 + x_{16}
p_{18}
            -x_{19}x_1 + x_{19} + x_{15} -
            -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
             -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
             -x_{24}x_1 + x_{20}
p_{24}
       = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
             x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
             -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
       = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
       = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
             -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
       = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
       = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
             x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
            x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
      = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
```

1.31 Triangulation, step 31

Choosing variable: Trying the variable with index 3.

Variable x_3 selected: The number of polynomials with this variable, with indexes from 1 to 3, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_1 . No reduction needed.

The triangular system has not been changed.

1.32 Triangulation, step 32

Choosing variable: Trying the variable with index 2.

Variable x_2 selected: The number of polynomials with this variable, with indexes from 1 to 2, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_2 from all other polynomials by reducing them with polynomial p_1 from previous step.

```
p_1 = -2x_1 + 1
     = -x_2 - x_1 + 1
     = x_3 -
 p_3
      = -x_4 + x_1
      = -x_5 + x_1
 p_5
      = 2x_6 -
 p_6
      = x_7 + x_6 -
 p_7
      = -x_8 + x_6
      = x_9 + x_6 -
      = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
      = -x_{11}x_1 + x_{10}x_2
p_{11}
p_{12} = -x_{12}x_1 + x_{10}x_3
     = -x_{13}x_7 + x_{11}
p_{13}
      = -x_{14}x_1 + x_{10}
p_{14}
      = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
            -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
      = -x_{18}x_7 + x_{16}
p_{18}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
      = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
      = -x_{22}x_1 + x_{20}x_3
p_{22}
     = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
      = -x_{24}x_1 + x_{20}
p_{24}
      = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
            x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
            -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
      = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
      = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
p_{27}
      = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
      = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
      = x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{30}
      = x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{31}
     = x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{32}
p_{33} = x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10}
```

1.33 Triangulation, step 33

Choosing variable: Trying the variable with index 1.

Variable x_1 selected: The number of polynomials with this variable, with indexes from 1 to 1, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_1 . No reduction needed.

The triangular system has not been changed.

The triangular system is:

```
p_1 = -2x_1 + 1
     = -x_2 - x_1 + 1
      = x_3 -
      = -x_4 + x_1
 p_4
      = -x_5 + x_1
      = 2x_6 -
      = x_7 + x_6 -
 p_7
      = -x_8 + x_6
      = x_9 + x_6 -
      = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
p_{10}
      = -x_{11}x_1 + x_{10}x_2
p_{11}
      = -x_{12}x_1 + x_{10}x_3
p_{12}
      = -x_{13}x_7 + x_{11}
p_{13}
      = -x_{14}x_1 + x_{10}
p_{14}
      = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
            -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
      = -x_{18}x_7 + x_{16}
p_{18}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
      = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
      = -x_{22}x_1 + x_{20}x_3
p_{22}
      = -x_{23}x_7 + x_{23} + x_{21} -
      = -x_{24}x_1 + x_{20}
p_{24}
      = -x_{25}x_7x_3x_1 + x_{25}x_7x_3 + x_{25}x_3x_1 - x_{25}x_3 - x_{25}x_2x_1 +
p_{25}
           x_{25}x_2 + x_{25}x_1 - x_{25} + x_7x_3x_1 - x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7
            -x_3x_1 + x_3 + x_2x_1 - x_2 - x_1^2 + x_1
      = -x_{26}x_1 + x_{26} + x_{25}x_2 - x_{25} - x_2 + x_1
p_{26}
      = -x_{27}x_1 + x_{27} + x_{25}x_3 - x_3
      = -x_{28}x_7 + x_{28} + x_{26} -
p_{28}
      = -x_{29}x_1 + x_{29} + x_{25} -
p_{29}
```

$$\begin{array}{rcl} p_{30} & = & x_{30} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12} \\ p_{31} & = & x_{31} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{32} & = & x_{32} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{33} & = & x_{33} + x_{32}x_{12} + x_{31}x_{11} + x_{30}x_{10} \end{array}$$

2 Final Remainder

2.1 Final remainder for conjecture geothm_zadatak

Calculating final remainder of the conclusion:

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

with respect to the triangular system.

1. Pseudo remainder with p_{33} over variable x_{33} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

2. Pseudo remainder with p_{32} over variable x_{32} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

3. Pseudo remainder with p_{31} over variable x_{31} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

4. Pseudo remainder with p_{30} over variable x_{30} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

5. Pseudo remainder with p_{29} over variable x_{29} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

6. Pseudo remainder with p_{28} over variable x_{28} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

7. Pseudo remainder with p_{27} over variable x_{27} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

8. Pseudo remainder with p_{26} over variable x_{26} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

9. Pseudo remainder with p_{25} over variable x_{25} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

10. Pseudo remainder with p_{24} over variable x_{24} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

11. Pseudo remainder with p_{23} over variable x_{23} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

12. Pseudo remainder with p_{22} over variable x_{22} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

13. Pseudo remainder with p_{21} over variable x_{21} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

14. Pseudo remainder with p_{20} over variable x_{20} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

15. Pseudo remainder with p_{19} over variable x_{19} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

16. Pseudo remainder with p_{18} over variable x_{18} :

$$g = x_{16}x_6 - x_{16} - x_{15}x_7 + x_7$$

17. Pseudo remainder with p_{17} over variable x_{17} :

$$g = x_{16}x_6 - x_{16} - x_{15}x_7 + x_7$$

18. Pseudo remainder with p_{16} over variable x_{16} :

$$g = x_{15}x_7x_1 - x_{15}x_7 - x_{15}x_6x_2 + x_{15}x_2 - x_7x_1 + x_7 + x_6x_2 - x_2$$

19. Pseudo remainder with p_{15} over variable x_{15} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

20. Pseudo remainder with p_{14} over variable x_{14} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

21. Pseudo remainder with p_{13} over variable x_{13} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

22. Pseudo remainder with p_{12} over variable x_{12} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

23. Pseudo remainder with p_{11} over variable x_{11} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

24. Pseudo remainder with p_{10} over variable x_{10} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

25. Pseudo remainder with p_9 over variable x_9 :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

26. Pseudo remainder with p_8 over variable x_8 :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

27. Pseudo remainder with p_7 over variable x_7 :

$$g = -x_6^2 x_2 x_1^2 + 2x_6^2 x_2 x_1 - x_6^2 x_2 - x_6^2 x_1^3 + 3x_6^2 x_1^2 - 3x_6^2 x_1 + x_6^2 + 2x_6 x_2 x_1^2 - 4x_6 x_2 x_1 + 2x_6 x_2 + 2x_6 x_1^3 - 6x_6 x_1^2 + 6x_6 x_1 - 2x_6 - x_2 x_1^2 + 2x_2 x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

28. Pseudo remainder with p_6 over variable x_6 :

$$g = -x_2x_1^2 + 2x_2x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

29. Pseudo remainder with p_5 over variable x_5 :

$$g = -x_2x_1^2 + 2x_2x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

30. Pseudo remainder with p_4 over variable x_4 :

$$g = -x_2x_1^2 + 2x_2x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

31. Pseudo remainder with p_3 over variable x_3 :

$$g = -x_2x_1^2 + 2x_2x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

32. Pseudo remainder with p_2 over variable x_2 :

$$g = 0$$

33. Pseudo remainder with p_1 over variable x_1 :

$$g = 0$$

3 Prover results

Status: Theorem has been proved.

Space Complexity: The biggest polynomial obtained during prover execution contains 21 terms.

Time Complexity: Time spent by the prover is 0.171 seconds.

4 NDG Conditions

NDG Conditions in readable form

• Failed to translate NDG Conditions to readable form