OpenGeoProver Output for conjecture "geothm_zadatak"

Wu's method used

September 30, 2016

1 Invoking the theorem prover

The used proving method is Wu's method. The input system is:

```
= -x_4 + x_1
 p_1
      = -x_4 + x_2
      = -x_5 + x_1
      = x_5 + x_2 -
 p_4
      = x_3 -
            -x_8 + x_6
      = -x_8 + x_7
      = x_9 + x_6 -
           -x_9 + x_7
      = -x_{13}x_7 + x_{11}
p_{10}
      = x_{13} + x_{12} -
           -x_{14}x_1 + x_{10}
p_{12}
p_{13}
     = -x_{14}x_2 + x_{11}
      = -x_{14}x_3 + x_{12}
p_{14}
      = -x_{18}x_7 + x_{16}
p_{15}
      = x_{18} + x_{17} -
p_{16}
      = -x_{19}x_1 + x_{19} + x_{15} -
      = -x_{19}x_2 + x_{16}
p_{18}
      = -x_{19}x_3 + x_{17}
p_{19}
      = -x_{23}x_7 + x_{23} + x_{21} -
p_{20}
      = x_{23} + x_{22} -
p_{21}
     = -x_{24}x_1 + x_{20}
p_{22}
     = -x_{24}x_2 + x_{24} + x_{21} -
p_{23}
     = -x_{24}x_3 + x_{22}
p_{25} = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
```

```
\begin{array}{llll} p_{26} & = & x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{27} & = & x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{28} & = & x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10} \\ p_{29} & = & -x_{32}x_{25} + x_{29} - x_{1} \\ p_{30} & = & -x_{32}x_{26} + x_{30} - x_{2} \\ p_{31} & = & -x_{32}x_{27} + x_{31} - x_{3} \\ p_{32} & = & x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28} \end{array}
```

1.1 Triangulation, step 1

Choosing variable: Trying the variable with index 32.

Variable x_{32} selected: The number of polynomials with this variable, with indexes from 1 to 32, is 3.

Minimal degrees: 3 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{32} from all other polynomials by reducing them with polynomial p_{29} from previous step.

```
= -x_4 + x_1
p_1
 p_2
     = -x_4 + x_2
     = -x_5 + x_1
 p_3
     = x_5 + x_2 -
 p_4
     = x_3 -
     = -x_8 + x_6
 p_6
     = -x_8 + x_7
 p_7
      = x_9 + x_6 -
      = -x_9 + x_7
p_9
     = -x_{13}x_7 + x_{11}
p_{10}
      = x_{13} + x_{12} -
p_{11}
     = -x_{14}x_1 + x_{10}
p_{12}
     = -x_{14}x_2 + x_{11}
p_{13}
     = -x_{14}x_3 + x_{12}
p_{14}
     = -x_{18}x_7 + x_{16}
p_{15}
     = x_{18} + x_{17} -
p_{16}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{17}
     = -x_{19}x_2 + x_{16}
p_{18}
     = -x_{19}x_3 + x_{17}
p_{19}
     = -x_{23}x_7 + x_{23} + x_{21} -
     = x_{23} + x_{22} -
p_{21}
     = -x_{24}x_1 + x_{20}
p_{22}
```

```
p_{23}
            -x_{24}x_2 + x_{24} + x_{21} -
              -x_{24}x_3 + x_{22}
p_{24}
              x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
              x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
              x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
              x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
        = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{29}
        = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
              -x_{31}x_{25} + x_{29}x_{27} - x_{27}x_1 + x_{25}x_3
p_{31}
        = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.2 Triangulation, step 2

Choosing variable: Trying the variable with index 31.

Variable x_{31} selected: The number of polynomials with this variable, with indexes from 1 to 31, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{31} from all other polynomials by reducing them with polynomial p_{29} from previous step.

```
= -x_4 + x_1
p_1
     = -x_4 + x_2
     = -x_5 + x_1
 p_3
     = x_5 + x_2 -
 p_4
      = x_3 -
      = -x_8 + x_6
 p_6
     = -x_8 + x_7
 p_7
      = x_9 + x_6 -
      = -x_9 + x_7
p_9
     = -x_{13}x_7 + x_{11}
p_{10}
      = x_{13} + x_{12} -
p_{11}
     = -x_{14}x_1 + x_{10}
p_{12}
     = -x_{14}x_2 + x_{11}
p_{13}
      = -x_{14}x_3 + x_{12}
p_{14}
     = -x_{18}x_7 + x_{16}
p_{15}
     = x_{18} + x_{17} -
p_{16}
     = -x_{19}x_1 + x_{19} + x_{15} -
p_{18}
     = -x_{19}x_2 + x_{16}
p_{19} = -x_{19}x_3 + x_{17}
```

```
p_{20}
            -x_{23}x_7 + x_{23} + x_{21} -
        = x_{23} + x_{22} -
p_{21}
              -x_{24}x_1 + x_{20}
p_{22}
              -x_{24}x_2 + x_{24} + x_{21} -
p_{23}
            -x_{24}x_3 + x_{22}
p_{24}
              x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
        = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
        = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
        = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
        = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{29}
        = x_{30}x_{26}x_{25} + x_{29}x_{27}^2 + x_{29}x_{25}^2 + x_{28}x_{25} - x_{27}^2x_1 +
p_{30}
              x_{27}x_{25}x_3
        = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
       = -x_{32}x_{25} + x_{29} - x_1
```

1.3 Triangulation, step 3

Choosing variable: Trying the variable with index 30.

Variable x_{30} selected: The number of polynomials with this variable, with indexes from 1 to 30, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{30} from all other polynomials by reducing them with polynomial p_{29} from previous step.

$$\begin{array}{rcl} p_1 & = & -x_4 + x_1 \\ p_2 & = & -x_4 + x_2 \\ p_3 & = & -x_5 + x_1 \\ p_4 & = & x_5 + x_2 - \\ p_5 & = & x_3 - \\ p_6 & = & -x_8 + x_6 \\ p_7 & = & -x_8 + x_7 \\ p_8 & = & x_9 + x_6 - \\ p_9 & = & -x_9 + x_7 \\ p_{10} & = & -x_{13}x_7 + x_{11} \\ p_{11} & = & x_{13} + x_{12} - \\ p_{12} & = & -x_{14}x_1 + x_{10} \\ p_{13} & = & -x_{14}x_2 + x_{11} \\ p_{14} & = & -x_{14}x_3 + x_{12} \\ p_{15} & = & -x_{18}x_7 + x_{16} \end{array}$$

```
= x_{18} + x_{17} -
        = -x_{19}x_1 + x_{19} + x_{15} -
p_{17}
              -x_{19}x_2 + x_{16}
p_{18}
              -x_{19}x_3 + x_{17}
             -x_{23}x_7 + x_{23} + x_{21} -
p_{20}
              x_{23} + x_{22} -
p_{21}
             -x_{24}x_1 + x_{20}
p_{22}
        = -x_{24}x_2 + x_{24} + x_{21} -
p_{23}
        =
             -x_{24}x_3 + x_{22}
p_{24}
        = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
        = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
        = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
        = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
        = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
              x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
        = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
        = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
              -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.4 Triangulation, step 4

Choosing variable: Trying the variable with index 29.

Variable x_{29} selected: The number of polynomials with this variable, with indexes from 1 to 29, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{29} . No reduction needed.

The triangular system has not been changed.

1.5 Triangulation, step 5

Choosing variable: Trying the variable with index 28.

Variable x_{28} selected: The number of polynomials with this variable, with indexes from 1 to 28, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{28} . No reduction needed.

The triangular system has not been changed.

1.6 Triangulation, step 6

Choosing variable: Trying the variable with index 27.

Variable x_{27} selected: The number of polynomials with this variable, with indexes from 1 to 27, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{27} . No reduction needed.

The triangular system has not been changed.

1.7 Triangulation, step 7

Choosing variable: Trying the variable with index 26.

Variable x_{26} selected: The number of polynomials with this variable, with indexes from 1 to 26, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{26} . No reduction needed.

The triangular system has not been changed.

1.8 Triangulation, step 8

Choosing variable: Trying the variable with index 25.

Variable x_{25} selected: The number of polynomials with this variable, with indexes from 1 to 25, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{25} . No reduction needed.

The triangular system has not been changed.

1.9 Triangulation, step 9

Choosing variable: Trying the variable with index 24.

Variable x_{24} selected: The number of polynomials with this variable, with indexes from 1 to 24, is 3.

Minimal degrees: 3 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{24} from all other polynomials by reducing them with polynomial p_{22} from previous step.

Finished a triangulation step, the current system is:

 $p_1 = -x_4 + x_1$

 $p_2 = -x_4 + x_2$

 $p_3 = -x_5 + x_1$

 $p_4 = x_5 + x_2 -$

 $p_5 = x_3 -$

 $p_6 = -x_8 + x_6$

 $p_7 = -x_8 + x_7$

 $p_8 = x_9 + x_6 -$

```
-x_9 + x_7
            -x_{13}x_7 + x_{11}
p_{10}
              x_{13} + x_{12} -
p_{11}
              -x_{14}x_1 + x_{10}
              -x_{14}x_2 + x_{11}
p_{13}
              -x_{14}x_3 + x_{12}
p_{14}
        = -x_{18}x_7 + x_{16}
p_{15}
        = x_{18} + x_{17} -
p_{16}
              -x_{19}x_1 + x_{19} + x_{15} -
p_{17}
        = -x_{19}x_2 + x_{16}
        = -x_{19}x_3 + x_{17}
p_{19}
              -x_{23}x_7 + x_{23} + x_{21} -
p_{20}
        = x_{23} + x_{22} -
p_{21}
        = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{22}
        = -x_{22}x_1 + x_{20}x_3
p_{23}
        = -x_{24}x_1 + x_{20}
p_{24}
        = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
        = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
        = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
        = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
        = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
              x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
       = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
       = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.10 Triangulation, step 10

Choosing variable: Trying the variable with index 23.

Variable x_{23} selected: The number of polynomials with this variable, with indexes from 1 to 23, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{23} from all other polynomials by reducing them with polynomial p_{20} from previous step.

$$p_1 = -x_4 + x_1$$

$$p_2 = -x_4 + x_2$$

$$p_3 = -x_5 + x_1$$

$$p_4 = x_5 + x_2 - x_3$$

```
p_5
       = x_3 -
            -x_8 + x_6
 p_6
             -x_8 + x_7
 p_7
       = x_9 + x_6 -
       = -x_9 + x_7
 p_9
             -x_{13}x_7 + x_{11}
p_{10}
       = x_{13} + x_{12} -
            -x_{14}x_1 + x_{10}
p_{12}
             -x_{14}x_2 + x_{11}
p_{13}
       = -x_{14}x_3 + x_{12}
       = -x_{18}x_7 + x_{16}
p_{15}
       =
             x_{18} + x_{17} -
p_{16}
             -x_{19}x_1 + x_{19} + x_{15} -
            -x_{19}x_2 + x_{16}
p_{18}
              -x_{19}x_3 + x_{17}
p_{19}
            -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{20}
            -x_{22}x_1 + x_{20}x_3
p_{21}
              -x_{22}x_7 + x_{22} - x_{21} + x_7
p_{22}
            -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
             x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
       = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
       = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
             x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
       = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
      = -x_{32}x_{25} + x_{29} - x_1
```

1.11 Triangulation, step 11

Choosing variable: Trying the variable with index 22.

Variable x_{22} selected: The number of polynomials with this variable, with indexes from 1 to 22, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{22} from all other polynomials by reducing them with polynomial p_{21} from previous step.

```
-x_4 + x_1
 p_1
             -x_4 + x_2
              -x_5 + x_1
 p_3
             x_5 + x_2 -
 p_4
             x_3 -
              -x_8 + x_6
 p_6
            -x_8 + x_7
 p_7
        = x_9 + x_6 -
              -x_9 + x_7
 p_9
        = -x_{13}x_7 + x_{11}
p_{10}
        = x_{13} + x_{12} -
p_{11}
              -x_{14}x_1 + x_{10}
p_{12}
       = -x_{14}x_2 + x_{11}
p_{13}
        = -x_{14}x_3 + x_{12}
p_{14}
p_{15}
              -x_{18}x_7 + x_{16}
        = x_{18} + x_{17} -
p_{16}
             -x_{19}x_1 + x_{19} + x_{15} -
p_{17}
              -x_{19}x_2 + x_{16}
p_{18}
             -x_{19}x_3 + x_{17}
p_{19}
             -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{20}
              x_{21}x_1 + x_{20}x_7x_3 - x_{20}x_3 - x_7x_1
p_{21}
             -x_{22}x_1 + x_{20}x_3
p_{22}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
              -x_{24}x_1 + x_{20}
p_{24}
        = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
        = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
        = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
        = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
              x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
       = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
             -x_{32}x_{25} + x_{29} - x_1
```

1.12 Triangulation, step 12

Choosing variable: Trying the variable with index 21.

Variable x_{21} selected: The number of polynomials with this variable, with indexes from 1 to 21, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{21} from all other polynomials by reducing them with polynomial p_{20} from previous step.

```
p_1
       = -x_4 + x_1
 p_2
       = -x_4 + x_2
       = -x_5 + x_1
 p_3
       = x_5 + x_2 -
       = x_3 -
 p_5
       = -x_8 + x_6
 p_6
       = -x_8 + x_7
       = x_9 + x_6 -
 p_8
       = -x_9 + x_7
 p_9
       = -x_{13}x_7 + x_{11}
p_{10}
       = x_{13} + x_{12} -
p_{11}
       = -x_{14}x_1 + x_{10}
p_{12}
       = -x_{14}x_2 + x_{11}
       = -x_{14}x_3 + x_{12}
p_{14}
       = -x_{18}x_7 + x_{16}
p_{15}
       = x_{18} + x_{17} -
p_{16}
             -x_{19}x_1 + x_{19} + x_{15} -
p_{17}
       = -x_{19}x_2 + x_{16}
p_{18}
       = -x_{19}x_3 + x_{17}
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
p_{22}
       = -x_{22}x_1 + x_{20}x_3
p_{23}
       = -x_{23}x_7 + x_{23} + x_{21} -
       = -x_{24}x_1 + x_{20}
p_{24}
p_{25}
       = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
       = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
       = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
             x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
       = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
      = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.13 Triangulation, step 13

Choosing variable: Trying the variable with index 20.

Variable x_{20} selected: The number of polynomials with this variable, with indexes from 1 to 20, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{20} . No reduction needed.

The triangular system has not been changed.

1.14 Triangulation, step 14

Choosing variable: Trying the variable with index 19.

Variable x_{19} selected: The number of polynomials with this variable, with indexes from 1 to 19, is 3.

Minimal degrees: 3 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{19} from all other polynomials by reducing them with polynomial p_{17} from previous step.

```
p_1
     = -x_4 + x_1
     = -x_4 + x_2
 p_2
     = -x_5 + x_1
      = x_5 + x_2 -
 p_5
     = x_3 -
     = -x_8 + x_6
 p_6
      = -x_8 + x_7
     = x_9 + x_6 -
 p_8
     = -x_9 + x_7
      = -x_{13}x_7 + x_{11}
p_{10}
     = x_{13} + x_{12} -
p_{11}
     = -x_{14}x_1 + x_{10}
p_{12}
      = -x_{14}x_2 + x_{11}
p_{13}
     = -x_{14}x_3 + x_{12}
p_{14}
     = -x_{18}x_7 + x_{16}
p_{15}
      = x_{18} + x_{17} -
p_{16}
     = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{17}
     = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{18}
     = -x_{19}x_1 + x_{19} + x_{15} -
     = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
     = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
```

```
-x_{22}x_1 + x_{20}x_3
            -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
              -x_{24}x_1 + x_{20}
p_{24}
        = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
        = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
        = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
        = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
              x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
        = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
        = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
        = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.15 Triangulation, step 15

Choosing variable: Trying the variable with index 18.

Variable x_{18} selected: The number of polynomials with this variable, with indexes from 1 to 18, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{18} from all other polynomials by reducing them with polynomial p_{15} from previous step.

$$\begin{array}{rcl} p_1 & = & -x_4 + x_1 \\ p_2 & = & -x_4 + x_2 \\ p_3 & = & -x_5 + x_1 \\ p_4 & = & x_5 + x_2 - \\ p_5 & = & x_3 - \\ p_6 & = & -x_8 + x_6 \\ p_7 & = & -x_8 + x_7 \\ p_8 & = & x_9 + x_6 - \\ p_9 & = & -x_9 + x_7 \\ p_{10} & = & -x_{13}x_7 + x_{11} \\ p_{11} & = & x_{13} + x_{12} - \\ p_{12} & = & -x_{14}x_1 + x_{10} \\ p_{13} & = & -x_{14}x_2 + x_{11} \\ p_{14} & = & -x_{14}x_3 + x_{12} \\ p_{15} & = & -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2 \\ p_{16} & = & -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3 \\ p_{17} & = & -x_{17}x_7 - x_{16} + x_7 \end{array}$$

```
p_{18}
        = -x_{18}x_7 + x_{16}
        = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
              -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
             -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
        = -x_{22}x_1 + x_{20}x_3
p_{22}
              -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
        = -x_{24}x_1 + x_{20}
p_{24}
        = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
        = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
        = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
        = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
              x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
       = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
       = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.16 Triangulation, step 16

Choosing variable: Trying the variable with index 17.

Variable x_{17} selected: The number of polynomials with this variable, with indexes from 1 to 17, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{17} from all other polynomials by reducing them with polynomial p_{16} from previous step.

$$\begin{array}{rcl} p_1 & = & -x_4 + x_1 \\ p_2 & = & -x_4 + x_2 \\ p_3 & = & -x_5 + x_1 \\ p_4 & = & x_5 + x_2 - \\ p_5 & = & x_3 - \\ p_6 & = & -x_8 + x_6 \\ p_7 & = & -x_8 + x_7 \\ p_8 & = & x_9 + x_6 - \\ p_9 & = & -x_9 + x_7 \\ p_{10} & = & -x_{13}x_7 + x_{11} \\ p_{11} & = & x_{13} + x_{12} - \\ p_{12} & = & -x_{14}x_1 + x_{10} \\ p_{13} & = & -x_{14}x_2 + x_{11} \end{array}$$

```
p_{14}
        = -x_{14}x_3 + x_{12}
        = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{15}
             x_{16}x_1 - x_{16} + x_{15}x_7x_3 - x_7x_3 - x_7x_1 + x_7
            -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
        = -x_{18}x_7 + x_{16}
p_{18}
             -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
        = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
        = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
             -x_{22}x_1 + x_{20}x_3
p_{22}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
        = -x_{24}x_1 + x_{20}
p_{24}
        = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
        = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
        = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
        = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
        = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
             x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
       = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
      = -x_{32}x_{25} + x_{29} - x_1
```

1.17 Triangulation, step 17

Choosing variable: Trying the variable with index 16.

Variable x_{16} selected: The number of polynomials with this variable, with indexes from 1 to 16, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{16} from all other polynomials by reducing them with polynomial p_{15} from previous step.

$$p_{1} = -x_{4} + x_{1}$$

$$p_{2} = -x_{4} + x_{2}$$

$$p_{3} = -x_{5} + x_{1}$$

$$p_{4} = x_{5} + x_{2} - x_{5} + x_{2} - x_{5} + x_{2} - x_{5} + x_{5} - x_{5} + x_{6}$$

$$p_{6} = -x_{8} + x_{6} - x_{5} + x_{6} - x_{5} + x_{6} - x_{5} + x_{7}$$

$$p_{8} = x_{9} + x_{6} - x_{5} + x_{7}$$

```
p_{10}
       = -x_{13}x_7 + x_{11}
       = x_{13} + x_{12} -
p_{11}
             -x_{14}x_1 + x_{10}
p_{12}
       = -x_{14}x_2 + x_{11}
       = -x_{14}x_3 + x_{12}
p_{14}
             -x_{15}x_{7}x_{3}x_{1} + x_{15}x_{7}x_{3} - x_{15}x_{2}x_{1} + x_{15}x_{2} + x_{7}x_{3}x_{1}
              -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
       = -x_{18}x_7 + x_{16}
       = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
             -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
       = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
       = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
             x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
      = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
       = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.18 Triangulation, step 18

Choosing variable: Trying the variable with index 15.

Variable x_{15} selected: The number of polynomials with this variable, with indexes from 1 to 15, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{15} . No reduction needed.

The triangular system has not been changed.

1.19 Triangulation, step 19

Choosing variable: Trying the variable with index 14.

Variable x_{14} selected: The number of polynomials with this variable, with indexes from 1 to 14, is 3.

Minimal degrees: 3 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{14} from all other polynomials by reducing them with polynomial p_{12} from previous step.

```
= -x_4 + x_1
      = -x_4 + x_2
 p_2
      = -x_5 + x_1
 p_3
       = x_5 + x_2 -
       = x_3 -
 p_5
       = -x_8 + x_6
 p_6
       = -x_8 + x_7
       = x_9 + x_6 -
 p_8
       = -x_9 + x_7
 p_9
       = -x_{13}x_7 + x_{11}
       = x_{13} + x_{12} -
p_{11}
       = -x_{11}x_1 + x_{10}x_2
p_{12}
       = -x_{12}x_1 + x_{10}x_3
       = -x_{14}x_1 + x_{10}
p_{14}
       = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
             -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
       = -x_{18}x_7 + x_{16}
       = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
       = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
       = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
            x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
      = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
      = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
      = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.20 Triangulation, step 20

Choosing variable: Trying the variable with index 13.

Variable x_{13} selected: The number of polynomials with this variable, with indexes from 1 to 13, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{13} from all other polynomials by reducing them with polynomial p_{10} from previous step.

```
p_1 = -x_4 + x_1
      = -x_4 + x_2
      = -x_5 + x_1
      = x_5 + x_2 -
 p_4
      = x_3 -
 p_5
      = -x_8 + x_6
 p_6
      = -x_8 + x_7
      = x_9 + x_6 -
 p_8
      = -x_9 + x_7
      = -x_{11}x_1 + x_{10}x_2
p_{10}
      = -x_{12}x_1 + x_{10}x_3
p_{11}
      = -x_{12}x_7 - x_{11} + x_7
p_{12}
      = -x_{13}x_7 + x_{11}
p_{13}
      = -x_{14}x_1 + x_{10}
p_{14}
      = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
            -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
       = -x_{18}x_7 + x_{16}
      = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
      = -x_{22}x_1 + x_{20}x_3
p22
      = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
      = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
      = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
      = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{28}
      = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
      = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
```

$$x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2$$

$$p_{30} = -x_{30} x_{25} + x_{29} x_{26} - x_{26} x_1 + x_{25} x_2$$

$$p_{31} = x_{31} x_{27} + x_{30} x_{26} + x_{29} x_{25} + x_{28}$$

$$p_{32} = -x_{32} x_{25} + x_{29} - x_1$$

1.21 Triangulation, step 21

Choosing variable: Trying the variable with index 12.

Variable x_{12} selected: The number of polynomials with this variable, with indexes from 1 to 12, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{12} from all other polynomials by reducing them with polynomial p_{11} from previous step.

```
= -x_4 + x_1
p_1
 p_2
     = -x_4 + x_2
     = -x_5 + x_1
 p_3
     = x_5 + x_2 -
 p_4
      = x_3 -
      = -x_8 + x_6
 p_6
     = -x_8 + x_7
      = x_9 + x_6 -
      = -x_9 + x_7
p_9
     = -x_{11}x_1 + x_{10}x_2
      = x_{11}x_1 + x_{10}x_7x_3 - x_7x_1
     = -x_{12}x_1 + x_{10}x_3
p_{12}
     = -x_{13}x_7 + x_{11}
p_{13}
      = -x_{14}x_1 + x_{10}
      = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
           -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
      = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
      = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
      = -x_{18}x_7 + x_{16}
p_{18}
      = -x_{19}x_1 + x_{19} + x_{15} -
      = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
     = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
      = -x_{22}x_1 + x_{20}x_3
     = -x_{23}x_7 + x_{23} + x_{21} -
     = -x_{24}x_1 + x_{20}
p_{24}
```

$$\begin{array}{lllll} p_{25} & = & x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12} \\ p_{26} & = & x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{27} & = & x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{28} & = & x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10} \\ p_{29} & = & -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 + \\ & & & x_{27}^2x_{25}x_{1} - x_{27}x_{25}^2x_{3} + x_{26}^2x_{25}x_{1} - x_{26}x_{25}^2x_{2} \\ p_{30} & = & -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_{1} + x_{25}x_{2} \\ p_{31} & = & x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28} \\ p_{32} & = & -x_{32}x_{25} + x_{29} - x_{1} \end{array}$$

1.22 Triangulation, step 22

Choosing variable: Trying the variable with index 11.

Variable x_{11} selected: The number of polynomials with this variable, with indexes from 1 to 11, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_{11} from all other polynomials by reducing them with polynomial p_{10} from previous step.

```
p_1 = -x_4 + x_1
     = -x_4 + x_2
     = -x_5 + x_1
     = x_5 + x_2 -
 p_{4}
     = x_3 -
 p_5
     = -x_8 + x_6
     = -x_8 + x_7
 p_7
     = x_9 + x_6 -
     = -x_9 + x_7
     = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
p_{10}
     = -x_{11}x_1 + x_{10}x_2
p_{11}
     = -x_{12}x_1 + x_{10}x_3
     = -x_{13}x_7 + x_{11}
p_{13}
     = -x_{14}x_1 + x_{10}
p_{14}
     = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
           -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
     = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
     = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{18}
     = -x_{18}x_7 + x_{16}
    = -x_{19}x_1 + x_{19} + x_{15} -
```

$$\begin{array}{rclcrcl} p_{20} & = & -x_{20}x_{7}x_{3}x_{1} + x_{20}x_{3}x_{1} - x_{20}x_{2}x_{1} + x_{20}x_{1} + x_{7}x_{1}^{2} - x_{1}^{2} \\ p_{21} & = & -x_{21}x_{1} + x_{20}x_{2} - x_{20} + x_{1} \\ p_{22} & = & -x_{22}x_{1} + x_{20}x_{3} \\ p_{23} & = & -x_{23}x_{7} + x_{23} + x_{21} - \\ p_{24} & = & -x_{24}x_{1} + x_{20} \\ p_{25} & = & x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12} \\ p_{26} & = & x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{27} & = & x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{28} & = & x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10} \\ p_{29} & = & -x_{29}x_{27}^{2}x_{25} - x_{29}x_{26}^{2}x_{25} - x_{29}x_{25}^{3} - x_{28}x_{25}^{2} + \\ & & & x_{27}^{2}x_{25}x_{1} - x_{27}x_{25}^{2}x_{3} + x_{26}^{2}x_{25}x_{1} - x_{26}x_{25}^{2}x_{2} \\ p_{30} & = & -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_{1} + x_{25}x_{2} \\ p_{31} & = & x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28} \\ p_{32} & = & -x_{32}x_{25} + x_{29} - x_{1} \end{array}$$

1.23 Triangulation, step 23

Choosing variable: Trying the variable with index 10.

Variable x_{10} selected: The number of polynomials with this variable, with indexes from 1 to 10, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_{10} . No reduction needed.

The triangular system has not been changed.

1.24 Triangulation, step 24

Choosing variable: Trying the variable with index 9.

Variable x_9 selected: The number of polynomials with this variable, with indexes from 1 to 9, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_9 from all other polynomials by reducing them with polynomial p_8 from previous step.

$$p_1 = -x_4 + x_1$$

$$p_2 = -x_4 + x_2$$

$$p_3 = -x_5 + x_1$$

$$p_4 = x_5 + x_2 - x_3$$

$$p_5 = x_3 - x_4$$

$$p_6 = -x_8 + x_6$$

```
= -x_8 + x_7
       = x_7 + x_6 -
       = x_9 + x_6 -
       = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
       = -x_{11}x_1 + x_{10}x_2
p_{11}
       = -x_{12}x_1 + x_{10}x_3
p_{12}
       = -x_{13}x_7 + x_{11}
p_{13}
       = -x_{14}x_1 + x_{10}
p_{14}
       = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
             -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
       = -x_{18}x_7 + x_{16}
       = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
            -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
       = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
       = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
             x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
      = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
      = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.25 Triangulation, step 25

Choosing variable: Trying the variable with index 8.

Variable x_8 selected: The number of polynomials with this variable, with indexes from 1 to 8, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_8 from all other polynomials by reducing them with polynomial p_6 from previous step.

$$p_1 = -x_4 + x_1$$

```
p_2
       = -x_4 + x_2
           -x_5 + x_1
 p_3
            x_5 + x_2 -
 p_4
       = x_3 -
       = x_7 + x_6 -
 p_6
       = -x_7 + x_6
 p_7
       = -x_8 + x_6
       = x_9 + x_6 -
            -x_{10}x_7x_3x_1-x_{10}x_2x_1+x_7x_1^2
p_{10}
       = -x_{11}x_1 + x_{10}x_2
       = -x_{12}x_1 + x_{10}x_3
p_{12}
       = -x_{13}x_7 + x_{11}
p_{13}
       = -x_{14}x_1 + x_{10}
       = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
             -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
           -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
             -x_{18}x_7 + x_{16}
p_{18}
       = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
             -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
       = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
       = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
P27
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
             x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
      = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
           -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.26 Triangulation, step 26

Choosing variable: Trying the variable with index 7.

Variable x_7 selected: The number of polynomials with this variable, with indexes from 1 to 7, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_7 from all other polynomials by reducing them with polynomial p_6 from previous step.

```
= -x_4 + x_1
      = -x_4 + x_2
 p_2
      = -x_5 + x_1
 p_3
       = x_5 + x_2 -
      = x_3 -
 p_5
       = 2x_6 -
 p_6
       = x_7 + x_6 -
       = -x_8 + x_6
 p_8
       = x_9 + x_6 -
 p_9
       = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
       = -x_{11}x_1 + x_{10}x_2
p_{11}
       = -x_{12}x_1 + x_{10}x_3
p_{12}
       = -x_{13}x_7 + x_{11}
       = -x_{14}x_1 + x_{10}
p_{14}
       = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
             -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
       = -x_{18}x_7 + x_{16}
       = -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
       = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
       = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
            x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
      = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
      = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
      = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.27 Triangulation, step 27

Choosing variable: Trying the variable with index 6.

Variable x_6 selected: The number of polynomials with this variable, with indexes from 1 to 6, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_6 . No reduction needed.

The triangular system has not been changed.

1.28 Triangulation, step 28

Choosing variable: Trying the variable with index 5.

Variable x_5 selected: The number of polynomials with this variable, with indexes from 1 to 5, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_5 from all other polynomials by reducing them with polynomial p_3 from previous step.

```
p_1 = -x_4 + x_1
     = -x_4 + x_2
 p_2
     = x_3 -
 p_3
     = -x_2 - x_1 + 1
     = -x_5 + x_1
 p_5
     = 2x_6 -
 p_6
     = x_7 + x_6 -
     = -x_8 + x_6
 p_8
     = x_9 + x_6 -
     = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
     = -x_{11}x_1 + x_{10}x_2
p_{11}
p_{12} = -x_{12}x_1 + x_{10}x_3
     = -x_{13}x_7 + x_{11}
     = -x_{14}x_1 + x_{10}
p_{14}
     = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
           -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
     = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
     = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
     = -x_{18}x_7 + x_{16}
     = -x_{19}x_1 + x_{19} + x_{15} -
p_{20} = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
```

```
= -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
        = -x_{22}x_1 + x_{20}x_3
p_{22}
             -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
        = -x_{24}x_1 + x_{20}
        = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
        = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
        = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
        = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
              x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
      = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
       = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

1.29 Triangulation, step 29

Choosing variable: Trying the variable with index 4.

Variable x_4 selected: The number of polynomials with this variable, with indexes from 1 to 4, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_4 from all other polynomials by reducing them with polynomial p_1 from previous step.

```
p_1
     = x_3 -
     = -x_2 - x_1 + 1
     = -x_2 + x_1
 p_3
     = -x_4 + x_1
 p_4
     = -x_5 + x_1
     = 2x_6 -
 p_6
     = x_7 + x_6 -
     = -x_8 + x_6
     = x_9 + x_6 -
 p_9
     = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
     = -x_{11}x_1 + x_{10}x_2
p_{11}
     = -x_{12}x_1 + x_{10}x_3
p_{12}
     = -x_{13}x_7 + x_{11}
p_{13}
     = -x_{14}x_1 + x_{10}
p_{14}
p_{15}
     = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
           -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
```

$$\begin{array}{lll} p_{16} & = & -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2 \\ p_{17} & = & -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3 \\ p_{18} & = & -x_{18}x_7 + x_{16} \\ p_{19} & = & -x_{19}x_1 + x_{19} + x_{15} - \\ p_{20} & = & -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2 \\ p_{21} & = & -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1 \\ p_{22} & = & -x_{22}x_1 + x_{20}x_3 \\ p_{23} & = & -x_{23}x_7 + x_{23} + x_{21} - \\ p_{24} & = & -x_{24}x_1 + x_{20} \\ p_{25} & = & x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12} \\ p_{26} & = & x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12} \\ p_{27} & = & x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11} \\ p_{28} & = & x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10} \\ p_{29} & = & -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 + \\ & & x_{27}^2x_{25}x_1 - x_{27}x_{25}^2x_3 + x_{26}^2x_{25}x_1 - x_{26}x_{25}^2x_2 \\ p_{30} & = & -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2 \\ p_{31} & = & x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28} \\ p_{32} & = & -x_{32}x_{25} + x_{29} - x_{1} \end{array}$$

1.30 Triangulation, step 30

Choosing variable: Trying the variable with index 3.

Variable x_3 selected: The number of polynomials with this variable, with indexes from 1 to 3, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_1 . No reduction needed.

The triangular system has not been changed.

1.31 Triangulation, step 31

Choosing variable: Trying the variable with index 2.

Variable x_2 selected: The number of polynomials with this variable, with indexes from 1 to 2, is 2.

Minimal degrees: 2 polynomial(s) with degree 1.

Polynomial with linear degree: Removing variable x_2 from all other polynomials by reducing them with polynomial p_1 from previous step.

$$p_1 = -2x_1 + 1$$

$$p_2 = -x_2 - x_1 + 1$$

```
p_3
           x_3 -
             -x_4 + x_1
 p_4
            -x_5 + x_1
 p_5
       = 2x_6 -
       = x_7 + x_6 -
 p_7
       = -x_8 + x_6
       = x_9 + x_6 -
       = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
p_{10}
       = -x_{11}x_1 + x_{10}x_2
p_{11}
       = -x_{12}x_1 + x_{10}x_3
       = -x_{13}x_7 + x_{11}
p_{13}
       = -x_{14}x_1 + x_{10}
p_{14}
       = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
             -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
p_{16}
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
       = -x_{18}x_7 + x_{16}
p_{18}
            -x_{19}x_1 + x_{19} + x_{15} -
p_{19}
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
            -x_{22}x_1 + x_{20}x_3
p22
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
p_{25}
       = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
       = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       = -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
             x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
       = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
       = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
      = -x_{32}x_{25} + x_{29} - x_1
```

1.32 Triangulation, step 32

Choosing variable: Trying the variable with index 1.

Variable x_1 selected: The number of polynomials with this variable, with indexes from 1 to 1, is 1.

Single polynomial with chosen variable: Chosen polynomial is p_1 . No reduction needed.

The triangular system has not been changed.

The triangular system is:

```
p_1 = -2x_1 + 1
     = -x_2 - x_1 + 1
      = x_3 -
 p_3
      = -x_4 + x_1
      = -x_5 + x_1
       = 2x_6 -
 p_6
       = x_7 + x_6 -
       = -x_8 + x_6
       = x_9 + x_6 -
p_9
       = -x_{10}x_7x_3x_1 - x_{10}x_2x_1 + x_7x_1^2
p_{10}
       = -x_{11}x_1 + x_{10}x_2
p_{11}
      = -x_{12}x_1 + x_{10}x_3
p_{12}
      = -x_{13}x_7 + x_{11}
p_{13}
      = -x_{14}x_1 + x_{10}
p_{14}
       = -x_{15}x_7x_3x_1 + x_{15}x_7x_3 - x_{15}x_2x_1 + x_{15}x_2 + x_7x_3x_1
p_{15}
             -x_7x_3 + x_7x_1^2 - 2x_7x_1 + x_7 + x_2x_1 - x_2
       = -x_{16}x_1 + x_{16} + x_{15}x_2 - x_2
       = -x_{17}x_1 + x_{17} + x_{15}x_3 - x_3
p_{17}
       = -x_{18}x_7 + x_{16}
p_{18}
       = -x_{19}x_1 + x_{19} + x_{15} -
       = -x_{20}x_7x_3x_1 + x_{20}x_3x_1 - x_{20}x_2x_1 + x_{20}x_1 + x_7x_1^2 - x_1^2
p_{20}
       = -x_{21}x_1 + x_{20}x_2 - x_{20} + x_1
p_{21}
       = -x_{22}x_1 + x_{20}x_3
p_{22}
       = -x_{23}x_7 + x_{23} + x_{21} -
p_{23}
       = -x_{24}x_1 + x_{20}
p_{24}
       = x_{25} - x_{22}x_{16} + x_{22}x_{11} + x_{21}x_{17} - x_{21}x_{12} - x_{17}x_{11} + x_{16}x_{12}
       = x_{26} + x_{22}x_{15} - x_{22}x_{10} - x_{20}x_{17} + x_{20}x_{12} + x_{17}x_{10} - x_{15}x_{12}
p_{26}
       = x_{27} - x_{21}x_{15} + x_{21}x_{10} + x_{20}x_{16} - x_{20}x_{11} - x_{16}x_{10} + x_{15}x_{11}
p_{27}
       = x_{28} + x_{27}x_{12} + x_{26}x_{11} + x_{25}x_{10}
p_{28}
       \hspace*{35pt} = \hspace*{35pt} -x_{29}x_{27}^2x_{25} - x_{29}x_{26}^2x_{25} - x_{29}x_{25}^3 - x_{28}x_{25}^2 +
p_{29}
            x_{27}^2 x_{25} x_1 - x_{27} x_{25}^2 x_3 + x_{26}^2 x_{25} x_1 - x_{26} x_{25}^2 x_2
      = -x_{30}x_{25} + x_{29}x_{26} - x_{26}x_1 + x_{25}x_2
p_{30}
      = x_{31}x_{27} + x_{30}x_{26} + x_{29}x_{25} + x_{28}
p_{31}
      = -x_{32}x_{25} + x_{29} - x_1
p_{32}
```

2 Final Remainder

2.1 Final remainder for conjecture geothm_zadatak

Calculating final remainder of the conclusion:

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

with respect to the triangular system.

1. Pseudo remainder with p_{32} over variable x_{32} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

2. Pseudo remainder with p_{31} over variable x_{31} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

3. Pseudo remainder with p_{30} over variable x_{30} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

4. Pseudo remainder with p_{29} over variable x_{29} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

5. Pseudo remainder with p_{28} over variable x_{28} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

6. Pseudo remainder with p_{27} over variable x_{27} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

7. Pseudo remainder with p_{26} over variable x_{26} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

8. Pseudo remainder with p_{25} over variable x_{25} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

9. Pseudo remainder with p_{24} over variable x_{24} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

10. Pseudo remainder with p_{23} over variable x_{23} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

11. Pseudo remainder with p_{22} over variable x_{22} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

12. Pseudo remainder with p_{21} over variable x_{21} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

13. Pseudo remainder with p_{20} over variable x_{20} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

14. Pseudo remainder with p_{19} over variable x_{19} :

$$g = -x_{18}x_6 + x_{18} + x_{15} -$$

15. Pseudo remainder with p_{18} over variable x_{18} :

$$g = x_{16}x_6 - x_{16} - x_{15}x_7 + x_7$$

16. Pseudo remainder with p_{17} over variable x_{17} :

$$g = x_{16}x_6 - x_{16} - x_{15}x_7 + x_7$$

17. Pseudo remainder with p_{16} over variable x_{16} :

$$g = x_{15}x_7x_1 - x_{15}x_7 - x_{15}x_6x_2 + x_{15}x_2 - x_7x_1 + x_7 + x_6x_2 - x_2$$

18. Pseudo remainder with p_{15} over variable x_{15} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

19. Pseudo remainder with p_{14} over variable x_{14} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

20. Pseudo remainder with p_{13} over variable x_{13} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

21. Pseudo remainder with p_{12} over variable x_{12} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

22. Pseudo remainder with p_{11} over variable x_{11} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

23. Pseudo remainder with p_{10} over variable x_{10} :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

24. Pseudo remainder with p_9 over variable x_9 :

$$g = -x_7^2 x_1^3 + 3x_7^2 x_1^2 - 3x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 -2x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2x_7 x_2 x_1 - x_7 x_2$$

25. Pseudo remainder with p_8 over variable x_8 :

$$\begin{array}{ll} g & = & -x_7^2 x_1^3 + 3 x_7^2 x_1^2 - 3 x_7^2 x_1 + x_7^2 + x_7 x_6 x_2 x_1^2 \\ & & -2 x_7 x_6 x_2 x_1 + x_7 x_6 x_2 - x_7 x_2 x_1^2 + 2 x_7 x_2 x_1 - x_7 x_2 \end{array}$$

26. Pseudo remainder with p_7 over variable x_7 :

$$g = -x_6^2 x_2 x_1^2 + 2x_6^2 x_2 x_1 - x_6^2 x_2 - x_6^2 x_1^3 + 3x_6^2 x_1^2 - 3x_6^2 x_1 + x_6^2 + 2x_6 x_2 x_1^2 - 4x_6 x_2 x_1 + 2x_6 x_2 + 2x_6 x_1^3 - 6x_6 x_1^2 + 6x_6 x_1 - 2x_6 - x_2 x_1^2 + 2x_2 x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

27. Pseudo remainder with p_6 over variable x_6 :

$$g = -x_2x_1^2 + 2x_2x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

28. Pseudo remainder with p_5 over variable x_5 :

$$g = -x_2x_1^2 + 2x_2x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

29. Pseudo remainder with p_4 over variable x_4 :

$$g = -x_2x_1^2 + 2x_2x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

30. Pseudo remainder with p_3 over variable x_3 :

$$g = -x_2x_1^2 + 2x_2x_1 - x_2 - x_1^3 + 3x_1^2 - 3x_1 + 1$$

31. Pseudo remainder with p_2 over variable x_2 :

$$g = 0$$

32. Pseudo remainder with p_1 over variable x_1 :

$$g = 0$$

3 Prover results

Status: Theorem has been proved.

Space Complexity: The biggest polynomial obtained during prover execution contains 21 terms.

Time Complexity: Time spent by the prover is 0.155 seconds.

4 NDG Conditions

NDG Conditions in readable form

• Failed to translate NDG Conditions to readable form