

Title	Animal cellulases with a focus on aquatic invertebrates
Author(s)	Tanimura, Aya; Liu, Wen; Yamada, Kyohei; Kishida, Takushi; Toyohara, Haruhiko
Citation	Fisheries Science (2013), 79(1): 1-13
Issue Date	2013-01
URL	<a href="http://hdl.handle.net/2433/169682">http://hdl.handle.net/2433/169682</a>
Right	The final publication is available at <a href="http://www.springerlink.com">www.springerlink.com</a>
Type	Journal Article
Textversion	author

1		
2		
3	1	<b>Title</b>
4		
5		
6	2	<b>Animal Cellulases with a Focus on Aquatic Invertebrates</b>
7		
8		
9	3	
10		
11		
12	4	<b>Aya Tanimura<sup>1</sup> • Wen Liu<sup>1</sup> • Kyohei Yamada<sup>1</sup> • Takushi Kishida<sup>2</sup> • Haruhiko Toyohara<sup>1*</sup></b>
13		
14		
15	5	
16		
17		
18	6	<sup>1</sup> Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University,
19		
20		
21		
22	7	Kyoto 606-8502, Japan
23		
24		
25	8	<sup>2</sup> Primate Research Institute, Kyoto University, Inuyama, Aichi 484-8506, Japan
26		
27		
28	9	
29		
30		
31	10	<b>Corresponding Author</b>
32		
33		
34		
35	11	Haruhiko Toyohara
36		
37		
38	12	Tel/Fax: 81-75-753-6446
39		
40		
41	13	<b>Email Address</b>
42		
43		
44	14	Haruhiko Toyohara: toyohara@kais.kyoto-u.ac.jp
45		
46		
47		
48	15	Aya Tanimura: ayatani@kais.kyoto-u.ac.jp
49		
50		
51	16	Wen Liu: liuwen@kais.kyoto-u.ac.jp
52		
53		
54	17	Kyohei Yamada: kyohei1216@gmail.com
55		
56		
57	18	Takushi Kishida: takushi@zoo.zool.kyoto-u.ac.jp
58		
59		
60	19	
61		
62		
63		
64		
65		

**Abstract**

Cellulose is utilized as a nutritional source by various organisms. It had been long believed that only protozoa, bacteria and fungi, in addition to plants and photo-synthetic bacteria, are able to synthesize cellulases encoded by their own genes. However, the wide spread distribution of cellulases throughout the animal kingdom has been recently recognized. Conventionally, animals digest cellulose by utilizing cellulases derived from symbiotic bacteria in the digestive organs. However, recent molecular biological studies have shown that some cellulase genes are actually encoded on animal chromosomes. In addition, the homologous primary structure of cellulases obtained from various phyla of invertebrates indicates the possible vertical transfer of the cellulase gene from ancient organisms that are now extinct. Studies on cellulase with unique enzymatic properties are expected to be applied to bioethanol production and aquaculture. In the present review, we describe cellulases, with a primary focus on aquatic invertebrates in which both endogenous and exogenous cellulases are involved in the breakdown of cellulose in the digestive organs.

**Keywords**

Breakdown • Cellulase • Cellulose • Endo- $\beta$ -1,4-glucanase • Endogenous • GHF9 • Invertebrate • Symbiosis

## Introduction

Cellulose synthesized by plants and phototrophic bacteria is the most abundant organic substance on the earth. Cellulose is chemically stable, and thereby plays an important role as a major component in the cell wall of plants and bacteria by providing physical strength [1]. This physical strength is attributed to the primary structure of cellulose, which consists of monomeric chains of D-glucopyranose bound by  $\beta$ -1,4-glycoside linkages that form cellulose microfibrils interconnected with hydrogen bonds (Fig. 1) [2]. In addition to cellulose, the cell wall contains lignin and various hemicelluloses, including mannan, xylan, and laminarin. The contents of the hemicelluloses differ across plant species [3].

Enzymes that degrade cellulose are collectively called cellulases, and are classified according to a range of characteristics. First, cellulases are classified according to the cleavage site on cellulose; comprising (1)  $\beta$ -1,4-endoglucanase (EC 3.2.1.4.), which cleaves cellulose at random sites, and (2)  $\beta$ -1,4-exocellobiohydrolase (EC 3.2.1.91.), which cleaves off glucose dimers from the terminal end of cellulose. Subsequently,  $\beta$ -glucosidase (EC 3.2.1.21.) cleaves glucose from the breakdown products formed by  $\beta$ -1,4-endoglucanase and  $\beta$ -1,4-exocellobiohydrolase (Fig. 2).

Second, cellulases are classified by the presence of the carbohydrate-binding module (CBM) within their molecules. Some carbohydrate degrading enzymes have a CBM that is independent of the catalytic site, which binds to substrates and stabilizes the enzymatic reaction. For example, cellulases that have a cellulose-binding domain (CBD) on a CBM constantly bind to cellulose, with the enzyme molecules

continuously moving to a subsequent cleavage site after each reaction. In comparison, cellulases that do not have a CBD detach from the cellulose after every cleaving reaction, and search for the cleaving site of the next hydrolytic reaction [2]. Cellulases that have a CBD are assumed to hydrolyze cellulose more efficiently than those that do not have (Fig. 3).

Third, cellulases are classified according to their primary structure. For instance, glycoside hydrolases are classified into the glycoside hydrolase family (GHF) by Henrissat et al., according to the amino acid sequence [4, CazyWeb: <http://www.cazy.org/> “Accessed 6 May 2012”]. At present, 130 families are registered, with cellulases being classified into families 1, 3, 5, 6, 7, 8, 9, 10, 12, 19, 26, 30, 44, 45, 48, 51, 61, 74, 116, and 124 (Table 1).

Cellulose is utilized as a nutritional source by various organisms. It has long been believed that only protozoa, bacteria, and fungi, in addition to plants and photo-synthetic bacteria, are able to synthesize cellulases encoded by their own genes [1, 5-7]. Before 1998, it was assumed that metazoans degraded cellulose using cellulases derived from symbiotic protozoa and bacteria in their digestive organs [1]. However, a novel gene encoding cellulase (GHF9;  $\beta$ -1,4-endoglucanase) was identified in *Reticulitermes speratus* (Arthropoda, Insecta) in 1998 [7]. Subsequently, the presence of endogenous  $\beta$ -1,4-endoglucanases, belonging to various GHF families, has been reported in various insects, crustaceans, mollusks, echinoderms, and nematodes [8-16].

The origin of cellulase in metazoans may be explained by two alternative hypotheses [15, 16]. The

1  
2  
3 72 first hypothesis is the horizontal transfer of cellulase genes from symbiotic protozoa. The second  
4  
5  
6 73 hypothesis is the vertical transfer of cellulase genes from ancient organisms that are now extinct. In this  
7  
8  
9 74 second hypothesis, cellulase genes have been inherited for a long period of time from the ancestor to the  
10  
11  
12 75 offspring. According to the reports on GHF9 [15, 16], which is the most intensively studied cellulases,  
13  
14  
15 76 vertical transfer is considered more likely. The amino acid sequence of GHF9 is very similar in several  
16  
17  
18 77 organisms, indicating the presence of a shared common cellulase ancestor, from which the GHF9 gene  
19  
20  
21 78 has been inherited for a hundred million years (Fig. 4). Unfortunately, evolutionary evidence of other  
22  
23  
24 79 GHFs is not available, and information about their primary structures remains fragmentary [15].  
25  
26  
27

28  
29 80 Table 2 classifies cellulases according to origin (endogenous or exogenous) and habitat (terrestrial or  
30  
31  
32 81 aquatic). Endogenous origin means that the cellulase gene is encoded on the chromosomes of the  
33  
34  
35 82 organisms, whereas an exogenous origin means that the cellulase gene is encoded on the chromosomes of  
36  
37  
38 83 symbiotic microorganisms. As shown in Table 2, enzymes from archaea, eubacteria, fungi, and plants are  
39  
40  
41 84 all classified as endogenous, whereas cellulases of both endogenous and exogenous origin are found in  
42  
43  
44 85 invertebrates. The exogenous origin of cellulases in invertebrates is assumed to compensate for cellulases  
45  
46  
47 86 of endogenous origin. In contrast, cellulases of endogenous origin have yet to be reported in vertebrates.  
48  
49  
50

51 87 In the present review, we describe cellulases, with a primary focus on aquatic invertebrates in which  
52  
53  
54 88 both endogenous and exogenous cellulases contribute to the breakdown of cellulose in the digestive  
55  
56  
57 89 organs.  
58  
59  
60  
61  
62  
63  
64  
65

**Cellulases derived from symbiotic microorganisms in terrestrial organisms**

It had long been doubted as to whether herbivorous terrestrial animals actually digest cellulose and utilize it as a nutritional source. For instance, there have been substantial efforts to validate whether herbivorous mammals utilize cellulose during digestion [1]. Studies on symbiotic microorganisms in the ruminant stomach of herbivorous animals, which are involved in cellulose breakdown, are documented in Table 3. In 1942, Hungate reported the ability of the genera *Diplodinium* and *Entodinium* to breakdown protozoan cellulose, and the possible implication of these microorganisms in the breakdown of cellulose in the lumen of herbivorous mammals [17]. Various microorganisms in the digestive organs of sheep have also been investigated. For example, in 1982, Wood et al. isolated an anaerobic symbiotic microorganism *Ruminococcus albus* from the sheep lumen that was able to degrade cellulose, and successfully purified the cellulase from the extracts [18]. In 1986, Coleman et al. recorded the cellulose-degrading ability of the protozoa *Entodinium caudatum* in the sheep lumen [19]. Subsequently, in 1992, Bernaler et al. reported the cellulose-degrading ability of the anaerobic fungus *Neocallinastix frontalis* in the sheep lumen [20]. In addition to sheep, Varel et al. reported the cellulase degrading ability of the gram-positive bacillus *Bacteroides succinogenes* and the gram-negative coccus *Ruminococcus flavefaciens* in the porcine colon [21].

In addition to herbivorous mammals, there have been extensive studies of symbiotic bacteria

implicated in the breakdown of cellulose in herbivorous insects. The cellulose-degrading ability of symbiotic bacteria in the termite *Reticulitermes flavipes* was first reported in 1924 [22]. Subsequently, in 1932, Trager et al. reported the cellulase activity of flagellates in the digestive organs of the wood roach [23]. Recently, Delalibera et al. found symbiotic cellulose-degrading bacteria and fungi in the digestive organs of the wood borer and bark beetle [24]. The presence of symbiotic microorganisms has been extensively studied in termites. For example, Wenzel et al. successfully isolated symbiotic aerobic and anaerobic bacteria from damp wood termites (*Zootermopsis angusticollis*; Arthropoda, Insecta) in 2002 [25]. In 2006, Watanabe et al. isolated archaeobacteria exhibiting cellulase activity from the digestive tract of the giant northern termite (*Mastotermes darwiniensis*; Arthropoda, Insecta) [26]. Interestingly, Martin et al. detected cellulase activity in the midgut extract of the fungus-growing termite, *Macrotermes natalensis* (Arthropoda, Insecta) in 1978 [27]. This termite species cultures cellulose-degrading bacteria in its hive, and obtains cellulase from these bacteria. A similar example was reported for another termite, *Macrotermes mulleri* (Arthropoda, Insecta) [28]. Hence, there are a large number of terrestrial animals that are associated with symbiotic microorganisms, allowing them to effectively utilize cellulose.

### **Cellulases derived from symbiotic microorganisms in aquatic organisms**

The shipworm (Bivalvia, Teredinidae) is a major pest that bores holes in the hulls of wooden ships and other wooden marine structures, occasionally resulting in the sinking of ships. Symbiotic bacteria



were first found in granules of “the Gland of Deshayes” from one shipworm species *Bankia australis* (Mollusca, Bivalvia) in 1973 [29]. In 1983, Waterbury et al. found some species of bacteria that had both cellulose-degrading activity and nitrogen-fixing ability in shipworms, indicating that this species of shipworm utilizes cellulose and fixes nitrogen as carbon and nitrogen sources, respectively [30]. In 1991, the authors also reported the presence of a symbiotic bacteria belonging to Proteobacteria in various species of shipworm [31]. This was the first report demonstrating the widespread distribution of symbiotic bacteria belonging to the same phylum being present in a variety of shipworm species. After 2002, *Teredinibacter turnerae* classified into Proteobacterium, was found to be distributed in 24 species belonging to 9 distinct genera out of the total 14 genera of shipworm [32]. This observation indicates the ubiquitous distribution of specific bacteria among various species of shipworm.

Studies on symbiotic bacteria with cellulase activity have extended to the deep sea ecology of animals inhabiting wood that has fallen to the ocean floor at the sea bottom. In 1997, bacteria morphologically resembling those found in shipworms were discovered in the digestive organs of a bivalve species *Xylophaga washingtona* (Mollusca, Bivalvia), which belongs to the same class as the shipworm [33]. New symbiotic bacteria were also found in the gills of a sunken wood-associated mussel (Mytilidae) in 2008 using the fluorescence in situ hybridization (FISH) technique, which employs specific molecular probes to identify different species of bacteria [34]. In 2010, another species of symbiotic bacteria was found in the gills and digestive organs of *Pectinodonta* sp., belonging to the

Gastropoda [35]. On the basis of these findings, the mechanism of cellulose breakdown in the wood ecological system of the deep sea is gradually being revealed.

In comparison, Cary et al. reported the vertical transfer of symbiotic bacteria from “mother to baby” in related species of *Calymene* (cold-seep clams) and *Solemya reidi* [36]. In a later study of *Bankia setacea*, a type of shipworm, Sipe et al. reported the vertical transfer of symbiotic bacteria from “mother to baby” via the eggs [37]. Since the cellulose-degrading activity of these bacteria has yet to be demonstrated, further studies on the occurrence of cellulase activities in these bacterial species are required.

The gribble worm is an isopod that bores holes in wooden ships, similar to shipworm. The presence of symbiotic bacteria in the gribble worm has long been disputed [38]; however, in 2010, one gribble worm species, *Limnoria quadripunctata* (Arthropoda, Malacostracea), was confirmed to have endogenous cellulases, the genes of which are encoded on the chromosomes of the species [39].

**Endogenous cellulases in terrestrial and aquatic animals**

Table 3 summarizes the reported endogenous  $\beta$ -1,4-endoglucanases belonging to GHF9 from terrestrial and aquatic invertebrates. Most species containing endogenous cellulases belong to the Arthropoda and Nematoda. The first species reported to have endogenous cellulase was the termite *Reticulitermes speratus*, which is widely distributed across Japan, and has been studied intensively as a

harmful insect that decomposes wooden houses [7]. In 1998, Watanabe et al. succeeded in cloning the cDNA of  $\beta$ -1,4-endoglucanase from *R. speratus*, and the enzyme was classified as GHF9, according to the deduced amino acid sequence. The authors confirmed its endogenous origin by using PCR and Southern blot analysis of DNA extracted from termites. These procedures revealed the presence of an intron in the gene of  $\beta$ -1,4-endoglucanase. Later, the cDNA of another cellulase was cloned from *Coptotermes formosanus* (Arthropoda, Insecta), a termite species related to *R. speratus*, with it being classified as an endogenous GHF9 cellulase [40]. This cellulase was expressed in the foregut and midgut, with a symbiotic flagellate expressing a GHF7 cellulase in the hindgut [40]. These findings indicate that *C. formosanus* first degrades cellulose in the foregut and midgut via endogenous cellulases, and then the degraded products of cellulose are further digested by cellulases from the symbiotic flagellate in the hindgut, facilitating the effective digestion of cellulose. The presence of endogenous cellulases was subsequently reported for herbivorous arthropods, including the cockroach, which inhabits forests [41], and the well-known flour beetle *Tribolium castaneum* (Arthropoda, Insecta), which has a worldwide distribution [42].

In addition to arthropods, cellulases from Nematoda have been studied intensively. In 1998, the cDNA of cellulase was cloned from the plant-pathogenic cyst nematodes *Globodera rostochiensis* and *Heterodera glycines* (Nematoda, Tylenchida), and was classified as an endogenous GHF5 family enzyme [11]. In addition to cellulase, *Heterodera glycines* has endogenous genes encoding chitinase [43].

Cellulase genes were assumed to be horizontally transferred from bacteria or fungi to the ancestors of these two nematodes [44]. These nematodes infest the roots of host plants through an aperture that is formed by using their cellulases to degrade the plant cell walls [45]. In Japan, the gene of endogenous  $\beta$ -1,4-endoglucanase belonging to GHF45 was found in a major pine wood nematode *Bursaphelenchus xylophilus* (Nematoda, Tylenchida) which is parasitic to pine trees, causing pine tree death. These nematodes are assumed to bore into pine trees by degrading the cell wall in a similar way to the other two described nematode species. The  $\beta$ -1,4-endoglucanase gene of this nematode was demonstrated to be horizontally transferred from fungi. Thus, this species may have acquired the ability to degrade cellulose independently from other plant parasitic nematodes [46].

A small number of aquatic invertebrate species with endogenous cellulases have been reported. However, the phylum of the species found to have cellulases is diverse: Arthropoda, Mollusca, Annelida, Echinodermata, and Chordata. In aquatic animals, endogenous cellulase ( $\beta$ -1,4 endoglucanase) was first identified in *Cherax quadricarinatus* (common crayfish: Arthropoda, Crustacea). The primary structure of their cellulases shows homology with those of the termite, a terrestrial arthropod [47]. Although this finding indicates that cellulases distribute widely in arthropod, further studies on the distribution of cellulases among the other subphylums as Cheliceriformes, Myriopoda, or Trilobitomorpha is needed to confirm the widespread distribution of cellulases in arthropods.

The presence of cellulases has been most intensively studied in mollusks. At present, GHF9

cellulases have, for example, been reported in *Halliotis discus hannai* (Mollusca, Gastropoda) [48],  
*Halliotis discus discus* (Mollusca, Gastropoda) [49], *Corbicula japonica* (Mollusca, Bivalvia) [12],  
*Ampullaria crossean* (Mollusca, Gastropoda) [50], and *Mizuhopecten yessoensis* (Mollusca, Bivalvia)  
[CazyWeb: <http://www.cazy.org/> “Accessed 6 May 2012”]. In addition, GHF45 cellulase has been  
reported in *Mytilus edulis* (Mollusca, Bivalvia) [51], *Corbicula japonica* [52], and *Ampullaria crossean*  
[53]. There are species differences in the feeding habits of these mollusks. For instance, *H. discus hannai*  
and *H. discus discus* feed on diatoms in the larval stage and macroalgae in the adult stage. *M. edulis* is an  
epifaunal suspension feeder in coastal marine areas. *C. japonica* preferentially feeds on terrestrial  
particulate organic matter over phytoplankton in brackish waters [49, 54–58]. To the best of our  
knowledge, the diet of *A. crossean* has not been reported. Nevertheless, *Pomacea canaliculata*, which  
belongs to the same family as *A. crossean*, is known to digest fresh leaves. In 2011, Qiu et al. reported  
that *P. canaliculata* (Mollusca, Gastropoda) feeds on both fresh and decayed leaves of a variety of  
macrophytes [59]. Qiu et al. also mentioned that several species of Ampullariidae (Pomacea) eat leaves of  
macrophytes in wetlands [59].

The widespread distribution of cellulases among various mollusks with different feeding habits may  
also indicate the importance of this enzyme in biochemical cellulose breakdown, particularly for *C.*  
*japonica*, which is thought to feed on detritus, including decaying plants. The detritus that accumulates on  
the substrate of brackish areas includes a large amount of plant fragments, mainly composed of cellulose,

and is considered an important food resource for estuarine benthos [60, 61]. A recent study revealed that *C. japonica* contains endogenous  $\beta$ -glucosidase, which degrades the digested products of cellulose formed by  $\beta$ -1,4-glucanase into free glucose [62]. In addition, immunological analysis using an antibody to GHF9 endo- $\beta$ -1,4-glucanase from *C. japonica* confirmed the production of GHF9 endo- $\beta$ -1,4-glucanase in the digestive gland [63]. Because of the synergistic action of  $\beta$ -1,4-glucanases [12, 52] and  $\beta$ -glucosidase, *C. japonica* is assumed to utilize cellulose efficiently as a nutritional resource. Furthermore, *C. japonica* is also able to degrade hemicellulose. Sakamoto et al. revealed that *C. japonica* contains xylanase, which degrades xylan, one of the hemicellulases [64]. A comparative study of cellulase and hemicellulase activities in bivalves confirmed that *C. japonica* exhibits significantly higher mannanase activity than other bivalves [65]. Hence, *C. japonica* appears to be well adapted to brackish environments rich in plant-derived detritus.

Furthermore, recent studies have confirmed that other mollusks also utilize multiple enzymes. For example, *H. discus hannai* contains mannanase,  $\beta$ -1,3-glucanase, and alginate lyase [66-68], and *M. edulis* contains mannanase [69]. More recently, Kumagai and Zahura found that *Aplysia kurodai* (Mollusca, Gastropoda), which is a sea hare that eats seaweed, contains  $\beta$ -1,3-glucanases and mannanase [70, 71]. The presence of endogenous cellulose-degrading enzymes, including  $\beta$ -1,4-endoglucanase, in these studied species indicates that these enzymes are widespread among mollusks.

Cellulase activity has also been well documented in other aquatic invertebrates, such as polychaetes

and crabs inhabiting a wide range of environments. Niiyama recorded high cellulase and hemicellulase activities in a variety of temperate region macrobenthos [72]. Toyohara et al. suggested that either meiobenthos (small annelids or nematodes) or sediments in temperate areas exhibit cellulase activity [73]. Furthermore, Yamada and Toyohara confirmed the presence of cellulase activity in the meiobenthos and sediments of the subantarctic region, which was dependent on climatic and sediment features [74]. A recent study by Liu and Toyohara proposed that sediment complexes harbor enzymes including cellulase [75], and it is suggested that plant-degrading enzymes are widely distributed in sediments and the environment.

$\beta$ -1,4-Endoglucanases of GHF9 have been reported in the echinoderm *Storonylocentrotus nudus* (Echinodermata, Echinoidea), which typically feeds on macroalgae [10], and *Ciona intertinalis* (Chordata, Ascidiacea) [76], for which the genome structure has been intensively studied. Although *C. intertinalis* belongs to the Chordata, which also includes humans, it is known to synthesize cellulose to protect its body. Genomic analyses have shown that the genome of this species harbors endogenous enzymes that synthesize cellulose, which were horizontally transferred from bacteria 530 million years ago. This species has evolved a specific biochemical system to synthesize cellulose that is distinct from that of plants [77]. While the  $\beta$ -1,4-endoglucanases of this species share a homologous structure with other GHF9 cellulases that have been reported in various aquatic invertebrates, they have a distinct function [76]. *C. intertinalis* contains cellulases to degrade cellulose in the tunic, not to degrade cellulose ingested

into the digestive organs. Hence, *C. intertinalis* may serve as a good model to study the horizontal gene transfer mechanism from prokaryotes to eukaryotes, although supplementary genetic information must be collected for other sea squirts.

255

#### Utilization of animal cellulases

Cellulose is the most abundant organic material on the earth. A variety of animals that have flourished on this planet are dependent on this widespread resource. Cellulose is even essential for humans as an industrial material for generating paper and clothes, as well as nutritional food fiber materials. The degradation and reconstitution of cellulose has been extensively studied with the aim of producing desirable industrial materials or food, in addition to synthesizing various cellulose derivatives. The modification of cellulose has been studied intensively, and cellulase is probably one of the most industrially utilized enzymes [2, 78, 79].

Recently, cellulases have attracted attention as potential energy sources, such as bioethanol. Bioethanol is synthesized from glucose derived from plant cellulose and hemicellulose. Bioethanol differs from conventional fossil fuels, such as petroleum and coal, in that it is a carbon-neutral fuel, which does not increase the amount of CO<sub>2</sub> in the air. This is because the amount of CO<sub>2</sub> released into the air when consuming bioethanol is equal to that fixed by plants used for bioethanol production [80]. An important aspect of bioethanol production is to reduce energy expenditure when degrading cellulose. At present,



microbial cellulases with heat-stable and acid-stable properties are available for the breakdown of cellulose. To date, anaerobic bacteria have been intensively studied for this purpose, as described in a review by Demain, in which the co-culture of anaerobic bacteria was shown to be effective for producing bioethanol under moderate conditions, and at low cost by using crude substrates [81]. Animal cellulases are potential candidates for bioethanol production, because they are expected to be equipped with specific enzymatic properties that are different to microbial cellulases. In 2011, Xu et al. reported that endogenous  $\beta$ -1,4 endoglucanase from a gastropod *Ampullaria crosseana* showed acid and heat stability [82]. Yanagisawa et al. reported that drips containing cellulase and amylase from the mid-gut gland of scallops saccharified sea lettuce and suggested that these enzymes could be used to produce ethanol when combined with yeast [83].

The digestive efficiency of termite enzymes is reported to be as high as 99 % for cellulose and 87 % for hemicelluloses, driving researchers to investigate potential industrial applications. It has been suggested that the ability of chewing and biting plants led to the high efficiency of cellulose breakdown by termites [2, 84]. Aquatic invertebrates, such as gastropods, also scrape plants using radula. Hence, further studies on both insect and aquatic animal cellulases might contribute toward improving the efficiency of bioethanol production.

Information about aquatic animal cellulases might also be beneficial for aquaculture. The feeding habits of algae eaters that are important to fisheries, such as *Haliotis discus hannai*, *Mytilus*

*edulis*, and *Mizuhopecten yessoensis*, have been intensively studied [54–56, 85]. For example, abalone is known to feed on diatoms in the larval stage, and feeds on macroalgae in the adult stage. It has also been reported that climatic events, such as tidal streams, may affect ecological circumstances by changing levels of competition or predation stress [54, 55]. Hence detailed information about the feeding habits of species important to fisheries may help in the development of efficient aquaculture systems. Other useful studies in associated fields, including protoplast preparation [86] and improving fruit yields [87], are also advancing.

Recent studies on cellulases have revealed that these enzymes are present in many organisms, reflecting the widespread distribution of cellulose on the earth. Organisms, including bacteria, fungi, and invertebrates, have developed a system to digest cellulose, which may originally have evolved in the bacteria or plants for cell wall construction [2]. Compared to studies of cellulases in microorganisms, such as bacteria and fungi, limited information is available about aquatic invertebrates. In conclusion, further studies on the cellulases of aquatic invertebrates are anticipated; these will contribute toward improving the efficient use of plants by human beings.

**Acknowledgements**

This study was partly supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan (No. 21380131).

1  
2  
3 306  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 307 **References**  
4  
5

- 6 308 1 Watanabe H and Tokuda G (2001) Animal cellulases. Cell Mol Life Sci 58: 1167-1178  
7  
8  
9 309 2 Watanabe H and Tokuda G (2010) Cellulolytic systems in insects. Annu Rev Entomol Annu  
10  
11  
12 310 Rev 55: 609-632  
13  
14  
15  
16 311 3 Cosgrove DJ (2005) Growth of the plant cell wall. Nat Rev Mol Cell Bio 6: 850-861  
17  
18  
19 312 4 Henrissat B, Claeyssens M, Tomme P, Lemesle L and Mornon JP (1989) Cellulase families  
20  
21  
22 313 revealed by hydrophobic cluster-analysis. Gene 81: 83-95  
23  
24  
25 314 5 Karrer P, Schubert P and Wehrli W (1925) Polysaccharides XXXIII on the enzymatic  
26  
27  
28 315 breakdown of artificial silk and native cellulose. Helv Chim Acta 8: 797-810  
29  
30  
31  
32 316 6 Lenski RE (2003) Life's solution: Inevitable humans in a lonely universe. Nature 425: 767-768  
33  
34  
35 317 7 Watanabe H, Noda H, Tokuda G and Lo N (1998) A cellulase gene of termite origin. Nature  
36  
37  
38 318 1998; 394: 330-331  
39  
40  
41 319 8 Martin MM (1991) The evolution of cellulose digestion in insects. Philos Trans Roy Soc B  
42  
43  
44 320 333: 281-288  
45  
46  
47 321 9 Sugimura M, Watanabe H, Lo N and Saito H (2003) Purification, characterization, cDNA  
48  
49  
50 322 cloning and nucleotide sequencing of a cellulase from the yellow-spotted longicorn beetle, Psacotha  
51  
52  
53 323 hilaris. Eur J Biochem 270: 3455-3460  
54  
55  
56  
57 324 10 Nishida Y, Suzuki K, Kumagai Y, Tanaka H, Inoue A and Ojima T (2007) Isolation and primary  
58  
59  
60  
61  
62  
63  
64  
65

- 1
- 2
- 3 325 structure of a cellulase from the Japanese sea urchin *Strongylocentrotus nudus*. *Biochimie* 89: 1002-1011
- 4
- 5
- 6 326 11 Smant G, Stokkermans J, Yan YT, de Boer JM, Baum TJ, Wang XH, Hussey RS, Gommers FJ,
- 7
- 8
- 9 327 Henrissat B, Davis EL, Helder J, Schots A and Bakker J (1998) Endogenous cellulases in animals:
- 10
- 11
- 12 328 Isolation of beta-1,4-endoglucanase genes from two species of plant-parasitic cyst nematodes. *Proc Natl*
- 13
- 14 329 *Acad Sci USA* 95: 4906-4911
- 15
- 16
- 17
- 18
- 19 330 12 Sakamoto K, Touhata K, Yamashita M, Kasai A and Toyohara H (2007) Cellulose digestion by
- 20
- 21
- 22 331 common Japanese freshwater clam *Corbicula japonica*. *Fish Sci* 2007; 73: 675-683
- 23
- 24
- 25 332 13 Imjongjirak C, Amparyup P and Sittipraneed S (2008) Cloning, genomic organization and
- 26
- 27
- 28 333 expression of two glycosyl hydrolase family 10 (GHF10) genes from golden apple snail (*Pomacea*
- 29
- 30 334 *canaliculata*). *DNA Sequence* 19: 224-236
- 31
- 32
- 33
- 34
- 35 335 14 Kostanjsek R, Milatovic M and Srus J (2010) Endogenous origin of endo-beta-1,4-glucanase in
- 36
- 37
- 38 336 common woodlouse *Porcellio scaber* (Crustacea, Isopoda). *J Comp Physiol B* 180: 1143-1153
- 39
- 40
- 41 337 15 Lo N, Watanabe H and Sugimura M (2003) Evidence for the presence of a cellulase gene in the
- 42
- 43
- 44 338 last common ancestor of bilaterian animals. *Proc Roy Soc Lond B Bio* 270: S69-S72
- 45
- 46
- 47
- 48 339 16 Davison A and Blaxter M (2005) Ancient origin of glycosyl hydrolase family 9 cellulase genes.
- 49
- 50
- 51 340 *Mol Biol Evol* 22: 1273-1284
- 52
- 53
- 54 341 17 Hungate RE (1946) The cellulose-decomposing bacteria in the rumen of cattle. *J Bacteriol* 51:
- 55
- 56
- 57 342 28-28
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1
- 2
- 3 343 18 Wood TM, Wilson CA and Stewart CS (1982) Preparation of the cellulase from the cellulolytic
- 4
- 5
- 6 344 anaerobic rumen bacterium *Ruminococcus-albus* and its release from the bacterial-cell wall. *Biochem J*
- 7
- 8
- 9 345 205: 129-137
- 10
- 11
- 12 346 19 Coleman GS (1986) The distribution of carboxymethylcellulase between fractions taken from
- 13
- 14
- 15 347 the rumens of sheep containing no protozoa or one of 5 different protozoal populations. *J Agr Sci* 106:
- 16
- 17
- 18 348 121-127
- 19
- 20
- 21
- 22 349 20 Bernalier A, Fonty G, Bonnemoy F and Gouet P (1992) Degradation and fermentation of
- 23
- 24
- 25 350 cellulose by the rumen anaerobic fungi in axenic cultures or in association with cellulolytic bacteria. *Curr*
- 26
- 27
- 28 351 *Microbiol* 25: 143-148
- 29
- 30
- 31
- 32 352 21 Varel VH, Fryda SJ and Robinson IM (1984) Cellulolytic bacteria from pig large-intestine. *Appl*
- 33
- 34
- 35 353 *Environ Microb* 47: 219-221
- 36
- 37
- 38 354 22 Cleveland LR (1924) The physiological and symbiotic relationships between the intestinal
- 39
- 40
- 41 355 protozoa of termites and their host, with special reference to *Reticulitermes flavipes* Kollar. *Biol Bull-US*
- 42
- 43
- 44 356 46: 178-201
- 45
- 46
- 47
- 48 357 23 Trager W (1932) A cellulase from the symbiotic intestinal flagellates of termites and of the
- 49
- 50
- 51 358 roach, *Cryptocercus punctulatus*. *Biochem J* 26: 1762-1771
- 52
- 53
- 54 359 24 Delalibera I, Handelsman J and Raffa KF (2005) Contrasts in cellulolytic activities of gut
- 55
- 56
- 57 360 microorganisms between the wood borer, *Saperda vestita* (Coleoptera: Cerambycidae), and the bark
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

361 beetles, *Ips pini* and *Dendroctonus frontalis* (Coleoptera: Curculionidae). *Environ Entomol* 34: 541-547  
 362 25 Wenzel M, Schonig I, Berchtold M, Kampfer P and Konig H (2002) Aerobic and facultatively  
 363 anaerobic cellulolytic bacteria from the gut of the termite *Zootermopsis angusticollis*. *J Appl Microbiol*  
 364 92: 32-40  
 365 26 Watanabe H, Takase A, Tokuda G, Yamada A and Lo N (2006) Symbiotic "archaezoa" of the  
 366 primitive termite *Mastotermes darwiniensis* still play a role in cellulase production. *Eukaryot Cell* 5:  
 367 1571-1576  
 368 27 Martin MM and Martin JS (1978) Cellulose digestion in midgut of fungus-growing termite  
 369 *Macrotermes natalensis* - Role of acquired digestive enzymes. *Science* 199: 1453-1455  
 370 28 Rouland C, Civas A, Renoux J and Petek F (1988) Purification and properties of cellulases  
 371 from the termite *Macrotermes mulleri* (Termitidae, Macrotermitinae) and its symbiotic fungus  
 372 *Termitomyces* sp. *Comp Biochem Phys B* 91: 449-458  
 373 29 Popham JD and Dickson MR (1973) Bacterial associations in the teredo *Bankia australis*  
 374 (Lamellibranchia: Mollusca). *Mar Biol* 19: 338-340  
 375 30 Waterbury JB, Calloway CB and Turner RD (1983) A cellulolytic nitrogen-fixing bacterium  
 376 cultured from the Gland of Deshayes in shipworms (Bivalvia: Teredinidae). *Science* 221: 1401-1403  
 377 31 Distel DL, DeLong EF and Waterbury JB (1991) Phylogenetic characterization and in situ  
 378 localization of the bacterial symbiont of shipworms (Teredinidae: Bivalvia) by using 16S rRNA sequence

- 1
- 2
- 3 379 analysis and oligodeoxynucleotide probe hybridization. AEM 57: 2376-2382
- 4
- 5
- 6 380 32 Distel DL, Morrill W, Maclaren-Toussaint N, Franks D and Waterbury J (2002) *Teredinibacter*
- 7
- 8
- 9 381 *turnerae* gen. nov., sp. nov., a dinitrogen-fixing, cellulolytic, endosymbiotic  $\gamma$ -proteobacterium isolated
- 10
- 11
- 12 382 from the gills of wood-boring molluscs (Bivalvia: Teredinidae). IJSEM 52: 2261-2269
- 13
- 14
- 15
- 16 383 33 Distel DL and Roberts SJ (1997) Bacterial endosymbionts in the gills of the deep-sea
- 17
- 18
- 19 384 wood-boring bivalves *Xylophaga atlantica* and *Xylophaga washingtona*. Biol Bull 192: 253-261
- 20
- 21
- 22 385 34 Duperron S, Laurent MCZ, Gaill F and Gros O (2008) Sulphur-oxidizing extracellular bacteria
- 23
- 24
- 25 386 in the gills of Mytilidae associated with wood falls. FEMS Microbiol Ecol 63: 338-349
- 26
- 27
- 28 387 35 Zbinden M, Pailleret M, Ravaux J, Gaudron SM, Hoyoux C, Lambourdiere J, Waren A, Lorion
- 29
- 30
- 31 388 J, Halary S and Duperron S (2010) Bacterial communities associated with the wood-feeding gastropod
- 32
- 33
- 34 389 *Pectinodonta* sp. (Patellogastropoda, Mollusca). FEMS Microbiol Ecol 74: 450-463
- 35
- 36
- 37
- 38 390 36 Cary SC (1994) Vertical transmission of a chemoautotrophic symbiont in the protobranch
- 39
- 40
- 41 391 bivalve, *Solemya reidi*. Mol Mar Biol Biotech 3: 121-130
- 42
- 43
- 44 392 37 Sipe AR, Wilbur AE and Cary SC (2000) Bacterial symbiont transmission in the wood-boring
- 45
- 46
- 47 393 shipworm *Bankia setacea* (Bivalvia: Teredinidae). Environ Microbiol 66: 1685-1691
- 48
- 49
- 50
- 51 394 38 Boyle J. P (1978) Absence of Microorganism in Crustacean Digestive Tracts. Science 200:
- 52
- 53
- 54 395 1157-1159
- 55
- 56
- 57 396 39 King AJ (2010) Molecular insight into lignocelluloses digestion by a marine isopod in the
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65



1  
2  
3 397 absence of gut microbes. Proc Natl Acad Sci USA 107: 5345-5350  
4  
5  
6 398 40 Nakashima K, Watanabe H, Saitoh H, Tokuda G and Azuma JI (2000) Dual cellulose-digesting  
7  
8  
9 399 system of the wood-feeding termite, *Coptotermes formosanus* Shiraki. Insect Biochem Molec 32:  
10  
11  
12 400 777-784  
13  
14  
15  
16 401 41 Lo N, Tokuda G, Watanabe H, Rose H, Slaytor M, Maekawa K, Bandi C and Noda H (2000)  
17  
18  
19 402 Evidence from multiple gene sequences indicates that termites evolved from wood-feeding cockroaches.  
20  
21  
22 403 Curr Biol 10: 801-804  
23  
24  
25  
26 404 42 Willis JD, Oppert B, Oppert C, Klingeman WE and Jurat-Fuentes JL (2011) Identification,  
27  
28  
29 405 cloning, and expression of a GHF9 cellulase from *Tribolium castaneum* (Coleoptera: Tenebrionidae). J  
30  
31  
32 406 Insect Physiol 57: 300-306  
33  
34  
35 407 43 Gao BL, Allen R, Maier T, McDermott JP, Davis EL, Baum TJ and Hussey RS (2002)  
36  
37  
38 408 Characterisation and developmental expression of a chitinase gene in *Heterodera glycines*. Int J Parasitol  
39  
40  
41 409 32: 1293-1300  
42  
43  
44  
45 410 44 Yan YT, Smant G, Stokkermans J, Qin L, Helder J, Baum T, Schots A and Davis E (1998)  
46  
47  
48 411 Genomic organization of four beta-1,4-endoglucanase genes in plant-parasitic cyst nematodes and its  
49  
50  
51 412 evolutionary implications. Gene 220: 61-70  
52  
53  
54 413 45 Jones JT, Furlanetto C and Kikuchi T (2005) Horizontal gene transfer from bacteria and fungi  
55  
56  
57 414 as a driving force in the evolution of plant parasitism in nematodes. Nematology 7: 641-646  
58  
59  
60  
61  
62  
63  
64  
65

- 1
- 2
- 3 415 46 Kikuchi T, Jones JT, Aikawa T, Kosaka H and Ogura N (2004) A family of glycosyl hydrolase
- 4
- 5
- 6 416 family 45 cellulases from the pine wood nematode *Bursaphelenchus xylophilus*. *FEBS Lett* 572: 201-205
- 7
- 8
- 9 417 47 Byrne KA, Lehnert SA, Johnson SE and Moore SS (1999) Isolation of a cDNA encoding a
- 10
- 11
- 12 418 putative cellulase in the red claw crayfish *Cherax quadricarinatus*. *Gene* 239: 317-324
- 13
- 14
- 15
- 16 419 48 Suzuki K, Ojima T and Nishita K (2003) Purification and cDNA cloning of a cellulase from
- 17
- 18
- 19 420 abalone *Haliotis discus hannai*. *Eur J Biochem* 270: 771-778
- 20
- 21
- 22 421 49 Nikapitiya C, Oh C, De Zoysa M, Whang I, Kang DH, Lee SR, Kim SJ and Lee J (2010)
- 23
- 24
- 25 422 Characterization of beta-1,4-endoglucanase as a polysaccharide-degrading digestive enzyme from disk
- 26
- 27
- 28 423 abalone, *Haliotis discus discus*. *Aquacult Int* 18: 1061-1078
- 29
- 30
- 31
- 32 424 50 Li YH, Yin QY, Ding M and Zhao FK (2009) Purification, characterization and molecular
- 33
- 34
- 35 425 cloning of a novel endo-beta-1,4-glucanase AC-EG65 from the mollusc *Ampullaria crosseana*. *Comp*
- 36
- 37
- 38 426 *Biochem Phys B* 153: 149-156
- 39
- 40
- 41 427 51 Xu BZ, Janson JC and Sellos D (2001) Cloning and sequencing of a molluscan
- 42
- 43
- 44 428 endo-beta-1,4-glucanase gene from the blue mussel, *Mytilus edulis*. *Eur J Biochem* 268: 3718-3727
- 45
- 46
- 47
- 48 429 52 Sakamoto K and Toyohara H (2009) Molecular cloning of glycoside hydrolase family 45
- 49
- 50
- 51 430 cellulase genes from brackish water clam *Corbicula japonica*. *Comp Biochem Phys B* 152: 390-396
- 52
- 53
- 54 431 53 Guo R, Ding M, Zhang SL, Xu GJ and Zhao FK (2008) Molecular cloning and characterization
- 55
- 56
- 57 432 of two novel cellulase genes from the mollusc *Ampullaria crosseana*. *J Comp Physiol B* 178: 209-215
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1
- 2
- 3 433 54 Won NI, Kawamura T, Takami H, Hoshikawa H and Watanabe Y (2011) Comparison of
- 4
- 5
- 6 434 abalone (*Haliotis discus hannai*) catches in natural habitats affected by different current systems:
- 7
- 8
- 9 435 Implication of climate effects on abalone fishery. *Fish Res* 110: 84-91
- 10
- 11
- 12 436 55 Xing RL, Wang CH, Cao XB and Chang YQ (2008) Settlement, growth and survival of abalone,
- 13
- 14
- 15 437 *Haliotis discus hannai*, in response to eight monospecific benthic diatoms. *J Appl Phycol* 20: 47-53
- 16
- 17
- 18 438 56 Hawkins AJS, Navarro E and Iglesias JIP (1990) Comparative allometries of gut-passage time,
- 19
- 20
- 21 439 gut content and metabolic fecal loss in *Mytilus edulis* and *Cerastoderma edule*. *Mar Biol* 105: 197-204
- 22
- 23
- 24 440 57 Antonio ES, Kasai A, Ueno M, Kurikawa Y, Tsuchiya K, Toyohara H, Ishihi Y, Yokoyama H
- 25
- 26
- 27 441 and Yamashita Y (2010) Consumption of terrestrial organic matter by estuarine molluscs determined by
- 28
- 29
- 30 442 analysis of their stable isotopes and cellulase activity. *Estuar Coast Shelf S* 86: 401-407
- 31
- 32
- 33 443 58 Kasai A and Nakata A (2005) Utilization of terrestrial organic matter by the bivalve *Corbicula*
- 34
- 35 444 *japonica* estimated from stable isotope analysis. *Fish Sci* 71: 151-158
- 36
- 37
- 38 445 59 Qiu JW, Chan MT, Kwong KL and Sun J (2011) Consumption, Survival and growth in the
- 39
- 40
- 41 446 invasive freshwater snail *Pomacea Canaliculata*: Does food freshness matter? *J Mollus Stud* 77: 189-195
- 42
- 43
- 44 447 60 Kanaya G, Suzuki T and Kikuchi E (2011) Spatio-temporal variations in macrozoobenthic
- 45
- 46
- 47 448 assemblage structures in a river-affected lagoon (Idoura Lagoon, Sendai Bay, Japan): Influences of
- 48
- 49
- 50 449 freshwater inflow. *Estuar Coast Shelf S* 92: 169-179
- 51
- 52
- 53 450 61 Kanaya G, Takagi S and Kikuchi E (2008) Spatial dietary variations in *Laternula marilina*
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

451 (Bivalva) and Hediste spp. (Polychaeta) along environmental gradients in two brackish lagoons. Mar  
 452 Ecol-Prog Ser 359: 133-144  
 453 62 Sakamoto K, Uji S, Kurokawa T and Toyohara H (2009) Molecular cloning of endogenous  
 454 beta-glucosidase from common Japanese brackish water clam Corbicula japonica. Gene 435: 72-79  
 455 63 Sakamoto K, Uji S, Kurokawa T and Toyohara H (2008) Immunohistochemical, in situ  
 456 hybridization and biochemical studies on endogenous cellulase of Corbicula japonica. Comp Biochem  
 457 Phys B 150: 216-221  
 458 64 Sakamoto K and Toyohara H (2009) Putative endogenous xylanase from brackish-water clam  
 459 Corbicula japonica. Comp Biochem Phys B 154: 85-92  
 460 65 Sakamoto K and Toyohara H (2009) A comparative study of cellulase and hemicellulase  
 461 activities of brackish water clam Corbicula japonica with those of other marine Veneroida bivalves. J Exp  
 462 Biol 212: 2812-2818  
 463 66 Ootsuka S, Saga N, Suzuki K, Inoue A and Ojima T (2006) Isolation and cloning of an  
 464 endo-beta-1,4-mannanase from Pacific abalone Haliotis discus hannai. J Biotechnol 125: 269-280  
 465 67 Shimizu E, Ojima T and Nishita K (2003) cDNA cloning of an alginate lyase from abalone,  
 466 Haliotis discus hannai. Carbohydr Res 338: 2841-2852  
 467 68 Kumagai Y and Ojima T (2009) Enzymatic properties and the primary structure of a  
 468 beta-1,3-glucanase from the digestive fluid of the Pacific abalone Haliotis discus hannai. Comp Biochem

469 Phys B 154: 113-120

470 69 Xu BZ, Hagglund P, Stalbrand H and Janson JC (2002) endo-beta-1,4-mannanases from blue  
471 mussel, *Mytilus edulis*: purification, characterization, and mode of action. J Biotechnol 92: 267-277

472 70 Zahura UA, Rahman MM, Inoue A, Tanaka H and Ojima T (2011) cDNA cloning and bacterial  
473 expression of an endo-beta-1,4-mannanase, AkMan, from *Aplysia kurodai*. Comp Biochem Phys B 159:  
474 227-235

475 71 Kumagai Y and Ojima T (2010) Isolation and characterization of two types of  
476 beta-1,3-glucanases from the common sea hare *Aplysia kurodai*. Comp Biochem Phys B 155: 138-144

477 72 Niiyama T and Toyohara H (2011) Widespread distribution of cellulase and hemicellulase  
478 activities among aquatic invertebrates. Fish Sci 77: 649-655

479 73 Toyohara H, Park Y, Tsuchiya K and Liu W (2012) Cellulase activity in meiobenthos in  
480 wetlands. Fish Sci 78: 133-137

481 74 Yamada K and Toyohara H (2012) Functions of meiobenthos and microorganisms in cellulose  
482 breakdown in sediments of wetlands with different origins in Hokkaido. Fish Sci 78: 699-706

483 75 Liu W and Toyohara H (2012) Sediment-complex-binding cellulose breakdown in wetlands of  
484 rivers. Fish Sci 78: 661-665

485 76 Dehal P, Satou Y, Campbell RK, Chapman J et al (2002) The draft genome of *Ciona*  
486 intestinalis: Insights into chordate and vertebrate origins. Science 298: 2157-2167

487 77 Nakashima K, Yamada L, Satou Y, Azuma J and Satoh N (2004) The evolutionary origin of  
 488 animal cellulose synthase. Dev Genes Evol 214: 81-88  
 489 78 Bhat MK (2000) Cellulases and related enzymes in biotechnology. Biotechnol Adv 18: 355-383  
 490 79 Murad HA and Azzaz HH (2010) Cellulase and dairy animal feeding. Biotechnology 9: 1-19  
 491 80 Demirbas A (2007) Progress and recent trends in biofuels. Prog Energ Combust 33: 1-18  
 492 81 Demain AL (2009) Biosolutions to the energy problem. J Ind Microbiol Biol 36: 319-332  
 493 82 Yin QY, Teng YG, Li YH, Ding M and Zhao FK (2011) Expression and characterization of  
 494 full-length *Ampullaria crosseana* endoglucanase EG65s and their two functional modules. Biosci Biotech  
 495 Biochem 75: 240-246  
 496 83 Yanagisawa M, Ojima T and Nakasaki K (2011) Bioethanol from sea lettuce with the use of  
 497 crude enzymes derived from waste. J Mater Cycl Waste Manag 13: 321-326  
 498 84 Ohkuma M (2003) Termite symbiotic systems: efficient bio-recycling of lignocellulose.  
 499 Appl. Microbiol Biot 61: 1-9  
 500 85 Silina AV and Zhukova NV. Growth variability and feeding of scallop *Patinopecten yessoensis*  
 501 on different bottom sediments: Evidence from fatty acid analysis (2007) J Exp Mar Biol Ecol 348:  
 502 46-59  
 503 86 Inoue A, Mashino C, Kodama T and Ojima T (2011) Protoplast preparation from *Laminaria*  
 504 *japonica* with recombinant alginate lyase and cellulase. Mar Biotechnol 13: 256-263

505 87 Zhao FY, Lin JF, Zeng XL, Guo LQ, Wang YH and You LR (2010) Improvement in fruiting  
 506 body yield by introduction of the *Ampullaria crosseana* multi-functional cellulase gene into *Volvariella*  
 507 *volvacea*. *Biores Technol* 101: 6482-6486  
 508 87 Zhao FY, Lin JF, Zeng XL, Guo LQ, Wang YH and You LR (2010) Improvement in fruiting  
 509 body yield by introduction of the *Ampullaria crosseana* multi-functional cellulase gene into *Volvariella*  
 510 *volvacea*. *Biores Technol* 101: 6482-6486  
 511 88 Nozaki M, Miura C, Tozawa Y and Miura T (2009) The contribution of endogenous cellulase to  
 512 the cellulose digestion in the gut of earthworm (*Pheretima hilgendorfi*: Megascolecidae). *Soil Biol*  
 513 *Biochem* 41: 762-769  
 514 88 Shimada K and Maekawa K (2008) Correlation between social structure and nymphal  
 515 wood-digestion ability in the xylophagous cockroaches *Salgallea esakii* and *Panesthia angustipennis*  
 516 (Blaberidae: Panesthiinae). *Sociobiology* 52: 417-427  
 517 90 Tokuda G, Lo N, Watanabe H, Slaytor M, Matsumoto T, and Noda H (1999) Metazoan  
 518 cellulase genes from termites: intron/exon structures and site of expression. *BBA-Struct expr* 1447:  
 519 146-159  
 520 91 Tokuda G, Lo N, Watanabe H, Arakawa G, Matsumoto T and Noda H (2004) Major alternation  
 521 of the expression site of endogenous cellulases in members of an apical termite lineage. *Mol Ecol* 13:  
 522 3219-3228

523 92 Zhang DH, Lax AR, Bland JM and Allen AB (2011) characterization of a new endogenous  
 524 endo-beta-1,4-glucanase of Formosan subterranean termite (*Coptotermes formosanus*). Insect Biochem  
 525 Molec 41: 211-218  
 526 93 Zhang DH, Lax AR, Raina AK and Bland JM (2009) Differential cellulolytic activity of  
 527 native-form and C-terminal tagged-form cellulase derived from *Coptotermes formosanus* and expressed  
 528 in *E. coli*. Insect Biochem Molec 39: 516-522  
 529 94 Li L, Frohlich J, Pfeiffer P and Konig H (2003) Termite gut symbiotic archaezoa are becoming  
 530 living metabolic fossils. Eucaryot Cell 2: 1091-1098  
 531 95 Scharf ME, Wu-Scharf D, Zhou X, Pittendrich BR and Bennett GW (2005) Gene expression  
 532 profiles among immature and adult reproductive castes of the termite *Reticulitermes flavipes*. Insect Mol  
 533 Biol 14: 31-44  
 534 96 Itakura S, Masuta T, Tanaka H and Enoki A (2006) Identification of two subterranean termite  
 535 species (Isoptera: Rhinotermitidae) using cellulase genes. J Econ Entomol 99: 123-128  
 536 97 Kim N, Choo YM, Lee KS, Hong SJ, Seol KY, Je YH, Sohn HD, Jin BR (2008) Molecular  
 537 cloning and characterization of a glycosyl hydrolase family 9 cellulase distributed throughout the  
 538 digestive tract of the cricket *Teleogryllus emma*. Comp Biochem Phys B 150: 368-376  
 539 98 Ito K, Nozakli M, Ohta T, Miura C, Tozawa Y and Miura T (2011) Differences of two  
 540 po;ychaete specis reflected in enzyme activities. Mar Biol 158: 1211-1221



541 99 Davison A and Blaxter ML (2005) An expression sequence tag survey of gene expression in the  
 542 pond snail *Lymnaea stagnails*, an intermediate vector of *Fasciola hepatica*. Parasitology 130: 539-552  
 543 100 Clark NL, Gasper J, Sekino M, Springer SA, Aquadro CF and Swanson WJ (2009) Coevolution  
 544 of interacting Fertilization Proteins. Plos Genet 5: e1000570  
 545 101 Crawford AC, Kricker JA, Anderson AJ, Richardson NR and Mather PB (2004) Structure and  
 546 function of a cellulase gene in redclaw crayfish, *Cherax quadricarinatus*. Gene 340: 267-274  
 547 102 Zhu XD, Mahairas G, Illies M, Cameron RA, Davidson EH and Etensohn CA (2001) A  
 548 large-scale analysis of mRNAs expressed by primary mesenchyme cells of the sea urchin embryo.  
 549 Development 128: 2615-2627  
 550 103 Edvardsen RB, Seo HC, Jensen MF, Mialon A, Mikhaleva J, Bjordal M, Cartry J, Reinhardt R,  
 551 weissenbach J, Wincker P and Chourrout D (2005) Remodeling of the homeobox gene complement in the  
 552 tunicate *Oikopleura dioica*. Curr Biol 15: R12-R13  
 553 104 Seo HC, Edvardsen RB, Maelnd AD, Bjordal M, Jansen MF, Hansen A, Flaar M, Weissenbach  
 554 J, Lahrach H, Wincker P, Reinhardt R and Chourrout D (2004) Hox cluster disintegration with persistent  
 555 anteroposterior order of expression in *Oikopleura dioica*. Nature 431: 67-71  
 556 105 Schloss PD, Delalibera I, Handelsman J and Radda KF (2006) Bacteria associated with the guts  
 557 of two wood-boring beetles: *Anoplophora glabripennis* and *Saperda vestita* (Cerambycidae). Environ  
 558 Entomol 35: 625-629

## Figure legends

**Fig.1** Three types of cellulose hydrolysis enzymes collectively named cellulase (Exocellubiohydrolase, Endo- $\beta$ -1,4-glucanase, and  $\beta$ -glucosidase) are involved in the cellulose degradation process

Exocellubiohydrolases (exocellulase) cut cellulose from reducing (R) or nonreducing (NR) termini to release cellubiose (dotted circle). Endo- $\beta$ -1,4-glucanases cut the cellulose randomly from the internal  $\beta$ -1,4 linkages (broken circle).  $\beta$ -Glucosidases hydrolyze cellubiose or cellu-oligomers from the reducing termini to produce glucoses (double circle). Scissors represents the cutting site of each enzyme.

**Fig.2** Schematic view of the crystalline structure of cellulose chains

Glucose units are joined by  $\beta$ -1,4 linkages. Different cellulose chains are linked by hydrogen bonds (marked by dotted lines) to form cellulose crystals.

**Fig.3** Two different types of cellulase act on the cellulose binding domain (CBD); multi-domain cellulases and single-domain cellulases

Multi-domain cellulases comprise a catalytic domain that is linked to a CBD by a peptide strand, termed a linker. In this instance, CBDs are assumed to bind to cellulose, which improves the efficiency of the hydrolytic process of multi-domain cellulases by continuing catalytic action on the cellulose surface, while following CBD movement. In contrast, single-domain cellulases leave the cellulose surface after

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

577 catalytic action is complete, and subsequently approach a new linkage to participate in another catalytic  
578 action.

579

580 **Fig.4** Phylogenetic tree of GHF9 endo- $\beta$ -1,4-glucanase based on a nucleotide sequence (CazyWeb:  
581 <http://www.cazy.org/> “Accessed 6 May 2012”.)

582 This tree was prepared according to the method of Davidson et al. (2005) [16]. Please  
583 note that this tree was prepared according to the nucleotide sequence of part of the endogenous  
584 endo- $\beta$ -1,4-glucanases reported to date.

585

586

和文要旨

動物セルラーゼ ― 水生無脊椎由来酵素に注目して

谷村 彩，劉 文，山田 京平（京大院農），岸田 拓士（京大霊長研），豊原 治彦（京大院農）

本総説では，動物セルラーゼに関する研究について 1900 年代半ばから最新のものに至る論文を渉猟し，陸生と水生，共生と内源性の観点から分類した。特に自身の染色体 DNA 上にコードされる内源性セルラーゼについては詳述した。これらの酵素について，糖加水分解酵素ファミリーにおける分類や一次構造上の類似性についても述べた。なかでも，近年著しく研究が進んだ軟体動物，節足動物，棘皮動物などの水生生物については，それらの食性や生態との関連について概説するとともに，水生生物由来のセルラーゼの今後の応用の可能性についても論じた。

キーワード

エンド- $\beta$ -1,4-グルカナーゼ，共生，セルラーゼ，セルロース，内源性，分解，無脊椎動物，

GHF9

Fig 1




Fig 2




Fig 3





Fig 4





**Table 1** List of members of the Gglycoside Hhydrolase Ffamily (GHF) \*

GHF	Cellulase Type	Archaea	Eukaryote			
			Eubacteria	Fungus	Plant	Animal(Invertebrate)
1	$\beta$ -glucosidase	+	+	+	+	+
3	$\beta$ -glucosidase	+	+	+	+	+
5	Endoglucanase, $\beta$ -glucosidase	+	+	+	+	+
6	Endoglucanase, Cellobiohydrolase	-	+	+	-	-
7	Endoglucanase, Cellobiohydrolase	-	-	+	+	+
8	Endoglucanase	+	+	-	-	-
9	Endoglucanase, Cellobiohydrolase, $\beta$ -glucosidase	+	+	+	+	+
10	Endoglucanase	-	+	-	-	-
12	Endoglucanase	+	+	+	+	-
19	Endoglucanase	+	+	-	-	-
26	Endoglucanase	-	+	-	-	-
30	$\beta$ -glucosidase	+	+	+	+	+
44	Endoglucanase	-	+	-	-	-
45	Endoglucanase	-	+	+	-	+
48	Endoglucanase	-	+	+	-	-
51	Endoglucanase	+	+	+	+	-
61	Endoglucanase	-	-	+	+	-
74	Endoglucanase	-	+	+	-	-
116	$\beta$ -glucosidase	+	+	+	+	+
124	Endoglucanase	-	+	-	-	-

\* The GHF includes comprises all of glycoside hydrolases, including cellulases. Cellulases are further classified into 20 families [CazyWeb:

<http://www.cazy.org/> “Accessed 6 May 2012”]., most of which belongs to endo- $\beta$ -1, 4-glucanase. Only GHF5, 9, 10, and 45 are found in metazoans, while and GHF9 and 45 are exclusively found in aquatic invertebrates.  $\beta$ -Glycosidases have beenare reported infrom most animal phyla phylum of animals, because since this enzyme hydrolyzes various dimers of sugars besidein addition to cellulbiose.

**Table 2** Classification of cellulases based on habitatliving area and enzyme genetic origin\*

Habitat type	Organism	Exogenous/Endogenous
Terrestrial	Archaea	Endo
	Eubacteria	Endo
	Fungus	Endo
	Plant	Endo
	Invertebrate Animal	Exo/Endo
	Vertebrate Animal	Exo
Aquatic	Archaea	Endo
	Eubacteria	Endo
	Fungus	Endo
	Plant	Endo
	Invertebrate Animal	Exo/Endo
	Vertebrate Animal	Exo

\* Animals are classified as into “aquatic” and “terrestrial.”. Then, cCellulases are classified into “endogenous” and “exogenous.”. Note,It should be noted that only exogenous cellulases have been so far reported for in vertebrates to date.

**Table 3** List of invertebrate cellulases\*

Exogenous/ Endogenous	Terrestrial/ Aquatic	Organism					Reference
		Phylum	Class	Order	Species	Common Name	
Endogenous	Terrestrial	Platyhelminthe	Trematoda	Strigeidida	<i>Schistosoma mansoni</i>		[16]
		Annelida	Oligochaeta	Haplotaxida	<i>Eisenia andrei</i>		CazyWeb
					<i>Eisenia andrei</i>		CazyWeb
					<i>Eisenia fetida</i>	branding worm	CazyWeb
					<i>Lumbricus rubellus</i>	humus earthworm	[16]
					<i>Pheretima hilgendorfi</i>	earthworm	[88]
		Arthropoda	Insecta	Blattaria	<i>Panesthia angustipennis</i>	cockroach	[89]
					<i>Panesthia cribrata</i>	cockroach	[41]
					<i>Salganea esakii</i>	cockroach	[89]
				Coleoptera	<i>Timarcha balearica</i>	beetle	[16]
				Copelata	<i>Tribolium castaneum</i>	red flour beetle	[42]
				Hymenoptera	<i>Apis mellifera</i>	honeybee	[16]
				Isopoda	<i>Glyptotermes sp. Wyong</i>	termite	[90]
					<i>Hodotermopsis sjoestdi</i>	termite	[91]
					<i>Sinocapritermes mushae</i>	termite	[91]
				Isoptera	<i>Coptotermes acinaciformis</i>	termite	CazyWeb
					<i>Coptotermes formosanus</i>	Formosan subterranean termite	[92]
					<i>Coptotermes formosanus</i>	Formosan subterranean termite	[93]
					<i>Coptotermes formosanus Shiraki</i>	termite	[40]
					<i>Mastotermes darwiniensis</i>	termite	[94]
					<i>Nasutitermes takasagonesis</i>	termite	[91]
					<i>Nasutitermes takasagonesis</i>	termite	[90]
					<i>Nasutitermes walkeri</i>	termite	[90]
					<i>Neotermes koshunensis</i>	termite	[91]
					<i>Odontotermes formosanus</i>	termite	CazyWeb
					<i>Odontotermes formosanus</i>	termite	[91]
					<i>Reticulitermes flavipes</i>	eastern subterrenean termite	[95]
					<i>Reticulitermes speratus</i>	termite	[7]
					<i>Reticulitermes speratus</i>	termite	[90]
					<i>Reticulitermes speratus</i>	termite	[96]
				Orthoptera	<i>Teleogryllus emma</i>	emma field cricket	[97]
			Malacostraca	Mysida	<i>Neomysis intermedia</i>		CazyWeb

Exogenous/ Endogenous	Terrestrial/ Aquatic	Organism					Reference
		Phylum	Class	Order	Species	Common Name	
Endogenous	Aquatic	Annelida	Polychaeta	Aciculata	<i>Perinereis nuntia brevicirris</i>		[98]
		Mollusca	Bivalvia	Ostreoidea	<i>Crassostrea virginica</i>	eastern oyster	[16]
				Pectinoida	<i>Argopecten irradians</i>	bay scallop	[16]
					<i>Argopecten irradians</i>	bay scallop	[16]
					<i>Mizuhopecten yessoensis</i>	ezo giant scallop	CazyWeb
				Veneroida	<i>Corbicula japonica</i>		[12]
			Gastropoda	Architaenioglossa	<i>Ampullaria crosseana</i>		[50]
				Pulmonata	<i>Biomphalaria glabrata</i>	bloodfluke planorb	[16]
					<i>Lymnaea stagnalis</i>	great pond snail	[99]
				Vetigastropoda	<i>Haliotis corrugata</i>	pink abalone	[100]
					<i>Haliotis cracherodii</i>	black abalone	[100]
					<i>Haliotis discus discus</i>	disc abalone	[49]
					<i>Haliotis discus hannai</i>	ezo abalone	[48]
					<i>Haliotis fulgens</i>	green abalone	[100]
					<i>Haliotis kamtschatkana</i>		[100]
					<i>Haliotis rufescens</i>		[100]
					<i>Haliotis sorenseni</i>	white abalone	[100]
					<i>Haliotis walallensis</i>	flat abalone	[100]
		Arthropoda	Branchiopoda	Diplostraca	<i>Daphnia magna</i>	water flea	[16]
			Insecta	Isoptera	<i>Macrotermes barnevi</i>		CazyWeb
			Malacostraca	Amphipoda	<i>Gammarus pulex</i>	shrimp	[16]
				Decapoda	<i>Austrothelphusa transversa</i>		CazyWeb
					<i>Callinectes sapidus</i>	blue crab	[16]
					<i>Cherax quadricarinatus</i>	crayfish	[47]
					<i>Cherax quadricarinatus</i>	crayfish	[101]
					<i>Euastacus sp. SL-2005</i>		CazyWeb
					<i>Homarus americanus</i>	American lobster	[16]
				Isopoda	<i>Limnoria quadripunctata</i>		[39]
					<i>Porcellio scaber</i>		[14]
		Echinodermata	Echinoidea	Echinacea	<i>Strongylocentrotus nudus</i>	sea urchin	[10]
					<i>Strongylocentrotus purpuratus</i>	purple sea urchin	[102]

continued

Exogenous/ Endogenous	Terrestrial/ Aquatic	Organism					Reference
		Phylum	Class	Order	Species	Common Name	
Endogenous	Aquatic	Chordata	Appendiculari	Copelata	<i>Oikopleura dioica</i>	sea squirt	[103]
					<i>Oikopleura dioica</i>	sea squirt	[104]
			Ascidiacea	Enterogona	<i>Ciona intestinalis</i>	sea squirt	[76]
					<i>Ciona savignyi</i>	sea squirt	[16]
				Stolidobranchia	<i>Botryllus schlosseri</i>	sea squirt	[16]
					<i>Halocynthia roretzi</i>	sea squirt	[16]
					<i>Molgula tectiformis</i>	sea squirt	[16]
Exogenous	Terrestrial	Arthropoda	Insecta	Blattaria	<i>Cryptocercus punctulatus</i>	brown-hooted cockroach	[23]
				Coleoptera	<i>Dendroctonus frontalis</i>	southern pine beetle	[24]
					<i>Ips pini</i>	North American pine engraver	[24]
					<i>Saperda vestita</i>	Wood Borer	[105]
				Isoptera	<i>Macrotermes mulleri</i>	termite	[28]
					<i>Macrotermes natalensis</i>	termite	[27]
					<i>Mastotermes darwiniensis</i>	Giant Northern Termite	[26]
					<i>Reticulitermes flavipes</i>	eastern subterranean termite	[22]
					<i>Zootermopsis angusticollis</i>	Pacific dampwood termite	[25]
		Chordata	Mammalia	Artiodactyla	<i>Sus sp.</i>	pig	[21]
				Catartiodactyla	<i>Bos primigenius</i>	aurochs	[17]
					<i>Ovis aries</i>	sheep	[19]
					<i>Ovis aries</i>	sheep	[18]
					<i>Ovis aries</i>	sheep	[20]
	Aquatic	Mollusca	Bivalvia	Teredinidae	<i>Bankia gouldi</i>	shipworm	[30]
					<i>Lyrodus pedicellatus</i>	shipworm	[32]
					<i>Psiloteredo healdi</i>	shipworm	[30]
					<i>Teredo bartschi</i>	shipworm	[30]
					<i>Teredo furcifera</i>	shipworm	[30]
					<i>Teredo navalis</i>	shipworm	[30]
			Gastropoda	Patellogastropoda	<i>Pectinodonta sp.</i>		[35]

\* Invertebrate cellulases are classified according to the standard described in Table 2. Please note that endogenous cellulases include only GHF9, while exogenous cellulases include some of the cellulases reported to date. Animal classification and common names are presented according to the Species Dictionary: [http://bm2.genes.nig.ac.jp/Integrated\\_BodyMap/species.php](http://bm2.genes.nig.ac.jp/Integrated_BodyMap/species.php) “Accessed 6 May 2012”. Endogenous GHF9s are summarized according to the CazyWeb: <http://www.cazy.org/> “Accessed 6 May 2012”. and Davison et al (2005) [16].