

DECOLORIZATION OF ORANGE II DYE USING BACTERIA IMMOBILIZED ONTO MULTI-WALLED CARBON NANOTUBE CEMENT COMPOSITE

MOHAMMAD VOJDANI

A dissertation submitted in partial fulfillment of the
requirements for the award of the degree of
Master of Science (Biotechnology)

Faculty of Biosciences and Bioengineering
Universiti Teknologi Malaysia

JULY 2012

I would like to dedicate this
thesis to my ever-supportive family
and my supervisor Dr. Shafnaz Shahir who
have been my great inspiration in completing this thesis

ACKNOWLEDGEMENT

First and foremost, I want to thank my dear supervisor Dr. Shafinaz Shahir and dear co-supervisor Dr. Zaiton Abdul Majid for their kindness and guidance throughout my entire research. Their encouragement, advices and semi daily observations on my work gave me the inspiration to keep on the right direction during my research project. Without them I could never accomplish my project smoothly. I would also like to thank all those who were involved directly or indirectly in the completion of this project. Last but not least, I would like to express my utmost appreciation for my lovely Mom and Dad who have given me all that I have and my lovely and kind siblings for their love and support of my decision and the path I choose. Indeed they are the best people in my life.

ABSTRACT

In this study, decolorization of Orange II dye using *Enterococcus faecalis* immobilized onto multi-walled carbon nanotube (MWCNT) cement composite was assessed. Multi-walled carbon nanotube (MWCNT) cement composites were made by mixing zeolite (40g), cement (12g), water (8ml) and MWCNT (0.6g) for attachment of biofilm. Biofilm was allowed to develop on the biocomposites for 7 days. The efficiency of decolorization of orange II dye was examined in the presence and absence of biofilm. By comparing the system with and without biofilm, the system without biofilm (using free suspended cells) achieved the highest percentage removal of orange II dye (97.69%). Unexpectedly the lowest percentage of dye decolorization was achieved for the system using cells immobilized onto Hardened Cement Paste + MWCNT (72.74%). This was most likely due to the cytotoxic effect of MWCNT on the bacterial cells. As expected maximum decolorization for all the systems occurred in anaerobic condition and after entering the aerobic condition decolorization had reached a steady state because of the competition between oxygen and dye.

ABSTRAK

Dalam kajian ini, penyahwarnaan pewarna *Orange II* menggunakan *Enterococcus faecalis* yang disekat gerak pada tiub nano karbon berdingg ganda (*MWCNT*) simen komposit telah dikaji. *MWCNT* simen komposit telah dibuat dengan mencampurkan zeolit (40 g), simen (12 g), air (8 mL) dan *MWCNT* (0.6 g). Lapisan biofilem telah dibangunkan pada komposit selama 7 hari. Kecekapan penyahwarnaan pewarna *Orange II* telah dikaji dalam kehadiran dan ketiadaan biofilem. Kajian mendapati bahawa sistem tanpa biofilem (menggunakan sel bebas) mencapai peratusan penyingkiran tertinggi pewarna *Orange II* (97,69%) manakalaperatusan terendah telah dicapai untuk sistem menggunakan sel-sel yang telah disekat gerak dan membentuk biofilem pada *MWCNT* simen komposit (72,74%). Pencapaian peratusan penyahwarnaan yang rendah ini mungkin disebabkan oleh kesan sitotoksik *MWCNT* pada sel-sel bakteria. Sebagaimana yang dijangka proses penyahwarnaan maksimum telah dicapai dalam keadaan anaerobik. Walaubagaimanpun proses ini telah mencapai keadaan mantap dalam keadaan aerobik akibat persaingan antara oksigen dan pewarna.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	x
	LIST OF FIGURES	xi
1	INTRODUCTION	
	1.1 Introduction	1
	1.2 Background of Study	2
	1.3 Objectives of the Study	3
	1.4 Scope of the Study	4
2	LITERATURE REVIEW	5
	2.1 Textile wastewater	5
	2.2 Dyes	6
	2.2.1 Dye classification	7
	2.3 Azodyes	8
	2.4 Methods of color removal	9
	2.4.1 Physico-chemical methods	9
	2.4.2 Physical methods	10
	2.4.3 Chemical methods	10
	2.4.4 Biological methods	12
	2.5 Decolorization by Fungi	13
	2.6 Mechanism of microbial color removal	14

	2.6.1	Aerobic decolorization	14
	2.6.2	Anaerobic decolorization	15
	2.7	Mixed and single(pure) bacterial cultures	16
	2.8	Aerobic fate of aromatic amines	17
	2.9	Biodegradation of Azo Dyes	18
	2.10	Orange II	23
	2.11	Biodegradation of Orange II Dye	24
	2.12	Immobilized microorganisms	25
	2.13	Carbon Nanotube	25
3		MATERIALS AND METHODS	27
	3.1	Experimental Design	27
	3.2	Microorganism	28
	3.3	Preparation of Stock Solutions	28
	3.4	Medium preparation	28
	3.5	Preparation of starter culture	28
	3.6	Multi-walled carbon nanotube and cement	29
	3.7	Hardened cement paste (HCP) preparation	29
	3.8	Immobilization and Development of Biofilm	29
	3.9	Analysis of Dye Decolorization	30
	3.10	COD Determination	31
	3.11	pH Measurement	31
	3.12	FESEM	31
4		RESULTS AND DISCUSSION	32
	4.1	Introduction	32
	4.2	Development of bioparticles	32
	4.3	Effect of pH on dye decolorization	34
	4.4	Analysis of Dye Decolorization	39
	4.5	Analysis of Chemical Oxygen Demand (COD)	42
	4.6	Microscopy Examination of Biofilm (FESEM)	45
5		CONCLUSION	48
	5.1	Conclusion	48
	5.2	Future works	49
		REFERENCES	50

LIST of TABLES

TABLE NO.	TITLE	PAGE
2.1	properties of dyes classified based on their usage	8
2.2	Advantages and disadvantages of some of the current non-biological methods of dye removal	12
2.3	Some of the advantages and disadvantages of anaerobic treatment	16
4.1	Different ratio and combination of materials to develop bioparticles	33
4.2	Highest decolorization in different systems	35
4.3	Highest COD removal rate among treatment systems	42

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	General view of the fate of azo dyes and aromatic amines in anaerobic-aerobic systems	19
2.2	Proposed mechanism for reduction of azo dyes by whole bacterial cells	22
2.3	structure of Orange II	23
4.1	Bioparticles which used in treatment	33
4.2	Changes in pH for free suspended cells grown in Orange II dye medium at anaerobic condition (37°C) and aerobic condition (25°C)	34
4.3	Changes in pH for HCP, IHCP, HCP + MWCNT and IHCP + MWCNT grown in Orange II dye medium at anaerobic condition (37°C) and aerobic condition (25°C)	35
4.4	Comparison between pH and decolorization of the free suspended cell system	36
4.5	Comparison between pH and decolorization of the HCP system	37
4.6	Comparison between pH and decolorization of the IHCP system	37
4.7	Comparison between pH and decolorization of the HCP + MWCNT system	38
4.8	Comparison between pH and decolorization of the IHCP + MWCNT system	38
4.9	Decolorization of Orange II dye in free suspended cells and immobilized hardened cement pastes with MWCNT in an anaerobic-aerobic system	40
4.10	Decolorization of Orange II dye in hardened cement pastes and immobilized hardened cement pastes in an anaerobic-aerobic system	41
4.11	Decolorization of Orange II dye in hardened cement pastes with MWCNT and immobilized hardened cement pastes with MWCNT in an anaerobic-aerobic system	41

4.12	Comparison of COD removal percentage between hardened cement pastes and immobilized hardened cement pastes	43
4.13	Comparison of COD removal percentage between hardened cement pastes with MWCNT and immobilized hardened cement pastes with MWCNT	44
4.14	Comparison of COD removal percentage between immobilized hardened cement pastes and immobilized hardened cement pastes with MWCNT	44
4.15	FESEM image of HCP with 5K magnification	46
4.16	FESEM image of HCP + MWCNT with 10K magnification	46
4.17	FESEM image of IHCP with biofilm after 7 days of treatment with 5K magnification	47
4.18	FESEM image of IHCP + MWCNT and biofilm after 7 days of treatment with 10k magnification	47

CHAPTER 1

INTRODUCTION

1.1 Introduction

Textile wastewater as one of the industrial wastes always draws attention of researchers and scientists. Textile industries are described by high water and chemical consumption as a result of different processes which give rise to noticeable amounts of colored wastewater. Their conventional treatment like the other waste has some steps and is mostly physico-chemical and biological. Textile companies may face a lack of available water sources due to water insufficiency and restrictions of ground water use. In the near future, many textile companies will have to improve wastewater quality to the fresh water standards for reuse purpose.

Tendency to use biological methods has increased in recent years, because of their ability for removing the organic materials and the problems that chemicals cause to nature. Furthermore for using microorganisms which can degrade these waste, some methods have been suggested but most of them are time consuming, expensive and low efficient versus chemicals.

Dyes are the visual part of textile wastewater and because of that take most attention at first glance and the usual methods are not efficient in decolorizing them and it seems that microorganisms are more important in this part. Types of the dyes

according to their application and chemical composition are diverse and due that we need to use different kind of microorganisms.

Because of that, biological methods should be consist either of one species of special microorganisms or of various kinds of different microorganisms that can establish an ecosystem suitable for dye omitting. At the end dyes can be degraded into simpler compounds and are finally mineralized to water and carbon dioxide by a wide variety of aerobic or anaerobic organisms.

Microorganisms need time, stability and physical support to work efficiently on wastes from effluent and for this matter one of the method is immobilizing enzyme or cell to greatly restrict the freedom of movement of them. So immobilization provides a physical support for cells. Therefore the first consideration is to decide on the support material and then the main method of immobilization.

Lately the use of nanomaterials in bio-related research has increased sharply. Nanomaterials due to their dimension have specific characteristics which make them valuable and provide new ideas for novel research.

1.2 Background of Study

No ideal support material or method of immobilization has emerged to provide a standard for each type of immobilization. Selection of support material and method of immobilization is made by measuring the various attributes of the enzyme or cell application against the properties and limitations of the combined immobilization and support. Several practical aspects should be considered before starting laboratory work to ensure that the final immobilized enzyme or cell preparation is suitable for the planned purpose or application and will operate at optimum effectiveness.

Different methods of immobilization can influence the activity and half-life of a cell-based biotransformation. There are five principal methods for immobilization of enzymes or cells: adsorption, covalent binding, entrapment, encapsulation, and cross-linking.

There are many different support materials such as calcium alginate, polyacrylamide, DEAE-cellulose, glutaraldehyde and agar. Moreover in some research solid and porous materials like sponge have been used. Recently by the emerging and development of nanotechnology, the interest of using them in environmental research regarding to their unique properties has increased.

Azo dyes are the most important groups of synthetic dyes. It had been estimated that about 10% of the dye stuff used during the dyeing processes does not bind to the fibers and is released into the effluents. Because all the industrially produced azo dyes are xenobiotic compounds, it is not surprising that they are recalcitrant in conventional treatment methods (Steffan *et al.*, 2005).

1.3 Objectives of the Study

- To immobilize dye degrading bacterial cells onto hardened cement paste with multi-walled carbon nanotubes (MWCNT).
- To compare the ability of freely suspended and immobilized hardened cement paste with MWCNT to degrade Orange II dye.

1.4 Scope of the Study

The study was concentrated on the decolorization of pure dye solution of Orange II. Composite of zeolite, cement and MWCNT were used to immobilize bacterial cells (*Enterococcus faecalis*) which are able to degrade Orange II. Dye decolorization, pH, optical density (OD) and chemical oxygen demand (COD) were determined and analyzed by standard methods.

REFERENCES

- Abramian, L. & El-Rassy, H. (2009). Adsorption kinetics and thermodynamics of azo-dye Orange II onto highly porous titania aerogel. *Chemical Engineering Journal*, 150, 403-410.
- Adebajo, M.O., Frost, R.L., Klopogge, J.T., Carmody, O., Kokot, S. (2003). Porous materials for oil spill cleanup: a review of synthesis and absorbing properties. *J. Porous Mat.*, 10, 159-170.
- Akasaka, T., Watari, F. (2009). Capture of bacteria by flexible carbon nanotubes. *Acta Biomater.*, 5, 607-12.
- Aksu, Z., Cagatay, S.S. (2006). Investigation of biosorption of Gemazol Turquoise Blue-G reactive dye by dried *Rhizopus arrhizus* in batch and continuous systems. *Sep. Purif. Technol.*, 48, 24-35.
- Alaton, I.A., Balcioglu, I.A., Bahnemann, D.W. (2002). Advanced oxidation of a reactive dyebath effluent: comparison of O₃, H₂O₂/UV-C and TiO₂/UV-A processes. *Water Res.*, 36, 1143-1154.
- Alpat, S.K., Ozbayrak, O., Alpat, S., Akcay, H. (2008). The adsorption kinetics and removal of cationic dye, Toluidine Blue O, from aqueous solution with Turkish zeolite. *J. Hazard. Mater.*, 151, 213-220.

- Anjaneyulu, Y., Sreedhara Chary, N., Raj, D.S.S. (2005). Decolourization of industrial effluents – available methods and emerging technologies – a review. *Rev. Environ. Sci. Bio/Technol.*, 4, 245–273.
- Bakshi, D.K., Sharma, P. (2003). Genotoxicity of textile dyes evaluated with Ames test and rec-assay. *J. Environ. Pathol. Toxicol. Oncol.*, 22 (2), 101–109.
- Banat IM, Nigam P, Singh D, Marchant R. (1996). Microbial decolourization of textile dyes containing effluents: a review. *Bioresour Technol*, 58, 217–27.
- Bansal, R.C., Goyal, M. (2005). Activated Carbon Adsorption. Taylor & Francis Group, Boca Raton.
- Barragan, B.E., Costa, C., Carmen Marquez, M. (2007). Biodegradation of azo dyes by bacteria inoculated on solid media. *Dyes Pigments*, 75, 73–81.
- Bayramoglu, G., Bektas_, S., Arica, M.Y. (2003). Biosorption of heavy metal ions on immobilized white-rot fungus *Trametes versicolor*. *J. Hazard. Mater.*, B101, 285–300.
- Beydilli, M.I., Pavlostathis, S.G., Tincher, W.C. (1998). Decolorization and toxicity screening of selected reactive azo dyes under methanogenic conditions. *Water Sci. Technol.*, 38, 225–232.
- Bhattacharyya, K.G., Sarma, A. (2003). Adsorption characteristics of the dye, Brilliant Green, on Neem leaf powder. *Dyes Pigments*, 57, 211–222.
- Bras, R., Ferra, I.A., Pinheiro, H.M., Goncalves, I.C. (2001). Batch tests for assessing decolourisation of azo dyes by methanogenic and mixed cultures. *Journal of Biothechnology*, 89, 155–62.

- Bromley-Challenor, K.C.A., Knapp, J.S., Zhang, Z., Gray, N.C.C., Hetheridge, M.J., Evans, M.R. (2000). Decolorization of an azo dye by unacclimated activated sludge under anaerobic conditions. *Water Res.*, 34, 4410–4418.
- Bumpus, J. A. & Ved Pal, S. (1995). Microbial degradation of azo dyes. *Progress in Industrial Microbiology*. Elsevier.
- Carliell, C.M., Barclay, S.J., Naidoo, N., Buckley, C.A., Mulholland, D.A., Senior, E. (1995). Microbial decolourisation of a reactive azo dye under anaerobic conditions. *Water SA.*, 21 (1), 61-69.
- Carliell, C.M., Barclay, S.J., Buckley, C.A., 1996. Treatment of exhausted reactive dye bath effluent using anaerobic digestion: laboratory and full scale trials. *Water S.A.*, 22, 225-233.
- Champagne, P.P., Ramsay, J.A. (2005). Contribution of manganese peroxidase and laccase to dye decoloration by *Trametes versicolor*. *Appl. Microbiol. Biotechnol.*, 69, 276–285.
- Chang J.S, Chou C, Chen S.Y. (2001a). Decolorization of azo dyes with immobilized *Pseudomonas luteola*. *Proc Biochem*, 36, 757–63.
- Chang, J.S, Chou, C., Lin, Y.C., Lin, P.J., Ho, J.Y, Hu, T.L. (2001b). Kinetic characteristics of bacterial azo-dye decolorization by *Pseudomonas luteola*. *Water Research*, 35(12), 2841–50.
- Cheetham, P.S.J., Bucke, C. (1984). Immobilisation of microbial cells and their use in waste water treatment. In: Grainger JM, Lynch JM, editors. *Microbial methods for environmental biotechnology*. London, UK: Academic Press Inc., p. 219.

- Chen K-C, Wu J-Y, Liou D-J, Hwang S-CJ. (2003). Decolorization of textile dyes by newly isolated bacterial strains. *J Biotechnol*, 10, 57–68.
- Chen, B.Y., Chen, S.Y., Chang, J.S. (2005). Immobilized cell fixed-bed bioreactor for wastewater decolorization. *Process Biochem.*, 40, 3434–3440.
- Christie, R.M. (2007). Environmental Aspects of Textile Dyeing. Woodhead, Boca Raton, Cambridge.
- Chung, K.T., Fulk, G.E., Egan, M. (1978). Reduction of azo dyes by intestinal anaerobes. *Appl. Environ. Microbiol.*, 35 (3), 558–562.
- Chung, K.T., Stevens, S.E.J., Cerniglia, C.E. (1992). The reduction of azo dyes by the intestinal microflora. *Crit. Rev. Microbiol.*, 18, 175–197.
- Claus, H., Faber, G., Koenig, H. (2002). Redox-mediated decolorization of synthetic dyes by fungal laccases. *Appl. Microbiol. Biotechnol.*, 59, 672–678.
- Coughlin, M.F., Kinkle, B.K., Bishop, P.L. (2002). Degradation of acid orange 7 in an anaerobic biofilm. *Chemosphere*, 46, 11–19.
- Coughlin, M.F., Kinkle, B.K., Bishop, P.L. (2003). High performance degradation of azo dye Acid Orange 7 and sulfanilic acid in a laboratory scale reactor after seeding with cultured bacterial strains. *Water Res.*, 37 (11), 2757–2763.
- Couto, S.R., Sanroman, M.A., Hofer, D., Guñbitz, G.M. (2004). Stainless steel sponge: a novel carrier for the immobilization of the white-rot fungus *Trametes hirsuta* for decolourization of textile dyes. *Bioresour. Technol.*, 95, 67–72.

- Crini, G. (2006). Non-conventional low-cost adsorbents for dye removal: a review. *Bioresour. Technol.*, 97, 1061–1085.
- Cruz, A., Buitron, G. (2001). Biodegradation of Disperse Blue 79 using sequenced anaerobic/aerobic biofilters. *Water Sci. Technol.* 44 (4), 159–166.
- Delee, W., O' Neill, C., Hawkes, F.R., Pinheiro, H.M. (1998). Anaerobic treatment of textile effluents: a review. *J. Chem. Technol. Biotechnol.*, 73, 323–335.
- Dilek FB, Taplamacioglu HM, Tarlan E. (1999). Color and AOX removal from pulping effluents by algae. *Appl Microbiol Biotechnol*, 52, 585–91.
- Dos Santos, A.B., Madrid, M.P., de Bok, F.A.M., Stams, A.J.M., Van Lier, J.B., Cervantes, F.J. (2006). The contribution of fermentative bacteria and methanogenic archaea to azo dye reduction by thermophilic anaerobic consortium. *Enzyme Microbiol. Technol.*, 39, 38–46.
- Dos Santos, A. B., Cervantes, F. J. & Van Lier, J. B. (2007). Review paper on current technologies for decolourisation of textile wastewaters: Perspectives for anaerobic biotechnology. *Bioresource Technology*, 98, 2369–2385.
- Duran, N., Rosa, M.A., D'Annibale, A., Gianfreda, L. (2002). Applications of laccases and tyrosinases (phenoloxidases) immobilized on different supports: a review. *Enzyme Microbiol. Technol.*, 31, 907–931.
- Fang, H., Wenrong, H., Yuezhong, L. (2004). Investigation of isolation and immobilization of a microbial consortium for decoloring of azo dye 4BS. *Water Res.*, 38, 3596–3604.

- Field, J.A., Stams, A.J.M., Kato, M., Schraa, G. (1995). Enhanced biodegradation of aromatic pollutants in cocultures of anaerobic and aerobic bacterial consortia. *Antonie van Leeuwenhoek Int. J. Gen. Mol. Microbiol.*, 67, 47–77.
- Field, J.A., Cervantes, F.J., Van der Zee, F.P., Lettinga, G. (2000). Role of quinones in the biodegradation of priority pollutants: a review. *Water Sci. Technol.*, 42, 215–222.
- FitzGerald, S.W., Bishop, P.L. (1995). Two stage anaerobic/aerobic treatment of sulfonated azo dyes. *J. Environ. Sci. Health*, A30, 1251–76.
- Frijters, C.T.M.J., Vos, R.H., Scheffer, G., Mulder, R. (2006). Decolorizing and detoxifying textile wastewater, containing both soluble and insoluble dyes, in a full scale combined anaerobic/aerobic system. *Water Res.*, 40, 1249–1257.
- Fu, Y.Z., Viraraghavan, T. (2001a). Fungal decolorization of dye wastewaters: a review. *Bioresour. Technol.*, 79, 251–262.
- Gingell, R., Walker, R. (1971). Mechanisms of azo reduction by *Streptococcus faecalis*. II. The role of soluble flavins. *Xenobiotica*, 1(3), 231–9.
- Goncalves, I., Gomes, A., Brás, R., Ferra, M.I.A., Amorim, M.T.P., Porter, R.S. (2000). Biological treatment of effluent containing textile dyes. *J Soc Dyers Colour*, 6, 393–7.
- Gupta, V. K. & Suhas. (2009). Application of low-cost adsorbents for dye removal - A review. *Journal of Environmental Management*, 90, 2313–2342.
- Hai, F.I., Yamamoto, K., Fukushima, K. (2007). Hybrid treatment systems for dye wastewater. *Crit. Rev. Env. Sci. Technol.*, 37, 315–377.

- Henriques, I. D. S., Love, N. G. (2007). The role of extracellular polymeric substances in the toxicity response of activated sludge bacteria to chemical toxins. *Water Research*, 41, 4177-4185.
- Hu TL. (1994). Decolorization of reactive azo dyes by transformation with *Pseudomonas luteola*. *Biores Technol*, 49: 47–51.
- Hu TL. (2001). Kinetics of azoreductase and assessment of toxicity of metabolic products from azo dyes by *Pseudomonas luteola*. *Water Sci Technol*, 43(2), 261–9.
- Hunger, K. (2003). Industrial Dyes: Chemistry, Properties, Applications. Wiley-VCH, Weinheim; [Cambridge].
- Husain, Q. (2006). Potential applications of the oxidoreductive enzymes in the decolorization and detoxification of textile and other synthetic dyes from polluted water: a review. *Crit. Rev. Biotechnol.*, 26, 201–221.
- Imamura, K., Ikeda, E., Nagayasu, T., Sakiyama, T., Nakanishi, K. (2002). Adsorption behavior of methylene blue and its congeners on a stainless steel surface. *J. Colloid Interface Sci.*, 245, 50–57.
- Isik, M., Sponza, D.T. (2004b). Decolorization of azo dyes under batch anaerobic and sequential anaerobic/aerobic conditions. *J. Environ. Sci. Health, Part A-Toxic/Hazard. Subst. Environ. Eng.*, 39 (4), 1107–1127.
- Klibanov, A.M., Morris, E.D. (1981). Horsheradish peroxidise for the removal of carcinogenic aromatic amines from water. *Enzyme Microb. Technol.*, 3, 119–122.

- Kalyuzhnyi, S., Sklyar, V. (2000). Biomineralisation of azo dyes and their breakdown products in anaerobic–aerobic hybrid and UASB reactors. *Water Sci. Technol.*, 41 (12), 23–30.
- Kang, S., Herzberg, M., Rodrigues, D.F., Elimelech, M. (2008). Antibacterial effects of carbon nanotubes: size does matter. *Langmuir.*, 24, 6409–13.
- Kapdan, I.K., Tekol, M., Sengul, F. (2003). Decolorization of simulated textile wastewater in an anaerobic–aerobic sequential treatment system. *Process Biochem.*, 38 (7), 1031–1037.
- Kapdan, I.K., Alparslan, S. (2005). Application of anaerobic–aerobic sequential treatment system to real textile wastewater for color and COD removal. *Enzyme Microb. Technol.*, 36 (2–3), 273–279.
- Keck, A., Klein, J., Kudlich, M., Stolz, A., Knackmuss, H-J., Mattes, R. (1997). Reduction of azo dyes by redox mediators originating in the naphthalenesulfonic acid degradation pathway of *Sphingomonas sp.* strain BN6. *Applied and Environmental Microbiology*, 63(9), 3684–3690.
- Khare, U. K., P. Bose, and P. S. Vankar. (2007). Impact of Ozonation on Subsequent Treatment of Azo Dye Solution. *J. Chem. Technol. Biotechnol.*, 82, 1012.
- Khehra, M. S., Saini, H. S., Sharma, D. K., Chadha, B. S., Chimni, S. S. (2005). Decolorization of various azo dyes by bacterial consortium. *Dyes and Pigments*, 67, 55-61.
- Kirby, N., Marchant, R., McMullan, G. (2000). Decolourisation of synthetic textile dyes by *Phlebia tremellosa*. *FEMS Microbiol. Lett.*, 188, 93–96.

- Knackmuss, H.J. (1996). Basic knowledge and perspectives of bioelimination of xenobiotic compounds. *J. Biotechnol.*, 51, 287–295.
- Kudlich, M., Keck, A., Klein, J., Stolz, A. (1997). Localization of the enzyme system involves in anaerobic reduction of azo dyes by *Sphingomonas sp.* strain BN6 and effect of artificial redox mediators on the rate of azo dye reduction. *Applied and Environmental Microbiology*, 63, 3691–3694.
- Kudlich, M., Hetheridge, M.J., Knackmuss, H.J., Stolz, A. (1999). Autoxidation reactions of different aromatic oaminohydroxynaphthalenes that are formed during the anaerobic reduction of sulfonated azo dyes. *Environ. Sci. Technol.*, 33 (6), 896–901.
- Kulla, H.G. (1981). Aerobic bacterial degradation of azo dyes. *FEMS Microbiol. Lett.*, 12, 387–399.
- Lee, J.-W., Choi, S.-P., Thiruvengkatachari, R., Shim, W.-G., Moon, H. (2006). Evaluation of the performance of adsorption and coagulation processes for the maximum removal of reactive dyes. *Dyes Pigments*, 69, 196–203.
- Libra, J. A., M. Borchert, L. Vigelahn, and T. Storm. (2004). Two Stage Biological Treatment of a Diazo Reactive Textile Dye and the Fate of the Dye Metabolites. *Chemosphere*, 56, 167.
- Li, G. Y., Wang, P. M. & Zhao, X. (2005). Mechanical behavior and microstructure of cement composites incorporating surface-treated multi-walled carbon nanotubes. *Carbon*, 43, 1239–1245.

- Lourenco, N.D., Novais, J.M, Pinheiro, H.M. (2000). Reactive textile dye colour removal in a sequencing batch reactor. *Water Science and Technology*, 42(5-6), 321–8.
- Madigan, M.T., Martinko, J.M., Parker, J. (2003). Brock Biology of Microorganisms, 10th ed. Prentice-Hall Inc., Simon & Schuster/A Viacom Company, Upper Saddle River, New Jersey, USA.
- Manu, B., Chaudhari, S. (2002). Anaerobic decolorisation of simulated textile wastewater containing azo dyes. *Bioresource Technology*, 82, 225-231.
- Maurya, N.S., Mittal, A.K., Cornel, P., Rother, E. (2006). Biosorption of dyes using dead macro fungi: effect of dye structure, ionic strength and pH. *Bioresour. Technol.*, 97, 512–521.
- Mauter, S.M., Elimelech, M. (2008). Environmental applications of carbon based nanomaterials. *Environmental Science & Technology J.*, 42, 5843–5859.
- Menon, V.C., Komarneni, S. (1998). Porous adsorbents for vehicular natural gas storage: a review. *J. Porous Mat.*, 5, 43-58.
- Metcalf and Eddy. (2003). Wastewater Engineering: Treatment and Reuse, 4th ed. McGraw-Hill, New York, USA.
- Mezohegyi, G., Kolodkin, A., Castro, U.I., Bengoa, C., Stuber, F., Font, J., Fabregat, A., Fortuny, A.. (2007). Effective anaerobic decolorization of azo dye Acid Orange 7 in continuous up flow packed-bed reactor using biological activated carbon system. *Ind. Eng. Chem. Res.*, 46, 6788–6792.

- Moawad, H., El-Rahim, W.M., Khalafallah, M. (2003). Evaluation of biotoxicity of textile dyes using two bioassays. *J. Basic Microbiol*, 43 (3), 218–229.
- Myers, C.R., Myers, J.M. (1992). Localization of cytochromes to the outer membrane of anaerobically grown *Shewanella putrefaciens* MR-1. *Journal of Bacteriology*, 174(11), 3429–38.
- Nepal, D., Balasubramanian, S., Simonian, A.L., Davis, V.A. (2008). String antimicrobial coatings: single walled carbon nanotubes armored with biopolymers. *Nano Lett.*, 8, 1896–902.
- Nigam, P., Banat, I.M., Singh, D., Marchant, R. (1996). Microbial process for the decolorization of textile effluent containing azo, diazo and reactive dyes. *Process Biochemistry*, 31(5), 435–442.
- Niu, H., Xu, X.S., Wang, J.H. (1993). Removal of lead from aqueous solutions by penicillin biomass. *Biotechnol. Bioeng.*, 42, 785–787.
- Nur, H., Manan, A.F.N.A., Wei, L.K., Muhid, M.N.M., Hamdan, H. (2005). Simultaneous adsorption of a mixture of paraquat and dye by NaY zeolite covered with alkylsilane. *J. Hazard. Mater.*, 117, 35–40.
- O'Neill, C., Freda, R.H., Dennis, L.H., Nidia DL, Helena MP, Wouter D. (1999). Colour in textile effluents, sources, measurements, discharge contents and simulation: a review. *J Chem Technol Biotechnol*, 74, 1009–1018.
- O'Neill, C., Lopez, A., Esteves, S., Hawkes, F.R., Hawkes, D.L., Wilcox, S. (2000). Azo-dye degradation in an anaerobic-aerobic treatment system operating on simulated textile effluent, *Appl. Microbiol. Biotechnol.*, 53, 249–254.

- Ong, S.-A., Toorisaka, E., Hirata, M. & Hano, T. (2005a). Decolorization of azo dye (Orange II) in a sequential UASB-SBR system. *Separation and Purification Technology*.
- Ong, S.-A., Toorisaka, E., Hirata, M. & Hano, T. (2005b). Treatment of azo dye Orange II in aerobic and anaerobic-SBR systems. *Process Biochemistry*, 40, 2907-2914.
- Pagga, U., Brown, D. (1986). The degradation of dyestuffs: Part II. Behaviour of dyestuffs in aerobic biodegradation tests. *Chemosphere*, 15, 479-491.
- Panswad, T., Iamsamer, K., Anotai, J. (2001a). Decolorisation of azo-reactive dye by polyphosphate and glycogen-accumulating organisms in an anaerobic-aerobic sequencing batch reactor. *Bioresour Technol.*, 76, 151-159.
- Pavlidis, I., Tsoufis, T., Gournis, D. & Stamatis, H. (2009). Enzyme immobilization on chemically functionalized multi-walled carbon nanotubes. *New Biotechnology*, 25, S131-S131.
- Pazarlioglu, N.K., Urek, R.O., Ergun, F. (2005). Biodecolourization of Direct Blue 15 by immobilized *Phanerochaete chrysosporium*. *Process Biochem.*, 40, 1923-1929.
- Pearce, C. I., Lloyd, J. R. & Guthrie, J. T. (2003). The removal of colour from textile wastewater using whole bacterial cells: a review. *Dyes and Pigments*, 58, 179-196.
- Pearce, C.I., Christie, R., Boothman, C., Von Canstein, H., Guthrie, J.T., Lloyd, J.R. (2006). Reactive azo dye reduction by *Shewanella* Strain J18 143. *Biotechnol. Bioeng.*, 95, 692-703.

- Pinheiro, H.M., Touraud, E., Thomas, O. (2004). Aromatic amines from azo dye reduction: status review with emphasis on direct UV spectrophotometric detection in textile industry wastewaters. *Dyes Pigm.*, 61 (2), 121–139.
- Plumb, J.J., Bell, J., Stuckey, D.C. (2001). Microbial populations associated with treatment of an industrial dye effluent in an anaerobic baffled reactor. *Applied and Environmental Microbiology*, 67(7), 3226–35.
- Rai, H.S., Bhattacharyya, M.S., Singh, J., Bansal, T.K., Vats, P., Banerjee, U.C. (2005). Removal of dyes from the effluent of textile and dyestuff manufacturing industry: a review of emerging techniques with reference to biological treatment. *Crit. Rev. Env. Sci. Technol.*, 35, 219–238.
- Rajaguru, P., Kalaiselvi, K., Palanivel, M., Subburam, V. (2000). Biodegradation of azo dyes in a sequential anaerobic–aerobic system. *Appl. Microbiol. Biotechnol.*, 54, 268–273.
- Ramakrishna, K.R., Viraraghavan, T. (1997). Dye removal using low cost adsorbents. *Water Sci. Technol.*, 36, 189–196.
- Rau, J., Knackmuss, H.J., Stolz, A. (2002). Effects of different quinoid redox mediators on the anaerobic reduction of azo dyes by bacteria. *Environ. Sci. Technol.*, 36, 1497–1504.
- Rau, J., Stolz, A. (2003). Oxygen-sensitive nioreductases NfsA and NsfB of *Escherichia coli* function under anaerobic conditions as lawsone-dependent azo reductases. *Appl. Environ. Microbiol.*, 69 (6), 3448–3455.

- Robinson, T., McMullan, G., Marchant, R., Nigam, P. (2001). Remediation of dyes in textile effluent: a critical review on current treatment technologies with a proposed alternative. *Bioresource Technology*, 77, 247-255.
- Russ, R., Rau, J., Stolz, A. (2000). The function of cytoplasmic flavin reductases in the reduction of azodyes by bacteria. *Applied and Environmental Microbiology*, 66(4), 1429–1434.
- Sani, R., Banerjee, U. (1999b). Decolorization of triphenylmethane dyes and textile and dye-stuff effluent by *Kurthia sp.* *Enzyme Microb. Technol.*, 24, 433–437.
- Sarsour, J., Janitza, J., Gahr, F. (2001). Biological degradation of dye-containing wastewater (Biologische Abbau farbstoffhaltiger Abwässer). *Wasser Luft Boden*, 6, 44–46.
- Sharma DK, Saini HS, Singh M, Chimni SS, Chadha BS. (2004). Isolation and characterization of microorganisms capable of decolorizing various triphenylmethane dyes. *J Basic Microbiol*, 44(1), 59–65.
- Sharma, P., Singh, L. (2009). biodegradation of orange II dye by *Phanerochaete chrysosporium* in simulated wastewater. *Journal of Scientific and Industrial Research*, 68 157-161.
- Shaw, C.B., Carliell, C.M., Wheatley, A.D. (2002). Anaerobic/aerobic treatment of coloured textile effluents using sequencing batch reactors. *Water Res.*, 36 (8), 1993–2001.

- Silva, A. C., PIC, J. S., Sant'anna Jr, G. L. & Dezotti, M. (2009) Ozonation of azo dyes (Orange II and Acid Red 27) in saline media. *Journal of Hazardous Materials*, 169, 965-971.
- Simmons, P., Tobin, J.M., Singleton, I. (1995). Consideration of the use of commercially available yeast biomass for the treatment of metal containing effluents. *J. Ind. Microbiol.*, 14, 240–246.
- Sosath, F., Libra, J.A. (1997). Purification of wastewaters containing azo dyes (Biologische Behandlung von synthetischen Abwässern mit Azofarbstoffen). *Acta Hydrochim. Hydrobiol.*, 25 (5), 259–264.
- Specht, K., Platzek, T. (1995). Textile dyes and finishes-remarks to toxicological and analytical aspects. *Dtsch. Lebensm. Rundsch.*, 91, 352–359.
- Sponza, D.T. (2002). Necessity of toxicity assessment in Turkish industrial discharges (examples from metal and textile industry effluents). *Environ. Monit. Assess.*, 73 (1), 41–66.
- Sponza, D.T., Isik, M. (2002). Decolorization and azo dye degradation by anaerobic/aerobic sequential process. *Enzym. Microb. Technol.*, 31, 102–110.
- Steffan, S., Bardi, L. & Marzona, M. (2005) Azo dye biodegradation by microbial cultures immobilized in alginate beads. *Environment International*, 31, 201-205.
- Stolz A. (2001). Basic and applied aspects in the microbial degradation of azo dyes. *Appl Microbiol Biotechnol.*, 56, 69–80.

- Supaka, N., Juntongjin, K., Damronglerd, S., Delia, M.L., Strehaiano, P. (2004). Microbial decolorization of reactive azo dyes in a sequential anaerobic–aerobic system. *Chem. Eng. J.*, 99, 169–176.
- Sutherland, I. W. (2001). The biofilm matrix – an immobilized but dynamic microbial environment. *Trends in Microbiology*, 9, 222–227.
- Tamburlini, G., Ehrenstein, O.V., Bertolini, R.. (2002). Children's health and environment: a review of evidence. Environmental Issue Report No. 129, WHO/European Environment Agency, WHO Geneva, 223.
- Tan, N.C.G., Field, J.A. (2000). Biodegradation of sulfonated aromatic compounds. In: Lens, P., Hulshoff-Pol, L.W. (Eds.), *Environmental Technologies to Treat Sulfur Pollution. Principles and Engineering*. IWA Publishing, London, pp. 377–392.
- Tan, N.C.G. (2001). Integrated and sequential anaerobic/aerobic biodegradation of azo dyes. Ph.D. Thesis, Agrotechnology and Food Sciences, Sub-department of Environmental Technology, Wageningen University, Wageningen, The Netherlands.
- Tsai, Y.-C., Chen, S.-Y. & Liaw, H.-W. (2007). Immobilization of lactate dehydrogenase within multiwalled carbon nanotube-chitosan nanocomposite for application to lactate biosensors. *Sensors and Actuators B: Chemical*, 125, 474-481.
- Vandevivere, P.C., Bianchi, R., Verstraete, W. (1998). Treatment and reuse of wastewater from the textile wet-processing industry: review of emerging technologies. *J. Chem. Technol. Biotechnol.*, 72, 289–302.

- Van der Zee, F.P., Lettinga, G., Field, J.A. (2001b). Azo dye decolourisation by anaerobic granular sludge. *Chemosphere*, 44 (5), 1169–1176.
- Van der Zee, F.P., Bisschops, I.A.E., Lettinga, G., Field, J.A. (2003). Activated carbon as an electron acceptor and redox mediator during the anaerobic biotransformation of azo dyes. *Environ. Sci. Technol.*, 37, 402–408.
- Van der Zee, F. P., Villaverde, S. (2005). Combined anaerobic–aerobic treatment of azo dyes—A short review of bioreactor studies. *Water Research*, 39, 1425–1440.
- Vaseashta A., Mihailescu I. N. (2008). Functionalized Nanoscale Materials. *Devices and Systems*, Springer, 978-1-4020-8902-2 (Print) 978-1-4020-8903-9 (Online).
- Vijayaraghavan, K. and Y. S. Yun. (2007). Utilization of Fermentation Waste (*Corynebacterium glutamicum*) for Biosorption of Reactive Black 5 from Aqueous Solution. *J. Hazard. Mater.*, 141, 45.
- Wesenberg, D., Kyriakides, I., Agathos, S.N. (2003). White rot fungi and their enzymes for the treatment of industrial dye effluents. *Biotechnol. Adv.*, 22, 161–187.
- Willmott, NJ. (1997). The use of bacteria–polymer composites for the removal of colour from reactive dye effluents. PhD thesis, UK: University of Leeds.
- Wingender, J., Neu, T. R., & Flemming, H.-C. (1999). What are bacterial extracellular polymeric substances? In J. Wingender, T. R. Neu, & H.-C. Flemming (Eds.), *Microbial extracellular polymeric substances – Characterization, structure and function* (pp. 1–19).

- Wuhrmann K, Mechsner K, Kappeler T. (1980). Investigations on rate determining factors in the microbial reduction of azo dyes. *Eur J Appl Microbiol Biotechnol*, 9, 325–38.
- Yoo, E.S., Libra, J., Adrian, L. (2001). Mechanism of decolorization of azo dyes in anaerobic mixed culture. *Journal of Environmental Engineering*, 127(9), 844–9.
- Zhang, X.Q., Bishop, P.L. (2003). Biodegradability of biofilm extracellular polymeric substances. *Chemosphere*, 50, 63–9.
- Zhang, S.J., Yang, M., Yang, Q.X., Zhang, Y., Xin, B.P., Pan, F. (2003). Biosorption of reactive dyes by the mycelium pellets of a new isolate of *Penicillium oxalicum*. *Biotechnol. Lett.*, 25, 1479–1482.
- Zheng Z, Levin RE, Pinkham JL, Shetty K. (1999). Decolorization of polymeric dyes by a novel *Penicillium* isolate. *Proc Biochem*, 34, 31–7.
- Zhou W, Zimmermann W. (1993). Decolorization of industrial effluents containing reactive dyes by *actinomycetes*. *FEMS Microbiol Lett*, 107, 157–62.
- Zimmermann, T., Kulla, H., Leisinger, T. (1982). Purification and properties of orange II-azoreductase from *Pseudomonas KF46*. *Experientia*, 38, 1380.
- Zimmermann, T., Gasser, F., Kulla, H., Leisinger, T. (1984). Comparison of two bacterial azoreductases acquired during adaptation to growth on azo dyes. *Arch. Microbiol.*, 138, 37–43.