

# Non-convulsive status epilepticus in adults – an overview

■ S. Rüegg

Division of Clinical Neurophysiology, Department of Neurology, University Hospital Basel

## Summary

Rüegg S. Non-convulsive status epilepticus in adults – an overview. *Schweiz Arch Neurol Psychiatr.* 2008;159:53–83.

Status epilepticus (SE) is the most frequent neurological emergency requiring admission on the intensive care unit (ICU). The classification of status epilepticus may be dichotomised into the “convulsive” and the “non-convulsive” form. Whereas the diagnosis of generalised convulsive status epilepticus (GCSE) is easily made in view of the evident manifestation, the diagnosis of non-convulsive status epilepticus (NCSE) may become very difficult according to its non-spectacular and protean presentation. Non-convulsive status epilepticus is a common cause of altered mental status and delirium; and a substantial number of comatose patients in the intensive care unit may suffer from NCSE. The most important part of making a diagnosis of NCSE is to think of it at all. The definite diagnosis of NCSE is dependent on electroencephalographic (EEG) confirmation. Repetitive or, optimally, continuous EEG recording may help to closely monitor seizure activity and to guide the therapy in order to protect the patient from under- and/or over-treatment. The interpretation of EEG recordings of NCSE may become challenging because of numerous artifacts caused by the patient, caregivers and devices, and because of trace elements of ambiguous significance and of waveforms mimicking epileptic activity.

Non-convulsive status epilepticus was historically subdivided into the groups of complex-partial status epilepticus (focal; CPSE) and absence status (generalised; AS). Most recently, a classification

proposed by the International League Against Epilepsy (ILAE) has subdivided focal NCSE into aura continua (non-convulsive simple partial SE with maintained consciousness) and into dyscognitive SE (with impaired consciousness) of

## abbreviations

AED	antiepileptic drug
AS	absence status
BD	benzodiazepines
CISE	critical illness status epilepticus
CPR	cardiopulmonary resuscitation
CPSE	complex-partial status epilepticus
CSE	convulsive status epilepticus
CSF	cerebrospinal fluid
CT	computer tomography
CYP450	cytochrome P 450
DTI	diffusion tensor imaging
DWI	diffusion-weighted imaging
EEG	electroencephalography
GCSE	generalised convulsive status epilepticus
HIV	human immunodeficiency virus
ICU	intensive care unit
IGE	idiopathic generalised epilepsy
LEV	levetiracetam
LZP	lorazepam
MDL	midazolam
MRI	magnetic resonance imaging
MSE	myoclonic status epilepticus
NCSE	non-convulsive status epilepticus
PET	positron emission tomography
PHT	phenytoin
PLED	periodic lateralised epileptiform discharges
RSE	refractory status epilepticus
SE	status epilepticus
SGSE	secondarily generalised status epilepticus
SPECT	single-photon emission computer tomography
TPW	triphasic waves
VPA	valproic acid

Correspondence:  
Stephan Rüegg, MD  
Division of Clinical Neurophysiology  
Department of Neurology  
University Hospital  
Petersgraben 4  
CH-4031 Basel  
e-mail: srueegg@uhbs.ch

mesial temporal or neocortical origin. Several patients with focal NCSE or AS were known for pre-existing epilepsy. In addition, other types of NCSE have been reported, like “subtle” SE where convulsions in patients with GCSE clinically stop, but the continuous epileptic activity electroencephalographically still persists. Another variant belongs to those patients with critical illness where the serious disorder and/or medications may trigger an acute, prolonged and all too often very-hard-to-treat epileptic condition (“critical illness SE” [CISE]). The majority of these patients will never have had seizures before. Of note, none of the currently proposed classifications has gained broad acceptance and official approval.

Since the need for immediate and intensive treatment of GCSE is beyond debate, the insidiously calm presentation of NCSE may mislead to a much less aggressive therapeutic approach. However, experimental and clinical data suggest that continuous epileptic activity in the brain during NCSE associated with consecutive (glutamatergic) hyperexcitation triggers neurotoxicity and proapoptotic mechanisms which eventually may lead to irreversible brain damage and the onset of a chronic epileptic disorder or to an irreversible neurological deficit. Thus, recent national and international guidelines and reviews recommend immediate and intensive treatment for NCSE as well as for GCSE. With a few exceptions, the first drug is an intravenous benzodiazepine, mainly lorazepam; intravenous phenytoin or valproate should be started without delay. Persistence of NCSE after administration of these two classes of drugs heralds refractory SE (RSE) where single or combinations of drugs in anaesthetic dosages requiring intubation and enteral antiepileptics may be added except for the cases of AS or an underlying terminal disease. The outcome of NCSE is mainly determined by the type, duration, cause and the severity of concomitant diseases.

*Keywords: non-convulsive status epilepticus; intensive care unit; diagnosis; electroencephalography; treatment; benzodiazepines*

## Introduction

Status epilepticus (SE) is the most common neurological emergency requiring treatment in the intensive care unit (ICU). After the first depiction almost 3000 years ago, SE has only gained increased attention since a few decades because of the improved availability of neurophysiological techniques, the better understanding of the mechanisms of seizure propagation and termination,

the advances in intensive care medicine and the growing number of antiepileptic drugs available for treatment. From an operational and clinical viewpoint, SE is divided into two main entities, convulsive SE (CSE) and non-convulsive SE (NCSE). Convulsive SE includes the focal and generalised (GCSE) forms, while NCSE encompasses all other non-convulsive forms of prolonged, not self-limited focal and generalised epileptic manifestations. While a widely accepted definition of GCSE exists, such a definition for NCSE is still lacking. The uncertainties in NCSE considerably augment with respect to the yet unresolved issues of types representing NCSE, the almost unknown epidemiology, the immense number of causes, the difficulties to determine the prognosis and, especially, to the debates about best treatment [1]. Although guidelines approved by specialty boards and neurological societies exist, they lack high-level evidence due to the almost absence of high-quality (level of evidence class I- or II-) studies. Thus, several important issues regarding NCSE urgently need further investigation: How to improve awareness of NCSE? Which clinical signs point to the presence of NCSE? Is there a valid score to build? Which is the best treatment? What is the role of the newer antiepileptic drugs? The value of steroids or other immunotherapeutics in the treatment of NCSE is still unclear, but increasing, mainly experimental, evidence points to an important contribution in RSE. On the more pathophysiological level, the crucial question “Does non-convulsive status epilepticus damage the brain?” is still waiting for a conclusive answer. Basic and clinical research on the mechanisms and treatment of NCSE may enormously profit from the formation of a worldwide and interdisciplinary network. The present article tries to address these issues in the light of the current literature; it has to be considered as a momentarily valuable, but not definitively evident and conclusive statement.

## History

The term “status epilepticus” was introduced in 1824 by Calmeil as the French expression “état de mal” when he was writing his thesis on the experiences he had made at the Salpêtrière and Charenton Asylum [2]. While the ancient Greek-Roman medicine assumed that epilepsy and seizures may be an impressive but not lethal disease, the probably very first account of SE and its serious prognosis can even earlier be found on the Neo-Babylonian Akkadian cuneiform in the 25/26 Sakikku (718–612 BC) [3]. Almost half a

century after Cameil, the term “status epilepticus” appeared for the first time in a textbook of clinical medicine when Bazire translated Trousseau’s book “Leçons en Médecine Clinique” [4]. At that time, it became clear that there were different types of SE, the obvious form with convulsions as well as the more “silent” variants of the “dreamy state” or the “fugue épileptique” or the prepsychotic “twilight states”, well depicted by the eminent French (Charcot, Bourneville, Trousseau) and British (Jackson, Gowers) neurologists. The advent of clinical electroencephalography (EEG) [5] allowed for better differentiation of the clinically almost indistinguishable states of purely postictal alterations and non-convulsive status epilepticus. The typical EEG picture of absence status (AS) was discovered by Lennox and co-workers [6]. The occurrence of long-lasting, continuous neurological signs as a result of epileptic activity recorded by EEG was presented in a hallmark paper by Scott and Masland in 1953 [7]. Penfield and Jasper reported the existence of continuous somato-sensory SE for which they coined the term “aura continua” [8]. Landolt emphasised the usefulness of EEG for the discrimination between AS and complex-partial SE (CPSE) which clinically often are barely to distinguish [9]. Karbowski having organised a symposium on psychomotor status (i.e. CPSE from temporal origin) in 1979 compiled 36 already published and eight own cases, all supported by EEG, in the conference monograph [10]. The Xth Marseille Colloquium on Epilepsy held in 1962 was the first conference devoted to classify the various forms of SE. However, there were “as many forms of SE as types of seizures exist” as Gastaut stated, and the classification was only published 8 years later and received almost no attention [11]. Despite the Santa Monica Meeting on SE in 1980 [12], SE was not included in the still valid “Classification of Epilepsy and Epileptic Syndromes” of the International League Against Epilepsy (ILAE) of 1989 [13]. The most recent attempts of a classification of SE have come from the ILAE and the Cleveland Clinic Group [14, 15], and, with special regard to NCSE, from Shorvon [16, 17].

This lack of a broadly accepted classification is the more unsatisfying, as it impedes research efforts for better epidemiological treatment and prognostic data, and as it obscures prognosis and stronger evidence for the treatment of particularly NCSE.

## Definition

There is no accepted definition of NCSE. One carefully elaborated definition recently proposed

by Shorvon states: “Non-convulsive status epilepticus is a term used to denote a range of conditions in which electrographic seizure activity is prolonged and results in non-convulsive clinical symptoms.” [16, 17] These clinical symptoms mainly include focal deficits like aphasia and amnesia, an impairment of consciousness or an altered behaviour ranging from stupor and coma to delirium and frantic psychosis. Occasionally, automatisms or subtle facial, periorbicular, abdominal and limb twitches, tonic eye deviation or spontaneous (mostly horizontal) nystagmus-like movements, and extensor response(s) of the big toe(s) may be present [18].

How long is prolonged? Thirty minutes were the historically determined duration of GCSE [19]; however, this view has been challenged after the hallmark Veterans Affairs-study published in 1998, evaluating the best first-line treatment of GCSE [20], by Lowenstein et al. who proposed a substantially shorter duration of five minutes [21]. This proposal was adopted by several groups and incorporated into national guidelines, like the Swiss one [22]. In the case of NCSE, the situation is much more equivocal as an expert panel first recommended 60 minutes [19], whereas Jordan six years later proposed 15 to 30 minutes [23] and the most recent comprehensive overview has reiterated a duration of 30 minutes [24]. The author favours a much shorter duration of five minutes identical to the one used in GCSE. This opinion relies on two pivotal facts: first, from the brain’s perspective, it does not make a big difference whether a motor neuron or a non-motor neuron is exposed to prolonged continuous epileptic discharges – both neurons risk to be damaged by excitotoxic mechanisms leading to neuronal death and impaired neurological function. Second, sound data supports the observation that 80% of seizures last less than one minute and more than 90% are terminated within two minutes, i.e., self-terminating seizures persisting for five minutes or more may be very rare and indicate that the usually self-restricting, inherent antiepileptic mechanisms were insufficient and fail to stop the epileptic activity [25–27]. It is also important to underscore that the shorter the time of diagnosis and the start of treatment of NCSE are, the faster NCSE can be stopped and the better its outcome may be. Conversely, delayed diagnosis and therapy of NCSE may result in a vicious circle of progressive brain damage and increasing resistance to the antiepileptic drugs applied which all together then may prolong the duration of NCSE and impair its treatment and prognosis [28–36].

## Types, clinical and electroencephalographic manifestations of NCSE

In general, NCSE in adults is subdivided into the two main groups of focal and generalised NCSE, where simple-partial SE (SPSE) and CPSE, and AS are the principal types, respectively [37]. The most recent proposal of classification has been created by a group of foremost experts in the field of epilepsy on behalf of the ILAE [14]. However, this group too was unable to unanimously settle with their own proposal what may reflect the difficulties encountered in establishing a useful and correct classification of seizure, epilepsy and status epilepticus. The group introduced the term “aura continua” globally encompassing all forms of non-convulsive SPSE where maintained consciousness is a prerequisite. Thereby, they left the more restricted definition of “aura continua” of Penfield who assigned only the forms with continuous somatosensory sensations to “aura continua”. The ILAE also replaced CPSE by the term “dyscognitive” SE making an additional subdivision into those forms attributed to derive from the “mesial temporal” and those from the “neocortical” brain areas. Both forms essentially require impaired consciousness. While the forms originating from the mesial temporal lobe regions may be summarised as “limbic status” manifesting with limbic sensations, the manifestations of the neocortical forms reflect the region of origin involved – in the case of aphasia, for example, the inferior frontal lobe or the superior temporal gyrus.


With respect to the advent of advanced life support and the astounding progress of (critical care)

medicine the author of this review would like to introduce a third group of subtypes of NCSE concerning seriously affected patients in intensive care units (ICU). This group will be called “critical illness SE” (CISE) and more extensively delineated below. An overview of the different types of NCSE is given in figure 1.

The primary generalised NCSE may occur as typical or atypical AS, where the former is present in adults with a known history of idiopathic generalised epilepsy (IGE). The onset of typical AS is almost always precipitated by a medication error, malcompliance or provocative factors, like fever and infection or sleep disturbances; clinically the patients show a confusional state, but often were able to perform complex activities of daily living. Duration can last from half an hour up to weeks [38–40]. The EEG shows the classical generalised 3/sec spike-wave pattern; the interictal background activity is normal. Atypical AS in adults may manifest either in patients with complex epileptic syndromes or “de novo” in patients without history of epilepsy. Clinical differentiation of atypical from typical AS may be difficult; however, the presence of additional subtle signs like eyelid myocloni, perioral automatisms has been proposed to rather point to atypical AS. Some authors suggest that the level of impaired consciousness and the deficits in performing complex tasks may be more pronounced [24]. The EEG shows less regular spike-wave activity of 2.5–4 Hz and the interictal background activity is slow [41]. The “de novo” AS in adults results from benzodiazepine withdrawal in the majority of cases, but it may also occur without any precipitating factor [42]. The EEG is characterised by spike-wave activity of markedly unstable frequency ranging from 0.5 to 4 Hz (fig. 2).

The clinical manifestations of aura continua are as various as there are distinct focal neurological non-convulsive deficits. Aura continua manifests with the typical “plus” symptoms of the involved cortical area and its adjacent tissues in the form of somatosensory perceptions, illusions or hallucinations. The quality of the experience is determined by the localisation of the epileptic focus, for instance, aura continua originating in the temporal lobe manifests by gustatory, olfactory, auditory or rising epigastric sensations, while somatosensory and visual illusions or hallucinations point to a parieto-(temporo-)occipital focus. The unspectacular appearance of aura continua may have led to only a few of publications; nevertheless, these forms of NCSE may be by far more frequently present in clinical practice. Other rare manifestations of aura continua include the opercular type with pharyngeal myoclonus and anarthria [43], the inhibitory

**Figure 1** Types of non-convulsive status epilepticus. The three main groups are the (primary) generalised (red), the focal (blue) and the group of acutely and severely affected patients with markedly impaired consciousness (green).






(paretic, aphasic and amaurotic), emetic, pilomotor and the gelastic types [44–50]. The surface EEG shows focal slowing with repetitive, regular and/or irregular spike-wave discharges of various frequencies (fig. 3 and 4).

Patients with cognitive, purely amnesic limbic or isolated fear SE were at the borderline between aura continua and dyscognitive SE of the mesial temporal subgroup according to the ILAE classification [51–55]. The surface EEG often eludes

**Figure 2** EEG in absence status.

35-year-old computer engineer, having acute phases of confusion, but during them is able to drive a car. No intake of medication nor illicit drugs. Emergency EEG about 5 hours after start of confusional state. Patient walks around and talks intelligibly, however, he was unable to add  $5 + 7$ . Note the aperiodic, but almost continuously generalised epileptic spikes and spike-wave discharges (1–2.5 Hz), pronounced over the frontal regions (fig. 2a). After administration of 5 mg lorazepam, the epileptic activity disappears and is replaced by an almost normal background alpha activity (fig. 2b).



**Figure 3** Aura continua with tonic spasms of the right leg.


38-year-old business man, operated for parietal meningioma six months ago. Levetiracetam tapered to 500 mg/d. Start of feelings of “something is wrong in my head” followed by permanent spasms and short subtle myocloni of the left leg two days before admission. The EEG shows right parieto-central almost continuous beta activity typical of tonic seizures, intermingled with small spike-like discharges with phase reversal over electrode C4, the motor cortex region of the left leg.






**Figure 4** Aura continua with repetitive adverse seizures.

82-year-old woman without pre-existing medical history presenting with speech arrest and prolonged adverse head version to the left. EEG showing slightly irregular/aperiodic spike-wave discharges mainly on the right lateral brain regions with extension to closer to the midline interrupted by short episodes of rhythmic delta waves (fig. 4a–c). This pattern may resemble periodic lateralised epileptiform discharges (PLEDs), but is sometimes slightly irregular (aperiodic) and – together with the clearly ictal semiology – was viewed to reflect epileptic activity which responded well to the application of lorazepam (fig. 4d). Repetitive brain imaging and CSF examinations did not reveal a causative process and the dyscognitive status epilepticus was considered to be of kryptogenic origin.



**Figure 5** Symptomatic dyscognitive status epilepticus.

74-year-old retired high-school teacher suddenly experiencing sensorimotor aphasia, stupor and right-upper-sided eye deviation. EEG showed continuous periodic abortive spike-wave discharges, i.e. PLED, over the left hemisphere with predominance in the parieto-temporo-occipital region and also extending to the contralateral right side (fig. 5a). Administration of lorazepam immediately led to abolition of stupor, eye deviation and global aphasia; however, profound sensory (Wernicke's) aphasia persisted (fig. 5b). Eventually, diffusely expansive glioblastoma multiforme originating from the left parieto-occipital territory was found.






recognition because of the localisation of the focus in the deeply “covered” limbic structures [53] or the ictal activity is represented by rhythmic mid-temporal high-amplitude delta waves [56].

Dyscognitive SE (formerly CPSE) belongs to the group of focal NCSE in which an impairment up to the complete loss of consciousness is mandatory [14, 57]. In addition to the key cognitive and


**Figure 6** Subtle status epilepticus / symptomatic dyscognitive status epilepticus.

55-year-old man suffering from pneumococcal sepsis with meningo-encephalitis, undergoing repetitive tonic-clonic generalised seizures, but later showed periodic episodes of tonic head version to the left. Continuous EEG shows short phases of slow low-amplitude theta-delta activity (fig. 6a) with series of repetitive seizures starting from the right temporo-central region with low-amplitude high-(beta) frequency discharges typical of initiation of tonic seizures (fig. 6b). This activity evolves into higher-amplitude slower epileptic discharges over the next ca 90 sec (fig. 6c–e) with attenuation over the next ca 80 sec (fig. 6e–h) when a new seizure starts from the right parietal region (fig. 6h–l). The patient did not regain consciousness within the seizures. These episodes were considered to represent “subtle” SE. The patient was aggressively treated with midazolam anaesthesia, intravenous phenytoin and levetiracetam and high-dose topiramate by nasogastric tube and recovered well.



**Figure 7** Drug-induced generalised critical illness status epilepticus.

89-year-old man admitted to another hospital for abdominal pain. Sigmoid diverticulitis was suspected and treated with the 4th-generation cephalosporine cefepime. The patient slowly became comatose and he was referred to the tertiary care centre. Head CT and CSF were normal. The EEG showed almost absent background activity replaced by generalised partially rhythmic delta activity mixed with a few multifocal epileptic spikes (fig. 7a). After administration of 1 mg midazolam, the patient immediately regained consciousness and started to talk relentlessly; theta background activity reappeared (fig. 7b). Some days later he returned home in a good condition.






behavioural changes, some subtle clinical signs like discrete nystagmus, automatisms, muscle twitching, (unilateral) mydriasis or extensor response upon plantar stimulation may be present. Dyscognitive

SE usually originates from the temporal and less frequently from the frontal lobes [58–61], rarely it originates from the parietal or occipital lobes. The long-lasting episodes of dyscognitive SE often


**Figure 8A** Triphasic waves or epileptic discharges? Stimulus-sensitive triphasic waves.

57-year-old woman with subarachnoid haemorrhage (Fischer Grade IV) and clipping of aneurysms of the left anterior and medial cerebral artery. The patient remained at GCS 4–6 without sedation. The EEG revealed a biphasic curve pattern with slowed background activity and focal slowing in the delta/subdelta range in the left frontal region (fig. 8A-a) which spontaneously or after acoustic stimulation (fig. 8A-b) changed into high-amplitude serial triphasic waves (fig. 8A-c).




**Figure 8B** Triphasic waves are sensitive to benzodiazepines.

45-year-old woman with end-stage metastatic leiomyosarcoma of stomach undergoing chemotherapy with high-dose ifosfamide. The patient was comatose. The EEG showed almost continuous generalised, frontally accentuated triphasic waves and a few multifocal epileptic spikes (fig. 8B-a). These triphasic waves were markedly reduced in amplitude after administration of 5 mg of lorazepam (fig. 8B-b).



**Figure 8C** Triphasic waves resulting from accumulation of opioids – response to the opioid antagonist naloxone.

74-year-old man undergoing pancreatojejunostomy for pancreatic cancer. Postoperatively, he had multiorgan failure. Analgetic management included fentanyl. The patient did not regain consciousness and NCSE was suspected. The EEG revealed permanent rhythmically generalised, frontally accentuated triphasic waves mixed with some multifocal epileptic discharges (fig. 8C-a); the triphasic waves almost completely disappeared after administration of naloxone while the epileptic discharges persisted (fig. 8C-b).







are also called “psychomotor status”, emphasising the “strange” coincidence of severely impaired consciousness and altered behaviour together with an almost perfectly preserved functioning of the

**Figure 9** Postanoxic myoclonic status epilepticus – differentiation from generalised convulsive status epilepticus.


79-year-old man undergoing aortocoronary bypass surgery. Some hours later on the same day he had cardiac arrest and was electromechanically resuscitated during 10 minutes. He had repetitive subtle generalised myoclonic seizures. The EEG revealed a slow low-amplitude background activity (fig. 9a) alternatively changing with generalised epileptic spike-wave discharges of 1 Hz (fig. 9b) which ceased after the administration of lorazepam and resulting in a diffusely spreading, non-reactive, frontally accentuated theta activity, a so-called theta coma often seen in postanoxic encephalopathy (fig. 9c). The patient did not regain consciousness and died 13 days later from septic pneumonia.



**a**



**b**



**c**

motor system allowing for extensive and often seemingly purposeful activities. It is well understandable that these episodes often were believed to be of “psychogenic” origin by the lay people already a long time ago [62]. In fact, the differentiation of this type of dyscognitive SE from non-epileptic conditions without EEG and by pure clinical means may become challenging even for the skilled epileptologist [63, 64]. The differentiation of NCSE from non-epileptic psychogenic pseudostatus is very important since the treatment of pseudostatus as SE may put these patients at life-threatening risks [65, 66]. Furthermore, it may become especially difficult to diagnose dyscognitive SE in elderly patients in whom it is as frequent as of 40% [67–69]. Typical psychiatric manifestations of dyscognitive SE include delirium [67, 70, 71], stupor or catatonia [72, 73], mental slowing [74], cognitive decline [75], aggressive behaviour [76] and psychotic depression [77]. When spreading to the neocortical areas of the temporal lobes, auditory or visual hallucinations may occur [78, 79]. The EEG of dyscognitive SE is characterised by irregular or regular focal spikes or spike-wave activity similar to the one observed in aura continua; however, the ictal activity in dyscognitive SE tends to involve a larger area which increases the likelihood to detect it by surface EEG (fig. 5) [80, 81].

In this context, the significance of “periodic lateralised epileptiform discharges” (PLED), as they are present in figures 4a and b and figure 5a, should be discussed. The EEG pattern of PLED was first described by Chatrian et al. [82] and denotes periodic lateralised spike(-wave)-like activity of about 0.5–1.5 Hz which may occur after a variety of cerebral events like tumours, bleedings and ischaemic strokes. They often persist for weeks and months and may poorly respond to antiepileptic treatment; the authors already doubted their true epileptic nature by choosing the term “epileptiform”. However, he also occasionally observed PLED of epileptic nature. Pohlmann-Eden et al. in their seminal review proposed that PLED may not represent a specific single entity of pathological EEG activity, but a continuum ranging from rather benign epileptiform activity to clear-cut continuous ictal activity, i.e. SE [83]. This view was supported by later studies which stated that PLED following GCSE or serial epileptic seizures have to be considered as an ictal pattern; these PLED also responded to antiepileptic treatment [84]. The study of Assal et al. by using SPECT elegantly showed that PLED following SE or repetitive seizures were coupled with hypermetabolism which switched to hypo/(normo-)metabolism as


soon as PLED ceased, again pointing to PLED as an ictal pattern [85].

The last group includes the “critical illness SE” (CISE) where patients experience NCSE because of (multi-)organ failure and/or often complex polypharmacotherapy [18]. The most frequent type observed clinically and electroencephalographically resembles dyscognitive SE with often secondary generalisation. The background activity in the EEG is usually very slow, and serial epileptic seizures without interictal recovery of the patient (instead of continuous epileptic activity) might be observed (fig. 6). Some of the CISE were induced by drugs, notably those targeting the CNS, but also antibiotics, like carbapenems, chinolones

(gyrase inhibitors) and most prominently, the fourth-generation cephalosporine cefepime [86–88]. These CISE mostly show a generalised ictal pattern with a very polymorphous background slowing, rhythmic delta activity and multifocal or generalised spike-wave activity. The EEG may also display triphasic waves (TPW) (fig. 7) which may become very rhythmic and sharply contoured, mimicking spike-wave activity of (NC)SE. However, these TPW were not ictal, but reflect encephalopathy [81, 89]. Although first described in the context of hepatic encephalopathy [90, 91], TPW may occur in any other type of encephalopathy, like renal, postanoxic and drug-induced (cytostatic drugs, opiates) encephalopathy. Importantly, TPW

**Figure 10** Postanoxic myoclonic status epilepticus with subclinical seizures.

81-year-old man undergoing cardiac arrest and cardiopulmonary resuscitation. He was treated with midazolam anaesthesia for 3 days, but did not awake after stopping anaesthesia. The EEG showed a non-reactive, very low amplitude (sensitivity of 2  $\mu$ V/mm) fast beta activity of 20–25 Hz (fig. 10a). This background activity was interrupted by extensive bursts of generalised myocloni, most probably originating from the reticular brain-stem region (“reticular myoclonus”) (fig. 10b). But there were also generalised seizures with evolving delta activity, sometimes intermingled with the myoclonic bursts (fig. 10c–f). This activity eventually responded to anaesthesia with propofol and midazolam (fig. 10g). Nevertheless, the patient died 2 days later of non-epileptic complications.





may disappear after administration of benzodiazepines (fig. 8a and 8b) (or naloxone in the case of opiate-overdose [fig. 8c]) [92, 93]. Thus, the response of this EEG pattern to benzodiazepines does not help to distinguish TPW from ictal spike-wave activity. A more detailed analysis for the differentiation of NCSE from TPW is given by a recent study by Boulanger et al. [94].

**Figure 11** Generalised subtle myoclonic status epilepticus.

64-year-old woman undergoing cardiopulmonary resuscitation after ventricular fibrillation. She remained comatose and had bilateral eyelid myoclonus, tonic left upper gaze deviation and irregular body myocloni. The EEG revealed generalised (slightly a-)periodic epileptiform discharges (GPED) which – together with the subtle ictal semiology – were considered to be of epileptic nature (fig. 11a). The initial response to intravenous midazolam led to the suppression of the epileptic discharges (fig. 11b). However, the generalised epileptic discharges reappeared some hours later (fig. 11c) and neither high-dose midazolam nor propofol did abort this activity and barbiturates could not be administered because of the cardiac frailty. The patient died 3 days later.



Another subtype is called “subtle” SE, denoting those cases of NCSE which smoothly develop from preceding overt GCSE, a concept built upon the experimental work of Treiman et al. [29] and clinically supported by DeLorenzo and colleagues [95].

The third subtype encompasses the postanoxic myoclonic SE (MSE) where patients experience bursts, clusters or continuous, focal or generalised, symmetric, rhythmic myocloni [96, 97]. Clinically and electrographically this state may be sometimes difficult to separate from the putatively non-epileptic, often symmetric reticular myoclonus [98]. Occasionally, stimulus-sensitive focal or generalised subtle myocloni associated with epileptic discharges in the EEG can be observed in patients after cardiopulmonary resuscitation (CPR); evolution from severely suppressed background activity into an ictal pattern and attenuation following administration of i/v midazolam represent seizure activity and not a purely encephalopathic pattern (fig. 9). Alternating reticular brain-stem and epileptic cortical activity can occasionally be recorded in EEG of patients after CPR (fig. 10). Subtle generalised MSE initially may respond to midazolam, but can become increasingly resistant to further treatment (fig. 11).

Several authors do not consider postanoxic myoclonic SE as a form of NCSE because its treatment very often clinically does not improve the general serious condition of the patient. However, continuous clinical and electrographical postanoxic (subtle) myoclonic epileptic activity *is* SE, irrespective of the lack of substantial response to treatment. This resistance to therapy may rather reflect the very severe global brain damage in these patients making recovery implausible (cf. also paragraph on prognosis).

## Epidemiology

Sound epidemiological data about the incidence of NCSE and its subtypes are not available yet because of the lack of an accepted definition and as a result from using different criteria including and excluding some forms of NCSE. Additionally, it is likely that the protean, often unspectacular and calm presentation of NCSE is frequently missed, leading to a substantially underestimated incidence of NCSE when compared to GCSE with its self-evident clinical manifestation. There are five large population-based studies and an overview about the incidence of SE in general [99–104]. Their data are summarised in table 1a and 1b after being pooled and weighted according to their size; information about the incidence of specifically NCSE

**Table 1a** Incidence of NCSE: synopsis of five population-based studies on the epidemiology of status epilepticus.

types of status epilepticus	Bologna (I) <sup>a, e</sup> (Vignatelli 2003) (n = 336.876) <4% non-whites	Marburg (D) <sup>a, e</sup> (Knake 2001) (n = 743.285) <4% non-whites	Geneva (CH) <sup>a</sup> (Coeytaux 2000) (n = 1.735.420) 4% non-whites	Rochester, MN (USA) <sup>b</sup> (Hesdorffer 1998) (n = ~544.215) 4% non-whites	Richmond, VA (USA) <sup>a</sup> (DeLorenzo 1996) (n = 202.774) 57% non-whites
absence status	2.0	6.0	3.5	0.6 (3.3) <sup>d</sup>	1.0
CPSE	16.0	43.3	26.7	7.1 <sup>c</sup> (38.9) <sup>d</sup>	3.0 (?) <sup>*</sup>
subtle	–	–	1.2	–	–
myoclonic	16.0	–	–	1.9 (10.4) <sup>d</sup>	2.0
(others)	7.0	4.0	6.4	1.2 (6.5) <sup>d</sup>	1.0
GCSE	9.0	14.0	33.1	4.6 (25.1) <sup>d</sup>	29
SGSE	41.0	19.3	13.9	2.9 (15.8) <sup>d</sup>	43 (?) <sup>*</sup>
other	9.0	13.4	15.2	–	21 (?)
total <sup>b</sup>	10.7	17.1	10.3	18.3	50
gender: ♂/♀ <sup>b</sup>	9.7/11.5	26.1/13.7	12.1/7.8	23.2/13.1	not available

<sup>a</sup> Values in per cent.<sup>b</sup> Incidences per 100 000.<sup>c</sup> Classification “partial” only, no further subdivision into “simple” or “complex”.<sup>d</sup> A substantial proportion of CPSE may be included in the category “partial, secondarily generalised”.<sup>e</sup> Adults > age 20 only.<sup>\*</sup> Some cases of NCSE probably included in the group of SGSE.

and its subtypes is extracted as much as possible. The incidence of NCSE may be 6/100 000, representing about 40% of all SE; however, when looking at this proportion in a tertiary care centre where critically ill patients are more frequent, it may be markedly increased. About 8% of all comatose patients in a tertiary-care-centre ICU had NCSE [105]. It is important to use the EEG as an additional tool to examine patients in coma of unknown

origin and know the various patterns of EEG changes associated with coma of both NCSE and metabolic encephalopathy, respectively [81, 89]. About two thirds of NCSE are dyscognitive SE. The incidence of NCSE in males is twice that of females.

## Causes

Two general conditions have to be considered when looking at the causes of NCSE: first, NCSE in those patients with known epilepsy and, second, in those patients with no history of epilepsy.

The main reasons for NCSE in epileptic patients are changes in AED levels. They include lowering of levels by purposeful or inadvertent reduction of AED dosages, or by non-compliance [106]. Toxic levels of AED too may cause NCSE [107]. Almost all AED may provoke worsening of seizures or even of an epileptic syndrome in some patients [108]. The AED tiagabine, inhibiting the re-uptake of GABA from the synaptic cleft by blocking the GABA transporter-1, may induce NCSE by interference with the pre- and postsynaptic GABA<sub>A/B</sub> receptors and other mechanisms modulating transmitter release [109, 110]. Antiepileptic drug levels impaired by co-administration of other enzyme-inducing drugs may also facilitate the onset of NCSE. Other reasons may be inappropriate AED selection, for example, carbamazepine or phenytoin (PHT) for an IGE syndrome [111, 112]. Ex-

**Table 1b** Incidence of different types of NCSE according to the results of population-based studies on the epidemiology of status epilepticus.

	all studies percentual [%]	all studies, population-weighted percentual [%]	absolute per 100 000 p.	
non-convulsive forms				
absence status	3.16	3.71	0.531	
CPSE	25.58	29.67	4.219	
myoclonic	5.68	3.22	0.510	
subtle	0.24	0.58	0.060	
others	4.98	5.66	0.745	
total	39.64	42.84	6.065	39.80%
convulsive forms				
GCSE	22.04	25.38	3.647	
SGSE	26.60	19.54	3.559	
others	11.72	11.48	2.340	
total	60.36	56.40	9.173	60.20%
total	100.00	99.24	15.238	100.00%



ternal factors, like fever, drugs, sleep shifts and deprivation, photic stimulation and hormonal changes also may trigger NCSE in epileptic patients [32].

Structural and metabolic-toxic changes are the principal causes of NCSE in patients without a history of epilepsy. Non-convulsive status epilepticus resulting from structural alterations include acute events like intracranial haemorrhages, traumatic brain injuries and ischaemic strokes [113–116]. Subacutely, primary brain tumours and metastases may provoke NCSE [117–119]. Infections, like herpes encephalitis, neurosyphilis and Creutzfeldt-Jakob disease, may be associated with NCSE [120–122]. Chronically progressing conditions causing seizures encompass all the different forms of neurodegenerative disorders, like Alzheimer's disease, frontotemporal dementia spectrum, cerebral amyloid angiopathy, etc. [123, 124]. Several metabolic conditions may be associated with NCSE as an epileptic reaction to the changed cerebral physiological environment. Both hypoglycaemic and hyperglycaemic states may be associated with NCSE [125, 126]. Hyperthyroidism and hypothyroidism are causes of NCSE; while the first is known in thyrotoxic crisis [127], the latter occurs in myxoedema [128] or steroid-responsive encephalopathy associated with autoimmune thyroiditis (the former "Hashimoto encephalopathy") where NCSE occurs with a frequency of 10 to 15% [129–133]. Disturbances of sodium, potassium and calcium homoeostasis may be associated with NCSE [37, 78, 134–136]. Hypomagnesaemia induced by therapies with cis-platin, amphotericin B and so on can cause NCSE [137]. Numerous other (illicit and approved) drugs are known to lower seizure threshold (table 2). The most important drug causing NCSE is alcohol withdrawal and only very rarely alcohol intoxication [138]. It is of special interest to the neurologist that – beyond drugs like antibiotics, anticancer and immunosuppressive drugs – some antipsychotic drugs, antidepressants and some antiepileptic drugs (tiagabine, vigabatrin, gabapentin, carbamazepine, phenytoin) may cause NCSE [139]. This list would be incomplete if increasing age as "risk factor" for NCSE was not mentioned [140, 141]; an epidemiological study in the Hong Kong population showed a 3- to 8-fold increase in the incidence of SE [142].

### Diagnosis and differential diagnosis

The diagnosis of NCSE relies on the combination of the patient's history, the clinical signs and the EEG. Sometimes, the clinical and electroen-

cephalographic response to the administration of benzodiazepines may help to confirm the diagnosis. The broad range of clinical signs and symptoms were mentioned in the section on the different types of NCSE. Their protean, often unspectacular manifestations make it very difficult to diagnose NCSE on clinical grounds alone. Nevertheless, a remote condition facilitating the onset of an epileptic disorder, severely impaired consciousness and spontaneous eye movements (i.e. horizontal nystagmus) were significantly associated with the presence of NCSE [143]. A history of epilepsy may facilitate to think of NCSE; absence of such a history will not exclude the presence of NCSE in patients, since NCSE may often be the first manifestation of an epileptic condition in patients admitted to hospitals [30, 144, 145]. However, NCSE does not exclusively start in ICUs or hospitals, but also at home, in psychiatric institutions, in nursing homes and asylums.

According to Jordan, the following conditions warrant further evaluation regarding a possible diagnosis of NCSE [23]:

- episodes of blank staring, automatisms, aphasia or perseverations of actions;
- unexplained onset of impaired consciousness, especially if its level fluctuates;
- fluctuating aphasia without structural lesion explaining the aphasic deficit;
- impaired consciousness or mentation associated with minimal clinical signs, like eyelid, facial or truncal subtle myoclonus, horizontal nystagmus and spontaneous extensor position of one or both big toes;
- prolonged postictal state or postictal unawareness of longer duration than 15 to 30 minutes;
- protracted state of reduced alertness after brain surgery or any other surgery where cerebral functions are at risk.

To confirm the diagnosis of NCSE warrants the exclusion of other diagnoses. The differential diagnosis of NCSE encompasses acute stroke, inflammation (like limbic or paraneoplastic encephalitis), infection (like herpes simplex encephalitis), primary brain tumours and metastases in the non-motor areas, "pure" psychiatric causes of delirium, stupor and delusions or hallucinations, and non-epileptic, psychogenic pseudostatus. Several approved and illicit drugs as well as metabolic and electrolyte alterations may produce neurological and behavioural states resembling NCSE without detectable ictal activity in the concurrently recorded EEG.

To conclude, the most important step in the diagnosis of NCSE is "to think of it at all!" [18]. In addition, the value of 24-hour availability of EEG

cannot be overestimated if NCSE is clinically suspected, but warrants definitive confirmation. The various EEG patterns of NCSE and its confounders were already presented in the “types of NCSE” section of this article; more detailed information is displayed in several excellent reviews elsewhere [37, 41, 56, 81, 89, 146].

The diagnostic value of other paraclinical examinations may depend on the very specific context. Thus, analysis of cerebrospinal fluid may


help to diagnose an underlying CNS infection; unspecifically and as a bystander phenomenon, there may be pleocytosis of up to about 30 cells and slightly elevated protein levels in the CSF of non-infectious NCSE [147]. An increase of neuron-specific enolase upon NCSE was suggested [148–150]. The various imaging modalities may be of lesser value in the emergency diagnosis of NCSE, but may reveal valuable localisatory, structural (CT, MRI, DTI), pathophysiological (DWI, SPECT) and

**Table 2** Drugs causing non-convulsive status epilepticus.

approved medications	
<b>CNS active drugs</b>	
neuroleptics	carbopenems
especially: clozapine	imipenem
chlorpromazine	meropenem
olanzapine	<b>anticancer drugs</b>
lithium	ifosfamide
antidepressants	busulphane
tricyclic (-“pramins”)	<b>immunosuppressants</b>
tetracyclic (maprotilin, mianserin)	cyclosporine
not (or even anticonvulsant): SSRI	mycophenolate mofetil
stimulants	tacrolimus
theophylline	<b>intravenous contrast mediums</b>
appetite moderators	<b>antiarrhythmics</b>
(methylphenidate; effect not definitely determined)	
mid- and high-dose opiates	<b>illicit drugs</b>
flumazenil	<b>tranquillisers/sedatives</b>
benzodiazepine withdrawal	benzodiazepine withdrawal
<b>antiepileptic drugs</b>	barbiturate withdrawal
tiagabine	<b>opiates</b>
vigabatrine	mid to high doses of all opiates
carbamazepine	(low doses probably anticonvulsive)
phenytoin	morphine
<b>antibiotics</b>	heroin
penicillin (derivatives)	buprenorphine
penicillin G	(hydr-)oxycodon
mezlocillin	<b>stimulants</b>
piperacillin	cocaine
calvulanic acid	amphetamine
cephalosporins	designer amphetamines
cefepime	paramethoxyamphetamine (PMA)
ceftazidime	paramethoxymethamphetamine (PMMA)
gyrase inhibitors	(benzyl-)piperazines
ciprofloxacin	<b>alcohol</b>
ofloxacin	withdrawal
enoxacin	(excessively high intake)
exception: norfloxacin (does not cross the BBB)	
	<b>Abbreviations:</b>
	BBB: blood-brain barrier; SSRI: selective serotonin re-uptake inhibitors.



**Figure 12** Pathophysiological aspects and sequelae of non-convulsive status epilepticus.



metabolic (PET) information about the involved brain areas, especially when used in a combined, complementary manner [151–154]. The readily availability, the ease of acquisition and the important information displayed in acute stroke placed DWI among the most performed imaging modalities, increasingly also in an emergency setting. In the context of NCSE, it is of note that hyperintense DWI changes (with or without reduced ADC maps) may also be observed. However, the hyperintensities in NCSE are very often less bright, show a very cortical pattern, often not exactly respecting a perfusion territory typical of a specific cerebral artery and they may disappear within 5 to 7 days – in contrast to the ischaemic DWI hyperintensities persisting for 2 to 6 weeks [155–158].

### Pathophysiological aspects

Pathophysiological aspects of NCSE are similar to those observed in the convulsive forms of SE on the cellular and local network (neuronal/astroglial) level, but they may be different on the level of the brain and the whole body [31, 159, 160].

Neurons, astroglial cells and small local networks affected by continuous epileptic activity of GCSE and almost all forms of NCSE (except for AS) display a cascade of electrochemical events triggered by the synchronous hyperexcitability. This hyperexcitability may result from a direct increase of excitatory mechanisms or a reduction of the local inhibitory network (i.e. disinhibition), or both of them. Hyperexcitability usually is linked with an excessive glutamatergic spillover of neu-


rons which also exceeds the capacity of glutamate-reuptake (“buffering”) by astroglial cells. Toxic amounts of extracellular glutamate overstimulate glutamate receptors and induce a disruption of the extra-/intracellular  $\text{Ca}^{2+}$ -homoeostasis with accumulation of cytotoxic intraneuronal  $\text{Ca}^{2+}$ -concentrations. Excess of the second messenger  $\text{Ca}^{2+}$ -ion paralyses or overdrives numerous downstream metabolic processes vital for the neurons and may induce acute cell death or apoptosis [161]. The high energy demands of the hyperexcited neurons exhaust the pool of energy sources of the brain which is almost exclusively dependent on glucose. Energy failure leads to a loss of function of the  $\text{Na}^+/\text{K}^+$ -ATPase which is followed by a breakdown of the ionic transmembrane homoeostasis and by subsequent concomitant water influx into the neurons and astroglial cells, cell swelling and death. As another consequence, toxic metabolites, nitric oxide formed upon activation of the neuronal and inducible forms of NO synthase, and radicals may accumulate intra- and extracellularly, and further act as cytotoxic substances, especially to cell membranes, or as chemoattractants for inflammatory cells. Eventually, there may be local destruction of functional brain tissue and gliosis with loss of function leading to persistent neurological deficits or the onset or aggravation of an epileptic disorder.

On the level of the whole body, GCSE has the additional consequence of exhaustive motor activity which will lead to profound metabolic changes such as lactic acidosis, hyperthermia and sympathetic overdrive in the phase of decompensation [162, 163]. When persisting, these factors may create an immediate life-threatening condition. In contrast, NCSE will not induce whole-body energy exhaustion with its metabolic sequelae because of the lack of excessive motor activity, but it is essential to note that continuous epileptic activity in non-motor brain regions, like in the temporal lobes or the insular and opercular cortex, may directly overstimulate autonomic sympathetic and parasympathetic centres which in turn may induce life-threatening autonomic hyper- or hypo-activity, most importantly ventricular arrhythmias or asystolia and bronchospasm [164]. An overview of these mechanisms is schematically drawn in figure 12.

### Prognosis

The prognosis of a disorder in general refers to the outcome with regard to survival or death (i.e. mortality), to deficits, disabilities or handicaps, and to the response to treatment. Prognosis is essential

**Figure 13** Prognosis of different forms of non-convulsive status epilepticus according to the impact of the underlying condition.



to determine the need, the immediacy and the intensity of the treatment; it also serves to handle the questions and expectations of the patients and their relatives.

Overall mortality of SE in general ranges within 10–33% in the population-based epidemiological studies [165]. Short-term mortality was 19% in a large population-based study irrespective of the type of SE; however, the data were collected from 1965 to 1984, at a time when treatment and ICU care of SE were not at the level of nowadays standards [166]. The long-term mortality over the first 10 years after SE in the same population was 40% and thus threefold higher than that of the matched general population [167]. Older patients with NCSE also share a higher mortality than those without [142]. Mortality of specifically NCSE has been 19% in a more recent study; mortality was significantly associated with symptomatic NCSE [168]; accordingly, an earlier study observed a much higher mortality of 57% in patients with exclusively symptomatic NCSE on an ICU [169]. Additionally, the prognosis of NCSE may not only include the issues of mortality and response to treatment, but also those of the mid- and long-term sequelae, like the likelihood of recurrence of NCSE and the onset of epilepsy after NCSE [136]. Recent data suggest that the risk of recurrence of (NC)SE ranges from 30 to 100% depending on the underlying cause of NCSE [170]. About 40% of patients will experience an unprovoked seizure within 10 years after one or repetitive episodes of (NC)SE [171].

For several reasons the prognosis of NCSE is one of its most equivocal and debated topics [172–174]. First, there is no broadly accepted defi-

nition which leads to the inclusion and/or exclusion of some forms of NCSE depending on the specific criteria of the investigators in these studies which in turn may have weakened the strength of the results and made it difficult to compare these studies. Second, it is important to respect the variety and heterogeneity of types of NCSE which render it impossible to make a universal prognosis for NCSE [175]. Third, it has become clear that any prognosis in any form of NCSE has to be considered within the specific context of the presence of underlying (causative) or concomitant (non-causative) disorders [1, 37, 174]. Fourth, the prognosis of NCSE is influenced by the time until adequate treatment will have been installed, i.e., delayed diagnosis and therapy will impact the outcome in a negative way [35, 37]. Fifth, there is increasing doubt as to whether the vast amount of experimental animal data on (NC)SE in fact strongly relates to NCSE in humans [176]. Many of these data were derived from specific animal models, often at different developmental stages or periods of postnatal life. The uncertainty of the reliability of the results of these studies was best reflected in one issue of *Progress in Brain Research* by addressing the yet unresolved question “Do seizures damage the brain?” [177]. In recent years, however, there has been increasing evidence from pathological and imaging data that all types of NCSE except for AS may damage the brain [178–183].

Keeping these issues in mind, the prognosis of the different types of NCSE will be discussed according to the classification shown in figure 1. Additionally, it seems helpful to place these types of NCSE into a Cartesian graph where the x-axis denotes non-epileptic factors increasingly responsible for the patient’s state and prognosis, and where the y-axis assigns increasing importance to the epileptic activity determining the patient’s global state and prognosis (fig. 13). Thus, it becomes clear that in the case of AS (left top) the epileptic activity is almost completely responsible for the patient’s state, while, at the other end (right bottom), the epileptic activity in postanoxic MSE may contribute only little to the patient’s global state of almost always deep coma. However, a recent study has shown that the occurrence of MSE in patients after cerebral anoxia represents an independent poor-outcome predictor, emphasising that MSE might be more than an innocent bystander of hypoxic brain damage [184]. Nevertheless, it follows that the prognosis of AS is closely linked to the response to antiepileptic treatment, whereas the prognosis of MSE is mainly dependent on the severity of global brain damage and the failure of other vital organs, and less on the result of the



antiepileptic treatment. Accordingly, the prognosis of CISE is mainly determined by the prognosis and response to treatment of the underlying critical illness.

As an approximate rule, the types of NCSE where the epileptic activity largely contributes to the patient's impairment share a better prognosis than those forms where patients suffer from critical illness. This fact has been corroborated by a simple clinical score for the prognosis of SE in adults, using consciousness, type of SE, age and history of seizures as items. The patients with the highest scores corresponding to the poorest prognosis were those without history of seizures, age over 65, NCSE and coma [185]. Absence status has an excellent prognosis with respect to response to treatment and outcome [37]. An exception might be *de novo* AS in adults caused by withdrawal of benzodiazepines; here, the prognosis is influenced by the interference of benzodiazepine addiction. The prognosis of atypical AS also is good, especially when atypical AS occurs *de novo* in non-epileptic patients; however, it has often to be considered with respect to an underlying more severe idiopathic generalised epileptic syndrome which *per se* may be associated with a more serious outcome. Aura continua often shares a good prognosis, but this is again dependent on the cause of aura continua and concomitant diseases; aura continua difficult to control may be caused by malformations, astroglial tumours, (residual) intracerebral haematomas or mitochondrial diseases. The prognosis of dyscognitive SE is good in general, but also determined by the presence or absence of structural lesions and concomitant disorders. The prognosis is much more serious in most forms of CISE because of the underlying disorder; however, the NCSE plays an important contributory role to the course of the disease as it may additionally worsen the global outcome of the patient. Subtle SE also shares a serious prognosis with respect to the long-lasting persisting epileptic activity which impairs response to treatment (inducing the vicious circle described earlier) which may increase the likelihood of irreversible brain damage. The worst prognosis must be attributed to postanoxic MSE, where about 80% of patients die and the surviving almost never return to a conscious life remaining in the permanent vegetative state. This outcome reflects the catastrophic brain damage caused by anoxia; MSE has to be considered not as a main causative factor, but as an indicator and epiphenomenon reflecting the disinhibitory state after abolition of inhibitory mechanisms of the brain and the global cerebral spillover with hyperexcitatory transmitters and excitotoxic metabolites [184, 186–188].

## Treatment

The treatment of NCSE basically follows the same principles as do therapeutic algorithms for GCSE, although there is an appalling lack of studies specifically devoted to the optimal treatment of NCSE which would fulfil the criteria of evidence classes I or II. Again, this absence of data may result from the absence of an accepted definition of NCSE, from its heterogeneity and the divergent opinions even of experts whether NCSE might damage the brain or not implicating a more or less aggressive treatment strategy [1, 172, 173]. Nevertheless, more recent reviews and guidelines emphasise the need for immediate, resolute and type-adapted treatment of NCSE [18, 22, 24, 33, 189]. It is important to note that the most studies providing data discussed below were predominantly carried out in patients with GCSE and not in patients with NCSE. However, their evidence may also apply to the patients having NCSE, at least not until novel specific data will provide different results. As mentioned in the previous paragraph, the antiepileptic treatment becomes the more important, the more the epileptic activity *per se* contributes to the patient's state, whereas in the symptomatic forms of NCSE like CISE the successful therapy of the underlying and concomitant non-epileptic disorders will mainly determine prognosis and outcome of the patient.

Once NCSE has been diagnosed, the patient should be checked and stabilised for the basic vital parameters. Then, thiamine 250 mg should be given intravenously before any glucose-containing fluids or higher concentrated glucose solutions for correction of hypoglycaemia will have been administered; this schedule is especially recommended in the case of suspected alcohol addiction or malnutrition, but may be performed in all patients with NCSE with regard to its harm-sparing potential and the low risk of adverse effects.

The initiation of the antiepileptic treatment without delay is one of the most important factors in the therapy of NCSE: the longer NCSE persists, the more difficult it is to be terminated.

The immediate administration of intravenous benzodiazepines (BD) is the unequivocal, evidence-based first step for the efficient treatment of NCSE; caregivers hereby should be prepared to ventilate the patient if respiratory depression will occur. The most frequently used BD are diazepam, lorazepam (LZP), midazolam (MDL) and clonazepam (only in Europe where an i/v-formulation is available). Although these substances share the same basic mode of action, they differ in many pharmacokinetic and pharmacodynamic aspects [190–192].

The most used BD diazepam is highly lipophilic, enters the brain very quickly and binds to the GABA<sub>A</sub> receptor. However, it then rapidly dissociates from this receptor subsequently equilibrating with fat tissue. This may explain the often observed recurrence (“breakthrough”) of (NC)SE after administration of diazepam. If additional doses of diazepam are given, the patient is at risk to be overdosed as soon as a new equilibrium level is reached and the diazepam accumulated in the fat tissue redistributes to the brain and its receptors [193]. This phenomenon and the fact that diazepam undergoes complex metabolism generating more than 40 active metabolites make this substance very demanding to handle and achieve predictable levels and effects [190, 191]. Thus – and against widely common practice – diazepam is no longer the BD of choice in SE.

The less lipophilic BD LZP which is not metabolised to further active compounds accordingly has become the BD of choice for the first-line treatment of (NC)SE [194]. Lorazepam additionally leads to less respiratory depression [195] when compared to other BD and – most important for a sustained antiepileptic effect and the avoidance of recurrence of SE – it seems to bind in a semi-covalent manner at the GABA<sub>A</sub> receptor site extending its effect up to about 24 hours despite falling blood levels [196–199]. The dominant role of LZP in the treatment of (NC)SE was definitively established by the hallmark multicentre prospective double-blind Veterans Affairs-Study published in the *New England Journal of Medicine* in 1998, comparing four different first-line regimens (phenytoin [PHT], diazepam followed by PHT, phenobarbital and LZP) where LZP showed to be significantly superior to PHT, but not to the other drugs; it was, however, easier to handle [20]. The pre-hospital use of LZP for the treatment of serial seizures or SE was safe and efficient in a large study in adults 3 years later. Unexpectedly (but calming), respiratory problems were significantly more frequent in the placebo group; it also seemed likely that LZP was superior to diazepam in terms of efficacy [200]. Unfortunately, lorazepam is still not approved by the Swiss authorities (Swissmedic) for the use in SE.

Midazolam (MDL) is a very rapidly acting BD with a short half-time [190, 191]; therefore, stable antiepileptic effect needs repetitive or continuous administration [201–203]. It also has markedly sedative and stronger respiratory depressive properties which make it less suitable for first-line treatment of SE, but it is widely used for the treatment of refractory SE (RSE, cf. below). Nevertheless, for pharmacological reasons and in

the specific ICU environment, the ultra-fast antiepileptic activity of MDL may be exploited by first-line administration of a bolus of 1 to 5 mg to induce the very immediate effect, while at the same time also lorazepam 2–8 mg is given which will start to exert its sustained effect just at the decline phase of MDL. Midazolam can also be safely administered intramuscularly and may be preferred in those SE where i/v access is not available or its installation at risk [203, 204].

Clonazepam may have a profile similar to that of LZP and is widely used in French-speaking countries; however, sound studies are lacking probably because of no available intravenous formulation in some Anglo-American countries [205–207].

While BD are being given as first-line treatment, the patient should undergo laboratory exams including haematological and chemical parameters as well as checking for thyroid hormone status, infectious diseases (especially herpes encephalitis) and an extensive drug screen because several “club drugs” like amphetamine and its (“designer-”)derivatives (paroxymethamphetamine [PMA], paramethoxymethamphetamine [PMMA], [benzyl-]piperazines and cocaine) may provoke NCSE [208–214].

Since BD are not the mid- or long-term treatment of epileptic disorders and these medications impair the patient’s awareness, memory and consciousness which may impede the judgement of the patient’s neurological state in the course of the disease, the concomitant administration of a “classic” AED should be put ahead. Actually, there are three different compounds with an intravenous formulation available on the market: PHT, valproic acid (VPA) and levetiracetam (LEV), but only PHT is approved for the treatment of (NC)SE in Switzerland.

Phenytoin was the first antiepileptic drug beyond the sedatives BD and barbiturates to be administered intravenously. The feasibility, safety and efficacy of i/v PHT was demonstrated in the early 1950s, however, not in studies which would satisfy standards of nowadays [215, 216]. The only study yielding Class-I evidence for i/v PHT in GCSE was the already cited VA study; it showed that PHT alone was significantly inferior to lorazepam to stop SE, but it was efficient when given together with i/v diazepam [20]. The use of the i/v formulation of the drug implicates to keep several cautions in mind: the solution has to be stabilised at a high, very basic pH, which necessitates a central line and a infusion rate of no more than 50 mg per minute. The substance by itself and the additives of the i/v formulation may induce serious cardiac and local cutaneous (“purple-glove syn-

drome") adverse effects [217–219]. In addition, phenytoin is highly protein bound and metabolised by the cytochrome P 450 (CYP450) system, mainly the 2C9 and 2C19 variants, which produces several pharmacokinetic and -dynamic interactions, especially to be observed when coumarins, dexamethasone, cytostatic drugs, other AED or some antibiotics were co-administered [220–224]. The non-linear pharmacokinetics of PHT bear a high risk of toxic levels; their close monitoring is essential. The prodrug fosphenytoin is stabilised at a physiological pH and free of cardiotoxic additives which allows for a much faster infusion rate and also for intramuscular administration [225], but there is no advantage in terms of faster reaching the therapeutic level because the prodrug has to be metabolised by plasma phosphatases into phenytoin [226, 227]. These properties, together with an about 10 times higher price than PHT, resulted in a lack of approval in almost all European countries, Switzerland included [228, 229].

Intravenous valproic acid (VPA) has been available in central European countries for more than 20 years and may be an alternative to PHT in the treatment of SE [230–232]. Valproic acid has the broadest spectrum off all AED to date and does not induce marked impairment of consciousness. It is a weak to moderate inhibitor of the CYP450 system and, as a short-chain fatty acid, metabolised by beta-oxidation in the liver and renally eliminated after glucuronidation [233–238]. The substance leads to the production of ammonia (sometimes dramatically exacerbated by the presence of a mitochondrial urea-cycle enzyme defect like ornithyl-carbamoyl-transferase deficiency) which should be closely controlled after fast loading; levels of ammonia up to 70  $\mu\text{mol}$  (normal 7–34) usually were well tolerated, whereas levels >90  $\mu\text{mol}$  may lead to substantial obtundation which warrants for a dose reduction [239–241]. During i/v therapy for SE, levels of free (unbound) VPA should be monitored, because this fraction (normally 10–15%) rises overproportionally in the case of high doses and hypoalbuminaemia [242, 243] and may induce toxicity (sedation, elevated liver enzymes, low platelet count, reversible parkinsonism, pancreatitis) [244–246]. The use of VPA in patients with intracranial bleedings is questionable because of the various effects of VPA on platelet number and function as well as on several clotting factors [247, 248]. But, in general, the safety of i/v VPA, even at high doses, fast infusion rates and in cardiac instable patients, was demonstrated in several studies [249–259]. However, prospective randomised well-(i.e. PHT-)controlled studies

which had provided the Class-I evidence necessary to gain approval by regulatory boards (including Swissmedic) are still lacking. Nevertheless, some countries (Norway, Canada, Singapore) approved the drug on "summative, use-proven evidence"; i/v VPA was lately approved for the treatment of GCSE also in Germany after recommendations upon an extensive review of an expert panel [260]. A randomised double-blind pilot trial, using PHT or VPA as *first-line treatment before administration of benzodiazepines*, showed a trend to superiority of VPA against PHT; however, the study was underpowered and statistically flawed; thus, these results should be interpreted with caution and further studies are warranted [261, 262]. Most recently, a controlled randomised prospective trial comparing i/v VPA with i/v PHT (n = 50 in each group) after failure of BD has shown a significantly superior efficacy of VPA compared to PHT (p >0.05); the study also reiterated the importance of a treatment as soon as possible [263].

An intravenous formulation of levetiracetam (LEV) has recently been introduced, but has not gained approval for SE yet. This drug has a very favourable pharmacokinetic and -dynamic profile with no known interaction with any drug and only very few adverse effects (mainly slight somnolence and behavioural alterations which play no role in the acute treatment of SE) [264–266]. Levetiracetam has a substantial first-dose effect [267] and its fast i/v-application in healthy volunteers is safe and leads to high drug levels [268, 269]. After an experimental study had performed and shown an important anticonvulsant effect in SE [270], small studies used LEV as a third- or fourth-line treatment in refractory SE, often also given by nasogastric tube [271–275]. There are two reports of a few cases of NCSE probably induced by LEV, but these intriguing observations await further confirmation by other studies [276, 277].


The administration of LEV seems to be most favourable in critically ill, post-transplantation and HIV patients with (NC)SE on therapy with multiple drugs where the risk of significant adverse effects and interactions is especially high.

All the three substances discussed have the advantage that they can be switched 1:1 from the i/v-dose to oral administration without problems.

The success rate of the administration of one BD and an i/v AED (PHT/VPA/LEV) depends on the type of NCSE. It is almost 100% in all forms of AS and it is high in epileptic patients with aura continua or dyscognitive SE. The rate drops in patients with aura continua or dyscognitive SE resulting from structural lesions (tumours, haemorrhages) or in patients with CISE due to multiorgan failure. It



**Figure 14** Therapy algorithm of non-convulsive status epilepticus.



is lowest in patients with subtle SE and especially in postanoxic MSE [184, 185, 278].

Status epilepticus persisting after the administration of two AED is called refractory SE which occurs in about (20–)30% of all cases of SE [279–281]. It is treated by induction of anaesthesia after intubation of the patient with either midazolam, lorazepam, propofol and midazolam, or thiopental [282–298]. Studies evaluating these treatments are of small size and often not directly comparable because of different parameters used, especially electroencephalographic titration of depth of coma (suppression of epileptic activity vs burst suppression vs electrocerebral silence) [299–303]. A meta-analysis could not find a superiority of any of these treatments in terms of outcome [303].

Midazolam lowers blood pressure and undergoes tachyphylaxis because of alterations of the GABA<sub>A</sub> receptor subunit composition upon long-term stimulation like in SE [304]. It also bears the risk of accumulation accompanied by a very prolonged recovery from coma in patients with impaired renal function due to the pharmacologically active glucuronated metabolite  $\alpha$ -1-OH-midazolam [305].

Propofol is a highly lipophilic, very short-acting anaesthetic which functions as an anticonvulsant at different sites (GABA<sub>A</sub> receptor, NMDA receptor, voltage-gated Ca<sup>2+</sup>-channels) [294, 306–311]. When used as single anaesthetic compound, the substance may cause myocloni resembling seizures; however, this phenomenon is non-epileptic and most probably related to disinhibition of subcortical structures [312, 313]. To avoid the phenomenon or true “withdrawal” seizures, it is recommended to use propofol always combined with (a small dose of) a BD

[22]. Anaesthesia with propofol is associated with intake of 1200–1500 kcal of lipids due to the solvent based on soy oil. The drug should not be used longer than 5 days because it may cause the life-threatening “propofol-infusion syndrome” consisting of severe acidosis, liver failure and extensive rhabdomyolysis [314]. A pilot trial using propofol in 31 episodes of patients with refractory SE showed control of SE in two thirds of the patients without serious adverse effects [315].

When MDL and propofol fail, barbiturate coma with thiopental may be started. Barbiturates are very potent antiepileptic drugs and additionally downregulate global brain metabolism leading to a sort of “hibernating” brain state and, thus, reduce the risk of hyperexcitation and the accrual of toxic metabolites. However, they are markedly cardio-depressive and may induce severe hypotension. Barbiturate coma in SE is accompanied by a mortality of 30–50% [303]. Unfortunately, the barbiturate easiest to handle, pentobarbital, is no longer on the market in Europe. Thus, thiopental is the only available drug for this purpose; it is initially ultra-fast and -short acting because it is highly lipophilic, and it quickly crosses the blood-brain barrier, but it immediately accumulates also in the fatty tissues. From there, it will again equilibrate and redistribute to the brain structures. This eventually leads to very prolonged recovery from coma after stopping the drug [316, 317].

Other (additional) therapeutic options in RSE include the administration of all available i/v-AED (i.e. VPA and LEV, when PHT was first used) or enteral AED by nasogastric tube. Among those AED, high-dose topiramate was successfully used in small case series [318–320]. Lidocaine, ketamine, etomidate and clomethiazole were also “rescue” therapeutics [321–332], while others reported successful use of inhalative anaesthetics (“-fluranes”) despite reports of a possible proconvulsant effect of some compounds of this class [333–338].

Intravenous steroids may become another therapeutic alternative in RSE because there is increasing evidence that continuous epileptic activity stimulates the production of proinflammatory mediators like cytokines (especially interleukin-1 $\beta$  [339]) and helps to sustain epileptic activity. Steroids also are effective in some paediatric epilepsy syndromes, like infantile spasms, Landau-Kleffner syndrome and Rasmussen encephalitis [340–342]. The reported clinical experience in adult patients is small, but some cases with RSE responded to treatment with steroids [343, 344].

To summarise, a treatment algorithm of NCSE is proposed in figure 14 and dosages of the drugs discussed in this section are shown in table 3.

**Table 3** Drug dosing for non-convulsive status epilepticus.

drug		dosing	
A. first-line			
lorazepam	<60 y	4 mg i/v	repeat once
	60–80 y	2 mg i/v	repeat 1–3 times
	>80 y	1 mg i/v	repeat up to 5 times
clonazepam	<60 y	1 mg i/v	repeat 1–2 times
	60–80 y	0.75 mg i/v	repeat 1–3 times
	>80 y	0.50 mg i/v	repeat up to 3 times
midazolam	<60 y	5 mg i/v	repeat several times, if necessary
	60–80 y	2 mg i/v	repeat several times, if necessary
	>80 y	1 mg i/v	repeat several times, if necessary
diazepam	<60 y	10 mg i/v	repeat 1–2 times
	60–80 y	5 mg i/v	repeat 1–3 times
	>80 y	2.5 mg i/v	repeat up to 5 times
B. second-line			
phenytoin			
loading		15–18 mg / kg body weight at 50 mg/min	
maintenance	<70 kg	150 mg i/v	every 12 h; start 6 h after loading
	70–90 kg	175 mg i/v	every 12 h; start 6 h after loading
	>90 kg	200 mg i/v	every 12 h; start 6 h after loading
valproic acid			
loading	<60 y	30–45 mg i/v / kg body weight in 30 min	
	60–80 y	20–30 mg i/v / kg body weight in 30 min	
	>80 y	15–25 mg i/v / kg body weight in 30 min	
maintenance		start the same dose as the loading dose <i>concomitantly over 24 h</i>	
levetiracetam			
loading	<60 y	25–30 mg i/v / kg body weight in 15 min	
	60–80 y	15–25 mg i/v / kg body weight in 15 min	
	>80 y	10–20 mg i/v / kg body weight in 15 min	
maintenance*	<60 y	1000–1500 mg every 12 h	
	60–80 y	750–1000 mg every 12 h	
	>80 y	500–750 mg every 12 h	
C. third-line			
midazolam		(100–) 300–600 (–4000) µg/min	
propofol		(100–) 300–600 (–1000) mg/hour (+ midazolam or lorazepam!)	
thiopental			
bolus		100–250 mg	repeat until seizures stop
maintenance		3–5 mg / kg body weight / hour	
D. additional (“fourth-line”, “rescue”) therapy			
i/v			
lidocaine			
loading dose		1.5–2.0 mg/ kg body weight	repeat once after 20–30 min
maintenance		3–4 mg/ kg body weight/ hour	



ketamine		
loading dose	0.7–2.0 mg/ kg body weight	
maintenance	2–6 mg/ kg body weight/ hour	
clomethiazole		
loading dose	5–12 mg/ kg body weight	
maintenance	0.6–4.0 mg/ kg body weight/ hour	
etomidate		
loading dose	0.25 mg/ kg body weight as bolus within 30 sec	
maintenance	0.02 mg/ kg body weight per minute as continuous infusion	
<i>inhalative</i>		
isoflurane		
loading	0.5 (–3.0)% of end-tidal volume	
maintenance	0.8–2.0% of end-tidal volume	
<i>oral</i>		
topiramate	100–400 mg	every 12 hours
oxcarbazepine	600–1200 mg	every 12 hours; monitor sodium!
* dose must be adapted if renal function is impaired		

\* dose must be adapted if renal function is impaired

## Conclusions

Non-convulsive status epilepticus is one of the most important neurological emergencies requiring rapid diagnosis, being confirmed by EEG and treated without delay and with appropriate aggressiveness. The often unspectacular and un-specific clinical manifestation of NCSE makes it important to “think of it at all” in any patient presenting with unexplained new onset of behavioural changes, impaired consciousness and/or focal, non-convulsive neurological deficits. Treatment of NCSE closely follows the rules of the other forms of SE.

## References

- Kaplan PW. Assessing the outcomes in patients with nonconvulsive status epilepticus: nonconvulsive status epilepticus is underdiagnosed, potentially overtreated, and confounded by comorbidity. *J Clin Neurophysiol.* 1999;16:341–52.
- Calmeil LF. De l'épilepsie. Paris: Didot; 1824. p. 13.
- Wilson JVK, Reynolds EH. Translation and analysis of a cuneiform text forming part of a Babylonian treatise on epilepsy. *Med Hist.* 1990;34:185–98.
- Trousseau A. Lectures on Clinical Medicine. Volume 1. Translated by Bazier PV. London: New Sydenham Society; 1867. p. 53–4.
- Berger H. Über das Elektrenkephalogramm des Menschen. *Arch Psychiatrie.* 1929;87:527–70.
- Gibbs FA, Davis H, Lennox WG. The EEG in epilepsy and in the impaired states of consciousness. *Arch Neurol Psychiatry.* 1935;34:1133.
- Scott JS, Masland RL. Occurrence of “continuous symptoms” in epilepsy patients. *Neurology.* 1953;3:297–301.
- Penfield W, Jasper HH. Epilepsy and Functional Anatomy of the Human Brain. Boston: Little Brown; 1954.
- Landolt H. Die Dämmer- und Verstimmungszustände bei Epilepsie und ihre Elektroencephalographie. *Z Gesamte Inn Med.* 1963;185:411–30.
- Karbowski K. Status psychomotoricus. Klinische und elektroencephalographische Aspekte. In: Karbowski K, Herausgeber. Status psychomotoricus und seine Differentialdiagnose. Bern: Hans Huber; 1980. S. 39–71.
- Gastaut H. Clinical and electroencephalographic classification of epileptic seizures. *Epilepsia.* 1970;11:102–13.
- Delgado-Escueta AV, Wasterlain CG, Treiman DM, Porter RJ, editors. Status epilepticus. Mechanisms of brain damage and treatment. *Adv Neurol.* 1983;34:1–541.
- Commission on Classification and Terminology of the International League Against Epilepsy. Proposal for revised clinical and electroencephalographic classification of epileptic seizures. *Epilepsia.* 1989;30:389–99.
- Engel J Jr. Report of the ILAE Classification Core Group. *Epilepsia.* 2006;47:1558–68.
- Rona S, Rosenow F, Arnold S, Carreno M, Diehl B, Ebner A, et al. A semiological classification of status epilepticus. *Epileptic Disord.* 2005;7:5–12.
- Shorvon S. The classification of status epilepticus. *Epileptic Disord.* 2005;7:1–3.
- Shorvon S. What is non-convulsive status epilepticus, and what are its subtypes? *Epilepsia.* 2007;48(Suppl 8):S35–8.
- Rüegg SJ, Dichter MA. Diagnosis and treatment of nonconvulsive status epilepticus in an intensive care unit setting. *Curr Treat Options Neurol.* 2003;5:93–110.
- Working Group on Status Epilepticus. Treatment of convulsive status epilepticus. Recommendations of the Epilepsy Foundation of America's Working Group on Status Epilepticus. *JAMA.* 1993;270:854–9.

- 20 Treiman DM, Meyers PD, Walton NY, Collins JF, Colling C, Rowan AJ, et al. A comparison of four treatments for generalized convulsive status epilepticus. Veterans Affairs Status Epilepticus Cooperative Study Group. *N Engl J Med*. 1998;339:792–8.
- 21 Lowenstein DH, Bleck T, MacDonald RL. It's time to revise the definition of status epilepticus. *Epilepsia*. 1999;40:120–2.
- 22 Leppert D, Stöckli HR, Fuhr P. Richtlinien für die Behandlung des Status epilepticus. *Schweiz Ärztezeitung*. 2005;86:2097–100. Full e-text available by: [www.saez.ch/status\\_epilepticus\\_d.pdf](http://www.saez.ch/status_epilepticus_d.pdf).
- 23 Jordan KG. Continuous EEG monitoring in the neuroscience intensive care unit and emergency department. *J Clin Neurophysiol*. 1999;16:14–39.
- 24 Meierkord H, Holtkamp M. Non-convulsive status epilepticus in adults: clinical forms and treatment. *Lancet Neurol*. 2007;6:329–39.
- 25 Theodore WH, Porter RJ, Penry JK. Complex partial seizures: clinical characteristics and differential diagnosis. *Neurology*. 1983;33:1115–21.
- 26 Theodore WH, Porter RJ, Albert P, Kelley K, Bromfield E, Devinsky O, et al. The secondarily generalized tonic-clonic seizure: a videotape analysis. *Neurology*. 1994;44:1403–7.
- 27 Jenssen S, Gracely AJ, Sperling MR. How long do most seizures last? A systematic comparison of seizures recorded in the epilepsy monitoring unit. *Epilepsia*. 2006;47:1499–503.
- 28 Lowenstein D, Aminoff M, Simon R. Barbiturate anesthesia in the treatment of status epilepticus: clinical experience with 14 patients. *Neurology*. 1988;38:395–400.
- 29 Treiman DM, Walton NY, Kendrick C. A progressive sequence of electroencephalographic changes during generalized convulsive status epilepticus. *Epilepsy Res*. 1990;5:49–60.
- 30 Scholtes FB, Renier WO, Meinardi H. Generalized convulsive status epilepticus: causes, therapy, and outcome in 346 patients. *Epilepsia*. 1994;35:1104–12.
- 31 Fountain NB, Lothman EW. Pathophysiology of status epilepticus. *J Clin Neurophysiol*. 1995;12:326–42.
- 32 Fountain NB. Status epilepticus: risk factors and complications. *Epilepsia*. 2000;41(Suppl 2):S23–30.
- 33 Walker MC. Diagnosis and treatment of nonconvulsive status epilepticus. *CNS Drugs*. 2001;15:931–9.
- 34 Walker MC, White HS, Sander JW. Disease modification in partial epilepsy. *Brain*. 2002;125:1937–50.
- 35 Eriksson K, Metsäranta P, Huhtala H, Auvinen A, Kuusela AL, Koivikko M. Treatment delay and the risk of prolonged status epilepticus. *Neurology*. 2005;65:1316–8.
- 36 Sheth RD, Draskowski JF, Sirven JI, Gidal BE, Hermann BP. Prolonged ictal confusion in elderly patients. *Arch Neurol*. 2006;63:529–32.
- 37 Kaplan PW. The clinical features, diagnosis, and prognosis of nonconvulsive status epilepticus. *Neurologist*. 2005;11:348–61.
- 38 Tomson T, Lindbom U, Nilsson BY. Nonconvulsive status epilepticus in adults: thirty-two consecutive patients from a general hospital population. *Epilepsia*. 1992;33:829–35.
- 39 Gaitanis JN, Drislane FW. Status epilepticus: a review of different syndromes, their current evaluation, and treatment. *Neurologist*. 2003;9:61–76.
- 40 Tay SK, Hirsch LJ, Leary L, Jette N, Wittman J, Akman CI. Nonconvulsive status epilepticus in children: clinical and EEG characteristics. *Epilepsia*. 2006;47:1504–9.
- 41 Brenner RP. EEG in convulsive and nonconvulsive status epilepticus. *J Clin Neurophysiol*. 2004;21:319–31.
- 42 Thomas P, Beaumanoir A, Genton P, Dolisi C, Chatel M. “De novo” absence status of late onset: report of 11 cases. *Neurology*. 1992;42:104–10.
- 43 Thomas P, Borg M, Suisse G, Chatel M. Opercular myoclonic-anarthric status epilepticus. *Epilepsia*. 1995;36:281–9.
- 44 Raze A, Osborn MA, Vern BA, Molinari GF. Epileptic aphasia: first onset of prolonged monosymptomatic status epilepticus in adults. *Arch Neurol*. 1980;37:419–22.
- 45 Barry E, Sussman NM, Bosley TM, Harner RN. Ictal blindness and status epilepticus amauroticus. *Epilepsia*. 1985;26:577–84.
- 46 Roze E, Oubary P, Chédru F. Status-like recurrent pilomotor seizures: case report and review of the literature. *J Neurol Neurosurg Psychiatry*. 2000;68:647–9.
- 47 Armon C, Radtke RA, Friedman AH. Inhibitory simple partial (non-convulsive) status epilepticus after intracranial surgery. *J Neurol Neurosurg Psychiatry*. 2000;69:18–24.
- 48 Kramer RE, Lüders H, Goldstick LP, Dinner DS, Morris HH, Lesser RP, et al. Ictus emeticus: an electroclinical analysis. *Neurology*. 1988;38:1048–52.
- 49 DeToledo JC, Minagar A, Lowe MR. Persisting aphasia as the sole manifestation of partial status epilepticus. *Clin Neurol Neurosurg*. 2000;102:144–8.
- 50 Palmieri A, van Paesschen W, Dupont P, Van Laere K, van Driel G. Status gelasticus after temporal lobectomy: ictal FDG-PET findings and the question of dual pathology involving hypothalamic hamartomas. *Epilepsia*. 2005;46:1313–6.
- 51 McLachlan RS, Blume WT. Isolated fear in complex partial status epilepticus. *Ann Neurol*. 1980;8:639–41.
- 52 Sacquegnia T, Pazzaglia P, Baldrati A, De Carolis P, Gallassi R, Maccheroni M. Status epilepticus with cognitive symptomatology in a patient with partial complex epilepsy. *Eur Neurol*. 1981;20:319–25.
- 53 Wieser HG, Hailemariam S, Regard M, Landis T. Unilateral limbic epileptic status activity: stereo EEG, behavioural, and cognitive data. *Epilepsia*. 1985;26:19–29.
- 54 Lee BI, Lee BC, Hwang YM, Sohn YH, Jung JW, Park SC, et al. Prolonged ictal amnesia with transient focal abnormalities on magnetic resonance imaging. *Epilepsia*. 1992;33:1042–6.
- 55 Vuilleumier P, Despland PA, Regli F. Failure to recall (but not to remember): pure transient amnesia during nonconvulsive status epilepticus. *Neurology*. 1996;46:1036–9.
- 56 Kaplan PW. The EEG of status epilepticus. *J Clin Neurophysiol*. 2006;23:221–9.
- 57 Kaplan PW. Behavioral manifestations of nonconvulsive status epilepticus. *Epilepsy Behav*. 2002;3:122–39.
- 58 Wieser HG. Temporal lobe or psychomotor status epilepticus. A case report. *Electroencephalogr Clin Neurophysiol*. 1980;48:558–72.



- 59 Williamson PD, Spencer DD, Spencer SS, Novelly RA, Mattson RH. Complex partial status of frontal origin. *Ann Neurol.* 1985;18:497–504.
- 60 Williamson PD, Spencer DD, Spencer SS, Novelly RA, Mattson RH. Complex partial status epilepticus: a depth-electrode study. *Ann Neurol.* 1985;18:647–54.
- 61 Thomas P, Zifkin B, Migneco O, Lebrun C, Darcourt J, Andermann F. Nonconvulsive status epilepticus of frontal origin. *Neurology.* 1999;52:1174–83.
- 62 Ayres S. Sudden and temporary mental aberration – unconscious automatism – temporary irresponsible states. *JAMA.* 1902;38:315–6.
- 63 Drislane FW, Blum AS, Schomer DL. Focal status epilepticus: clinical features and significance of different EEG patterns. *Epilepsia.* 1999;40:1254–60.
- 64 Riggio S. Psychiatric manifestations of nonconvulsive status epilepticus. *Mt Sinai J Med.* 2006;73:960–6.
- 65 Reuber M, House AO. Treating patients with psychogenic non-epileptic seizures. *Curr Opin Neurol.* 2002;15:207–11.
- 66 Reuber M, Baker GA, Gill R, Smith DF, Chadwick DW. Failure to recognize psychogenic nonepileptic seizures may cause death. *Neurology.* 2004;62:834–5.
- 67 Towne AR. Epidemiology and outcomes of status epilepticus in the elderly. *Int Rev Neurobiol.* 2007;81:111–27.
- 68 Primavera A, Giberti L, Scotto P, Cocito L. Nonconvulsive status epilepticus as a cause of confusion in later life: report of 5 cases. *Neuropsychobiology.* 1994;30:148–52.
- 69 Fernandez-Torre JL, Diaz-Castroverde AG. Non-convulsive status epilepticus in elderly individuals: report of four representative cases. *Age Ageing.* 2004;33:78–81.
- 70 Kanemoto K, Tomikimi T, Kawasaki J. Prolonged post-ictal confusion as a manifestation of continuous complex partial status epilepticus: a depth EEG study. *Seizure.* 2000;9:151–5.
- 71 Cocito L, Audenino D, Primavera A. Altered mental state and nonconvulsive status epilepticus in patients with cancer. *Arch Neurol.* 2001;58:1310–5.
- 72 Lim J, Yagnik P, Schraeder P, Wheeler S. Ictal catatonia as a manifestation of nonconvulsive status epilepticus. *J Neurol Neurosurg Psychiatry.* 1986;49:833–6.
- 73 Swartz CM, Bottum KM, Salazar LS. Suppression of catatonia-like signs by lorazepam in nonconvulsive status epilepticus without seizure termination. *Am J Geriatr Psychiatry.* 2002;10:348–50.
- 74 McCarron MO, Tyagi A, Thomas SR, Sweeney BJ, Bone I. Non-convulsive status epilepticus of frontal origin following surgery for craniopharyngeoma. *Eur J Neurol.* 2002;9:178–9.
- 75 Jacobs DA, Fung KM, Cook NM, Schlaepfer WW, Goldberg HI, Stecker MM. Complex partial status epilepticus associated with anti-Hu paraneoplastic syndrome. *J Neurol Sci.* 2003;211:77–82.
- 76 Feddersen B, Bender A, Arnold S, Klopstock T, Noachtar S. Aggressive confusional state as a clinical manifestation of status epilepticus in MELAS. *Neurology.* 2003;61:1149–50.
- 77 Griffiths H, Farmer A. Psychotic depression presenting as status epilepticus. *Br J Psychiatry.* 1990;157:909–11.
- 78 Kümpfel T, Lechner C, Auer D, Kraft E, Lydtin H, Trenkwalder C. Non-convulsive status epilepticus with marked neuropsychiatric manifestations and MRI changes after treatment of hypercalcemia. *Acta Scand Neurol.* 2000;102:337–9.
- 79 Fernandez-Torre JL, Sanchez JM, Gonzalez C, Fernandez-Guinea O. Complex partial status epilepticus of extratemporal origin in a patient with systemic lupus erythematoses. *Seizure.* 2003;12:245–8.
- 80 Granner MA, Lee SJ. Nonconvulsive status epilepticus: EEG analysis in a large series. *Epilepsia.* 1994;35:42–7.
- 81 Brenner RP. The interpretation of the EEG in stupor and coma. *Neurologist.* 2005;11:271–84.
- 82 Chatrian GE, Shaw CM, Leffman H. The significance of periodic lateralized epileptiform discharges in EEG: an electrographic, clinical and pathological study. *Electroencephalogr Clin Neurophysiol.* 1964;17:177–93.
- 83 Pohlmann-Eden B, Hoch DB, Cochius JI, Chiappa KH. Periodic lateralized epileptiform discharges – a critical review. *J Clin Neurophysiol.* 1996;13:519–30.
- 84 Garzon E, Fernandez RM, Sakamoto AC. Serial EEG during human status epilepticus: evidence for PLED as an ictal pattern. *Neurology.* 2001;57:1175–83.
- 85 Assal F, Papazyan JP, Slosman DO, Jallon P, Goerres GW. SPECT in periodic lateralized epileptiform discharges (PLEDs): a form of partial status epilepticus? *Seizure.* 2001;10:260–4.
- 86 Dixit S, Kurlle P, Buyan-Dent L, Sheth RD. Status epilepticus associated with cefepime. *Neurology.* 2000;54:2153–5.
- 87 Barbey F, Bugnon D, Wauters JP. Severe neurotoxicity of cefepime in uremic patients. *Ann Intern Med.* 2001;135:1011.
- 88 Martinez-Rodriguez JE, Barriga FJ, Santamaria J, Iranzo A, Pareja JA, Revilla M, et al. Nonconvulsive status epilepticus associated with cephalosporins in patients with renal failure. *Am J Med.* 2001;111:115–9.
- 89 Kaplan PW. The EEG in metabolic encephalopathy and coma. *J Clin Neurophysiol.* 2004;21:307–18.
- 90 Foley JM, Watson CM, Adams RD. Significance of the electroencephalographic changes in hepatic coma. *Trans Am Neurol Assoc.* 1950;75:161–4.
- 91 Bickford RG, Butt HR. Hepatic coma: the encephalographic pattern. *J Clin Invest.* 1955;34:790–9.
- 92 Fountain NB, Waldman WA. Effects of benzodiazepines on triphasic waves. Implications for nonconvulsive status epilepticus. *J Clin Neurophysiol.* 2001;18:345–52.
- 93 Bertran F, Denise P, Letellier P. Nonconvulsive status epilepticus: the role of morphine and its antagonists. *Clin Neurophysiol.* 2000;30:109–12.
- 94 Boulanger JM, Deacon C, Lécuyer D, Gosselin S, Reiher J. Triphasic waves versus nonconvulsive status epilepticus: EEG distinction. *Can J Neurol Sci.* 2006;33:175–80.
- 95 DeLorenzo RJ, Waterhouse EJ, Towne AR, Boggs JG, Ko D, DeLorenzo GA, et al. Persistent nonconvulsive status epilepticus after the control of convulsive status epilepticus. *Epilepsia.* 1998;39:833–40.
- 96 Lowenstein DH, Aminoff MJ. Clinical and EEG features of status epilepticus in comatose patients. *Neurology.* 1992;42:100–4.
- 97 Jumao-as A, Brenner RP. Myoclonic status epilepticus: a clinical and electroencephalographic study. *Neurology.* 1990;40:1199–202.
- 98 Hallett M. Physiology of human posthypoxic myoclonus. *Mov Disord.* 2000;15(Suppl 1):S8–13.
- 99 DeLorenzo RJ, Hauser WA, Towne AR, Boggs JG, Pellock JM, Penberthy L, et al. A prospective, population-based epidemiologic study of status epilepticus in Richmond, Virginia. *Neurology.* 1996;46:1029–35.

- 100 Hesdorffer D, Logroscino G, Cascino G, Annegers JF, Hauser WA. Incidence of status epilepticus in Rochester Minnesota 1965–84. *Neurology*. 1998;50:735–41.
- 101 Coeytaux A, Jallon P, Galobardes B, Morabia A. Incidence of status epilepticus in French-speaking Switzerland (EPISTAR). *Neurology*. 2000;55:693–7.
- 102 Knake S, Rosenow F, Vescovi M, Oertel WH, Mueller HH, Wirbatz A, et al. Incidence of status epilepticus in adults in Germany: a prospective, population-based study. *Epilepsia*. 2001;42:714–8.
- 103 Vignatelli L, Tonon C, D'Alessandro R. Incidence and short-term prognosis of status epilepticus in adults in Bologna, Italy. *Epilepsia*. 2003;44:964–8.
- 104 Chin RF, Neville BG, Scott RC. A systematic review of the epidemiology of status epilepticus. *Eur J Neurol*. 2004;11:800–10.
- 105 Towne AR, Waterhouse EJ, Boggs JG, Garnett LK, Brown AJ, Smith Jr JT, et al. Prevalence of non-convulsive status epilepticus in comatose patients. *Neurology*. 2000;54:340–5.
- 106 Barry E, Hauser WA. Status epilepticus and antiepileptic medication levels. *Neurology*. 1994;44:47–50.
- 107 Spiller HA, Carlisle RD. Status epilepticus after massive carbamazepine overdose. *J Toxicol Clin Toxicol*. 2002;40:81–90.
- 108 Sazgar M, Bourgeois BF. Aggravation of epilepsy by antiepileptic drugs. *Pediatr Neurol*. 2005;33:227–34.
- 109 Kellinghaus C, Dziewas R, Lüdemann P. Tiagabine-related non-convulsive status epilepticus in partial epilepsy: three case reports and a review of the literature. *Seizure*. 2002;11:243–9.
- 110 Vinton A, Kornberg AJ, Cowley M, Matkovic Z, Kilpatrick C, O'Brien TJ. Tiagabine-induced generalised non convulsive status epilepticus in patients with lesional focal epilepsy. *J Clin Neurosci*. 2005;12:128–33.
- 111 Osorio L, Reed RC, Peltzer JN. Refractory idiopathic absence status epilepticus: a probable paradoxical effect of phenytoin and carbamazepine. *Epilepsia*. 2000;41:887–94.
- 112 Thomas P, Valton L, Genton P. Absence and myoclonic status epilepticus precipitated by antiepileptic drugs in idiopathic generalized epilepsy. *Brain*. 2006;129:1281–92.
- 113 Vespa PM, Nuwer MR, Nenov V, Ronne-Engstrom E, Hovda DA, Bergsneider M, et al. Increased incidence and impact of nonconvulsive and convulsive seizures after traumatic brain injury as detected by continuous electroencephalographic monitoring. *J Neurosurg*. 1999;91:750–60.
- 114 Velioglu SK, Ozmenoglu M, Boz C, Alioglu Z. Status epilepticus after stroke. *Stroke*. 2001;32:1169–72.
- 115 Afsar N, Kaya D, Aktan S, Aykut-Bingol C. Stroke and status epilepticus: stroke type, type of status epilepticus, and prognosis. *Seizure*. 2003;12:23–7.
- 116 Little AS, Kerrigan JF, McDougall CG, Zabramski JM, Albuquerque FC, Nakaji P, et al. Nonconvulsive status epilepticus in patients suffering spontaneous subarachnoid hemorrhage. *J Neurosurg*. 2007;106:805–11.
- 117 Hormigo A, Liberato B, Lis E, DeAngelis LM. Nonconvulsive status epilepticus in patients with cancer. Imaging abnormalities. *Arch Neurol*. 2004;61:362–5.
- 118 Blitshteyn S, Jaeckle KA. Nonconvulsive status epilepticus in metastatic CNS disease. *Neurology*. 2006;66:1261–3.
- 119 Cavaliere R, Farace E, Schiff D. Clinical implications of status epilepticus in patients with neoplasms. *Arch Neurol*. 2006;63:1746–9.
- 120 Gunduz A, Beskardes AF, Kutlu A, Ozkara C, Karaagac N, Yeni SN. Herpes encephalitis as a cause of non-convulsive status epilepticus. *Epileptic Disord*. 2006;8:57–60.
- 121 Ances BM, Shellhaus R, Brown MJ, Rios OV, Herman ST, French JA. Neurosyphilis and status epilepticus: case report and literature review. *Epilepsy Res*. 2004;59:67–70.
- 122 Cohen D, Kutluay E, Edwards J, Peltier A, Beydoun A. Sporadic Creutzfeldt-Jakob disease presenting with nonconvulsive status epilepticus. *Epilepsy Behav*. 2004;5:792–6.
- 123 Rosen I. Electroencephalography as a diagnostic tool in dementia. *Dement Geriatr Cogn Disord*. 1997;8:110–6.
- 124 Armon C, Peterson GW, Liwnicz BH. Alzheimer's disease underlies some cases of complex partial status epilepticus. *J Clin Neurophysiol*. 2000;17:511–8.
- 125 Manford M, Fuller GN, Wade JP. "Silent diabetes": non-ketotic hyperglycemia presenting as aphasic status epilepticus. *J Neurol Neurosurg Psychiatry*. 1995;59:99–100.
- 126 Kawai N, Miyake K, Kuroda Y, Yamashita S, Nishiyama Y, Monden T, et al. Magnetic resonance imaging and positron emission tomography findings in status epilepticus following severe hypoglycaemia. *Ann Nucl Med*. 2006;20:371–6.
- 127 Safe AF, Griffiths KD, Maxwell RT. Thyrotoxic crisis presenting as status epilepticus. *Postgrad Med J*. 1990;66:150–2.
- 128 Woods KL, Holmes GK. Myxoedema coma presenting in status epilepticus. *Postgrad Med J*. 1977;53:46–8.
- 129 Chong JY, Rowland LP, Utiger RD. Hashimoto encephalopathy: syndrome or myth? *Arch Neurol*. 2003;60:164–71.
- 130 Schäuble B, Castillo PR, Boeve BF, Westmoreland BF. EEG findings in steroid-responsive encephalopathy associated with autoimmune thyroiditis. *Clin Neurophysiol*. 2003;114:32–7.
- 131 McKeon A, McNamara B, Sweeney B. Hashimoto's encephalopathy presenting with psychosis and generalized absence status. *J Neurol*. 2004;251:1025–7.
- 132 Ferlazzo E, Raffaele M, Mazzu I, Pisani F. Recurrent status epilepticus as the main feature of Hashimoto's encephalopathy. *Epilepsy Behav*. 2006;8:328–30.
- 133 Tsai MH, Lee LH, Chen SD, Lu CH, Chen MT, Chuang YC. Complex partial status epilepticus as a manifestation of Hashimoto's encephalopathy. *Seizure*. 2007;16:713–6.
- 134 Primavera A, Fonti A, Giberti L, Cocito L. Recurrent absence status epilepticus and hyponatremia in a patient with polydipsia. *Biol Psychiatry*. 1995;38:189–91.
- 135 Kline CA, Esekogwu VI, Henderson SO, Newton KI. Non-convulsive status epilepticus in a patient with hypocalcemia. *J Emerg Med*. 1998;16:715–8.
- 136 Krumholz A. Epidemiology and evidence for morbidity of nonconvulsive status epilepticus. *J Clin Neurophysiol*. 1999;16:314–22.



- 137 Conolly RM, Doherty CP, Beddy P, O'Byrne K. Chemotherapy induced reversible posterior leukoencephalopathy syndrome. *Lung Cancer*. 2007;56:459–63.
- 138 Hillbom M, Pieninkeroinen I, Leone M. Seizures in alcohol-dependent patients: epidemiology, pathophysiology and management. *CNS Drugs*. 2003;17:1013–20.
- 139 Miyata H, Kubota F, Shibata N, Kifune A. Non-convulsive status epilepticus induced by antidepressants. *Seizure*. 1997;6:405–7.
- 140 Bottaro FJ, Martinez OA, Pardal MM, Bruetman JE, Reisin RC. Nonconvulsive status epilepticus in the elderly: a case-control study. *Epilepsia*. 2007;48:966–72.
- 141 Waterhouse EJ, DeLorenzo RJ. Status epilepticus in older patients: epidemiology and treatment options. *Drugs Aging*. 2001;18:133–42.
- 142 Sung CY, Chu NS. Status epilepticus in the elderly: etiology, seizure type, and outcome. *Acta Neurol Scand*. 1989;80:51–6.
- 143 Husain AM, Horn GJ, Jacobson MP. Non-convulsive status epilepticus: usefulness of clinical features in selecting patients for urgent EEG. *J Neurol Neurosurg Psychiatry*. 2003;74:189–91.
- 144 Dunne JW, Summers QA, Stewart-Wynne EG. Non-convulsive status epilepticus: a prospective study in an adult general hospital. *QJM*. 1987;238:117–26.
- 145 Haffey S, McKernan A, Pang K. Non-convulsive status epilepticus: a profile of patients diagnosed within a tertiary referral centre. *J Neurol Neurosurg Psychiatry*. 2004;75:1043–4.
- 146 Brenner RP. Is it status? *Epilepsia*. 2002;43(Suppl 3):S103–13.
- 147 Barry E, Hauser WA. Pleocytosis after status epilepticus. *Arch Neurol*. 1994;51:190–3.
- 148 DeGiorgio CM, Gott PS, Rabinowicz AL, Heck CN, Smith TD, Correale TD. Neuron-specific enolase, a marker for acute neuronal injury, is increased in complex partial status epilepticus. *Epilepsia*. 1996;37:606–9.
- 149 DeGiorgio CM, Correale JD, Gott PS, Ginsburg DL, Bracht KA, Smith T, et al. Serum neuron-specific enolase in human status epilepticus. *Neurology*. 1995;45:1134–7.
- 150 DeGiorgio CM, Heck CN, Rabinowicz AL, Gott PS, Smith T, Correale J. Serum neuron-specific enolase in the major subtypes of status epilepticus. *Neurology*. 1999;52:746–9.
- 151 Chung PW, Seo DW, Kwon JC, Kim H, Na DL. Nonconvulsive status epilepticus presenting as a sub-acute progressive aphasia. *Seizure*. 2002;11:449–52.
- 152 Oishi M, Otsubo H, Kameyama S, Wachi M, Tanaka K, Masuda H, et al. Ictal magnetoencephalographic discharges from elementary visual hallucinations of status epilepticus. *J Neurol Neurosurg Psychiatry*. 2003;74:525–7.
- 153 Kutulay E, Beattie J, Passaro EA, Edwards JC, Minecan D, Milling C, et al. Diagnostic and localizing value of ictal SPECT in patients with nonconvulsive status epilepticus. *Epilepsy Behav*. 2005;6:212–7.
- 154 Wiest R, von Bredow F, Schindler K, Schäuble B, Slotboom J, Brekenfeld C, et al. Detection of regional blood perfusion changes in epileptic seizures with dynamic brain perfusion CT – a pilot study. *Epilepsy Res*. 2006;72:102–10.
- 155 Lansberg MG, O'Brain MW, Norbash AM, Moseley ME, Morell M, Albers GW. MRI abnormalities associated with partial status epilepticus. *Neurology*. 1999;52:1021–7.
- 156 Lazeyras F, Blanke O, Zimine I, Delavelle J, Perrig SH, Seeck M. MRI, <sup>1</sup>H-MRS, and functional MRI during and after prolonged nonconvulsive seizure activity. *Neurology*. 2000;55:1677–82.
- 157 Hong K-S, Cho YJ, Lee SK, Jeong SW, Kima WK, Oh EJ. Diffusion changes suggesting predominant vasogenic oedema during partial status epilepticus. *Seizure*. 2004;13:317–21.
- 158 Szabo K, Poepel A, Pohlmann-Eden B, Hirsch J, Back T, Sedlaczek O, et al. Diffusion-weighted and perfusion MRI demonstrates parenchymal changes in complex partial status epilepticus. *Brain*. 2005;128:1369–76.
- 159 Chen JW, Wasterlain CG. Status epilepticus: pathophysiology and management in adults. *Lancet Neurol*. 2006;5:246–56.
- 160 Murdoch D. Mechanisms of status epilepticus: an evidence-based review. *Curr Opin Neurol*. 2007;20:213–6.
- 161 DeLorenzo RJ, Sun DA. Basic mechanisms in status epilepticus: role of calcium in neuronal injury and the induction of epileptogenesis. *Adv Neurol*. 2006;97:187–97.
- 162 Shorvon S. The management of status epilepticus. *J Neurol Neurosurg Psychiatry*. 2001;70(Suppl 2):S22–7.
- 163 Manno EM, Pfeifer EA, Cascino GD, Noe KH, Wijdicks EF. Cardiac pathology in status epilepticus. *Ann Neurol*. 2005;58:954–7.
- 164 Leung H, Kwan P, Elger CE. Finding the missing link between ictal bradyarrhythmia, ictal asystole, and sudden unexpected death in epilepsy. *Epilepsy Behav*. 2006;9:19–30.
- 165 Logroscino G, Hesdorffer DC, Cascino G, Hauser WA, Coeytaux A, Galobardes B, et al. Mortality after a first episode of status epilepticus in the United States and Europe. *Epilepsia*. 2005;46(Suppl 11):S46–8.
- 166 Logroscino G, Hesdorffer DC, Cascino G, Annegers JF, Hauser WA. Short-term mortality after a first episode of status epilepticus. *Epilepsia*. 1997;38:1344–9.
- 167 Logroscino G, Hesdorffer DC, Cascino GD, Annegers JF, Bagiella E, Hauser WA. Long-term mortality after a first episode of status epilepticus. *Neurology*. 2002;58:537–41.
- 168 Shneker BF, Fountain NB. Assessment of acute morbidity and mortality in nonconvulsive status epilepticus. *Neurology*. 2003;61:1066–73.
- 169 Young GB, Jordan KG, Doig GS. An assessment of nonconvulsive seizures in the intensive care unit using continuous EEG monitoring: an investigation of variables associated with mortality. *Neurology*. 1996;47:83–9.
- 170 Hesdorffer DC, Logroscino G, Cascino G, Annegers JF, Hauser WA. Risk of unprovoked seizure after acute symptomatic seizure: effect of status epilepticus. *Ann Neurol*. 1998;44:908–12.
- 171 Hesdorffer DC, Logroscino G, Cascino GD, Hauser WA. Recurrence of afebrile status epilepticus in a population-based study in Rochester, Minnesota. *Neurology*. 2007;69:73–8.
- 172 Aminoff M. Do nonconvulsive seizures damage the brain? – No. Controversies in neurology. *Arch Neurol*. 1998;55:119–21.

- 173 Young GG, Jordan KG. Do nonconvulsive seizures damage the brain? – Yes. *Arch Neurol.* 1998;55:117–20.
- 174 Kaplan PW. No, some types of nonconvulsive status epilepticus cause little permanent neurologic sequelae (or: “the cure may be worse than the disease”). *Neurophysiol Clin.* 2000;30:377–82.
- 175 Kaplan PW. Prognosis in nonconvulsive status epilepticus. *Epileptic Disord.* 2000;2:185–93.
- 176 Sloviter RS. The neurobiology of temporal lobe epilepsy: too much information, not enough knowledge. *C R Biol.* 2005;328:143–53.
- 177 Sutula T, Pitkänen A, editors. Do seizures damage the brain? *Prog Brain Res.* 2002;135:1–531.
- 178 Wasterlain CG, Fujikawa DG, Penix L, Sankar R. Pathophysiological mechanisms of brain damage from status epilepticus. *Epilepsia.* 1993;34(Suppl 1):S37–53.
- 179 Fujikawa DG, Itabashi HH, Wu A, Shinmei SS. Status epilepticus-induced neuronal loss in humans without systemic complications or epilepsy. *Epilepsia.* 2000;41:981–91.
- 180 Men S, Lee DH, Barron JR, Munoz DG. Selective neuronal necrosis associated with status epilepticus: MR findings. *AJNR.* 2000;21:1837–40.
- 181 Nixon J, Bateman D, Moss T. An MRI and neuropathological study of a case of fatal status epilepticus. *Seizure.* 2001;10:588–91.
- 182 Shiraska Y. Lack of neuronal damage in atypical absence status epilepticus. *Epilepsia.* 2002;43:1498–501.
- 183 Young GB. Status epilepticus and brain damage: pathology and pathophysiology. *Adv Neurol.* 2006;97:217–20.
- 184 Rossetti AO, Logroscino G, Liaudet L, Ruffieux C, Ribordy V, Schaller MD, et al. Status epilepticus. An independent outcome predictor after cerebral anoxia. *Neurology.* 2007;69:255–60.
- 185 Rossetti AO, Logroscino G, Bromfield EB. A clinical score for prognosis of status epilepticus in adults. *Neurology.* 2006;66:1736–8.
- 186 Krumholz A, Stern BJ, Weiss HD. Outcome from coma after cardiopulmonary resuscitation: relation to seizures and myoclonus. *Neurology.* 1988;38:401–5.
- 187 Wijdicks EF, Parisi JE, Sharbrough FW. Prognostic value of myoclonus status in comatose survivors of cardiac arrest. *Ann Neurol.* 1994;35:239–43.
- 188 Hui AC, Cheng C, Lam A, Mok V, Joynt GM. Prognosis following postanoxic myoclonus status epilepticus. *Eur Neurol.* 2005;54:10–3.
- 189 Walker MC. Treatment of nonconvulsive status epilepticus. *Int Rev Neurobiol.* 2007;81:287–97.
- 190 Arendt RM, Greenblatt DJ, deJong RH, Bonin JD, Abernethy DR, Ehrenberg BL, et al. In vitro correlates of benzodiazepine cerebrospinal fluid uptake, pharmacodynamic action and peripheral distribution. *J Pharmacol Exp Ther.* 1983;227:98–106.
- 191 Treiman DM. Pharmacokinetics and clinical use of benzodiazepines in the management of status epilepticus. *Epilepsia.* 1989;30(Suppl 2):S4–10.
- 192 Rey E, Treluyer JM, Pons G. Pharmacokinetic optimization of benzodiazepine therapy for acute seizures. Focus on delivery routes. *Clin Pharmacokinet.* 1999;36:409–24.
- 193 Walker MC, Tong X, Brown S, Shorvon SD, Patsalos PN. Comparison of single- and repeated-dose pharmacokinetics of diazepam. *Epilepsia.* 1998;39:283–9.
- 194 Lowenstein DH. Treatment options for status epilepticus. *Curr Opin Pharmacol.* 2005;5:334–9.
- 195 Terndrup TE, Fordyce WE. Respiratory drive during status epilepticus and its treatment: comparison of diazepam and lorazepam. *Epilepsy Res.* 1995;20:21–30.
- 196 Leppik IE, Derivan AT, Homan RW, Walker J, Ramsay E, Patrick B. Double-blind study of lorazepam and diazepam in status epilepticus. *JAMA.* 1983;249:1452–4.
- 197 Levy RJ, Krall RL. Treatment of status epilepticus with lorazepam. *Arch Neurol.* 1984;41:605–11.
- 198 Appleton R, Sweeney A, Choonara I, Robson J, Molyneux E. Lorazepam versus diazepam in the acute treatment of epileptic seizures and status epilepticus. *Dev Med Child Neurol.* 1995;37:682–8.
- 199 Cock HR, Schapira AHV. A comparison of lorazepam and diazepam as initial therapy in convulsive status epilepticus. *Q J Med.* 2002;95:225–31.
- 200 Alldredge BK, Gelb AM, Isaacs SM, Corry MD, Allen F, Ulrich S, et al. A comparison of lorazepam, diazepam, and placebo for the treatment of out-of-hospital status epilepticus. *N Engl J Med.* 2001;345:631–7.
- 201 Parent JM, Lowenstein DH. Treatment of refractory generalized status epilepticus with continuous infusion of midazolam. *Neurology.* 1994;44:1837–40.
- 202 Koul RL, Aithala GR, Chacko A, Joshi R, Elbualy MS. Continuous midazolam infusion as treatment of status epilepticus. *Arch Dis Child.* 1997;76:445–8.
- 203 Fountain NB, Adams RE. Midazolam treatment of acute and refractory status epilepticus. *Clin Neuropharmacol.* 1999;22:261–7.
- 204 Towne AR, DeLorenzo RJ. Use of intramuscular midazolam for status epilepticus. *J Emerg Med.* 1999;17:323–8.
- 205 Gastaut H, Courjon J, Poiré R, Weber M. Treatment of status epilepticus with a new benzodiazepine more active than diazepam. *Epilepsia.* 1971;12:197–214.
- 206 Singh AN, Le Morvan P. Treatment of status epilepticus with intravenous clonazepam. *Prog Neuropsychopharmacol Biol Psychiatry.* 1982;6:539–42.
- 207 Greenblatt DJ, Miller LG, Shader RI. Clonazepam pharmacokinetics, brain uptake, and receptor interactions. *J Clin Psychiatry.* 1987;48(Suppl):S4–11.
- 208 Merriam AE, Medalia A, Levine B. Partial complex status epilepticus associated with cocaine abuse. *Biol Psychiatry.* 1988;23:515–8.
- 209 Alldredge BK, Lowenstein DH, Simon RP. Seizures associated with recreational drug abuse. *Neurology.* 1989;39:1037–9.
- 210 Dhuna A, Pascual-Leone A, Langendorf F, Anderson DC. Epileptogenic properties of cocaine in humans. *Neurotoxicology.* 1991;12:621–6.
- 211 Refstad S. Paramethoxyamphetamine (PMA) poisoning: a “party drug” with lethal effects. *Acta Anaesthesiol Scand.* 2003;53:1298–9.
- 212 Staack RF. Piperazine designer drugs of abuse. *Lancet.* 2007;369:1411–3.
- 213 Wood DM, Dargan PI, Button J, Holt DW, Ovaska H, Ramsey J, et al. Collapse, reported seizure – and an unexpected pill. *Lancet.* 2007;369:1490.

- 214 Gee P, Richardson S, Woltersdorf W, Moore G. Toxic effects of BZP-based herbal party-pills in humans: a prospective study in Christchurch, New Zealand. *N Z Med J*. 2005;118:U1784.
- 215 McWilliam PK. Intravenous phenytoin sodium in continuous convulsions in children. *Lancet*. 1958;272:1147–9.
- 216 Wallis W, Kutt H, McDowell F. Intravenous diphenylhydantoin in treatment of acute repetitive seizures. *Neurology*. 1968;18:513–25.
- 217 Russell MA, Bousvaros G. Fatal results from diphenylhydantoin administered intravenously. *JAMA*. 1968;206:2118–9.
- 218 Burneo JG, Anandan JV, Barkley GL. A prospective study of the incidence of the purple glove syndrome. *Epilepsia*. 2001;42:1156–9.
- 219 O'Brien TJ, Meara FM, Matthews H, Vajda FJ. Prospective study of local cutaneous reactions in patients receiving IV phenytoin. *Neurology*. 2001;57:1508–10.
- 220 Nation RL, Evans AM, Milne RW. Pharmacokinetic drug interactions with phenytoin (part I). *Clin Pharmacokinet*. 1990;18:37–60.
- 221 Nation RL, Evans AM, Milne RW. Pharmacokinetic drug interactions with phenytoin (part II). *Clin Pharmacokinet*. 1990;18:131–50.
- 222 Cuttle L, Munns AJ, Hogg NA, Scott JR, Hooper WD, Dickinson RG, et al. Phenytoin metabolism by human cytochrome P450: involvement of P450 3A and 2C forms in secondary metabolism and drug-protein adduct formation. *Drug Metab Dispos*. 2000;28:1361–8.
- 223 Relling MV, Pui CH, Sandlund JT, Rivera GK, Hancock ML, Boyett JM, et al. Adverse effect of anticonvulsants on efficacy of chemotherapy for acute lymphoblastic leukaemia. *Lancet*. 2000;356:285–90.
- 224 Rüegg S. Dexamethasone/phenytoin interactions: neurooncological concerns. *Swiss Med Wkly*. 2002;132:425–6.
- 225 Pryor FM, Gidal B, Ramsay RE, DeToledo J, Morgan RO. Fosphenytoin: pharmacokinetics and tolerance of intramuscular loading doses. *Epilepsia*. 2001;42(2):245–50.
- 226 Browne TR. Fosphenytoin (Cerebyx). *Clin Neuropharmacol*. 1997;20:1–12.
- 227 Fischer JH, Patel TV, Fischer PA. Fosphenytoin: clinical pharmacokinetics and comparative advantages in the acute treatment of seizures. *Clin Pharmacokinet*. 2003;42:33–58.
- 228 DeToledo JC, Ramsay RE. Fosphenytoin and phenytoin in patients with status epilepticus. Improved tolerability versus increased costs. *Drug Safety*. 2000;22:459–66.
- 229 Touchette DR, Rhoney DH. Cost-minimization analysis of phenytoin and fosphenytoin in the emergency department. *Pharmacotherapy*. 2000;20:908–16.
- 230 Nitsche V, Mascher H. The pharmacokinetics of valproic acid after oral and parenteral administration in healthy volunteers. *Epilepsia*. 1982;23:153–62.
- 231 Hönack D, Löscher W. Intravenous valproate: onset and duration of anticonvulsant activity against a series of electroconvulsions in comparison with diazepam and phenytoin. *Epilepsy Res*. 1992;13:215–21.
- 232 Kaplan PW. Intravenous valproate treatment of generalized nonconvulsive status epilepticus. *Clin Electroencephalogr*. 1999;30:1–4.
- 233 Perucca E, Gatti G, Frigo GM, Crema A. Pharmacokinetics of valproic acid after oral and intravenous administration. *J Clin Pharmacol*. 1978;5:313–8.
- 234 Ramsay RE, Uthman B, Leppik IE, Pellock JM, Wilder BJ, Morris D, et al. The tolerability and safety of valproate sodium injection given as an intravenous infusion. *J Epilepsy*. 1997;10:187–93.
- 235 Löscher W. Basic pharmacology of valproate. A review after 35 years of clinical use for the treatment of epilepsy. *CNS Drugs*. 2002;16:669–94.
- 236 Perucca E. Pharmacological and therapeutic properties of valproate: a summary after 35 years of clinical experience. *CNS Drugs*. 2002;16:695–714.
- 237 Lindsay C, De Vane LC. Pharmacokinetics, drug interactions, and tolerability of valproate. *Psychopharmacol Bull*. 2003;37:25–42.
- 238 Cloyd JC, Dutta S, Cao G, Walch JK, Collins SD, Granneman GR. Valproate unbound fraction and distribution volume following rapid infusions in patients with epilepsy. *Epilepsy Res*. 2003;53:19–27.
- 239 Hamer HM, Knake S, Schomburg U, Rosenow F. Valproate-induced hyperammonemic encephalopathy in the presence of topiramate. *Neurology*. 2000;54:230–2.
- 240 Verrotti A, Trotta D, Morgese G, Chiarelli F. Valproate-induced hyperammonemic encephalopathy. *Metab Brain Dis*. 2002;17:367–73.
- 241 Segura-Bruna N, Rodriguez-Campello A, Puente V, Roquer J. Valproate-induced hyperammonemic encephalopathy. *Acta Neurol Scand*. 2007;114:1–7.
- 242 Gomez-Bellver MJ, Garcia-Sanchez MJ, Alonso-Gonzalez AC, Santos-Beluga D, Dominguez-Gil A. Plasma protein binding kinetics of valproic acid over a broad dosage range: therapeutic implications. *J Clin Pharm Ther*. 1993;18:191–7.
- 243 Hermida J, Tutor JC. A theoretical method for normalizing total serum valproic acid concentrations in hypoalbuminemic patients. *J Pharmacol Sci*. 2005;97:489–93.
- 244 Gidal B, Spencer N, Maly M, Pitterle M, Williams E, Collins MN, et al. Valproate-mediated disturbances of hemostasis: relationship to dose and plasma concentration. *Neurology*. 1994;44:1418–22.
- 245 Easterford K, Clough P, Kellett M, Fallon K, Duncan S. Reversible parkinsonism with normal  $\beta$ -CIT-SPECT in patients exposed to sodium valproate. *Neurology*. 2004;62:1435–7.
- 246 Gerstner T, Busing D, Bell N, Longin E, Kasper JM, Klostermann W, et al. Valproic acid-induced pancreatitis: 16 new cases and a review of the literature. *J Gastroenterol*. 2007;42:39–48.
- 247 Loiseau P. Sodium valproate, platelet dysfunction, and bleeding. *Epilepsia*. 1981;22:141–6.
- 248 Kreuz W, Linde R, Funk M, Meyer-Schrod R, Föll E, Nowak-Göttl U, et al. Valproate therapy induces von Willebrand disease Typ I. *Epilepsia*. 1992;33:178–84.
- 249 Devinsky O, Leppik I, Wilmore I, Pellock J, Dean C, Gates J. Safety of intravenous valproate. *Ann Neurol*. 1995;38:670–4.
- 250 Wheless J, Venkataraman V. Safety of high intravenous valproate loading doses in epilepsy patients. *J Epilepsy*. 1998;11:319–24.
- 251 Venkataraman V, Wheless JW. Safety of rapid intravenous infusion of valproate loading doses in epilepsy patients. *Epilepsy Res*. 1999;35:147–53.



- 252 Naritoku DK, Mued S. Intravenous loading of valproate for epilepsy. *Clin Neuropharmacol.* 1999;22:102–6.
- 253 Limdi NA, Faught E. The safety of rapid valproic acid infusion. *Epilepsia.* 2000;41:1342–5.
- 254 Sinha S, Naritoku DK. Intravenous valproate is well tolerated in unstable patients with status epilepticus. *Neurology.* 2000;55:722–4.
- 255 Dutta S, Cloyd JC, Granneman GR, Collins SD. Oral/intravenous maintenance dosing of valproate following intravenous loading: a simulation. *Epilepsy Res.* 2003;53:29–38.
- 256 Ramsay RE, Cantrell D, Collins SD, Walch JK, Naritoku DK, Cloyd JC, et al. Safety and tolerance of rapidly infused Depacon®. A randomized trial in subjects with epilepsy. *Epilepsy Res.* 2003;52:189–201.
- 257 Wheless JW, Vazquez BR, Kanner AM, Ramsay RE, Morton L, Pellock JM. Rapid infusion with valproate sodium is well tolerated in patients with epilepsy. *Neurology.* 2004;63:1507–8.
- 258 Limdi NA, Shimpi AV, Faught E, Gomez CR, Burneo JG. Efficacy of rapid IV administration of valproic acid for status epilepticus. *Neurology.* 2005;64:353–5.
- 259 Peters CNA, Pohlmann-Eden B. Intravenous valproate as an innovative therapy in seizure emergency situations including status epilepticus-experience in 102 adult patients. *Seizure.* 2005;14:164–9.
- 260 Krämer G, Bergmann A, Despland PA, König S, Kurlmann G, Kurth C, et al. Derzeitiger Stellenwert intravenöser Valproinsäure in der Therapie des generalisierten tonisch-klonischen Status epilepticus. *Akt Neurol.* 2005;32:263–74.
- 261 Misra UK, Kalita J, Patel R. Sodium valproate vs. phenytoin in status epilepticus: a pilot study. *Neurology.* 2006;67:340–2.
- 262 Rossetti AO. Sodium valproate vs. phenytoin in status epilepticus: a pilot study. *Neurology.* 2007;68:709.
- 263 Agarwal P, Kumar N, Chandra R, Gupta G, Antony AR, Garg N. Randomized study of intravenous valproate and phenytoin in status epilepticus. *Seizure.* 2007;16:527–32.
- 264 Perucca E, Gidal BE, Baltes E. Effects of antiepileptic comedication on levetiracetam pharmacokinetics: a pooled analysis of data from randomized adjunctive therapy trials. *Epilepsy Res.* 2003;53:47–56.
- 265 Sirsi D, Safdieh JE. The safety of levetiracetam. *Expert Opin Drug Saf.* 2007;6:241–50.
- 266 De Smedt T, Raedt R, Vonck K, Boon P. Levetiracetam: part II, the clinical profile of a novel anticonvulsant drug. *CNS Drug Rev.* 2007;13:57–78.
- 267 French J, Arrigo C. Rapid onset of action of levetiracetam in refractory epilepsy patients. *Epilepsia.* 2005;46:324–6.
- 268 Ramael S, De Smedt F, Toubanc N, Otoul C, Boulanger P, Riethuisen JM, et al. Single-dose bio-availability of levetiracetam intravenous infusion relative to oral tablets and multiple-dose pharmacokinetics and tolerability of levetiracetam intravenous infusion compared with placebo in healthy subjects. *Clin Ther.* 2006;28:734–43.
- 269 Ramael S, Daoust A, Otoul C, Toubanc N, Troenaru M, Lu Z, et al. Levetiracetam intravenous infusion: a randomized, placebo-controlled safety and pharmacokinetic study. *Epilepsia.* 2006;47:1128–35.
- 270 Mazarati AM, Baldwin R, Klitgaard H, Matagne A, Wasterlain CG. Anticonvulsant effects of levetiracetam and levetiracetam-diazepam combinations in experimental status epilepticus. *Epilepsy Res.* 2004;58:167–74.
- 271 Rossetti A, Bromfield EB. Levetiracetam in the treatment of status epilepticus in adults: a study of 13 episodes. *Eur Neurol.* 2005;54:34–8.
- 272 Rossetti AO, Bromfield EB. Determinants of success in the use of oral levetiracetam in status epilepticus. *Epilepsy Behav.* 2006;8:651–4.
- 273 Falip M, Carreno M, Amaro S, Donaire A, Delgado R, Toledo M, et al. Use of levetiracetam in hospitalized patients. *Epilepsia.* 2006;47:2186–8.
- 274 Rupprecht S, Franke K, Fitzek S, Witte OW, Hagemann G. Levetiracetam as a treatment option in non-convulsive status epilepticus. *Epilepsy Res.* 2007;73:238–44.
- 275 Schulze-Bonhage A, Hefft S, Oehl B. Termination of complex partial status epilepticus by intravenous levetiracetam – a case report. *J Neurol Neurosurg Psychiatry.* 2007; online first, published on March 12, 2007 as 10.1136/jnnp.2006.113951
- 276 Atefy R, Tettenborn B. Nonconvulsive status epilepticus on treatment with levetiracetam. *Epilepsy Behav.* 2005;6:613–6.
- 277 Nakken KO, Eriksson AS, Lossius R, Johannessen SI. A paradoxical effect of levetiracetam may be seen in both children and adults with refractory epilepsy. *Seizure* 2003;12:42–6.
- 278 Claassen J, Lokin JK, Fitzsimmons BF, Mendelsohn FA, Mayer SA. Predictors of functional disability and mortality after status epilepticus. *Neurology.* 2002;58:139–42.
- 279 Mayer SA, Claassen J, Lokin J, Mendelsohn F, Dennis LJ, Fitzsimmons BF. Refractory status epilepticus. Frequency, risk factors, and impact on outcome. *Arch Neurol.* 2002;59:205–10.
- 280 Rossetti AO, Logroscino G, Bromfield EB. Refractory status epilepticus: effect of treatment aggressiveness on prognosis. *Arch Neurol.* 2005;62:1698–702; Erratum in: *Arch Neurol.* 2006;63:1482.
- 281 Holtkamp M. The anaesthetic use and intensive care of status epilepticus. *Curr Opin Neurol.* 2007;20:188–93.
- 282 Kumar A, Bleck TP. Intravenous midazolam for the treatment of refractory status epilepticus. *Crit Care Med.* 1992;20:483–8.
- 283 Labar DR, Ali A, Root J. High-dose intravenous lorazepam for the treatment of refractory status epilepticus. *Neurology.* 1994;44:1400–3.
- 284 Claassen J, Hirsch LJ, Emerson RG, Bates JE, Thompson TB, Mayer SA. Continuous EEG monitoring and midazolam infusion for refractory nonconvulsive status epilepticus. *Neurology.* 2001;57:1036–42.
- 285 Denzel D, Burstein AH. Midazolam in refractory status epilepticus. *Ann Pharmacother.* 1996;30:1481–3.
- 286 Mackenzie SJ, Kapadia F, Grant IS. Propofol infusion for control of status epilepticus. *Anaesthesia.* 1990;45:1043–5.
- 287 Pitt-Miller PL, Elcock BJ, Maharaj M. The management of status epilepticus with a continuous propofol infusion. *Anesth Analg.* 1994;78:1193–4.

- 288 Stecker MM, Kramer TH, Raps EC, O'Meeghan R, Dulaney E, Skaar DJ. Treatment of refractory status epilepticus with propofol: clinical and pharmacokinetic findings. *Epilepsia*. 1998;39:18–26.
- 289 Begemann M, Rowan AJ, Tuhim S. Treatment of refractory complex-partial status epilepticus with propofol: case report. *Epilepsia*. 2000;41:105–9.
- 290 Prasad A, Worrall BB, Bertram EH, Bleck TP. Propofol and midazolam in the treatment of refractory status epilepticus. *Epilepsia*. 2001;42:380–6.
- 291 Wijedicks EFM. Propofol in myoclonus status epilepticus in comatose patients following cardiac resuscitation. *J Neurol Neurosurg Psychiatry*. 2002;73:94–5.
- 292 Niermeijer JM, Uiterwaal CS, van Donselaar CA. Propofol in status epilepticus: little evidence, many dangers? *J Neurol*. 2003;250:1237–40.
- 293 Rossetti AO, Reichhart MD, Schaller MD, Despland PA, Bogousslavsky J. Propofol treatment of refractory status epilepticus: a study of 31 episodes. *Epilepsia*. 2004;45:757–63.
- 294 Parviainen I, Usaro A, Kalviainen R, Mervälä E, Ruokonen E. Propofol in the treatment of refractory status epilepticus. *Intensive Care Med*. 2006;32:1075–9.
- 295 Lowenstein DH, Aminoff MJ, Simon RP. Barbiturate anesthesia in the treatment of status epilepticus: clinical experience with 14 patients. *Neurology*. 1988;38:395–400.
- 296 Osorio I, Reed RC. Treatment of refractory generalized tonic-clonic status epilepticus with pentobarbital anesthesia after high-dose phenytoin. *Epilepsia*. 1989;30:464–71.
- 297 Mirski MA, Williams MA, Hanley DF. Prolonged pentobarbital and phenobarbital coma for refractory generalized status epilepticus. *Crit Care Med*. 1995;23:400–4.
- 298 Parviainen I, Uusaro A, Kälviäinen R, Kaukanen E, Mervälä E, Ruokonen E. High-dose thiopental in the treatment of refractory status epilepticus in intensive care unit. *Neurology*. 2002;59:1249–51.
- 299 Van Ness PC. Pentobarbital and EEG burst suppression in treatment of status epilepticus refractory to benzodiazepines and phenytoin. *Epilepsia*. 1990;31:61–7.
- 300 Yaffe K, Lowenstein DH. Prognostic factors of pentobarbital therapy for refractory generalized status epilepticus. *Neurology*. 1993;43:895–900.
- 301 Krishnamurthy KB, Drislane FW. Relapse and survival after barbiturate anesthetic treatment of refractory status epilepticus. *Epilepsia*. 1996;37:863–7.
- 302 Krishnamurthy KB, Drislane FW. Depth of EEG suppression and outcome in barbiturate anesthetic treatment for refractory status epilepticus. *Epilepsia*. 1999;40:759–62.
- 303 Claassen J, Hirsch LJ, Emerson RG, Mayer SA. Treatment of refractory status epilepticus with pentobarbital, propofol, or midazolam: a systematic review. *Epilepsia*. 2002;43:146–53.
- 304 Jones-Davies DM, MacDonald RL. GABA(A) receptor function and pharmacology in epilepsy and status epilepticus. *Curr Opin Pharmacol*. 2003;3:12–8.
- 305 Bauer TM, Ritz R, Habarth C, Ha HR, Hunkeler W, Steig AJ, et al. Prolonged sedation due to accumulation of conjugated metabolites of midazolam. *Lancet*. 1995;346:145–7.
- 306 De Riu PL, Petrucci V, Testa C, Mulas M, Melis F, Caria MA, et al. Propofol anticonvulsant activity in experimental epileptic status. *Br J Anaesth*. 1992;69:177–81.
- 307 Orser BA, Wang LY, Penefather PS, MacDonald JF. Propofol modulates activation and desensitization of GABA<sub>A</sub> receptors in cultured murine hippocampal neurons. *J Neurosci*. 1994;14:7747–60.
- 308 Orser BA, Bertlik M, Wang LY, MacDonald JF. Inhibition by propofol (2.6 di-isopropylphenol) of the N-methyl-D-aspartate subtype of glutamate receptor in cultured hippocampal neurones. *Br J Pharmacol*. 1995;116:1761–8.
- 309 Holtkamp M, Tong X, Walker MC. Propofol in sub-anesthetic doses terminates status epilepticus in a rodent model. *Ann Neurol*. 2001;49:260–3.
- 310 Lingamaneni R, Hemmings HC. Differential interaction of anaesthetics and antiepileptic drugs with neuronal Na<sup>+</sup> channels, Ca<sup>2+</sup> channels, and GABA<sub>A</sub> receptors. *Br J Anaesth*. 2003;90:199–211.
- 311 Feng HJ, MacDonald RL. Multiple actions of propofol on alphabeta gamma and alphabeta delta GABA<sub>A</sub> receptors. *Mol Pharmacol*. 2004;66:1517–24.
- 312 Mäkelä JP, Iivanainen M, Pieninkeroinen IP, Waltimo O, Lahdensuu M. Seizures associated with propofol anesthesia. *Epilepsia*. 1993;34:832–5.
- 313 Walder B, Tramèr MR, Seeck M. Seizure-like phenomena and propofol: a systematic review. *Neurology*. 2002;58:1327–32.
- 314 Vasile B, Rasulo F, Cnadiani A, Latronico N. The pathophysiology of propofol infusion syndrome: a simple name for a complex syndrome. *Intensive Care Med*. 2003;29:1417–25.
- 315 Rossetti AO, Reichhart MD, Schaller MD, Despland PA, Bogousslavsky J. Propofol treatment of refractory status epilepticus: a study of 31 episodes. *Epilepsia*. 2004;45:757–63.
- 316 Stanski DR, Mihm FG, Rosenthal MH, Kalman SM. Pharmacokinetics of high-dose thiopental used in cerebral resuscitation. *Anesthesiology*. 1980;53:169–71.
- 317 Russo H, Bressolle F. Pharmacodynamics and pharmacokinetics of thiopental. *Clin Pharmacokinet*. 1998;35:95–134.
- 318 Reuber M, Evans J, Bamford JM. Topiramate in drug-resistant complex partial status epilepticus. *Eur J Neurol*. 2002;9:111–2.
- 319 Towne AR, Garnett LK, Waterhouse EJ, Morton LD, DeLorenzo RJ. The use of topiramate in refractory status epilepticus. *Neurology*. 2003;60:332–4.
- 320 Bensalem MK, Fakhoury TA. Topiramate and status epilepticus: report of three cases. *Epilepsy Behav*. 2003;4:757–60.
- 321 Bernhard CG, Böhm E, Höjeberg S. A new treatment of status epilepticus. Intravenous injections of a local anesthetic (lidocaine). *Arch Neurol Psychiatry*. 1955;74:208–14.
- 322 De Giorgi C, Altman K, Hamilton-Byrd E, Rabinowicz AL. Lidocaine in refractory status epilepticus: confirmation of efficacy with continuous EEG monitoring. *Epilepsia*. 1992;33:913–6.
- 323 Pascual J, Ciudad J, Berciano J. Role of lidocaine (lignocaine) in managing status epilepticus. *J Neurol Neurosurg Psychiatry*. 1992;55:49–51.

- 324 Agrawal P, Wali JP. Lidocaine in refractory status epilepticus: a forgotten drug in the emergency department. *Am J Emerg Med.* 1993;11:243–4.
- 325 Teng E, Wilkins P. Lidocaine in status epilepticus. *Ann Pharmacother.* 1994;28:1248–9.
- 326 Walker LA, Slovis CM. Lidocaine in the treatment of status epilepticus. *Acad Emerg Med.* 1997;4:918–22.
- 327 Sawaishi Y, Yano T, Enoki M, Takada G. Lidocaine-dependent early infantile status epilepticus with highly suppressed EEG. *Epilepsia.* 2002;43:201–4.
- 328 Thulasimani M, Ramaswamy S. The role of paraldehyde and lidocaine in the management of status epilepticus. *Anaesthesia.* 2002;57:99–100.
- 329 Fujikawa DG. Neuroprotective effect of ketamine administered after status epilepticus onset. *Epilepsia.* 1995;36:186–95.
- 330 Kofke WA, Bloom MJ, Van Cott A, Brenner RP. Electrographic tachyphylaxis to etomidate and ketamine used for refractory status epilepticus controlled with isoflurane. *J Neurosurg Anesthesiol.* 1997;9:269–72.
- 331 Sheth RD, Gidal BE. Refractory status epilepticus: response to ketamine. *Neurology.* 1998;51:1765–6.
- 332 Borris DJ, Bertram EH, Kapur J. Ketamine controls prolonged status epilepticus. *Epilepsy Res.* 2000;42:117–22.
- 333 Kofke WA, Young RS, Davis P, Woelfel SK, Gray L, Johnson D, et al. Isoflurane for refractory status epilepticus: a clinical series. *Anesthesiology.* 1989;71:653–9.
- 334 Schultz A, Schultz B, Grouven U, Beger FA, Korsch G. Sharp transients in the EEGs of non-epileptic adult patients receiving sevoflurane. *Pharm World Sci.* 2001;23:82–5.
- 335 Sharpe MD, Young GB, Mirsattari S, Harris C. Prolonged desflurane administration for refractory status epilepticus. *Anesthesiology.* 2002;97:261–4.
- 336 Jääskeläinen SK, Kaisti K, Suni L, Hinkka S, Scheinin H. Sevoflurane is epileptogenic in healthy subjects at surgical levels of anesthesia. *Neurology.* 2003;61:1073–8.
- 337 Mirsattari SM, Sharpe MD, Young GB. Treatment of refractory status epilepticus with inhalational anesthetic agents isoflurane and desflurane. *Arch Neurol.* 2004;61:1254–9.
- 338 Hentschke H, Schwarz C, Antkowiak B. Neocortex is the major target of sedative concentrations of volatile anaesthetics: strong depression of firing rates and increase of GABA<sub>A</sub> receptor-mediated inhibition. *Eur J Neurosci.* 2005;21:93–102.
- 339 Vezzani A, Granta T. Brain inflammation in epilepsy: experimental and clinical evidence. *Epilepsia.* 2005;46:1724–3.
- 340 Lux AL, Edwards SE, Hancock E, Johnson AL, Kennedy CR, Newton RW, et al.; United Kingdom Infantile Spasms Study. The United Kingdom Infantile Spasms Study (UKISS) comparing hormone treatment with vigabatrin on developmental and epilepsy outcomes to age 14 months: a multicentre randomised trial. *Lancet Neurol.* 2005;4:712–7.
- 341 Sinclair DB, Snyder TJ. Corticosteroids for the treatment of Landau-Kleffner syndrome and continuous spike-wave discharges during sleep. *Pediatr Neurol.* 2005;32:300–6.
- 342 Bahi-Buisson N, Villanueva V, Bulteau C, Delalande O, Dulac O, Chiron C, et al. Long-term response to steroid therapy in Rasmussen encephalitis. *Seizure.* 2007;16:485–92. Epub 2007 Apr 3.
- 343 Robakis TK, Hirsch LJ. Literature review, case report, and expert discussion of prolonged refractory status epilepticus. *Neurocrit Care.* 2006;4:35–46.
- 344 Bonati LH, Naegelin Y, Wieser HG, Fuhr P, Rüegg S. Beta activity in status epilepticus. *Epilepsia.* 2006;47:207–10.