



Several Blood Constituents of Five Breeds of Dairy Cattle in Guatemala

ROBERT L. SQUIBB, Ph.D.; J. EDGAR BRAHAM, M.S.; MIGUEL A. GUZMAN, M.S.; NEVIN S. SCRIMSHAW, Ph.D., M.D.

Guatemala, Central America

IN ORDER to increase the production of milk in countries of the American tropics, governments, agricultural organizations, and private breeders are importing increased numbers of purebred dairy cattle from the United States and Canada. Some of these cattle are maintained as purebred herds; others are crossed with selected native cattle.

Many of these imported animals die and,

and Piercy¹³ have stressed the value of blood data as an aid in determining the nutritional status of dairy animals. This report, a study of 12 essential blood constituents of dairy cattle considered representative of better managed herds in Guatemala, concludes the series initiated by Squibb *et al.*¹⁵ which were designed to provide reference data on farm animals of this tropical area.

TABLE 1—Levels of Several Blood Constituents of Five Breeds of Dairy Calves and Cows in Guatemala

| Constituent | | Native | | Jersey | | Brown Swiss | | Holstein-Friesian | | Ayrshire | |
|------------------------------|--------|--------|--------|----------|--------|-------------|--------|-------------------|-------|----------|--------|
| | | Mean | s.d. | Mean | s.d. | Mean | s.d. | Mean | s.d. | Mean | s.d. |
| SERUM | | | | | | | | | | | |
| No. of animals | Calves | 40 | | 5 | | 47 | | 17 | | 22 | |
| | Cows | 15 | | 18 | | 49 | | 30 | | 10 | |
| Total proteins | Calves | 6.21 | 0.62 | 6.33 | 0.21 | 6.19 | 0.28 | 6.75 | 0.11 | 6.50 | 2.00 |
| (Gm./100 ml.) | Cows | | | 7.27 | 0.18 | 7.23 | 0.13 | | | | |
| Riboflavin | Calves | 1.44 | 0.59 | 2.01 | 0.24 | 1.97 | 0.61 | 0.86 | 0.08 | 1.10 | 0.24 |
| (µg./100 ml.) | Cows | 1.39 | 1.03 | 1.67 | 0.57 | 3.00 | 1.02 | 2.56 | 0.74 | 1.59 | 0.97 |
| Ascorbic acid | Calves | 0.60 | 0.23 | 1.44 | 0.21 | 0.81 | 0.14 | 1.10 | 0.05 | 0.50 | 0.09 |
| (µg./100 ml.) | Cows | | | 1.20 | 0.08 | 1.06 | 0.11 | | | | |
| Carotene | Calves | 77.00 | 43.00 | 478.00 | 51.00 | 304.00 | 95.00 | 79.00 | 8.00 | 293.00 | 184.00 |
| (µg./100 ml.) | Cows | 900.00 | 225.00 | 1,097.00 | 101.00 | 666.00 | 152.00 | 886.00 | 31.00 | 893.00 | 214.00 |
| Vitamin A | Calves | 12.00 | 5.50 | 24.00 | 2.60 | 23.00 | 28.70 | 12.00 | 0.90 | 16.00 | 13.90 |
| (µg./100 ml.) | Cows | 51.00 | 26.60 | 58.00 | 9.70 | 34.00 | 12.80 | 47.00 | 7.00 | 40.00 | 14.40 |
| Total tocopherols | Calves | 0.45 | 0.16 | 0.27 | 0.02 | 0.50 | 0.12 | 0.16 | 0.03 | 0.33 | 0.18 |
| (µg. 100 ml.) | Cows | | | 0.60 | 0.34 | 0.91 | 0.04 | | | | |
| Alkaline phosphatase | Calves | 9.76 | 3.14 | 5.70 | 0.85 | 5.48 | 0.91 | 2.03 | 0.14 | 3.77 | 2.91 |
| (mM./1/hr.) | Cows | 2.30 | 0.46 | 1.54 | 0.58 | 2.31 | 1.00 | 2.62 | 2.10 | 4.62 | 3.21 |
| Calcium | Calves | 15.53 | 2.23 | | | 10.13 | 0.81 | | | 9.12 | 0.74 |
| (mg./100 ml.) | Cows | | | 6.89 | 0.75 | 10.61 | 1.58 | 10.15 | 1.21 | 9.32 | 1.14 |
| Phosphorus | Calves | 9.29 | 1.07 | | | 7.03 | 0.58 | | | 8.31 | 1.16 |
| (mg./100 ml.) | Cows | | | 4.15 | 0.34 | 6.17 | 1.04 | 6.90 | 0.85 | 6.03 | 0.86 |
| WHOLE BLOOD | | | | | | | | | | | |
| No. of animals | Calves | 27 | | | | 10 | | 17 | | | |
| Hemoglobin | Calves | 12.9 | 1.8 | | | 10.1 | 0.6 | 11.8 | 0.4 | | |
| (Gm./100 ml.) | | | | | | | | | | | |
| Erythrocytes | Calves | 11.5 | 2.4 | | | 8.4 | 0.5 | 9.7 | 0.5 | | |
| (millions/mm. ³) | | | | | | | | | | | |
| count | | | | | | | | | | | |
| Hematocrit (%) | Calves | 53.0 | 7.0 | | | 34.0 | 1.4 | 31.0 | 0.2 | | |

often, they produce less milk than the native cattle. Although the reasons for this high mortality and inefficient production are numerous and complicated, there is no doubt that the existing low plane of nutrition is an important factor. Rusoff

From the Instituto Agropecuario Nacional (Squibb) and the Instituto de Nutrición de Centro América y Panamá (INCAP, Braham, Guzmán, and Scrimshaw), Guatemala, C. A.

INCAP scientific publication I-85.

MATERIALS AND METHODS

From April, 1952, to May, 1954, blood samples were obtained from several herds of Jerseys, Holstein-Friesians, and Ayrshires of highland areas (elevation over 3,000 ft.) and Brown Swiss and native breeds of lowland areas (from sea level to 1,500 ft.). All cattle had access to either green soilage or pasture. The cows were 3 to 5 years of age and were in the third to fifth month of lactation. The calves ranged from 2 to 5 months of age and were all nursing. There was no rela-

tionship between the cows and the calves used in this study.

Approximately 15 ml. of blood was drawn from the jugular vein and permitted to coagulate to yield the serum. An additional 2 ml. was collected in tubes containing an anticoagulant. All samples were immediately placed in a small ice box and shipped by air to the laboratory. Within five hours after blood samples were collected, analyses were begun for protein,¹⁰ riboflavin,⁴ ascorbic acid,¹¹ carotenoids, and vitamin A,¹ tocopherols,¹² alkaline phosphatase,¹ calcium,⁸ and phosphorus.⁸

RESULTS AND DISCUSSION

Since only better managed herds were sampled, it was anticipated that the blood levels for the various constituents studied would be within normal limits. The serum protein values of the cows (table 1) would be even closer to those given by Dukes⁵ for plasma protein if the fibrinogen content of the plasma were added. The vitamin A and carotene blood levels were adequate and similar to those reported elsewhere.^{6, 14, 17} Obviously, these cattle, fed ample quantities of fresh green soilage or given access to green pastures, had an adequate intake of carotene. The relatively lower serum carotene and vitamin A levels of the native and Holstein-Friesian calves suggest that they were consuming poorer quality forage prior to the time the blood samples were drawn.

The ascorbic acid serum levels were, on the average, higher than those reported by Braun³ for dairy cows on pasture. The serum alkaline phosphatase and tocopherol values were similar to those previously reported^{9, 15} as were the levels of calcium and phosphorus.^{3, 13} The rumen and intestinal synthesis of riboflavin, which takes place in cattle, and dietary intake determine the serum levels (table 1) of this vitamin.

While the hemoglobin and hematocrit values and the erythrocyte counts of the purebred calves were similar to published values,^{13, 18} the native calves had comparatively higher values for these blood constituents.

SUMMARY

Chemical analyses were made of several blood constituents of dairy cows and calves of five breeds which were considered representative of better managed herds in Guatemala. The data indicate that blood serum levels of total proteins, ascorbic

acid, tocopherols, hemoglobin, erythrocytes, hematocrit, alkaline phosphatase, carotenoids, and vitamin A were similar to values published elsewhere as normal. Native calves raised in the tropic lowlands were observed to have higher hemoglobin and hematocrit values and erythrocyte counts.

References

- ¹ Bessey, O. A., Lowry, O. H., and Brock, M. J.: A Method for the Rapid Determination of Alkaline Phosphatase with Five Cubic Millimeters of Serum. *J. Biol. Chem.*, 164, (1946): 321-329.
- ² Bessey, O. A., Lowry, O. H., Brock, M. J., and Lopez, J. A.: Determination of Vitamin A and Carotene in Small Quantities of Blood Serum. *J. Biol. Chem.*, 166, (1946): 177-188.
- ³ Braun, W.: Average Levels of Various Constituents, Physical Properties, and Formed Elements of the Blood of Cows on Pasture. *Am. J. Vet. Res.*, 7, (Oct., 1946): 450-454.
- ⁴ Burch, H. B., Bessey, O. A., and Lowry, O. H.: Fluorometric Measurements of Riboflavin and Its Natural Derivatives in Small Quantities of Blood Serum and Cells. *J. Biol. Chem.*, 175, (1948): 457-470.
- ⁵ Dukes, H. H.: The Physiology of Domestic Animals. Comstock Publishing Co. Inc., Ithaca, N. Y., (1943): 45-50.
- ⁶ Eaton, H. D., Carpenter, C. A., Caverno, R. J., Johnson, R. E., and Elliott, F. I.: A Comparison of United States No. 2 Field-Cured Field-Baled Alfalfa Hay with Artificially Dried and Ground and Pelleted Alfalfa Hays as a Source of Carotenoids and Roughage for Holstein Calves. *J. Dai. Sci.*, 34, (1951): 124-135.
- ⁷ Goodland, R. L., Sealock, R. R., Scrimshaw, N. S., and Clark, L. C.: Interference with the Ultramicro Ascorbic Acid Method of Lowry, Lopez and Bessey. *Science*, 109, (1949): 494-495.
- ⁸ Hawk, P. B., Oser, B. L., and Summerson, W. H.: Practical Physiological Chemistry. 12th ed. Blakiston Co., Philadelphia, 1947.
- ⁹ Hibbs, J. W., Krauss, W. E., Monroe, C. F., and Pounden, W. D.: A Report on the Occurrence of Rickets in Calves Under Farm Conditions. *J. Dai. Sci.*, 28, (1945): 525-529.
- ¹⁰ Lowry, O. H., and Hunter, T. H.: Determination of Serum Protein Concentration with a Gradient Tube. *J. Biol. Chem.*, 159, (1945): 465-474.
- ¹¹ Lowry, O. H., Lopez, J. A., and Bessey, O. A.: Determination of Ascorbic Acid in Small Amounts of Blood Serum. *J. Biol. Chem.*, 160, (1945): 609-615.
- ¹² Quaife, M. L., Scrimshaw, N. S., and Lowry, O. H.: A Micromethod for Assay of Total Tocopherols in Blood Serum. *J. Biol. Chem.*, 180, (1949): 1229-1235.
- ¹³ Rusoff, L. L., and Piercy, P. L.: Blood Studies of Louisiana Dairy Cows. II. Inorganic Phosphorus, Hemoglobin Value, Erythrocyte Count, Leucocyte Count and Differential Leucocyte Percentages. *J. Dai. Sci.*, 19, (1946): 831-838.

¹⁴ Squibb, R. L., Cannon, C. Y., and Allen, R. S.: Effect of Raw Soybeans and of Soybean Oil on Plasma Carotene and on Vitamin A as Measured by Activated Glycerol Dichlorohydrin. *J. Dai. Sci.*, *31*, (1948): 421-427.

¹⁵ Squibb, R. L., Guzmán, M., Aguirre, F., and Scrimshaw, N. S.: Ten Constituents of the Blood Stream of Well-Fed White Rats, Chickens, Swine, Sheep, and Horses in Guatemala. *Am. J. Vet. Res.*, *14*, (July, 1953): 484-486.

¹⁶ Whiting, F., Loosli, J. K., Krukovsky, V. N., and Luck, K. L.: The Influence of Tocopherols and Cod-

Liver Oil on Milk and Fat Production. *J. Dai. Sci.*, *32*, (1949): 133-138.

¹⁷ Wise, G. H., Atkeson, F. W., Caldwell, M. J., Parrish, D. B., and Hughes, J. S.: Effects of High Vitamin A Intake on Milk and Fat Yields and on Vitamin A Constituents in Milk, Blood, and Livers of Dairy Cows. *J. Dai. Sci.*, *30*, (1947): 279-291.

¹⁸ Wise, G. H., Caldwell, M. J., Parrish, D. B., Flipse, R. J., and Hughes, J. S.: Changes in Cell Volume and in Concentration of Hemoglobin and Several Inorganic Constituents of the Blood of Calves During Early Postnatal Development. *J. Dai. Sci.*, *30*, (1947): 982-992.

