
Mimivirus: membrane baits find mostly viral prey (70/88 pairs); nucleoid and factory baits mostly host (305/461)

a Mimivirus: ten baits, 433 prey, by region

Bait: a tagged virus protein pulled down from infected amoeba cells (Acanthamoeba castellanii).
Prey: a protein found with a bait. L and R names are virus genes; ids of five digits, amoeba genes.



Ring and baits

- Bait: a tagged virus protein
- One bait pulled down the other
- Each pulled down the other
- Line to a dot: prey that bait pulled down
- Ground: the five baits of a subnetwork

Ring sections: region of prey

		viral	host
1	VF-IL only	48	123
2	Virion and VF-IL	4	0
3	Nucleoid, virion and VF-IL	7	0
4	No region listed	39	166
5	Nucleoid, not VF-IL	12	0
6	Virion only	9	0
7	Membrane, not in virion	7	0
8	Membrane, in virion	18	0

VF-IL: internal layer of the viral factory (VF).
Membrane: predicted transmembrane domain.

Strips on the ring


- organism: viral prey
- organism: host prey (amoeba protein)

Function of viral prey (number)

Uncharacterized	96
DNA, RNA and translation	
Transcription	5
DNA Replication and Repair	5
Translation and Ribosome Function	3
Nucleotide Metabolism	3
Proteins and enzymes	
Protein Modification and Folding	9
Hydrolases	2
Transferases	1
Other metabolism	
Carbohydrate Metabolism	5
Oxidation-Reduction	2
Amino Acid Metabolism	2
Lipid Metabolism	1
Structure and assembly	
Morphogenesis	3
Structural/Motifs/Protein binding	3
Structural/Motifs/DNA binding	1
No category: host 289, viral 3	

b Prey per bait


Dark: viral prey. White: host prey. Sorted by total within a subnetwork.
VM: virion-membrane baits. NVF: nucleoid and viral-factory baits.
Rows add up to 549 bait-prey pairs for 433 prey, as several baits can pull down one prey.
Counts include baits found by other baits.



		viral	host	all
VM	L323	34	3	37
	R595	16	2	18
	R443	14	1	15
	L330	4	11	15
	R387	2	1	3
NVF	R252	62	123	185
	R561	18	66	84
	L274	21	59	80
	R721	26	39	65
	L410	29	18	47

c Prey shared by two baits


Number: prey both baits pulled down; darker is more.
Triangle: one bait pulled down the other; two: each did.




d Proteins homologous to PDB complex chains

Line: the two PDB chains these proteins match are in contact (Methods: under 8 Å). PDB: Protein Data Bank.
Circle: protein, dark if viral prey, white if host prey. Maroon: bait. Square: viral protein that no bait pulled down.
Ids: 14831 is amoeba gene BAESF_14831.mRNA.1. The paper chose the PDB entries sharing most chains with these proteins.


7a09
human 43S pre-initiation complex
16 host proteins, 29 links




6zj3
Euglena gracilis ribosome
17 host proteins, 20 links




7dvq
human minor spliceosome
12 host proteins, 22 links



RNA polymerase
'core' set (file name)
8 viral proteins, 14 links



RNA polymerase
'not core' set (file name)
17 proteins incl. the 8; 35 links



140 further viral proteins were pulled down by no bait and are not shown.
R443: the Results give 9 viral membrane partners; the file flags 8 (L446 is not flagged).

RNA polymerase entries (legend): poxvirus 6rfl, 7amv; African swine fever virus 8q3k; human 7und, 7nvu, 7ena.

Every count the paper prints matches the network file and Source data Fig. 3. Ring: one section per region; inside it, prey follow the baits that pulled them down.

After Fig. 3 of Safi et al., EMBO J 2026, doi:10.1038/s44318-026-00770-8, CC BY 4.0. Data: Zenodo 17313812. Redesigned; values computed from published data.