

CycloPhaser: A Python Package for Detecting Extratropical Cyclone Life Cycles

Danilo Couto de Souza¹, Pedro Leite da Silva Dias¹, Carolina Barnez Gramscianinov², and Ricardo de Camargo¹

¹ Institute of Astronomy, Geophysics and Atmospheric Sciences of the São Paulo University, Rua do Matão, 226, Cidade Universitária, 05508-090, São Paulo, Brazil ² Institute for Coastal Systems – Analysis and Modeling, Helmholtz-Zentrum Hereon, Geesthacht, Germany

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

CycloPhaser is an open-source Python package designed to detect and analyze extratropical cyclone life cycles using central relative vorticity data. By automating the classification of key cyclone phases, such as intensification, decay, and maturity, CycloPhaser enables efficient processing of large datasets with minimal computational overhead. The package provides customizable options for pre-processing data through noise-reducing filters and smoothing techniques, ensuring accurate phase detection. Initially developed for use in the Southern Hemisphere, CycloPhaser can also be applied to the Northern Hemisphere with minor adjustments. Its flexible design makes it adaptable to various data types, including sea level pressure and geopotential height, broadening its utility for cyclone studies. CycloPhaser can support researchers in atmospheric sciences by enabling in-depth analysis of cyclone life cycles, improving model validation, and offering insights into the impacts of climate change on cyclone behavior.

Summary

CycloPhaser is a Python package designed to detect and analyze extratropical cyclone life cycles from central relative vorticity data. It enables researchers in meteorology and atmospheric sciences to automatically identify key stages of cyclone development, such as intensification, decay, and mature phases, using the vorticity series and its derivatives. By leveraging vorticity data, CycloPhaser helps scientists study cyclones across various regions and timeframes, contributing to improved understanding of cyclone energetics and behavior.

Statement of Need

Extratropical cyclones are key features of the climate system. In South America, they are especially important due to the presence of cyclogenesis hotspots in southeast Brazil (SE-BR), the La Plata River basin (LA-PLATA), and southeastern Argentina (ARG) (C. Gramscianinov et al., 2019). These cyclones can cause extreme precipitation, intense winds, high sea waves, and landslides, significantly impacting communities (Cardoso et al., 2022; C. B. Gramscianinov et al., 2023; D. C. de Souza et al., 2024; D. de Souza & Silva, 2021). Understanding their temporal and spatial development and evolution is crucial for improving forecasts, ultimately aiding in the adoption of mitigation and adaptation strategies.

Accurately identifying the regions where cyclones are positioned throughout their distinct life cycle stages remains a significant challenge in atmospheric sciences. Seminal works by (Bjerknes & Solberg, 1922), (Shapiro & Keyser, 1990), and (Neiman & Shapiro, 1993)

described extratropical cyclone life cycles in terms of structural changes and large-scale dynamics. However, these classifications were based on manual analysis of satellite imagery and synoptic charts, limiting their applicability to large datasets with multiple cyclone cases. Recent research has sought to objectively define cyclone life cycle stages using techniques such as normalizing the life cycle duration (Rudeva & Gulev, 2007; Schemm et al., 2018) or bisecting the cycle into “intensification” and “decay” phases by focusing on periods before and after peak vorticity or the lowest central pressure (Azad & Sorteberg, 2014; Booth et al., 2018; Dacre & Gray, 2009; Michaelis et al., 2017; ?). While these approaches support the study of cyclone intensification and decay, they tend to overlook critical phases such as the incipient stage — where environmental dynamics are still adapting to the developing low-level disturbance and surface isobars are not yet fully closed. Additionally, they treat the mature phase as a single time step, failing to account for the possibility that it may encompass multiple time steps during which the cyclone exhibits homogeneous features.

The pioneering work by (Couto de Souza et al., 2024) was the first to offer a comprehensive analysis of extratropical cyclone life cycles, dissecting systems into distinct life cycle phases and enabling the detection of multiple configurations across different systems. This study presents the Python package that facilitated such work. The method allows for an automated classification of cyclone life cycle stages, enabling the efficient processing of large datasets with minimal computational cost. This tool opens new avenues for research, such as analyzing cyclone life cycle behavior in climate change projections, enabling comparisons with present-day climates, and providing insights into how cyclone life cycles may evolve in response to climate variability. Additionally, it offers potential for assisting model validation by comparing the spatial positioning of life cycle phases across different models and reanalysis datasets. The package is both flexible and fully customizable, making it adaptable to a wide range of datasets and research needs.




Figure 1: Figure 1: Yearly cyclone track densities normalized for the three cyclogenesis regions along the South American coast (SE-BR, LA-PLATA, and ARG). Contours represent normalized track densities above 0.8, plotted individually for each region. Details regarding the genesis regions, tracking procedures, and analysis techniques are discussed in (Couto de Souza et al., 2024).

Features

The program includes optional pre-processing steps, such as applying a Lanczos filter to remove noise from the series and a Savitzky-Golay filter for smoothing, ensuring sinusoidal patterns in the data for phase detection. Key cyclone phases — intensification, decay, and mature — are identified through peaks and valleys in the vorticity time series. The intensification phase spans from a vorticity peak to the next valley, while the decay phase covers the opposite. The mature phase is defined as the period between a vorticity valley and neighboring derivative peaks. The pre-processing steps, as well as peaks and valleys detection in the vorticity series, are computed using Scipy’s package (Virtanen et al., 2020).

Thresholds for phase detection were rigorously calibrated using a representative set of cyclone tracks, ensuring accurate phase identification while filtering out noise. CycloPhaser also includes

a residual phase to account for tracking anomalies, such as post-decay re-intensification without returning to maturity. A post-processing step further refines the phase boundaries by correcting gaps and isolating single time-step phases. Finally, the incipient stage is detected by missing labels in the series or by selecting the initial time steps. More details are discussed in (Couto de Souza et al., 2024).




Figure 2: Representative example of a cyclone life cycle exhibiting an incipient-intensification-mature-decay configuration.

Although the package was initially devised for detecting life cycle phases using relative vorticity, it can be applied to any time series used as a proxy for cyclone detection, such as sea level pressure and geopotential height. Also, the program was designed for use in the Southern Hemisphere, but it can be applied to Northern Hemisphere vorticity series by multiplying by minus one.

References

- Azad, R., & Sorteberg, A. (2014). The vorticity budgets of north atlantic winter extratropical cyclone life cycles in MERRA reanalysis. Part i: Development phase. *Journal of the Atmospheric Sciences*, 71(9), 3109–3128.
- Bjerknes, J., & Solberg, H. (1922). *Life cycle of cyclones and the polar front theory of atmospheric circulation*. Grondahl.
- Booth, J. F., Naud, C. M., & Jeyaratnam, J. (2018). Extratropical cyclone precipitation life cycles: A satellite-based analysis. *Geophysical Research Letters*, 45(16), 8647–8654.
- Cardoso, A. A., Rocha, R. P. da, & Crespo, N. M. (2022). Synoptic climatology of subtropical cyclone impacts on near-surface winds over the south atlantic basin. *Earth and Space Science*, 9(11), e2022EA002482.
- Couto de Souza, D., Silva Dias, P. L. da, Gramscianinov, C. B., Silva, M. B. L. da, & Camargo,

- 98 R. de. (2024). New perspectives on south atlantic storm track through an automatic
99 method for detecting extratropical cyclones' lifecycle. *International Journal of Climatology*,
100 44(10), 3568–3588.
- 101 Dacre, H. F., & Gray, S. L. (2009). The spatial distribution and evolution characteristics of
102 north atlantic cyclones. *Monthly Weather Review*, 137(1), 99–115.
- 103 Gramcianinov, C. B., Camargo, R. de, Campos, R. M., Guedes Soares, C., & Silva Dias, P.
104 L. da. (2023). Impact of extratropical cyclone intensity and speed on the extreme wave
105 trends in the atlantic ocean. *Climate Dynamics*, 60(5-6), 1447–1466.
- 106 Gramcianinov, C., Hodges, K., & Camargo, R. de. (2019). The properties and genesis
107 environments of south atlantic cyclones. *Climate Dynamics*, 53, 4115–4140.
- 108 Michaelis, A. C., Willison, J., Lackmann, G. M., & Robinson, W. A. (2017). Changes in winter
109 north atlantic extratropical cyclones in high-resolution regional pseudo-global warming
110 simulations. *Journal of Climate*, 30(17), 6905–6925.
- 111 Neiman, P. J., & Shapiro, M. (1993). The life cycle of an extratropical marine cyclone. Part
112 i: Frontal-cyclone evolution and thermodynamic air-sea interaction. *Monthly Weather*
113 *Review*, 121(8), 2153–2176.
- 114 Rudeva, I., & Gulev, S. K. (2007). Climatology of cyclone size characteristics and their changes
115 during the cyclone life cycle. *Monthly Weather Review*, 135(7), 2568–2587.
- 116 Schemm, S., Sprenger, M., & Wernli, H. (2018). When during their life cycle are extratropical
117 cyclones attended by fronts? *Bulletin of the American Meteorological Society*, 99(1),
118 149–165.
- 119 Shapiro, M. A., & Keyser, D. (1990). *Fronts, jet streams and the tropopause*. Springer.
- 120 Souza, D. C. de, Crespo, N. M., Silva, D. V. da, Harada, L. M., Godoy, R. M. P. de, Domingues,
121 L. M., Luiz, R., Bortolozo, C. A., Metodiev, D., Andrade, M. R. M. de, & others. (2024).
122 Extreme rainfall and landslides as a response to human-induced climate change: A case
123 study at baixada santista, brazil, 2020. *Natural Hazards*, 1–26.
- 124 Souza, D. de, & Silva, R. R. da. (2021). Ocean-land atmosphere model (OLAM) performance
125 for major extreme meteorological events near the coastal region of southern brazil. *Climate*
126 *Research*, 84, 1–21.
- 127 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,
128 Burovski, E., Peterson, P., Weckesser, W., Bright, J., & others. (2020). SciPy 1.0:
129 Fundamental algorithms for scientific computing in python. *Nature Methods*, 17(3),
130 261–272.