

Lorenz Energy Cycle Climatology for the Southwestern Atlantic Cyclones

Danilo Couto de Souza^{1,2*}, Pedro Leite da Silva Dias¹,
Carolina Barnez Gramscianinov³, Ricardo Camargo¹

^{1*}Institute of Astronomy, Geophysics and Atmospheric Sciences, São
Paulo University, Rua do Matão, 266, São Paulo, 05508-090, São Paulo,
Brazil.

²Climate Risk Initiative, IRB(re), Av. República do Chile, 330 - 4o
andar, Rio de Janeiro, 20031-170, Rio de Janeiro, Brazil.

³Institute for Coastal Systems Analysis and Modeling,
Helmholtz-Zentrum Hereon, Max-Planck-Straße, 1, Geesthacht, 21502,
Schleswig-Holstein, Germany.

*Corresponding author(s). E-mail(s): danilo.oceano@gmail.com .

Contributing authors: pldsdias@gmail.com .

cbgramscianinov@gmail.com ; ricamarg@usp.br .

Abstract

This study presents a climatological assessment of the Lorenz Energy Cycle (LEC) applied to South Atlantic cyclones, using a Semi-Lagrangian framework. Over 6,700 cyclones were identified from ERA5 reanalysis (1979–2020), and LEC components were computed and averaged across four objectively defined life cycle phases: incipient, intensification, mature, and decay. Results reveal a coherent energy transfer structure: baroclinic conversions dominate during intensification, while barotropic conversions peak during the mature phase and reach magnitudes 2–3 times larger than baroclinic counterparts. Diabatic generation of eddy available potential energy plays a secondary but relevant role, particularly during intensification, occasionally surpassing baroclinic contributions. Eddy kinetic energy imports are most significant during the early phases, reinforcing development. Despite substantial variability among systems, an EOF analysis shows that most cyclones share a common energy structure, with variability manifesting as amplification or suppression of specific pathways. The leading EOFs are linked to differences in cyclone intensity, genesis region, and seasonality. Among the most intense systems, distinct clusters emerge with varying energetic

configurations, some dominated by baroclinic processes, others by barotropic conversions or enhanced diabatic generation. These findings demonstrate that South Atlantic cyclones encompass a spectrum of dynamical behaviors and can be classified based on their energy cycle characteristics. This study provides the first large-sample application of the Semi-Lagrangian LEC to extratropical cyclones in the South Atlantic and highlights the importance of barotropic processes in their development. The results offer a robust framework for interpreting cyclone energetics and establish a comprehensive baseline for classifying these systems.

Keywords: Cyclone energetics, Baroclinic conversion, Barotropic conversion, Latent heat release, South Atlantic cyclones

Acknowledgements

The first author would like to thank the Laboratory of Applied Meteorology for Regional Meteorological Systems (MASTER) for providing the data processing infrastructure. Special thanks are extended to Jean Peres and Djalma Vieira for their invaluable technical assistance and support in the laboratory, which contributed significantly to the success of this research. CBG acknowledges the Helmholtz European Partnership “Research Capacity Building for Healthy, Productive and Resilient Seas” (SEA-ReCap, grant no. PIE-0025).

1 Introduction

Cyclones play a crucial role in the Earth’s climate system by redistributing heat, moisture, and momentum across different regions. These dynamic weather systems, contribute to the global energy balance by transferring energy from the tropics to higher latitudes. Understanding cyclone behavior is essential for improving climate projections and assessing future climate change impacts. In South America, surface cyclones influence precipitation regimes across the continent (Reboita et al., 2010, 2018; de Souza et al., 2022) often contributing to extreme rainfall events (e.g., de Souza and da Silva, 2021; de Souza et al., 2024), and are associated with severe meteorological hazards such as intense winds (Cardoso et al., 2022; de Souza and da Silva, 2021), high sea waves (da Silva et al., 2025; Gramscianinov et al., 2023), and storm surges along coastal areas (Leal et al., 2023; Tecchio et al., 2024). These extreme events can have profound socioeconomic impacts, including damage to coastal infrastructure, disruptions to port operations, and interruptions in oil and gas exploration. This is particularly critical in the southeastern region of South America, where large metropolitan areas, such São Paulo, Rio de Janeiro and Buenos Aires are located, as well as the Port of Santos, the largest in Latin America.

The Lorenz Energy Cycle (LEC) is a valuable framework for understanding the conversion and flow of energy within the atmosphere, particularly in relation to synoptic and large-scale processes (Lorenz, 1967). It divides the atmospheric energy budget into four main reservoirs: kinetic and available potential energy, both for zonal (mean

flow) and eddy components. Lorenz (1955) formulated the concept of available potential energy (APE), building upon Margules (1903) concept of energy based on the hypothetical adiabatic redistribution of atmospheric mass. The LEC serves as a foundational tool for quantifying and understanding atmospheric dynamics, by tracking energy transformations and explaining how these forms of energy are generated by adiabatic processes and dissipated by friction.

Substantial efforts have been made to estimate the LEC components for the global circulation (e.g., Muench, 1965; Wiin-Nielsen, 1968; Hu et al., 2004; Li et al., 2007; Oort, 2018), to assess its projected changes under climate change scenarios (e.g., Hernández-Deckers and von Storch, 2010; Veiga and Ambrizzi, 2013; Pan et al., 2017; Michaelides, 2021), and to explore its relationship with phases of the El Niño–Southern Oscillation (Gutierrez et al., 2009). The LEC framework has also been applied in a range of contexts, from idealized baroclinic wave simulations (Kirshbaum et al., 2018; Rantanen et al., 2019), to rotating annulus experiments (Young, 2014), and even to the atmospheres of other planets (e.g., Read et al., 2020). In contrast, studies examining the LEC in relation to cyclonic systems are more limited and primarily consist of case studies. These have typically focused on tropical (Brennan and Vincent, 1980; Veiga et al., 2008), subtropical (Michaelides, 1987; Dias Pinto et al., 2013; Pezza et al., 2014; Cavicchia et al., 2018), and extratropical cyclones (Michaelides, 1992; Wahab et al., 2002; Bulic, 2006; Pezza et al., 2010; Dias Pinto and Rocha, 2011). Noteworthy contributions include the work of Black and Pezza (2013), which investigated explosive cyclones across all major ocean basins over a 32-year period and identified a universal signature of explosive cyclogenesis, and Okajima et al. (2021), which quantified the contribution of migratory cyclones to global energetics, highlighting their role in accelerating westerly winds worldwide.

Although the original formulation by Lorenz (1967) was intended for global-scale studies, analyzing specific atmospheric regions is valuable for understanding the mechanisms driving the development of individual systems, such as cyclones, and the energy conversions throughout their life cycle. The initial framework for such regional analysis was introduced by Muench (1965), who studied Northern Hemisphere stratospheric circulation during winter. Subsequent formulations by Dutton and Johnson (1967), Vincent and Chang (1973), and Smith (1980) refined the approach, but it was only with Brennan and Vincent (1980) that a generalized set of equations, including both eddy and zonal components of kinetic and APE for limited regions in the troposphere, was established. However, cyclones are mobile systems that often traverse large regions (Hoskins and Hodges, 2002; de Souza et al., 2024, e.g.). Since these studies employ the LEC within an Eulerian framework, the computational domains required to analyze cyclone energetics can become quite large (Black and Pezza, 2013, e.g.), potentially incorporating other atmospheric features (e.g., migratory anticyclones, upper-level troughs, large convective systems), which limits the interpretation of results.

In this context, Michaelides et al. (1999) developed a Semi-Lagrangian framework for analyzing cyclone energetics, where the computational domain is movable. This approach minimizes the impact of neighboring circulations on the system’s energy dynamics by employing a regional computational domain that follows the cyclone

116 track. In the present study, we present a LEC climatology applied to cyclonic sys-
 117 tems using such a framework, as well as the first LEC climatology (considering both
 118 Eulerian and Semi-Lagrangian frameworks) for cyclones in the Southwestern Atlantic.
 119 Over 6,700 cyclones with genesis in the Southwestern Atlantic region are analyzed,
 120 and their energy cycles are divided into distinct life cycle phases: incipient, intensifica-
 121 tion, mature, and decay. This division allows for a more comprehensive understanding
 122 of the energy flows and the dynamical mechanisms acting across each phase of the
 123 cyclone’s life cycle.

124 2 Materials and methods

125 2.1 Data

126 For both the cyclone tracking and energetics computations, we employed the fifth-
 127 generation reanalysis from the European Centre for Medium-Range Weather Forecasts
 128 (ECMWF), known as ERA5 (Hersbach et al., 2020). ERA5 data are produced using
 129 the Integrated Forecasting System (IFS) Cy41r2, which assimilates a wide range of
 130 observations, including satellite and in-situ measurements. It provides global coverage
 131 of numerous atmospheric, ocean-wave, and land-surface parameters at a horizontal
 132 resolution of approximately 31 km (0.25°). In this study, we utilized data from all 37
 133 pressure levels, ranging from 1000 hPa to 1 hPa, with a temporal interval of three
 134 hours.

135 The cyclone tracks used in this study were obtained from the ”Atlantic Extratrop-
 136 ical Cyclone Tracks Database” (Gramscianinov et al., 2020). This database tracks the
 137 central relative vorticity of cyclones at the 850 hPa level (ζ_{850}) using the TRACK algo-
 138 rithm (Hodges, 1994, 1995) and the method outlined by Hoskins and Hodges (2002).
 139 This algorithm has been employed in previous studies to assess cyclone climatologies
 140 in the South Atlantic region (Gramscianinov et al., 2019, 2020; de Souza et al., 2024).
 141 The database covers the period from 1979 to 2020 and spans the entire Atlantic Ocean
 142 within the spatial domain of 15°S – 55°S and 75°W – 20°E . The use of ERA5, instead of
 143 other reanalysis datasets, is justified by its higher spatial resolution, which improves
 144 cyclone detection in regions with complex orography and temperature gradients, such
 145 as the SESA region (Gramscianinov et al., 2020). A detailed description of the tracking
 146 methodology can be found in Gramscianinov et al. (2020).

147 The cyclone tracking procedure employed multiple criteria to ensure accurate
 148 cyclone detection. These criteria required cyclones to have a duration of at least
 149 24 hours and a minimum displacement of 1000 km, consistent with previous South
 150 Atlantic cyclone climatologies (Sinclair, 1995; Gramscianinov et al., 2019). Systems
 151 that spent over 80% of their life cycle over continental regions were excluded from
 152 the analysis to avoid counting thermal lows and lee troughs (e.g. Crespo et al., 2021).
 153 Notably, although the TRACK algorithm’s calibration and sensitivity are focused on
 154 extratropical cyclones — the majority of systems in this region (Marrafon et al., 2022)
 155 — the methodology does not explicitly exclude subtropical or tropical cyclones.

156 To focus on the Southwestern Atlantic Ocean, only cyclones with genesis near
 157 the South American coast were considered, specifically those originating within the
 158 eastern Argentina (ARG), La Plata River basin (LA-PLATA), and southeast Brazil

(SE-BR) genesis regions (Gramscianinov et al., 2019; de Souza et al., 2024) (Figure 1). After applying these selection criteria, the database initially comprised a total of 7931 cyclones. The spatial distribution of these genesis regions and the cyclone track density for all selected systems are presented in Figure 1. Noteworthy, in traditional cyclone studies, cyclone density is often presented for a single time step or normalized by individual tracks (e.g., Hoskins and Hodges, 2005; Gramscianinov et al., 2019). In contrast, Figure 1 displays cyclone density computed across all time steps within a 5° spherical cap. Cyclones may form within the cap, move into it from adjacent regions, remain for several time steps, or pass through it entirely. Each of these time-step occurrences is counted, meaning that the density values in the figure reflect multiple temporal passages rather than unique storm events.

Among the original 7,931 cyclones, 4,445 originated in ARG (56.0%), 1870 in LA-PLATA (23.6%), and 1616 in SE-BR (20.4%). However, due to data availability issues during the download process from the ERA5 database, not all cyclones could be considered in this study, resulting in a reduced total of 6,789 cyclones. These issues were related to the ERA5 API instability during the second half of 2023 and the first half of 2024, when ECMWF was migrating its data infrastructure and updating its API services. As a result, slow transfers and repeated interruptions caused incomplete or failed downloads, despite multiple attempts. This led to some cyclones being excluded from the analysis. The remaining dataset still includes a sufficiently large number of systems for robust analysis and generalization of the results. Of the 6,789 cyclones, 4,015 originated in ARG (59.1%), 1,288 in LA-PLATA (18.9%), and 1,486 in SE-BR (21.8%). The reduction corresponds to a 14.4% decrease in the number of cyclones, with region-specific reductions of 9.7% for ARG, 31.1% for LA-PLATA, and 8.0% for SE-BR.




Fig. 1 Cyclone track density for all systems analyzed in this study, highlighting the cyclogenesis regions near the South American coast (ARG, LA-PLATA, and SE-BR). The track density unit is cyclonic centers per 10^6 km^2 per month.

2.2 Lorenz Energy Cycle Computation

For computing the LEC, we used the the open-source Python application LorenzCycleToolKit (de Souza et al., 2024). The energy budget equations for the zonal and eddy forms of APE and kinetic energy are then expressed as:

$$\frac{\partial A_Z}{\partial t} = BA_Z - C_Z - C_A + G_Z \quad (1)$$

$$\frac{\partial A_E}{\partial t} = BA_E - C_E + C_A + G_E \quad (2)$$

$$\frac{\partial K_Z}{\partial t} = BK_Z + C_Z - C_K + B\Phi_Z - D_Z \quad (3)$$

$$\frac{\partial K_E}{\partial t} = BK_E + C_E + C_K + B\Phi_E - D_E \quad (4)$$

In these equations, the atmospheric energy reservoirs are separated into zonal and eddy components of available potential energy (A_Z , A_E) and kinetic energy (K_Z , K_E). Energy conversions between reservoirs are represented by the following terms: C_A , the conversion between zonal and eddy available potential energy ($A_Z \leftrightarrow A_E$); C_E , the conversion from eddy available potential energy to eddy kinetic energy ($A_E \leftrightarrow K_E$); C_K , the barotropic conversion from eddy kinetic energy to zonal kinetic energy ($K_E \leftrightarrow K_Z$); and C_Z , the conversion from zonal available potential energy to zonal kinetic energy ($A_Z \leftrightarrow K_Z$). The generation of available potential energy (G) and dissipation of kinetic energy (D) are indicated by subscripts Z and E , such that G_Z and G_E represent the generation of zonal and eddy available potential energy, while D_Z and D_E denote the dissipation of zonal and eddy kinetic energy, respectively. The boundary flux terms, BA_Z , BA_E , BK_Z , and BK_E , represent the import/export of available potential energy and kinetic energy across the computational domain boundaries. Additionally, the terms $B\Phi_Z$ and $B\Phi_E$ appear alongside C_Z and C_E , respectively, as both arise from deriving the kinetic energy balances. According to Muench (1965), these terms are challenging to interpret physically, but indicate processes generating kinetic energy resulting from work performed at the boundaries of the computational domain. The complete mathematical formulation of these terms can be found in the Appendix.

In these formulations, a global reference state is used to define the APE, meaning that the computed APE represents the contribution of local APE to global energetics (Smith, 1980). As a result, it does not account for regional variability, which can introduce bias in the APE computation (see Novak and Tailleux (2018) for a thorough discussion). Despite the limitations of this methodology, the formulations by Muench (1965) and Brennan and Vincent (1980) still provide a valuable framework due to their simplicity, robustness, and ability to yield consistent, interpretable results, enabling direct analysis of the dynamic processes driving cyclones development (Dias Pinto and Rocha, 2011; de Souza et al., 2025, e.g.). Moreover, while using a local APE definition (Novak and Tailleux, 2018, e.g.) offers more precise results, it often comes with higher computational costs and does not provide the same level of direct, practical interpretability.

218 Following the methodologies of Brennan and Vincent (1980); Michaelides (1987);
 219 Veiga et al. (2008); Pezza et al. (2010); Dias Pinto and Rocha (2011), the program
 220 was set to compute the dissipation and generation terms as residuals, defined as:

$$RK_Z = B\Phi_Z - D_Z + \epsilon_{KZ} \quad (5)$$

$$RK_E = B\Phi_E - D_E + \epsilon_{KE} \quad (6)$$

$$RG_Z = G_Z + \epsilon_{GZ} \quad (7)$$

$$RG_E = G_E + \epsilon_{GE} \quad (8)$$

221 Where RK_Z and RK_E are the residual dissipation terms (zonal and eddy, respec-
 222 tively), RG_Z and RG_E are the residual generation terms, $B\Phi_Z$ and $B\Phi_E$ represent
 223 the boundary pressure work terms, and ϵ accounts for numerical errors in the
 224 computational procedures. The set of budget equations becomes, then:

$$\frac{\partial A_Z}{\partial t} = -C_A - C_Z + RG_Z + BA_Z \quad (9)$$

$$\frac{\partial A_E}{\partial t} = C_A - C_E + RG_E + BA_E \quad (10)$$

$$\frac{\partial K_Z}{\partial t} = C_K + C_Z + BK_Z + RK_Z \quad (11)$$

$$\frac{\partial K_E}{\partial t} = -C_K + C_E + BK_E + RK_E \quad (12)$$

225 A complete depiction of the LEC, with arrows indicating positive energy fluxes,
 226 is presented in Figure 2. This figure also highlights three primary energetic pathways
 227 (baroclinic instability, barotropic conversion, and latent heat release) which will be
 228 discussed throughout the text.

229 Here, we used a Semi-Lagrangian framework (Michaelides et al., 1999), aiming to
 230 minimize interactions with other non-related circulations while capturing the main
 231 structure of the cyclone. A $15^\circ \times 15^\circ$ computational domain centered at the cyclone's
 232 central position, obtained from the TRACK database, was created for each time step.
 233 Given the impracticality of manually selecting an appropriate domain size for each
 234 cyclone in the dataset, a fixed size was employed. The choice of a $15^\circ \times 15^\circ$ computa-
 235 tional domain is justified as it is large enough to capture the effective radius of most
 236 cyclonic systems (Rudeva and Gulev, 2007). The effective cyclone radius is defined as
 237 a measure of cyclone size, determined by establishing a coordinate system centered
 238 on the cyclone and measuring the distance at which the radial pressure gradient first
 239 falls to zero. This domain size also is large enough for accommodating the cyclone's
 240 synoptic structure (e.g. Gramscianinov et al., 2019).

241 2.3 Analysis Methods

242 A key challenge in computing the LEC for large datasets lies in the variable life
 243 durations of cyclones. Averaging energy values across an entire life cycle can obscure




Fig. 2 Representation of the Lorenz Energy Cycle (LEC), with arrows denoting positive energy fluxes. The figure also indicates three main energetic pathways: baroclinic instability (purple), barotropic conversion (green), and latent heat release (red), which will be addressed in detail in the following sections.

distinct dynamical processes tied to different stages of cyclone development, while technical limitations exist in dissecting the life cycle into distinct periods. For example, Black and Pezza (2013) averaged cyclone energetics over periods of 48 hours before explosive cyclogenesis, during explosive deepening, and for 24 and 72 hours after it. However, cyclone life cycles can range from less than 24 hours to over 10 days (Trigo, 2006; Reboita et al., 2010; Gramscianinov et al., 2019), presenting significant physical limitations to such approaches.

To address these issues, we employed Cyclophaser, an open-source Python package designed to detect cyclone life cycle phases (de Souza et al., 2025). This program uses

time series of relative vorticity at the system’s central position and its first derivative to identify intensification, mature, and decay phases based on peaks and valleys in the vorticity and its derivative. In this study, we adopted the same life cycle detection approach as de Souza et al. (2024). After computing the LEC for each system, Cyclophaser was used to identify the life cycle phases for all analyzed cyclones, followed by the computation of mean LEC values for each phase. This approach facilitates the investigation of dynamical mechanisms specific to each development phase. For example, distinct energy fluxes are expected to dominate during the intensification and decay phases, particularly near the cyclone center. As shown by de Souza et al. (2024), approximately 60% of the analyzed cyclones follow a classical life cycle, consisting of incipient, intensification, mature, and decay phases, while over 95% display at least one intensification phase. A comprehensive description of the detection methodology, as well as spatial distributions and phase-specific statistics such as duration, displacement, speed, and intensity, can be found in de Souza et al. (2024).

To understand the dominant LEC variability patterns within the TRACK dataset, we performed an Empirical Orthogonal Function (EOF) analysis (Fukuoka, 1951; Lorenz, 1956) using the Python open-source library pyEOF (Zheng, 2021). This analysis reduces the dimensionality of the dataset, retaining the most significant variability, and is particularly useful for uncovering underlying structures and patterns in complex environmental data. Typically, each EOF represents a spatial pattern, while the associated time series, or principal component (PC), describes the temporal evolution of that pattern. However, in this study, each EOF reflects a mode of variability derived from all LEC terms, using their average values computed separately for each cyclone life cycle phase (incipient, intensification, mature, and decay). Consequently, the PCs represent the variation across individual cyclones rather than temporal evolution, and therefore are not displayed individually, as they do not depict any meaningful temporal or spatial feature. Performing EOF analyses separately for each life cycle phase enabled a detailed investigation of the variability in cyclone energetics across distinct stages of cyclone development, as well as throughout the entire energy cycle.

To associate cyclones with predominant EOFs, we first project each cyclone’s energetics onto the PCs obtained from an EOF analysis. This allows us to represent each system in terms of its contribution to the dominant modes of variability. Next, we classify cyclones into EOF(+) and EOF(-) groups, identifying systems where at least one PC reaches extreme values. Specifically, EOF(+) corresponds to cyclones where at least one PC exceeds the 90th percentile (q90), indicating a strong positive projection onto a specific mode. Meanwhile, EOF(-) are cyclones for which at least one PC is below the 10th percentile (q10), indicating strong negative projection onto a specific mode. For each cyclone meeting these criteria, we determine its predominant EOF as the one corresponding to the PC with the highest absolute value. This ensures that each system is classified based on the mode that most strongly characterizes its structure and dynamics.

To determine the LEC patterns of the most intense systems, we first selected only the cyclones in which the maximum central vorticity at 850 hPa exceeded the 90th percentile of the dataset. Subsequently, we applied the K-Means algorithm to the PCs of these intense cyclones to investigate whether distinct energetic characteristics could

298 be identified among them. This clustering analysis was not performed on all cyclones,
 299 as sensitivity tests indicated that applying it to the full dataset resulted in groups
 300 that differed mainly in intensity rather than exhibiting distinct energetic behaviors
 301 (not shown); thus, intensity-based selection was necessary.

302 K-Means is an unsupervised learning algorithm that partitions data into K clusters
 303 by minimizing intra-cluster variance while maximizing separation between groups
 304 based on sample similarity (MacQueen et al., 1967; Hartigan and Wong, 1979). The
 305 optimal number of clusters was determined using the Elbow Method, which identifies
 306 the most suitable number of clusters by plotting the within-cluster sum of squares for
 307 different values of K and locating the "elbow point," where the rate of decrease slows,
 308 indicating a balance between model complexity and performance. For this analysis, we
 309 employed the K-Means implementation from the `scikit-learn` Python package (Hao
 310 and Ho, 2019) and used the `yellowbrick` package for the Elbow Method (Bengfort
 311 and Bilbro, 2019).

312 3 Results

313 3.1 Climatological Features

314 The exploratory statistical metrics for all LEC terms, presented as averages across the
 315 entire cyclone lifecycle, are shown in Table 1, while their probability density functions
 316 (PDFs) are provided in Figure 3. Notably, all terms related to the zonal jets (K_Z ,
 317 BK_Z , and $\frac{\partial K_Z}{\partial t}$) frequently exhibit statistical metrics one order of magnitude higher
 318 than their eddy counterparts within the same group (K_E , BK_E , and $\frac{\partial K_E}{\partial t}$), indicating
 319 significantly greater energy amount concentrated on the jet streams. Although the
 320 energy budgets were computed with the generation and dissipation terms calculated
 321 as residuals, the generation terms can be directly computed using ERA5 data. There-
 322 fore, their probability density functions (PDFs) are also shown for a more physically
 323 based analysis (Figure 3e). Furthermore, the comparison between the directly com-
 324 puted generation values and those estimated by residual computation provides a useful
 325 metric for evaluating the accumulation of errors in LEC computation procedure. The
 326 dissipation terms were not directly computed in the present study.

327 All the energy terms (K_Z , A_Z , A_E , and K_E , Figure 3a) exhibit right-skewed distri-
 328 butions. Among these, A_Z shows the largest variability, as reflected in its long-tail
 329 distribution, standard deviation (std), interquartile range (IQR), and range (defined
 330 as the difference between maximum and minimum values), compared to A_E and K_E ,
 331 peaking at approximately $4 \times 10^5 J m^{-2}$. Conversely, A_E presents the lowest variabil-
 332 ity, peaking near $1 \times 10^5 J m^{-2}$, closer to its mean value. The kinetic energy terms, K_Z
 333 and K_E , display similar distributions, with K_Z peaking near $20 \times 10^5 J m^{-2}$ and K_E
 334 peaking near $2 \times 10^5 J m^{-2}$.

335 For the conversion terms, C_Z , C_K , and C_E exhibit high variability, in contrast
 336 to C_A . The former terms display long-tail distributions, peaking near zero, -1 , and
 337 $2 W m^{-2}$, respectively, while C_A presents a narrow distribution with a sharp peak near
 338 its mean value, close to zero (Figure 3b). The C_Z probability density function (PDF),
 339 along with the mean and quantile values, and its mean value being close to the median,
 340 indicate an overall tendency for both $A_Z \rightarrow K_Z$ and $K_Z \rightarrow A_Z$ conversions. However,

Table 1 Summary Statistics of Lorenz Energetics Components, computed as averages across the entire cyclone lifecycle: mean, median, standard deviation (std), 25th percentile (Q25), 75th percentile (Q75), interquantile range (IQR), and range. The IQR measures the range within which the central 50% of the values fall, while the range is computed as the difference between the maximum and minimum values. The units for the energy terms (A_Z , A_E , K_Z , and K_E) are $10^5 J m^{-2}$ and $W m^{-2}$ for the remaining terms.

Term	Mean	Median	Std Dev	Q25	Q75	IQR	Range
A_Z	5.55	4.68	3.71	2.77	7.49	4.72	36.46
A_E	1.62	1.24	1.20	0.77	2.09	1.32	11.07
K_Z	28.55	26.19	15.10	17.34	37.12	19.78	114.55
K_E	3.70	2.95	2.62	1.92	4.63	2.71	24.81
C_Z	0.52	0.38	4.85	-1.96	2.92	4.89	86.70
C_A	0.93	0.46	1.57	0.02	1.44	1.41	19.78
C_K	-3.56	-1.63	9.51	-5.77	0.47	6.24	206.43
C_E	3.84	2.47	4.92	0.63	5.84	5.21	54.74
BA_Z	3.54	2.03	7.67	-0.67	6.67	7.34	175.05
BA_E	0.58	0.16	4.40	-1.13	2.02	3.15	75.72
BK_Z	-7.93	-5.53	31.75	-23.75	7.83	31.58	445.63
BK_E	0.47	0.08	8.59	-2.73	3.22	5.95	146.87
$B\Phi_Z$	63.87	49.25	128.05	-9.12	132.35	141.47	1419.08
$B\Phi_E$	39.66	31.19	126.50	-28.84	109.47	138.32	1353.99
G_Z	-0.31	-0.17	2.66	-1.39	0.81	2.21	68.74
G_E	1.17	0.56	3.14	-0.26	2.21	2.46	58.39
$\frac{\partial A_Z}{\partial t}$	-0.07	-0.14	4.46	-2.04	1.68	3.71	106.39
$\frac{\partial A_E}{\partial t}$	-0.23	-0.10	1.92	-0.85	0.51	1.35	48.19
$\frac{\partial K_Z}{\partial t}$	1.59	0.75	13.38	-5.33	8.15	13.48	251.24
$\frac{\partial K_E}{\partial t}$	0.28	0.12	3.18	-1.13	1.58	2.71	61.50
RG_Z	-2.17	-1.00	7.60	-5.22	1.56	6.78	220.06
RK_Z	12.56	8.69	40.49	-8.68	32.71	41.39	561.24
RGE	2.10	1.20	5.15	-0.40	4.21	4.61	97.25
RKE	-7.59	-4.63	10.62	-11.48	-1.04	10.43	138.42

the positive mean and median values suggest a predominant $A_Z \rightarrow K_Z$ pathway. For C_A and C_E , the metrics presented in Table 1 indicate a consistent tendency for positive conversions, corresponding to $A_Z \rightarrow A_E \rightarrow K_E$ (baroclinic chain), but with more modest $A_Z \rightarrow A_E$ conversions than $A_E \rightarrow K_E$. In the case of the C_K term (barotropic conversion term), the metrics indicate a predominant negative conversion, i.e., $K_Z \rightarrow K_E$, although the positive Q75 value suggests the occurrence of events where $K_E \rightarrow K_Z$ conversions also take place.

For the boundary terms, BA_E and BK_Z exhibit similar distributions, as do BA_Z and BK_E (Figure 3c). The terms BA_E and BK_E display narrow distributions, peaking near zero, whereas BA_Z and BK_Z show nearly symmetrical long-tail distributions. Although for all four terms, both positive (influx) and negative (outflux) energy fluxes occur, as indicated by their negative Q25 and positive Q75 values, the peak near zero for BA_E and BK_Z suggests that boundary energy fluxes are typically low in most cases. In contrast, for BA_Z , the peak near $2 W m^{-2}$ indicates an overall tendency for energy influx, while for BK_E , the peak near $-2 W m^{-2}$ points to a predominant tendency for energy outflux. Meanwhile, the boundary pressure work terms ($B\Phi_Z$ and $B\Phi_E$) exhibit significantly higher values compared to the other boundary terms, with




Fig. 3 Probability density function (PDF) plots for distinct Lorenz Energetics terms, computed as averages across the entire cyclone lifecycle: (A) energy, (B) conversion, (C) boundary, (D) pressure work, (E) generation and residual, and (F) budget terms. The x-axis represents the energy values, while the y-axis represents density counts. The units for the energy terms (A_Z , A_E , K_Z , and K_E) are in $10^5 J m^{-2}$, and in $W m^{-2}$ for the remaining terms. Terms related to K_Z are divided by 10 (*) to account for their order of magnitude difference, which would otherwise distort the visualization.

both presenting nearly symmetric distributions (Figure 3d). These terms show tail distributions exceeding $\pm 200 W m^{-2}$, indicating substantial variability. The physical interpretation of these terms is discussed in Section 4.1.

Overall, both generation terms (G_Z and G_E) present nearly symmetrical distributions centered near zero (Figure 3e), suggesting that in most cases, either the zonally averaged diabatic heating (G_Z) and the zonal deviations (G_E) are low, or the atmospheric stability is high. However, there are instances where this behavior is amplified ($G < 0$) or reversed ($G > 0$). In contrast, the generation residual terms (RG_Z and RG_E) display distinct characteristics. The RG_Z term is left-skewed, with a long-tail distribution and higher variability compared to G_Z , as indicated by its lower Q25 and higher Q75 and IQR values (Table 1). Additionally, RG_Z exhibits more negative mean and median values than G_Z , suggesting a bias toward processes such as diabatic cooling at lower latitudes and/or heat distribution mechanisms that reduce zonal temperature gradients, such as cyclonic activity. On the other hand, RG_E is more symmetrical than RG_Z , though it still shows higher variability compared to G_E , as evidenced by its lower Q25 and higher Q75 and IQR values. Moreover, RG_E exhibits higher mean and median values than G_E , indicating a bias toward overestimating the eddy effects on latent heat release and meridional temperature gradients.

Therefore, the residuals suggest that the computed LEC increases the contribution of eddy motions to the atmospheric energy cycle through enhanced meridional heat transport, reducing zonal temperature gradients and enhancing convective activity within cyclone frontal structures.

Similar to the generation residual terms, each dissipation residual term displays distinct distributions (Figure 3e). The PDF for RK_Z mirrors that of RG_E , showing modest variability. Its negative Q25 values indicate instances of net dissipation of K_Z . However, the high positive mean, median, Q75, and IQR values suggest that in most cases, there is a net increase of K_Z related to this term, possibly due to the high $B\Phi Z$ value. In contrast, RK_E shows high variability with a left-skewed, long-tail distribution. Both its Q25 and Q75 are negative, indicating that dissipation of K_E frequently occurs.

3.2 Mean Energy Values Across Life Cycle Phases

The use of the CycloPhaser program allowed for the application of an objective criterion to dissect the cyclones into distinct life cycle phases, enabling an investigation of their energy cycles across these phases (Figure 4). Our analysis revealed that the mean energy flow directions rarely change across phases, although the magnitude of each term varies. However, the high standard deviation values indicate that energy fluxes frequently deviate from their mean direction.

The A_Z budget term exhibits mean positive values during the incipient phase, which turn negative during the intensification and mature phases, becoming positive again during the decay phase. The mean K_Z budget values are positive throughout all phases except during the mature phase. In contrast, the A_E budget remains negative across the entire lifecycle, while the K_E budget is positive during the incipient and intensification phases but turns negative during the mature and decay phases. However, it is important to note that the large variability observed in these terms indicates that numerous individual cases deviate significantly from these mean behaviors, including instances of opposite signs or substantially greater magnitudes.

Across the different cyclone phases, the mean behavior suggests that both baroclinic and barotropic conversions remain active, continuously providing energy to the eddies, while the adiabatic contribution from the G_E term is variable throughout the lifecycle. There is a marked enhancement of baroclinic conversions from the incipient to intensification phase, which then decrease in magnitude during the mature and decay phases. Conversely, barotropic conversion peaks during the mature phase, surprisingly with a higher magnitude than the $A_E \rightarrow K_E$ conversion. The behavior of the baroclinic chain is mirrored by the G_E term, which begins with negative values and, during the intensification phase, reaches a higher magnitude than the C_A term. The large variability observed in these energy conversion terms indicates that multiple cyclone configurations likely exist, where either baroclinic or barotropic conversions dominate, or where both pathways coexist with varying intensities. Furthermore, the variability of the G_E term suggests that, in some systems, adiabatic heating plays a significant role in cyclonic development, while in others, diabatic cooling hinders the development. The G_Z term remains negative throughout the entire life cycle, with progressively decreasing values.

Throughout the life cycle, the mean values indicate imports of A_Z , which decrease in the mature phase but rise again during the decay phase. The mean BA_E values exhibit a bimodal behavior, being positive during the incipient and intensification phases but turning negative during the mature and decay phases, though with negligible mean values in the latter. However, the large variability observed suggests that for individual cyclones, imports and exports of A_Z and A_E can differ substantially from these mean patterns, with some systems potentially exhibiting more pronounced or negligible boundary flux contributions.

Across all phases, the RK_Z term stands out with the highest mean values, also displaying the greatest variability, as indicated by the high standard deviation values. This could be influenced by the $B\Phi_Z$ term, as mentioned in the previous section. The BK_Z term also presents high mean values and variability across all phases, suggesting a compensatory effect for the excess energy from the RK_Z term. Lastly, the mean RK_E values are negative across all phases, peaking during the mature phase, indicating significant dissipation in this phase and presenting similar values during the incipient and decay phases.

3.3 Empirical Orthogonal Function Analysis

To better characterize the substantial variability observed in the LEC terms, we performed an Empirical Orthogonal Function (EOF) analysis to identify distinct energetic configurations throughout the cyclone lifecycle. The first EOF accounts for 28.3% of the total variance, followed by the second and third EOFs with 11.0% and 10.9%, respectively. The fourth to eighth EOFs contribute progressively less (8.2%, 7.4%, 5.9%, 5.2%, and 4.4%), with the first four modes explaining 58.4% of the total variance and the first eight reaching 81.3%. These results highlight the dominant patterns of variability, with diminishing contributions beyond the initial modes. This study focuses on the first four EOFs. Although EOFs can be interpreted in terms of both positive and negative contributions, we limit our analysis to the positive phase for simplicity.

Each cyclone can be described as a linear combination of the EOFs, with the corresponding principal components (PCs) indicating how strongly each mode contributes to the system's energetic structure. Consequently, the importance of a given EOF varies from system to system. The energy fluxes associated with each EOF represent deviations from the climatological mean (Figure 4), either reinforcing or weakening specific energy pathways. In this way, each EOF reflects the amplification or attenuation of a characteristic energy transfer pattern within the LEC. In section 3.4, we examine cyclones that project strongly onto each EOF individually.

Figure 5 illustrates the Lorenz Energy Cycle (LEC) terms associated with EOF 1. Across all life cycle phases, there is a consistent enhancement of moist baroclinic conversions, characterized by the positive values of the $A_Z \rightarrow A_E \rightarrow K_E$ conversions, in conjunction with an enhancement of the G_E term. Furthermore, there is an increase in the $K_Z \rightarrow K_E$ conversion (C_K becomes more negative), particularly during the mature and decay phases. These variations largely reinforce the mean behavior observed across the cyclone's life cycle. In addition, A_Z imports are enhanced, while a negative tendency is observed in its budget and generation terms. Conversely, the K_Z




Fig. 4 Mean values and standard deviation of the limited-area Lorenz Energy Cycle (LEC) terms for each life cycle phase : incipient (A), intensification (B), mature (C), and decay (D). Each box represents an energy component ($J m^{-2}$), with arrows showing the direction of energy flow ($W m^{-2}$). The numbers adjacent to the arrows indicate the magnitude and direction of the energy flux, with green indicating positive values and red indicating negative values

boundary, budget, and residual terms show a positive tendency. Lastly, the C_Z term presents a weak positive signal during the incipient and decay phases, and a negative tendency during the intensification and mature phases.

During the incipient (Figure 6A) and intensification phases (Figure 6B), EOF 2 displays an overall weakening of the C_A term, in contrast to EOF 1. Additionally, there is a reduction in barotropic conversions, indicating a $K_E \rightarrow K_Z$ tendency, alongside a weakening of K_E imports. This reduction is compensated by an enhancement in A_E imports, which appears to be the primary mechanism driving the increase in the C_E




Fig. 5 Lorenz Energy Cycle (LEC) terms for the first Empirical Orthogonal Function (EOF 1) across different life cycle phases: (A) incipient, (B) intensification, (C) mature, and (D) decay.

term. In the mature phase (Figure 6C), the the signal for A_E imports reverses in sign, and an enhancement of the moist baroclinic chain occurs, accompanied by a slight increase in $K_Z \rightarrow K_E$ conversions and a more pronounced increase in K_E imports. During the decay phase (Figure 6D), both the moist baroclinic chain and K_Z imports show a negative tendency. However, the enhanced barotropic conversions, coupled with a weakening of dissipation (positive tendency for RK_E), result in a slight yet positive tendency in K_E , suggesting a slower dissipation process.

For EOF 3 (Figure 7), the role of the moist baroclinic chain diminishes in eddy development. This is evident from the weakening of the C_E and G_E terms across the incipient (Figure 7A), intensification (Figure 7B), and mature (Figure 7C) phases,




Fig. 6 Lorenz Energy Cycle (LEC) terms for the second Empirical Orthogonal Function (EOF 2) across different life cycle phases: (A) incipient, (B) intensification, (C) mature, and (D) decay.

although the C_A term shows a slight enhancement during these phases. As a result, we can expect a weaker development for the incipient phase, indicated by the reduced K_E budget. However, during the intensification and mature phases, the dominant source of energy shifts to barotropic conversions. During the decay phase (Figure 7D), the marginal enhancement of the C_E term is driven by a slight increase in the G_E term and, more importantly, by an enhancement of A_E imports. Although this works in conjunction with increased K_E imports, the simultaneous enhancement of $K_E \rightarrow K_Z$ conversions leads to the dissipation of eddies.

During the incipient phase (Figure 8A) of EOF 4, the moist baroclinic chain weakens, as do the K_E imports, while the barotropic conversions are enhanced. As the




Fig. 7 Lorenz Energy Cycle (LEC) terms for the third Empirical Orthogonal Function (EOF 3) across different life cycle phases: (A) incipient, (B) intensification, (C) mature, and (D) decay.

492 system progresses to the intensification phase (Figure 8B), the barotropic conversion
 493 signal becomes neutral, and the weakening of the baroclinic chain lessens, accompa-
 494 nished by a strengthening of A_E imports. In the mature phase (Figure 8C), both the
 495 baroclinic chain and the A_E and K_E imports are slightly enhanced. However, the
 496 barotropic conversions reverse, signaling $K_E \rightarrow K_Z$ conversions, indicating a transfer
 497 of energy from eddy kinetic energy to zonal kinetic energy. In the decay phase (Figure
 498 8D), the signal of baroclinic conversions returns to its initial weakened state and A_E
 499 imports are reduced, while K_E are enhanced. Although the $K_E \rightarrow K_Z$ conversions
 500 are enhanced, they are counterbalanced by the K_E imports, leading to a slower eddy
 501 dissipation (higher tendency for K_E the budget).




Fig. 8 Lorenz Energy Cycle (LEC) terms for the fourth Empirical Orthogonal Function (EOF 4) across different life cycle phases: (A) incipient, (B) intensification, (C) mature, and (D) decay.

Here we present a summary of the energy pathways that were enhanced for each EOF, indicating the increased contribution to cyclone development (i.e., K_E increases). During EOF1, both the moist baroclinic and barotropic chains are enhanced, with increased imports of K_E during the incipient and intensification phases. For EOF2, cyclone development is characterized by the enhancement of the $BA_E \rightarrow A_E \rightarrow K_E$ chain during the incipient and intensification phases, followed by an increase in K_E imports during the mature phase. For EOF3, during the incipient phase, there is an enhancement of K_E imports, which is followed by an increase in barotropic conversions during the intensification and mature phases. For EOF4, barotropic conversions are enhanced during the incipient phase, with nearly neutral baroclinic and barotropic

512 conversions during intensification, and an enhanced baroclinic chain along with K_E
 513 imports during the mature phase. During cyclone decay, EOF1 shows an enhancement
 514 in dissipation, EOF2 and EOF4 present an enhancement in exports of K_E , and EOF3
 515 exhibits an increase in $K_E \rightarrow K_Z$ conversions.

516 3.4 Characteristics of Cyclones Based on EOF Signal

517 In this section, as the characteristics of the systems associated with each EOF are ana-
 518 lyzed, the EOF signal becomes relevant, necessitating a distinction between systems
 519 related to positive EOF signals — EOF(+) — and negative EOF signals — EOF(-),
 520 respectively. EOF1 accounts for the largest number of cyclones among the first four
 521 EOFs for both EOF(+) and EOF(-). For EOF(+), it is followed by EOF2, EOF3,
 522 and EOF4, while for EOF(-), it is followed by EOF3, EOF2, and EOF4, respectively
 523 (Figure S1a). The track densities for the cyclones associated with EOF(+) are pre-
 524 sented in Figure 9, while those for EOF(-) are shown in Figure 10. For EOF1(+),
 525 the track density maximum extends from SE-BR and LA-PLATA toward the ARG
 526 region, highlighting the significant contribution of systems originating in the former
 527 regions compared to the latter, as also indicated in Figure 11a. In contrast, for the
 528 remaining EOFs(+), the track density maximum is located near the ARG region, as
 529 as most systems originate there. An exception is observed in EOF3(+), where the track
 530 density maximum extends northeastward due to the increased contribution of SE-BR
 531 systems to this EOF.




Fig. 9 Cyclone track density for the systems associated with EOFs(+): (A) EOF1, (B) EOF2, (C) EOF3, and (D) EOF4. The track density unit is cyclones per 10^6 km² per month. The dashed rectangles mark the genesis regions shown in Figure 1.

532 Meanwhile, EOFs(-) exhibit a more variable behavior. EOF1(-) displays two track
 533 density maxima: one near SE-BR and another near ARG, as fewer systems originate
 534 from LA-PLATA for this EOF (Figure 11c). EOF2(-) is predominantly influenced by
 535 ARG systems, whereas EOF3(-) shows an increased contribution from LA-PLATA
 536 systems, with a primary track density maximum near ARG and a secondary maximum




Fig. 10 Cyclone track density for the systems associated to EOFs(-): (A) EOF1, (B) EOF2, (C) EOF3 and (D) EOF4. The track density unit is cyclonic centers per 10^6 km^2 per month. The track density unit is cyclones per 10^6 km^2 per month. The dashed rectangles mark the genesis regions shown in Figure 1.

over the La Plata River mouth. Lastly, EOF4(-) presents a track density maximum between LA-PLATA and SE-BR, reflecting the increased relative contribution of systems from these regions to this EOF.

Despite regional differences in track density maxima, nearly all EOFs exhibit a common pattern of cyclone tracks extending southeastward across the South Atlantic (Figures 9 and 10), with the exception of EOF3(+). Some systems display exceptional mobility, reaching as far as 150°E . This behavior aligns with the typical lifecycle of cyclonic systems originating near the South American coast, where cyclones generally develop near the coastal region, mature as they propagate southeastward, and decay closer to the Antarctic region (de Souza et al., 2024).

As shown in Figure 11b, EOF1(+) and EOF4(+) systems predominantly originate during JJA, while EOF2(+) and EOF3(+) genesis events are more evenly distributed across the seasons. For EOFs(-), EOF1(-) systems occur mainly during DJF, whereas EOF4(-) systems are more frequent during DJF and MAM. In contrast, EOF2(-) and EOF3(-) systems are predominantly observed during JJA and SON (Figure 11d). The seasonal distribution of EOF2(+) and EOF3(+) systems aligns with the expected behavior, as ARG systems are well-distributed throughout the year (Gramscianinov et al., 2019; Crespo et al., 2021). However, for the remaining EOFs, the observed behavior suggests that seasonal variations are linked to changes in the systems' energetics.

3.5 Groups of Intense Systems

While the previous sections examined the LEC for all cyclonic systems in the South Atlantic — analyzing both the mean behavior and its associated variability through EOF analysis — this section focuses on the energetic groups associated with the most intense cyclones in the dataset. The K-Means algorithm was applied to the first eight PCs of the selected cyclones, identifying five distinct LEC groups, as determined by




Fig. 11 Genesis proportion and seasonal occurrences of EOFs. (A) Genesis proportion for EOF(+), (B) Seasonal occurrences for EOF(+), (C) Genesis proportion for EOF(-), and (D) Seasonal occurrences for EOF(-). The colors represent different genesis regions (ARG, LA-PLATA, SE-BR) in (A) and (C), while in (B) and (D), the colors correspond to seasonal distributions (DJF, MAM, JJA, SON).

the Elbow Method. Following the cluster identification analysis, the energetic characteristics of the systems were reconstructed, with the mean behavior for each group represented in Figure 12.

Comparing the energy fluxes of the clusters with the climatological mean computed across all cyclones provides insight into the mechanisms that distinguish these systems as the most intense (Figure 12). Cluster 1 exhibits a strong enhancement of the moist baroclinic chain, particularly in the C_A and G_E terms, as well as increased barotropic conversion, with both C_A and C_K contributing comparably to the K_E budget. However, imports of K_E are below average. Cluster 2 shows modest increases across all baroclinic and barotropic terms, with the highest K_E imports among all clusters. Cluster 3 displays similar proportions to Cluster 1 for C_A , C_E , and C_K , though with slightly lower values. However, it presents the highest G_E among all clusters, and unlike Cluster 1, it exhibits positive but below-average K_E imports. Cluster 4 features the weakest energy conversions overall, despite showing nearly doubled values for baroclinic and barotropic terms relative to the full-sample average, while still presenting below-average K_E imports. Across all clusters, the residual term is not strictly proportional to the K_E budget: Cluster 1, for instance, shows the highest RK_E despite




Fig. 13 Cyclone track density for the five clusters (A–E) identified among the most intense cyclones in the dataset. These systems are defined as those with maximum central relative vorticity at 850 hPa exceeding the 90th percentile. The dashed rectangles mark the genesis regions shown in Figure 1.

the most intense systems on average, while Clusters 1 and 2 show similar mean intensity. Clusters 1 and 2 are more frequent during JJA, while Cluster 3 peaks in SON and Cluster 4 shows no strong seasonality (Figure 14c). In all clusters, the proportion of cyclones originating in LA-PLATA (Figure 14d) exceeds its baseline occurrence in the full dataset (18.9%), reinforcing the region’s role in the development of intense systems, as noted by Gramscianinov et al. (2019). The seasonal distributions are also consistent with prior findings for the LA-PLATA region (Crespo et al., 2021). The low frequency of cyclones forming in SE-BR is expected, as these tend to be weaker (Gramscianinov et al., 2019; de Souza et al., 2024). Although subtropical cyclones are most frequent over SE-BR during DJF (Evans and Braun, 2012; Gozzo et al., 2014), their limited presence in this intense-cyclone subset prevents them from significantly influencing the results.

4 Discussion

4.1 Key Features

The LEC diagrams (Figure 4) indicate that the mean energetic behavior is preserved across distinct life cycle phases. However, the high variability shown in Figure 3 reveals a wide range of behaviors among South Atlantic cyclones. While this section focuses on the mean energy cycle, it is important to recognize that energy budget terms serve primarily as a starting point. The energy conversions and fluxes offer more robust insight into the dynamical processes governing cyclone development. In addition to the phase-specific diagrams, a schematic of the mean Lorenz Energy Cycle for all




Fig. 14 Characteristics of the five clusters (A–E) identified among the most intense cyclones in the dataset. (A) Number of cyclones per cluster, (B) maximum intensity (defined as maximum central relative vorticity at 850 hPa), (C) seasonal distribution of occurrences, and (D) genesis region count. In (C), colors represent different seasons (DJF, MAM, JJA, SON), while in (D), colors correspond to genesis regions (ARG, LA-PLATA, SE-BR).

systems is shown in Figure 15, facilitating comparison across phases and highlighting dominant pathways.

The A_Z term increases during the early phases due to persistent influxes across the boundaries ($BA_Z > 0$), rather than local generation (G_Z). As baroclinic conversions intensify, the A_Z budget becomes negative during the intensification and mature phases, before returning to positive in the decay phase. This evolution suggests that A_Z variations are not driven by latitudinal diabatic heating contrasts. A similar pattern was reported by Dias Pinto and Rocha (2011), although their methodology excluded the incipient phase due to the use of sea-level pressure for cyclone tracking, instead of relative vorticity at 850 hPa (Sinclair, 1994; Hoskins and Hodges, 2002).

For K_Z , a general increase is observed except during the mature phase, when strong barotropic conversion ($K_Z \rightarrow K_E$) coincides with negative C_Z values. This indicates descending motion at warmer latitudes and ascending motion at colder latitudes as the cold front reaches the equatorward side of the cyclone and the warm front reaches the poleward side. Notably, BK_Z and RK_Z present high magnitudes: the former suggests export of zonal kinetic energy (e.g., jet stream outflow), while the latter aggregates dissipation, boundary pressure work ($B\Phi_Z$), subgrid transfers, and numerical errors.




Fig. 15 Schematic representation of the mean Lorenz Energy Cycle (LEC) for all analyzed cyclones in the South Atlantic, combining information from all life cycle phases. Arrows represent energy fluxes between reservoirs, with colors indicating the life cycle phases. The arrow thicknesses and symbol sizes for the balance terms ($\partial/\partial t$) are proportional to the absolute magnitude of each term, as presented in Figure 4.

As the Semi-Lagrangian framework does not account for changes in the background energetics along a moving domain, part of the apparent time evolution in A_Z and K_Z may simply reflect the cyclone's poleward drift (de Souza et al., 2024). Such drift carries the system into latitudes of larger climatological background APE, implying an increase in A_Z (Novak and Tailleux, 2018; Liu et al., 2024). Because the zonally integrated K_Z reaches its climatological maximum near 60°S (Novak and Tailleux, 2018; Liu et al., 2024), K_Z should decrease for storms that originate over ARG and migrate poleward, but increase for those formed in SE-BR or LA-PLATA. In practice,

however, both reservoirs evolve irregularly and non-monotonically, indicating that intrinsic cyclone dynamics, rather than passive domain advection, control the local energetics, corroborating Federer et al. (2025).

The A_E budget is negative throughout the life cycle, with minimal values during the incipient and intensification phases. This reflects a highly active and efficient baroclinic chain. Energy is transferred from A_Z via C_A , and from A_E to K_E via C_E , driven by meridional heat transport and vertical motions in frontal zones. During intensification, G_E becomes positive, associated with convective heating along the cold front (e.g., Govekar et al., 2011). Simultaneously, radiative heating in the warm sector due to long-wave absorption from mid- to high-level clouds (Lau and Crane, 1997; Keshtgar et al., 2023), combined with latent heat release from intense precipitation during this phase (McErlich et al., 2023), enhances G_E . As the system matures and convection weakens, BA_E reverses (export), and the A_E budget sharply decreases. During decay, with disorganized frontal structures and weak diabatic forcing, A_E returns to near-zero budget values.

The K_E reservoir increases during incipient and intensification phases, as baroclinic, barotropic, and boundary fluxes supply energy. In the mature phase, barotropic conversion peaks, but weakening of baroclinic forcing and reversal of boundary fluxes, along with increased RK_E , lead to a net energy loss. During the decay phase, diminishing energy inputs and rising exports further reduce K_E . Although RK_E includes numerical errors and the $B\Phi E$ term, the trend suggests dominant dissipative processes, in line with Smith (1980).

Both $B\Phi_Z$ and $B\Phi_E$ were directly computed and yielded large values (Table 1, Figure 3d). These terms, representing pressure work by zonal and meridional winds at the domain boundaries, are highly sensitive to small errors in the geopotential field (Brennan and Vincent, 1980). Despite quality control in ERA5, some inconsistencies may persist, and since the LECTK tool does not apply preprocessing filters, such errors likely propagate into the diagnostics. Future work should examine the physical interpretation of $B\Phi$ terms and the causes of their anomalous magnitudes, as they are seldom reported in literature.

4.2 Cyclone Groups

Substantial variability in the dataset, as revealed by the EOF analysis, is largely captured by EOF1(+) and EOF1(-), which together represent nearly 40% of all systems. EOF1(+) is linked to stronger cyclones, while EOF1(-) corresponds to weaker ones (Supplementary Figure S1b), indicating a bi-modal structure in cyclone intensity. This suggests that "day-to-day" systems tend to follow distinct energetic regimes. Traditional cyclone climatologies over the South Atlantic tend to group all cyclones together (Sinclair, 1994; Hoskins and Hodges, 2005; Reboita et al., 2010; Gramscianinov et al., 2019, e.g.). In contrast, the results presented here suggest that these systems exhibit distinct behaviors and may therefore be categorized into separate groups.

These intensity differences manifest in their energy cycles and physical characteristics. Figure 16 presents schematic LECs for the four leading EOF(+) groups. EOF1(+) displays a consistent enhancement across all energy pathways, while EOF1(-) shows a reduction. These differences extend to cyclone mobility: EOF1(+) systems follow

688 the South Atlantic storm track (Hoskins and Hodges, 2005; Gramscianinov et al.,
 689 2019), whereas EOF1(-) systems remain more confined near the South American
 690 coast (Figure 10a), particularly within the ARG and SE-BR regions (Figure 11d), and
 691 exhibit lower translational speeds (Figure S1).




Fig. 16 Schematic representation of the mean Lorenz Energy Cycle (LEC) for cyclones associated with the four leading positive Empirical Orthogonal Functions (EOF1(+) to EOF4(+)). Arrows represent energy fluxes between reservoirs, with colors indicating life cycle phases. The thickness of each arrow and the size of the balance terms ($\partial/\partial t$) symbols are proportional to the absolute magnitude of each term.

692 EOF2-EOF4 describe less frequent cyclone types. EOF2(+), EOF3(-), and
 693 EOF4(-) are associated with relatively stronger systems, while their counterparts cor-
 694 respond to weaker ones (Figure S1b). EOF2(+) is notable for lacking enhancements

in baroclinic/barotropic conversions or K_E imports during early phases. EOF3(−) and EOF4(−), in contrast, display a pronounced intensification of the $G_E \rightarrow A_E \rightarrow K_E$ chain, especially during the incipient and intensification phases. These patterns indicate that despite a shared dominant energy structure, South Atlantic cyclones express diverse configurations in their development. This variability is further reflected in differences in genesis region and seasonality (Figure 11), with EOF1(+) and EOF3(−) dominating among the most intense systems (Supplementary Figure S3).

For intense cyclones (above the 90th percentile in central vorticity), genesis is most frequent in LA-PLATA and during JJA—consistent with earlier studies identifying this region as favorable for strong development (Reboita et al., 2010; Gramscianinov et al., 2019; Crespo et al., 2021). These systems are primarily linked to moist baroclinic instability (Figure 12), particularly under the influence of a strong upper-level jet in austral winter. Meanwhile, SE-BR systems are less frequent, especially during DJF, when baroclinicity in that region is weaker (Gramscianinov et al., 2019; Crespo et al., 2021).

Among the four LEC-based clusters, Cluster 4 contains the largest number of systems but the lowest maximum central vorticity, making it the weakest group. Importantly, all clusters show enhanced LEC terms relative to the overall mean, reinforcing the connection between stronger energy fluxes and cyclone intensity. These findings support the robustness of the Semi-Lagrangian framework and highlight the existence of distinct energetic regimes. Even within the subset of intense cyclones, considerable diversity in energy cycle structures persists, indicating that multiple pathways can lead to strong cyclogenesis in the South Atlantic.

4.3 Mechanisms of Energy Transfer

Throughout this study, barotropic conversions (C_K) emerged as a key mechanism for cyclone development in the South American region. While extratropical cyclones are traditionally associated with baroclinic instability (Bjerknes and Solberg, 1922; Charney and Eliassen, 1964; Hoskins and Valdes, 1990), our results show that moist baroclinic processes dominate during the intensification phase, whereas barotropic conversions peak during the mature stage (Figure 4). Notably, C_K magnitudes are nearly three times larger than those of C_A , suggesting a stronger contribution from zonal kinetic energy to eddy development than from meridional temperature gradients.

Despite LEC-based studies dating back decades, few have examined South Atlantic cyclones, limiting direct comparisons. Case studies on tropical systems emphasize the $G_E \rightarrow A_E \rightarrow K_E$ chain and barotropic conversions (Brennan and Vincent, 1980; Veiga et al., 2008), while subtropical systems show mixed roles for both baroclinic and barotropic conversions (Michaelides, 1987; Dias Pinto et al., 2013; Pezza et al., 2014; Cavicchia et al., 2018). For extratropical systems, energy pathways vary: some are baroclinically dominated (Dias Pinto and Rocha, 2011; Black and Pezza, 2013), others barotropically (Michaelides, 1992; Dias Pinto and Rocha, 2011), and some combine both with varying degrees of K_E import (Wahab et al., 2002; Bulic, 2006; Pezza et al., 2010). Thus, although the relevance of barotropic instability in extratropical cyclones is not novel, this work provides the first large-sample LEC analysis for the South Atlantic, offering a consolidated view of this behavior. These findings are consistent

with recent studies that link storm track intensification in the Southern Hemisphere to enhanced barotropic growth (Chemke et al., 2022), associate storm growth time with barotropic conversion (Hadas and Kaspi, 2025), and identify barotropic conversions from K_Z to K_E within storm track regions (Liu et al., 2024).

It is also important to emphasize that, while previous studies investigated the LEC using an Eulerian framework, this study employed a Semi-Lagrangian framework. This approach has the advantage of isolating the energetics strictly related to the target system. However, the extent to which this methodology differs from the Eulerian framework in representing the magnitude and direction of energy fluxes has not been extensively assessed, apart from one study by Michaelides et al. (1999). Also, the extent to which our findings generalize to other regions remains uncertain. Given the stronger jets and meridional wind gradients in the Southern Hemisphere (e.g., Swart et al., 2019; Jain et al., 2023; Savita et al., 2023), barotropic conversions may be more pronounced in the South Atlantic than in the North Atlantic—a hypothesis requiring further study.

Another distinction is found in the decay phase. While most previous studies report $K_E \rightarrow K_Z$ conversions during cyclone decay (e.g., Dias Pinto and Rocha, 2011; Veiga et al., 2008; Pezza et al., 2014), this pattern was not observed here. In our analysis, the decay phase, the longest for these systems (de Souza et al., 2024), is averaged over multiple timesteps and features C_K values that gradually shift from strongly negative in the mature phase to near neutral at the end of the lifecycle (Figures 4C and S4). This suggests that barotropic conversion continues to sustain the system during decay, until dissipation dominates. Prior use of large Eulerian domains may have underestimated C_K , as seen in Black and Pezza (2013), where C_K remains positive across all phases.

Some limitations of our approach must be acknowledged. Although the Semi-Lagrangian framework follows the cyclone’s trajectory, it does not account for regional variations in background APE and kinetic energy. This may influence A_Z , K_Z , and their boundary terms as the system moves through varying energy environments. Furthermore, the LEC formulation relies on a global reference state for APE. In contrast, the local APE framework proposed by Novak and Tailleux (2018) defines APE relative to each air parcel’s individual reference state. While this approach can yield additional insights—such as the potential for cyclones to locally generate APE (Federer et al., 2024), the overall global energy fluxes remain consistent between the global and local formulations (Liu et al., 2024). Moreover, the modest BA_Z values observed here (Figure 4) may be partially explained by APE transport from the polar upper troposphere, as shown for the North Atlantic storm tracks by Federer et al. (2025).

5 Summary and Conclusions

In this study, we examined the Lorenz Energy Cycle (LEC) for 6,789 cyclonic systems originating in the Southwestern Atlantic, presenting what is, to the best of our knowledge, the first comprehensive climatology of cyclones’ energy cycles in this region. A notable exception is the work by Black and Pezza (2013), which focused solely on explosive cyclones and used large fixed domains, contrasting with the Semi-Lagrangian

approach adopted in this study. While studies such as Smith (1980) provide valuable reviews of the energetics of extratropical cyclones, focusing on broad averages and generalized mechanisms, and case studies have explored the dynamics of specific systems (Pezza et al., 2014; Dias Pinto and Rocha, 2011; Cavicchia et al., 2018, e.g.), our study adopts a climatological approach. Specifically, we track the energy cycles of individual cyclones across distinct life cycle phases, offering a more detailed investigation of the unique physical mechanisms that drive energy conversions in this region. Furthermore, the use of the Cyclophaser program to dissect the life cycles of cyclones has enabled a novel investigation of the energy cycle across different developmental stages, providing deeper insights into the variability and dynamism of these systems.

The results presented here demonstrate that, for most cyclonic systems in the South Atlantic, a clear pattern of energy flow is evident. This pattern is characterized by both barotropic and baroclinic conversions providing energy for eddy development and is preserved throughout the cyclone life cycle phases, despite variations in magnitude. Barotropic conversions tend to be 2 to 3 times larger in magnitude than baroclinic conversions, with the former peaking during the mature phase and the latter during the intensification phase. The generation of eddy potential energy and the imports of eddy kinetic energy play secondary roles. The former peaks during the intensification phase and remains positive through the mature and decay phases, while the latter occurs primarily during the incipient (cyclogenesis) and intensification phases.

Despite the high variability among the systems, the EOF analysis indicates that, in most cases, the main behavior is preserved. This variability is instead expressed as a strengthening and/or weakening of specific energy pathways, indicating that cyclonic development in the South Atlantic region can be attributed to distinct dynamical processes, as supported by the literature. Consequently, it is possible to hypothesize that these systems can be grouped into distinct types based on their characteristics and energy cycles. This heterogeneity is also evident among the most intense systems. For these systems, distinct groups exhibit varying relative contributions of energy from barotropic and baroclinic chains, as well as differences in eddy generation of APE and imports of eddy kinetic energy. Although most of these intense systems originate in the LA-PLATA region, the variability in the relative proportions of genesis regions among the different clusters is reflected in their energy cycles and mean characteristics.

The goal of this study was to investigate the energy cycle of cyclones in the South Atlantic region using a Semi-Lagrangian approach. As a pioneering study of this kind, several questions have emerged throughout its development. For instance, previous studies for this region have assessed the distinct synoptic and dynamical conditions related to cyclogenesis in the three primary genesis regions, as well as the seasonal variability in their development (Gramscianinov et al., 2019; Crespo et al., 2021). A distinction in the energy cycles among these genesis regions, along with their seasonal variability, would therefore be valuable. Additionally, given the prominence of barotropic conversions as an energy source for cyclone development, the question arises as to whether barotropic instability is truly occurring near the cyclone center. de Souza (2024) proposed a framework for classifying cyclones into distinct groups

based on their energy cycles. In an upcoming study, this method will be applied to address these questions and further investigate these topics.

Appendix

Mathematical expressions used for the calculation of the Lorenz Energy Cycle (LEC), adopting the notation from Michaelides (1987).

Firstly, we define the zonal mean of a variable X , between longitudes λ_1 and λ_2 :

$$[X]_\lambda = \frac{1}{\lambda_2 - \lambda_1} \int_{\lambda_2}^{\lambda_1} X d\lambda \quad (13)$$

The eddy component of this variable is its deviation from the zonal mean:

$$(X)_\lambda = X - [X]_\lambda \quad (14)$$

The domain mean of the variable X , defined over the computational domain bounded by longitudes λ_1 and λ_2 , and latitudes ϕ_1 and ϕ_2 , is given by:

$$[X]_{\lambda\phi} = \left(\frac{1}{\lambda_2 - \lambda_1} \right) \left(\frac{1}{\sin \phi_2 - \sin \phi_1} \right) \int_{\lambda_2}^{\lambda_1} X \cos \phi d\lambda d\phi \quad (15)$$

Similarly, we define the deviation of the zonal mean from the domain mean:

$$([X]_\lambda)_\phi = [X]_\lambda - [X]_{\lambda\phi} \quad (16)$$

From the definitions above, the four energy components used in the LEC computation are defined as follows:

$$A_Z = \int_{p_t}^{p_b} \frac{[(T)_\lambda]^2_{\lambda\phi}}{2[\sigma]_{\lambda\phi}} dp \quad (17)$$

$$A_E = \int_{p_t}^{p_b} \frac{[(T)_\lambda]^2_{\lambda\phi}}{2[\sigma]_{\lambda\phi}} dp \quad (18)$$

$$K_Z = \int_{p_t}^{p_b} \frac{[[u]_\lambda]^2 + [[v]_\lambda]^2_{\lambda\phi}}{2g} dp \quad (19)$$

$$K_E = \int_{p_t}^{p_b} \frac{[(u)_\lambda]^2 + [(v)_\lambda]^2_{\lambda\phi}}{2g} dp \quad (20)$$

where p is the atmospheric pressure, with subscripts b and t denoting the lower (base) and upper (top) pressure boundaries of the atmosphere, respectively. T represents temperature, g is the acceleration due to gravity, and u and v are the zonal and meridional wind components, respectively. The static stability parameter σ is defined as:

$$\sigma = \left[\frac{gT}{c_p} - \frac{pg}{R} \frac{\partial T}{\partial p} \right]_{\lambda\phi} \quad (21)$$

843 where c_p is the specific heat at constant pressure, and R is the ideal gas constant
 844 for dry air.

845 The four conversion terms are defined as follows, integrating over the atmospheric
 846 column from the base (p_b) to the top (p_t) pressures:

$$C_Z = \int_{p_t}^{p_b} -[([T]_\lambda)_\phi([w]_\lambda)_\phi]_{\lambda\phi} \frac{R}{gp} dp \quad (22)$$

$$C_E = \int_{p_t}^{p_b} -[(T)_\lambda(\omega)_\lambda]_{\lambda\phi} \frac{R}{gp} dp \quad (23)$$

$$C_A = \int_{p_t}^{p_b} -\left(\frac{1}{2a\sigma} \left[(v)_\lambda(T)_\lambda \frac{\partial([T]_\lambda)_\phi}{\partial\phi} \right]_{\lambda\phi} + \frac{1}{\sigma} \left[(\omega)_\lambda(T)_\lambda \frac{\partial([T]_\lambda)_\phi}{\partial p} \right]_{\lambda\phi} \right) dp \quad (24)$$

$$C_K = \int_{p_t}^{p_b} \frac{1}{g} \left(\left[\frac{\cos\phi}{a} (u)_\lambda(v)_\lambda \frac{\partial}{\partial\phi} \left(\frac{[u]_\lambda}{\cos\phi} \right) \right]_{\lambda\phi} + \left[\frac{(v)_\lambda^2}{a} \frac{\partial[v]_\lambda}{\partial\phi} \right]_{\lambda\phi} \right. \\ \left. + \left[\frac{\tan\phi}{a} (u)_\lambda^2 [v]_\lambda \right]_{\lambda\phi} + \left[(\omega)_\lambda(u)_\lambda \frac{\partial[u]_\lambda}{\partial p} \right]_{\lambda\phi} + \left[(\omega)_\lambda(v)_\lambda \frac{\partial[v]_\lambda}{\partial p} \right]_{\lambda\phi} \right) dp \quad (25)$$

847 where a is the Earth's radius and ω is the vertical velocity in isobaric coordinates.
 848 The APE generation and K dissipation terms are defined as:

$$G_Z = \int_{p_t}^{p_b} \frac{[(q)_\lambda]_\phi([T]_\lambda)_\phi]_{\lambda\phi}}{c_p[\sigma]_{\lambda\phi}} dp \quad (26)$$

$$G_E = \int_{p_t}^{p_b} \frac{[(q)_\lambda(T)_\lambda]_{\lambda\phi}}{c_p[\sigma]_{\lambda\phi}} dp \quad (27)$$

$$D_Z = - \int_{p_t}^{p_b} \frac{1}{g} [[u]_\lambda[F_\lambda]_\lambda + [v]_\lambda[F_\phi]_\lambda]_{\lambda\phi} dp \quad (28)$$

$$D_E = - \int_{p_t}^{p_b} \frac{1}{g} [(u)_\lambda(F_\lambda)_\lambda + (v)_\lambda(F_\phi)_\lambda]_{\lambda\phi} dp \quad (29)$$

849 Here, F_λ and F_ϕ represent the zonal and meridional frictional components,
 850 respectively, and q is the diabatic heating term, computed as a residual from the
 851 thermodynamic equation:

$$\frac{q}{c_p} = \frac{\partial T}{\partial t} - \vec{V}_H \cdot \vec{\nabla}_p T - S_p \omega \quad (30)$$

852 where $-\vec{V}_H \cdot \vec{\nabla}_p T$ represents the horizontal advection of temperature and S_p
 853 approximates the static stability, given by:

$$S_p \equiv -\frac{T}{\theta} \frac{\partial\theta}{\partial p} \quad (31)$$

854 where θ is the potential temperature.

The boundary terms are given by:

$$\begin{aligned} \text{BAZ} = & c_1 \int_{p_1}^{p_2} \int_{\varphi_1}^{\varphi_2} \frac{1}{2[\sigma]_{\lambda\varphi}} \left(2 ([T]_{\lambda})_{\varphi} (T)_{\lambda} u + ([T]_{\lambda\varphi})_{\varphi}^2 u \right)_{\lambda_1}^{\lambda_2} \\ & \times d\varphi dp + c_2 \int_{p_1}^{p_2} \frac{1}{2[\sigma]_{\lambda\varphi}} \left(2 [(v)_{\lambda} (T)_{\lambda}]_{\lambda} ([T]_{\lambda})_{\varphi} \cos \varphi + ([T]_{\lambda})_{\varphi}^2 [(v)_{\lambda} \cos \varphi] \right)_{\varphi_1}^{\varphi_2} dp \\ & - \frac{1}{2[\sigma]_{\lambda\varphi}} \left([2(\omega)_{\lambda} (T)_{\lambda}]_{\lambda} ([T]_{\lambda})_{\varphi} + [\omega]_{\lambda} ([T]_{\lambda})_{\varphi}^2 \right)_{\lambda\varphi}^{p_2} \end{aligned} \quad (32)$$

$$\begin{aligned} \text{BAE} = & c_1 \int_{p_1}^{p_2} \int_{\varphi_1}^{\varphi_2} \frac{1}{2[\sigma]_{\lambda\varphi}} [u(T)_{\lambda}^2]_{\lambda_1}^{\lambda_2} d\varphi dp \\ & + c_2 \int_{p_1}^{p_2} \frac{1}{2[\sigma]_{\lambda\varphi}} \left([(T)_{\lambda}^2 v]_{\lambda} \cos \varphi \right)_{\varphi_1}^{\varphi_2} dp \\ & - \left(\frac{[\omega(T)_{\lambda}^2]_{\lambda\varphi}}{2[\sigma]_{\lambda\varphi}} \right)_{p_1}^{p_2} \end{aligned} \quad (33)$$

$$\begin{aligned} \text{BKZ} = & c_1 \int_{p_1}^{p_2} \int_{\varphi_1}^{\varphi_2} \frac{1}{2g} \left(u [u^2 + v^2 - (u)_{\lambda}^2 - (v)_{\lambda}^2] \right)_{\lambda_1}^{\lambda_2} \\ & \times d\varphi dp + c_2 \int_{p_1}^{p_2} \frac{1}{2g} \left([v \cos \varphi [u^2 + v^2 - (u)_{\lambda}^2 - (v)_{\lambda}^2]] \right)_{\varphi_1}^{\varphi_2} dp \\ & - \left(\frac{1}{2g} [\omega [u^2 + v^2 - (u)_{\lambda}^2 - (v)_{\lambda}^2]]_{\lambda\varphi} \right)_{p_1}^{p_2} \end{aligned} \quad (34)$$

$$\begin{aligned} \text{BKE} = & c_1 \int_{p_1}^{p_2} \int_{\varphi_1}^{\varphi_2} \frac{1}{2g} \left(u [(u)_{\lambda}^2 + (v)_{\lambda}^2] \right)_{\lambda_1}^{\lambda_2} d\varphi dp \\ & + c_2 \int_{p_1}^{p_2} \frac{1}{2g} \left([v \cos \varphi [(u)_{\lambda}^2 + (v)_{\lambda}^2]] \right)_{\varphi_1}^{\varphi_2} dp \\ & - \left(\frac{1}{2g} [\omega [(u)_{\lambda}^2 + (v)_{\lambda}^2]]_{\lambda\varphi} \right)_{p_1}^{p_2} \end{aligned} \quad (35)$$

where $c_1 = -[a(\lambda_2 - \lambda_1)(\sin \varphi_2 - \sin \varphi_1)]^{-1}$, $c_2 = -[a \ x(\sin \varphi_2 - \sin \varphi_1)]^{-1}$.

Lastly, the terms $B\Phi Z$ and $B\Phi E$ are given by:

$$\begin{aligned} \text{B}\Phi\text{Z} = & c_1 \int_{p_1}^{p_2} \int_{\varphi_1}^{\varphi_2} \frac{1}{g} \left([v]_{\lambda} ([\Phi]_{\lambda})_{\varphi} \right)_{\lambda_1}^{\lambda_2} d\varphi dp \\ & + c_2 \int_{p_1}^{p_2} \frac{1}{g} \left(\cos \varphi [v]_{\lambda} ([\Phi]_{\lambda})_{\varphi} \right)_{\varphi_1}^{\varphi_2} dp \\ & - \frac{1}{g} \left([([\omega]_{\lambda})_{\varphi} ([\Phi]_{\lambda})_{\varphi}]_{\lambda\varphi} \right)_{p_1}^{p_2} \\ \text{B}\Phi\text{E} = & c_1 \int_{p_1}^{p_2} \int_{\varphi_1}^{\varphi_2} \frac{1}{g} ((u)_{\lambda} (\Phi)_{\lambda})_{\lambda_1}^{\lambda_2} d\varphi dp \end{aligned} \quad (36)$$

$$\begin{aligned}
& + c_2 \int_{p_1}^{p_2} \frac{1}{g} \left([(\nu)_\lambda(\Phi)_{\lambda_\lambda}]_\lambda \cos \varphi \right)_{\varphi_1}^{\varphi_2} dp \\
& - \frac{1}{g} \left([(\omega)_\lambda(\Phi)_{\lambda_\varphi}]_{p_1}^{p_2} \right)
\end{aligned} \tag{37}$$

Declarations

Funding

This study was partly financed by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) under Finance Code 001.

Competing interests

The authors have no relevant financial or non-financial interests to disclose.

Data Availability

The cyclone tracks used in this study were obtained from the *Atlantic Extratropical Cyclone Tracks Database*, available at <https://doi.org/10.17632/kwcvfr52hp.4>. All datasets, including the computed Lorenz Energy Cycle results for each cyclone, along with the scripts used to generate the figures and analyses presented in this study, are publicly available at the following GitHub repository: https://github.com/daniloceano/energetic_patterns_cyclones_south_atlantic. The source code of the **Cyclophaser** package, used for detecting cyclone life cycle phases, is available on PyPI at <https://pypi.org/project/cyclophaser/>. The **LorenzCycleToolkit** package, used for computing the Lorenz Energy Cycle components, is also available on PyPI at <https://pypi.org/project/LorenzCycleToolkit/>.

Author contributions

Conceptualization: Danilo Couto de Souza, Pedro Leite da Silva Dias, Ricardo de Camargo, Carolina Barnez Gramscianinov; Methodology: Danilo Couto de Souza, Pedro Leite da Silva Dias, Ricardo de Camargo; Formal analysis and investigation: Danilo Couto de Souza, Pedro Leite da Silva Dias; Writing - original draft preparation: Danilo Couto de Souza; Writing - review and editing: Danilo Couto de Souza, Pedro Leite da Silva Dias, Ricardo de Camargo, Carolina Barnez Gramscianinov; Funding acquisition: Pedro Leite da Silva Dias, Ricardo de Camargo; Resources: Pedro Leite da Silva Dias, Ricardo de Camargo; Supervision: Pedro Leite da Silva Dias, Ricardo de Camargo.

References

Andrade, H.N., A.B. Nunes, M.S. Teixeira, M.F.L. de Quadro, V.D. de Avila, F.S.C. de Oliveira, and R.d.C.M. Alves. 2025. Composite analysis of explosive cyclones in the southern atlantic ocean. *International Journal of Climatology* 45(4): e8737 .

- 889 Andrade, H.N., M.F.L.D. Quadro, A.B. Nunes, F.S.D. Oliveira, V.D.D. Avila, M.S.
890 Teixeira, and R.D.C.M. Alves. 2024. Explosive cyclones occurred between 2010 and
891 2020 in the south atlantic under the perspective of two detection schemes. *Anais*
892 *da Academia Brasileira de Ciências* 96(suppl 1): e20231051 .
- 893 Bengfort, B. and R. Bilbro. 2019. Yellowbrick: Visualizing the scikit-learn model
894 selection process. *Journal of Open Source Software* 4(35): 1075 .
- 895 Bjerknes, J. and H. Solberg. 1922. *Life cycle of cyclones and the polar front theory of*
896 *atmospheric circulation*. Grondahl.
- 897 Black, M.T. and A.B. Pezza. 2013. A universal, broad-environment energy conversion
898 signature of explosive cyclones. *Geophysical Research Letters* 40(2): 452–457 .
- 899 Brennan, F.E. and D.G. Vincent. 1980. Zonal and eddy components of the synoptic-
900 scale energy budget during intensification of hurricane carmen (1974). *Monthly*
901 *Weather Review* 108(7): 954–965. [https://doi.org/10.1175/1520-0493\(1980\)](https://doi.org/10.1175/1520-0493(1980)108<0954:ZAECOT>2.0.CO;2)
902 108<0954:ZAECOT>2.0.CO;2 .
- 903 Bulic, I. 2006. Limited area energy budget during a life cycle of genoa cyclone
904 (18-21 november 1999). *NUOVO CIMENTO-SOCIETA ITALIANA DI FISICA*
905 *SEZIONE C* 29(2): 167 .
- 906 Cardoso, A.A., R.P. da Rocha, and N.M. Crespo. 2022. Synoptic climatology of
907 subtropical cyclone impacts on near-surface winds over the south atlantic basin.
908 *Earth and Space Science* 9(11): e2022EA002482 .
- 909 Cavicchia, L., A. Dowdy, and K. Walsh. 2018. Energetics and dynamics of sub-
910 tropical australian east coast cyclones: Two contrasting cases. *Monthly Weather*
911 *Review* 146(5): 1511–1525 .
- 912 Charney, J.G. and A. Eliassen. 1964. On the growth of the hurricane depression.
913 *Journal of the Atmospheric Sciences* 21(1): 68–75 .
- 914 Chemke, R., Y. Ming, and J. Yuval. 2022. The intensification of winter mid-latitude
915 storm tracks in the southern hemisphere. *Nature climate change* 12(6): 553–557 .
- 916 Crespo, N.M., R.P. da Rocha, M. Sprenger, and H. Wernli. 2021. A potential vorticity
917 perspective on cyclogenesis over centre-eastern south america. *International Journal*
918 *of Climatology* 41(1): 663–678 .
- 919 da Silva, M.B.L., D.C. de Souza, F.T.C. Barreto, R. Tecchio, R. de Freitas
920 Pimentel dos Anjos, C.B. Gramscianinov, and R. De Camargo. 2025. Classifica-
921 tion of synoptic weather patterns associated with extreme wave events in different
922 regions of western south atlantic. *Natural Hazards*: 1–25 .

- 923 de Souza, D. and R.R. da Silva. 2021. Ocean-land atmosphere model (olam) perfor-
 924 mance for major extreme meteorological events near the coastal region of southern
 925 brazil. *Climate Research* 84: 1–21 .
- 926 de Souza, D.C. 2024. *Cyclones in the Southwestern Atlantic: Life Cycle and Ener-*
 927 *getics*. Doctoral thesis, Universidade de São Paulo, São Paulo, Brazil. Versão
 928 corrigida.
- 929 de Souza, D.C., N.M. Crespo, D.V. da Silva, L.M. Harada, R.M.P. de Godoy, L.M.
 930 Domingues, R. Luiz, C.A. Bortolozo, D. Metodiev, M.R.M. de Andrade, et al. 2024.
 931 Extreme rainfall and landslides as a response to human-induced climate change: a
 932 case study at baixada santista, brazil, 2020. *Natural Hazards*: 1–26 .
- 933 de Souza, D.C., P.L.S. da Dias, M.B.L. da Silva, and R. de Camargo. 2024. Lorenz-
 934 cycletoolkit: A comprehensive python tool for analyzing atmospheric energy cycles.
 935 *Journal of Open Source Software* 9(101): 7139 .
- 936 de Souza, D.C., P.L.S. da Dias, C.B. Gramscianinov, and R. de Camargo. 2025.
 937 Cyclophaser: A python package for detecting extratropical cyclone life cycles.
 938 *Journal of Open Source Software* 10(108): 7363 .
- 939 de Souza, D.C., P.L. da Silva Dias, C.B. Gramscianinov, M.B.L. da Silva, and
 940 R. de Camargo. 2024. New perspectives on south atlantic storm track through
 941 an automatic method for detecting extratropical cyclones’ lifecycle. *International*
 942 *Journal of Climatology* 44(10): 3568–3588 .
- 943 de Souza, D.C., R. Ramos da Silva, P. Gomes da Silva, A.F.H. Fetter Filho, F.J.
 944 Mendez, and D. Werth. 2022. A hybrid regional climate downscaling for the south-
 945 ern brazil coastal region. *International Journal of Climatology* 42(13): 6753–6770
 946 .
- 947 de Souza, D.C., V.A. Ranieri, P.L. da Silva Dias, A.R. Flores, and R. de Camargo.
 948 2025. Revisiting the rare transition of a south atlantic cyclone to
 949 tropical storm akar: Energy cycle and stratosphere-troposphere interac-
 950 tion. manuscript submitted to Climate Dynamics. Preprint available at
 951 <https://scity.org/articles/activity/10.21203/rs.3.rs-6536558/v1>.
- 952 Dias Pinto, J.R., M.S. Reboita, and R.P. da Rocha. 2013. Synoptic and dynam-
 953 ical analysis of subtropical cyclone anita (2010) and its potential for tropical
 954 transition over the south atlantic ocean. *Journal of Geophysical Research:*
 955 *Atmospheres* 118(19): 10–870. <https://doi.org/10.1002/jgrd.50830> .
- 956 Dias Pinto, J.R. and R.P. Rocha. 2011. The energy cycle and structural evolu-
 957 tion of cyclones over southeastern south america in three case studies. *Journal of*
 958 *Geophysical Research: Atmospheres* 116(D14) .

- 959 Dutton, J.A. and D.R. Johnson. 1967. The theory of available potential energy and a
960 variational approach to atmospheric energetics. *Advances in geophysics* 12: 333–436
961 .
- 962 Evans, J.L. and A. Braun. 2012. A climatology of subtropical cyclones in the south
963 atlantic. *Journal of Climate* 25(21): 7328–7340 .
- 964 Federer, M., L. Papritz, M. Sprenger, and C.M. Grams. 2025. Synoptic perspective on
965 the conversion and maintenance of local available potential energy in extratropical
966 cyclones. *Weather and Climate Dynamics* 6(1): 211–230 .
- 967 Federer, M., L. Papritz, M. Sprenger, C.M. Grams, and M. Wenta. 2024. On the local
968 available potential energy perspective of baroclinic wave development. *Journal of
969 the Atmospheric Sciences* 81(5): 871–886 .
- 970 Fukuoka, A. 1951. A study of 10-day forecast (a synthetic report). *The Geophysical
971 Magazine* XXII: 177–218 .
- 972 Govekar, P.D., C. Jakob, M.J. Reeder, and J. Haynes. 2011. The three-dimensional dis-
973 tribution of clouds around southern hemisphere extratropical cyclones. *Geophysical
974 Research Letters* 38(21). <https://doi.org/10.1029/2011GL049091> .
- 975 Gozzo, L.F., R.P. da Rocha, M.S. Reboita, and S. Sugahara. 2014. Subtropical
976 cyclones over the southwestern south atlantic: Climatological aspects and case
977 study. *Journal of Climate* 27(22): 8543–8562 .
- 978 Gramscianinov, C., R. Campos, R. De Camargo, K. Hodges, C.G. Soares, and
979 P. da Silva Dias. 2020. Analysis of atlantic extratropical storm tracks characteristics
980 in 41 years of era5 and cfsr/cfsv2 databases. *Ocean Engineering* 216: 108–111 .
- 981 Gramscianinov, C., K. Hodges, and R.d. Camargo. 2019. The properties and genesis
982 environments of south atlantic cyclones. *Climate Dynamics* 53: 4115–4140 .
- 983 Gramscianinov, C.B., R.M. Campos, R. de Camargo, K.I. Hodges, C. Guedes Soares,
984 and P.L. da Silva Dias. 2020. Atlantic extratropical cyclone tracks in 41 years of
985 ERA5 and CFSR/CFSv2 databases. *Mendeley Data* V4. [https://doi.org/10.17632/
986 kwcvfr52hp.4](https://doi.org/10.17632/kwcvfr52hp.4) .
- 987 Gramscianinov, C.B., R. de Camargo, R.M. Campos, C. Guedes Soares, and P.L.
988 da Silva Dias. 2023. Impact of extratropical cyclone intensity and speed on the
989 extreme wave trends in the atlantic ocean. *Climate dynamics* 60(5-6): 1447–1466 .
- 990 Gutierrez, E.R., P.S. Dias, J. Veiga, R. Camayo, and A. Dos Santos. 2009. Multivariate
991 analysis of the energy cycle of the south american rainy season. *International
992 Journal of Climatology* 29(15): 2256–2269 .

- 993 Hadas, O. and Y. Kaspi. 2025. A lagrangian perspective on the growth of midlatitude
994 storms. *AGU Advances* 6(3): e2024AV001555 .
- 995 Hao, J. and T.K. Ho. 2019. Machine learning made easy: a review of scikit-learn
996 package in python programming language. *Journal of Educational and Behavioral*
997 *Statistics* 44(3): 348–361 .
- 998 Hartigan, J.A. and M.A. Wong. 1979. Algorithm as 136: A k-means clustering algo-
999 rithm. *Journal of the Royal Statistical Society. Series C (Applied Statistics)* 28(1):
1000 100–108 .
- 1001 Hernández-Deckers, D. and J.S. von Storch. 2010. Energetics responses to increases
1002 in greenhouse gas concentration. *Journal of Climate* 23(14): 3874–3887 .
- 1003 Hersbach, H., B. Bell, P. Berrisford, S. Hirahara, A. Horányi, J. Muñoz-Sabater,
1004 J. Nicolas, C. Peubey, R. Radu, D. Schepers, et al. 2020. The era5 global reanalysis.
1005 *Quarterly Journal of the Royal Meteorological Society* 146(730): 1999–2049 .
- 1006 Hodges, K. 1995. Feature tracking on the unit sphere. *Monthly Weather*
1007 *Review* 123(12): 3458–3465 .
- 1008 Hodges, K.I. 1994. A general method for tracking analysis and its application to
1009 meteorological data. *Monthly Weather Review* 122(11): 2573–2586 .
- 1010 Hoskins, B.J. and K.I. Hodges. 2002. New perspectives on the northern hemisphere
1011 winter storm tracks. *Journal of the Atmospheric Sciences* 59(6): 1041–1061. [https://doi.org/10.1175/1520-0469\(2002\)059<1041:NPO TNH>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1041:NPO TNH>2.0.CO;2) .
- 1012
- 1013 Hoskins, B.J. and K.I. Hodges. 2005. A new perspective on southern hemisphere
1014 storm tracks. *Journal of Climate* 18(20): 4108–4129 .
- 1015 Hoskins, B.J. and P.J. Valdes. 1990. On the existence of storm-tracks. *Journal of*
1016 *Atmospheric Sciences* 47(15): 1854–1864 .
- 1017 Hu, Q., Y. Tawaye, and S. Feng. 2004. Variations of the northern hemisphere
1018 atmospheric energetics: 1948–2000. *Journal of Climate* 17(10): 1975–1986 .
- 1019 Jain, D., S.A. Rao, R. Dandi, P.A. Pillai, A. Srivastava, and M. Pradhan. 2023.
1020 Monsoon mission coupled forecast system version 2.0: model description and indian
1021 monsoon simulations. *Geoscientific Model Development Discussions* 2023: 1–29 .
- 1022 Keshtgar, B., A. Voigt, C. Hoose, M. Riemer, and B. Mayer. 2023. Cloud-
1023 radiative impact on the dynamics and predictability of an idealized extratropical
1024 cyclone. *Weather and Climate Dynamics* 4(1): 115–132. <https://doi.org/10.5194/wcd-4-115-2023> .
- 1025
- 1026 Kirshbaum, D., T. Merlis, J. Gyakum, and R. McTaggart-Cowan. 2018. Sensitivity
1027 of idealized moist baroclinic waves to environmental temperature and moisture

- 1028 content. *Journal of the Atmospheric Sciences* 75(1): 337–360 .
- 1029 Lau, N.C. and M.W. Crane. 1997. Comparing satellite and surface observa-
 1030 tions of cloud patterns in synoptic-scale circulation systems. *Monthly weather*
 1031 *review* 125(12): 3172–3189. [https://doi.org/10.1175/1520-0493\(1997\)125\(3172:](https://doi.org/10.1175/1520-0493(1997)125(3172:CSASOO)2.0.CO;2)
 1032 [CSASOO\)2.0.CO;2](https://doi.org/10.1175/1520-0493(1997)125(3172:CSASOO)2.0.CO;2) .
- 1033 Leal, K.B., L.E.d.S. Robaina, T.S. Körting, J.L. Nicolodi, J.D. da Costa, and V.G.
 1034 Souza. 2023. Identification of coastal natural disasters using official databases to
 1035 provide support for the coastal management: the case of santa catarina, brazil.
 1036 *Natural Hazards*: 1–18 .
- 1037 Li, L., A.P. Ingersoll, X. Jiang, D. Feldman, and Y.L. Yung. 2007. Lorenz energy
 1038 cycle of the global atmosphere based on reanalysis datasets. *Geophysical Research*
 1039 *Letters* 34(16) .
- 1040 Liu, Z., C.L. Franzke, L. Novak, R. Tailleux, and V. Lembo. 2024. A systematic
 1041 local view of the long-term changes in the atmospheric energy cycle. *Journal of*
 1042 *Climate* 37(23): 6413–6434 .
- 1043 Lorenz, E. 1967. The nature and theory of the general circulation of the atmosphere.
 1044 *World meteorological organization* 161 .
- 1045 Lorenz, E.N. 1955. Available potential energy and the maintenance of the general
 1046 circulation. *Tellus* 7(2): 157–167 .
- 1047 Lorenz, E.N. 1956. *Empirical orthogonal functions and statistical weather predic-*
 1048 *tion*, Volume 1. Massachusetts Institute of Technology, Department of Meteorology
 1049 Cambridge.
- 1050 MacQueen, J. et al. 1967. Some methods for classification and analysis of multivari-
 1051 ate observations. In *Proceedings of the fifth Berkeley symposium on mathematical*
 1052 *statistics and probability*, Volume 1, pp. 281–297. Oakland, CA, USA.
- 1053 Margules, M. 1903. Über die energie der stürme. *Jahrbuch der Zentralanstalt für*
 1054 *Meteorologie, Wien* 48: Anhang .
- 1055 Marrafon, V.H., M.S. Reboita, R.P. da Rocha, and E.M. de Jesus. 2022. Classificação
 1056 dos tipos de ciclones sobre o oceano atlântico sul em projeções com o regcm4 e
 1057 mcgs. *Revista Brasileira de Climatologia* 30: 1–25 .
- 1058 McErlich, C., A. McDonald, J. Renwick, and A. Schuddeboom. 2023. An assessment
 1059 of extra-tropical cyclone precipitation extremes over the southern hemisphere using
 1060 era5. *Geophysical Research Letters* 50(17): e2023GL104130 .
- 1061 Michaelides, S. 2021. Lorenz atmospheric energy cycle in climatic projections.
 1062 *Climate* 9(12): 180 .

- 1063 Michaelides, S.C. 1987. Limited area energetics of genoa cyclogenesis. *Monthly*
1064 *Weather Review* 115(1): 13–26 .
- 1065 Michaelides, S.C. 1992. A spatial and temporal energetics analysis of a baroclinic
1066 disturbance in the mediterranean. *Monthly weather review* 120(7): 1224–1243 .
- 1067 Michaelides, S.C., N.G. Prezerakos, and H.A. Flocas. 1999. Quasi-lagrangian ener-
1068 getics of an intense mediterranean cyclone. *Quarterly Journal of the Royal*
1069 *Meteorological Society* 125(553): 139–168 .
- 1070 Muench, H.S. 1965. On the dynamics of the wintertime stratosphere circulation.
1071 *Journal of Atmospheric Sciences* 22(4): 349–360 .
- 1072 Novak, L. and R. Tailleux. 2018. On the local view of atmospheric available potential
1073 energy. *Journal of the Atmospheric Sciences* 75(6): 1891–1907 .
- 1074 Okajima, S., H. Nakamura, and Y. Kaspi. 2021. Cyclonic and anticyclonic contribu-
1075 tions to atmospheric energetics. *Scientific reports* 11(1): 13202 .
- 1076 Oort, A.H. 2018. On estimates of the atmospheric energy cycle, *Renewable Energy*,
1077 Vol1.214–Vol1.235. Routledge.
- 1078 Pan, Y., L. Li, X. Jiang, G. Li, W. Zhang, X. Wang, and A.P. Ingersoll. 2017. Earth’s
1079 changing global atmospheric energy cycle in response to climate change. *Nature*
1080 *communications* 8(1): 14367 .
- 1081 Pezza, A.B., L.A. Garde, J.A.P. Veiga, and I. Simmonds. 2014. Large scale features
1082 and energetics of the hybrid subtropical low ‘duck’over the tasman sea. *Climate*
1083 *dynamics* 42: 453–466 .
- 1084 Pezza, A.B., J.A.P. Veiga, I. Simmonds, K. Keay, and M.d.S. Mesquita. 2010. Envi-
1085 ronmental energetics of an exceptional high-latitude storm. *Atmospheric Science*
1086 *Letters* 11(1): 39–45 .
- 1087 Rantanen, M., J. Räisänen, V.A. Sinclair, and H. Järvinen. 2019. Sensitivity
1088 of idealised baroclinic waves to mean atmospheric temperature and meridional
1089 temperature gradient changes. *Climate Dynamics* 52: 2703–2719 .
- 1090 Read, P., D. Kennedy, N. Lewis, H. Scolan, F. Tabataba-Vakili, Y. Wang, S. Wright,
1091 and R. Young. 2020. Baroclinic and barotropic instabilities in planetary atmo-
1092 spheres: energetics, equilibration and adjustment. *Nonlinear Processes in Geo-*
1093 *physics* 27(2): 147–173 .
- 1094 Reboita, M.S., R.P. Da Rocha, T. Ambrizzi, and S. Sugahara. 2010. South atlantic
1095 ocean cyclogenesis climatology simulated by regional climate model (regcm3).
1096 *Climate Dynamics* 35: 1331–1347 .

- 1097 Reboita, M.S., R.P. da Rocha, M.R. de Souza, and M. Llopart. 2018. Extratropi-
 1098 cal cyclones over the southwestern south atlantic ocean: Hadgem2-es and regcm4
 1099 projections. *International Journal of Climatology* 38(6): 2866–2879 .
- 1100 Reboita, M.S., M.A. Gan, R. Da Rocha, T. Ambrizzi, et al. 2010. Precipitation regimes
 1101 in south america: a bibliography review. *Revista Brasileira de Meteorologia* 25(2):
 1102 185–204 .
- 1103 Rudeva, I. and S.K. Gulev. 2007. Climatology of cyclone size characteristics and their
 1104 changes during the cyclone life cycle. *Monthly Weather Review* 135(7): 2568–2587 .
- 1105 Savita, A., J. Kjellsson, R.P. Kedzierski, M. Latif, T. Rahm, S. Wahl, and W. Park.
 1106 2023. Assessment of climate biases in openifs version 43r3 across model horizontal
 1107 resolutions and time steps. *Geoscientific Model Development Discussions* 2023:
 1108 1–25 .
- 1109 Sinclair, M.R. 1994. An objective cyclone climatology for the southern hemi-
 1110 sphere. *Monthly Weather Review* 122(10): 2239–2256. [https://doi.org/10.1175/
 1111 1520-0493\(1994\)122<2239:AOCCT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1994)122<2239:AOCCT>2.0.CO;2) .
- 1112 Sinclair, M.R. 1995. A climatology of cyclogenesis for the southern hemisphere.
 1113 *Monthly Weather Review* 123(6): 1601–1619 .
- 1114 Smith, P.J. 1980. The energetics of extratropical cyclones. *Reviews of Geo-*
 1115 *physics* 18(2): 378–386 .
- 1116 Swart, N.C., J.N. Cole, V.V. Kharin, M. Lazare, J.F. Scinocca, N.P. Gillett, J. Anstey,
 1117 V. Arora, J.R. Christian, S. Hanna, et al. 2019. The canadian earth system model
 1118 version 5 (canesm5. 0.3). *Geoscientific Model Development* 12(11): 4823–4873 .
- 1119 Tecchio, R., D.C. de Souza, M.B.L. da Silva, M.C. de Oliveria Costa, R. de Camargo,
 1120 and J. Harari. 2024. Mean sea level, tidal components and surges in guanabara bay
 1121 (rio de janeiro) from 1990 to 2021. *International Journal of Climatology* .
- 1122 Trigo, I.F. 2006. Climatology and interannual variability of storm-tracks in the euro-
 1123 atlantic sector: a comparison between era-40 and ncep/ncar reanalyses. *Climate*
 1124 *Dynamics* 26(2): 127–143 .
- 1125 Veiga, J.A.P. and T. Ambrizzi. 2013. A global and hemispherical analysis of the
 1126 lorenz energetics based on the representative concentration pathways used in cmip5.
 1127 *Advances in Meteorology* 2013(1): 485047 .
- 1128 Veiga, J.A.P., A.B. Pezza, I. Simmonds, and P.L. Silva Dias. 2008. An analysis of the
 1129 environmental energetics associated with the transition of the first south atlantic
 1130 hurricane. *Geophysical Research Letters* 35(15) .

- 1131 Vincent, D. and L. Chang. 1973. Some further considerations concerning energy
1132 budgets of moving systems. *Tellus* 25(3): 224–232 .
- 1133 Wahab, M.A., H.A. Basset, and A. Lasheen. 2002. On the mechanism of winter cyclo-
1134 genesis in relation to vertical axis tilt. *Meteorology and Atmospheric Physics* 81(1):
1135 103–127 .
- 1136 Wiin-Nielsen, A. 1968. On the intensity of the general circulation of the atmosphere.
1137 *Reviews of Geophysics* 6(4): 559–579 .
- 1138 Young, R. 2014. The lorenz energy cycle in simulated rotating annulus flows. *Physics*
1139 *of Fluids* 26(5) .
- 1140 Zheng, Z. 2021. zzheng93/pyeof: First release (v0.0.0).

1141 Supplementary information




Fig. S1 Statistics for the cyclones classified as EOF(+) (red) and EOF(-) (blue). (A) Cyclone counts; (B) maximum intensity, expressed by the *minimum* central relative vorticity at 850 hPa; (C) total life-cycle duration (days); and (D) mean translational speed (km h⁻¹).




Fig. S2 Boxplot of the total life cycle duration (in days) for the four clusters (1–4) identified among the most intense cyclones in the dataset.




Fig. S3 Stacked time series of the first four principal components (PC1–PC4) for each intense cyclone ($\zeta_{850, \text{max}} > q_{99}$). Positive values are plotted upward and negative values downward, so the coloured bars represent the signed contribution of each EOF mode to the energetics of every system. The x -axis lists the individual cyclones (sorted arbitrarily), while the y -axis gives the PC amplitude (dimensionless).




Fig. S4 Median barotropic-conversion term (C_k) at the start (blue) and end (red) of the decay phase, with the error bars representing the inter-quartile range.