

GEOMETRIC INVARIANT THEORY

Minicourse by Michel Brion (Université Grenoble Alpes, Institut Fourier, France),
CIMPA school Florianópolis 2025.

Notes at github.com/danimalabares/cimpa-floripa

Abstract. Geometric Invariant Theory (GIT) studies the actions of reductive algebraic groups on algebraic varieties and the constructions of quotients in algebraic geometry. One of the most relevant aspect of GIT is its application to moduli theory: we can indeed approach moduli problems trying to construct moduli spaces as GIT quotients. In this course we will present the basic notions of geometric invariant theory, we will illustrate its connections to moduli and provide concrete examples of moduli spaces obtained as GIT quotients.

CONTENTS

1. Introduction	1
2. Local structure of quotients	2
3. Linear algebraic groups	3
4. Quotients by linearly reductive groups	5
5. Hilbert-Mumford criterion	8
6. Hyperelliptic curves	9
7. Local structure of quotients	10
References	10

Plan.

- (1) Quotients by finite groups.
- (2) Linear algebraic groups.
- (3) GIT (affine case).
- (4) GIT (general case).

1. INTRODUCTION

Let G be a finite group of order N , $G \subset \text{Aut}(X)$, where X is an affine variety over a field $k = \bar{k}$ of arbitrary characteristic $p \geq 0$. G acts on $\Gamma(X, \mathcal{O}_X) = \mathcal{O}(X)$ by

$$(gf)(x) = f(g^{-1}x).$$

Define

$$\mathcal{O}(X)^G := \{f | gf = f \forall g \in G\} \subset \mathcal{O}(X)$$

Proposition 1.1. $\mathcal{O}(X)^G$ is finitely generated and $\mathcal{O}(X)$ is a finite module over $\mathcal{O}(X)^G$.

Proof. $A = \mathcal{O}(X) \supset \mathcal{O}(X)^G = B$, $\mathcal{O}(X) = k[f_1, \dots, f_n]$, $p_1(f) = \prod_{g \in G} (t - gf) \in A^G[t] = B[t]$. \square

Let $Y = \text{Spec} B$ and $\pi : X \rightarrow Y$ a dominant, finite morphism, which implies it is surjective.

Proposition 1.2. *The set-theoretic fibers of π are the orbits Gx*

Proof. π is G -invariant, i.e. constant on orbits. \square

Proposition 1.3. *The quotient map π is open and if $U \subset X$ is open, affine, G -stable, then the restriction $\pi|_U$ is the quotient map $U \rightarrow U/G$.*

Proposition 1.4. *$f : X \rightarrow Z$ G -invariant morphism, then there is a unique morphism $\varphi : Y \rightarrow Z$ such that $f = \varphi \circ \pi$. This says that π is the categorical quotient.*

Sketch of proof. It is clear that φ exists as a map of sets. It is continuous since π is open surjective. $\varphi^\sharp : \mathcal{O}_Z \rightarrow \varphi_*(\mathcal{O}_Y)$. \square

Example 1.5. (1) $G = \mu_n$ the roots of unity of order n (prime to p). G acts on \mathbb{A}^2 by $g(x, y) = (gx, gy)$. $\mathcal{O}(\mathbb{A}^2) = k[x, y]$ and $\mathcal{O}(\mathbb{A}^2)^G = k[x^n, x^{n-1}y, \dots, xy^{n-1}, y^n]$. \mathbb{A}^2/G is a singular surface.
(2) $G = S_n$ symmetric group, acts on \mathbb{A}^n permuting the coordinates. $\mathcal{O}(\mathbb{A}^n) = k[x_1, \dots, x_n]$, $\mathcal{O}(\mathbb{A}^n)^G = k[e_1, \dots, e_n]$ where e_i is the i -th symmetric function. The quotient $\pi : \mathbb{A}^n \xrightarrow{(e_1, \dots, e_n)} \mathbb{A}^n$ is finite, flat of degree $n!$.
(3) (Symmetric products.) $X = Z^n$, Z affine algebraic variety, $G = S_n$ acting by permuting copies of Z . Then $X/G := Z^{(n)}$, the n -th symmetric product of Z . Points of $Z^{(n)}$ = effective 0-cycles in Z of degree $n = z_1 + \dots + z_n$, ($z_i \in Z$) = $n_1 z_1 + \dots + n_k z_k$ where z_i are distinct and n_i are positive integers with some n .

2. LOCAL STRUCTURE OF QUOTIENTS

See [Mum70].

Definition 2.1. Points with trivial stabilizer.

The local structure of the symmetric product variety $Z^{(n)}$ at the point x is the stabilizer

$$G_x = S_{n_1} \times \dots \times S_{n_k} \subset S_n = G$$

and we get

$$Z^{(n_1)} \times \dots \times Z^{(n_k)} = X/G_x \rightarrow Z^{(n)} = X/G$$

which is étale at the image of x .

Proposition 2.2. *If Z is smooth then $Z^{(n)}$ is smooth.*

Proof. Using power series rings. \square

Proposition 2.3. *If $\dim Z \geq 2$ then $Z^{(n)}$ is singular.*

Proof. Done in lecture. \square

Then we have the *Hilbert scheme characteristic morphism*

$$\begin{aligned}\gamma : \text{Hilb}_n(Z) &\longrightarrow Z^{(n)} \\ W \subset Z &\longmapsto \sum_{z \in Z} \dim(\mathcal{O}_{W,z} z)\end{aligned}$$

where W must be **finite of length** n . See Binger, Linear determinants (older) or J. Bertin, The punctual Hilbert Scheme.

If Z is smooth of dimension 1 then γ is an isomorphism. If Z is of dimension 2 then γ is a resolution. In general γ is an isomorphism above $Z_{f_2}^{(n)}$.

3. LINEAR ALGEBRAIC GROUPS

Definition 3.1. An *algebraic group* is a variety G equipped with morphisms

$$\begin{aligned}m : G \times G &\longrightarrow G \\ (g, h) &\longmapsto gh\end{aligned}$$

and

$$\begin{aligned}i : G &\longrightarrow G \\ g &\longmapsto g^{-1}\end{aligned}$$

with an element $e \in G$ which satisfies the group axioms.

(In fact this is the same as a group scheme of finite type over k .)

Example 3.2. (1) Finite groups.

- (2) \mathbb{G}_a = additive group = $(\mathbb{A}^1, +)$ and \mathbb{G}_m = multiplicative group = $(\mathbb{A}^1 \setminus \{0\}, \times)$.
- (3) GL_n = general linear group. This is a principal open subset ($\det \neq 0$), an affine variety.
- (4) Classical groups $\text{SL}_n, \text{O}_n, \text{Sp}_{2n}$ closed in GL_n . Also $\text{PGL}_n = \text{GL}_n / \mathbb{G}_m : I_n \hookrightarrow \text{GL}_{n^2}$.
- (5) $(E, 0)$ elliptic curve; there is a unique algebraic group law $+$ with neutral element 0. (Abelian varieties.)

Proposition 3.3. G algebraic group, G^0 connected component of e : G^0 is a closed normal subgroup of G and the connected components of G are the gG^0 for $g \in G$. Moreover, G^0 is irreducible and G is smooth (which is obvious because it is smooth somewhere and we may just translate).

$$1 \longrightarrow G^0 \longrightarrow G \longrightarrow \pi_0(G) \longrightarrow 1$$

Definition 3.4. An *action* of an algebraic group G on a variety X is a morphism $a : G \times X \rightarrow X$ which satisfies the axioms of an action:

$$a(g, x) = g \cdot x, \quad g \cdot (h \cdot x) = gh \cdot x, \quad e \cdot x = x$$

The *orbit* of $x \in X$ is $G \cdot x = \{g \cdot x | g \in G\}$. The *stabilizer* G_x is $\{g \in G | g \cdot x = x\}$, which is a closed subgroup of G .

Proposition 3.5. (1) Each orbit $G \cdot x$ is locally closed, smooth of dimension $\dim(G) - \dim(G_x)$.

- (2) The orbit map $G \rightarrow G \cdot x, g \mapsto g \cdot x$ is faithfully flat.

- (3) $\overline{G \cdot x} \setminus G \cdot x$ is a union of orbits of smaller dimension.
- (4) Closed orbits exist.

Proof. Should be easy. For example, to see it's faithfully flat you just notice it is flat somewhere (if it is connected, is it connected?) and then use homogeneity again. \square

Proposition 3.6. *For any $n \geq 1$, the set $\{x \in X \mid \dim(G \cdot x) \geq n\}$ is open. That is, $\{x \in X \mid \dim(G_x) \leq n\}$ is open.*

In particular, the points with finite stabilizer form an open subset.

Proof. Use

$$\begin{aligned} \gamma : G \times X &\longrightarrow X \times X \\ (g, x) &\longmapsto (g \cdot x, x) \end{aligned}$$

\square

Lemma 3.7. *Let $f : G \rightarrow H$ be an homomorphism of algebraic groups. Then $\text{Ker}(f)$ is a normal subgroup of G . $\text{Im}(f)$ is a closed subgroup of H of dimension $\dim(G) - \dim \text{Ker } f$.*

Example 3.8. $G = \text{SL}_2$ acts on $k[x, y]_n := V_n$ (these are called the *binary forms of degree n*) by linear change of variables.

Definition 3.9. An algebraic group is *linear* if $G \hookrightarrow \text{GL}_n$ for some n .

Example 3.10. $\mathbb{G}_a = \left\{ \begin{pmatrix} 1 & t \\ 0 & q \end{pmatrix} \right\}$ is linear.

Proposition 3.11. *G linear algebraic group, X a G -variety, G acts on $\mathcal{O}(X) = \Gamma(X, \mathcal{O}_X)$ via $(g \cdot f)(x) = f(g^{-1} \cdot x)$ then $\mathcal{O}(X)$ is a union of f.d. G -stable subspaces on which G acts algebraically.*

Proof. From the action $a : G \times X \rightarrow X$ we get the *co-action*

$$\begin{aligned} a^\# : \mathcal{O}(X) &\longrightarrow \mathcal{O}(G \times X) \\ f &\longmapsto \sum_{i=1}^n (\varphi_i \otimes \psi_i) \end{aligned}$$

\square

Definition 3.12. A finite dimensional vector space V is called a *G -module* if V is equipped with an action of G via a homomorphism of algebraic groups $G \rightarrow \text{GL}(V)$.

A *G -module* is a vector space V equipped with a linear action of the group G such that $V = \text{union of finite dimensional } G\text{-submodules}$.

Proposition 3.13. (1) *X affine G -variety, G linear algebraic group, then $X \hookrightarrow V$ finite dimensional G -module as a closed G -stable subvariety.*
 (2) *Every affine algebraic group is linear.*

Let X be an affine G -variety. Then

$$\begin{aligned} \mathcal{O}(X) &\longrightarrow \mathcal{O}(G \times X) = \mathcal{O}(X)[t, t^{-1}] \\ f &\longmapsto \sum_{n \in \mathbb{Z}} f_n t^n \end{aligned}$$

In particular, every \mathbb{G}_m -module is semisimple: every nonzero G -module contains a nonzero fixed point. Indeed, let V be a finite-dimensional G -module. Let $v \in V$ be a \mathbb{G}_a -eigenvector. Then $g \cdot v = \gamma(g)v$ for $X \in \mathcal{O}(\mathbb{G}_a)^\times = k^\times$ taking $g = 0$ yields $X = 1$.

G connected algebraic group, then G is *unipotent* if it is an iterated extension of copies of \mathbb{G}_a . Then every nonzero G -module contains a nonzero fixed point.

As an example consider $G = \mathrm{SL}_2$. (We saw that it is not linearly reductive in the sense described in the next section.)

4. QUOTIENTS BY LINEARLY REDUCTIVE GROUPS

Definition 4.1. A linear algebraic group G is *linearly reductive* if every G -module is semi-simple.

Exercise 4.2. G is linearly reductive if and only if the functor $(G\text{-modules}) \rightarrow (\text{vector spaces}), V \mapsto V^G$ is exact.

Theorem 4.3. *If G is linearly reductive then every connected normal unipotent subgroup U of G is trivial. The converse is true in characteristic zero.*

Proof. (\implies) Let V be a G -module, $V \neq 0$ and U as in the statement. Then $V^U \neq 0$ is a G -module. Let W be a G -module. Then $V = V^U \oplus W$. [Missing]

(\impliedby) We show this for $k = \mathbb{C}$. View $G \supset K$ compact subgroup, Zariski dense (eg. $C^* \supset S^1$ or $\mathrm{GL}_n(\mathbb{C}) \supset \mathrm{U}_n$.) [Missing] \square

Definition 4.4. A linear algebraic group G is *reductive* if it has a nontrivial connected unipotent normal subgroup.

The following definition is from StacksProject, `brauer.tex`:

Definition 4.5. Let A be a k -algebra. We say an A -module M is *simple* if it is nonzero and the only A -submodules are 0 and M . We say A is *simple* if the only two-sided ideals of A are 0 and A .

Lemma 4.6. *G linearly reductive, V a G -module. Then*

- (1) *There is a unique projection of G -modules $R_V : V \rightarrow V^G$ called the Reynolds operator.*
- (2)

$$\begin{array}{ccc} V & \xrightarrow{f} & W \\ R_V \downarrow & & \downarrow R_V \\ V^G & \xrightarrow{f^G} & W^G \end{array}$$

commutes.

- (3) *A a G -algebra, $B := A^G$. Then $R_A(ab) = bR_A(a)$ for all $a \in A, b \in B$.*

Proof. (1) There is a unique decomposition $V = V^G \oplus V'$ where V' is the sum of nontrivial simple G -submodules.

- (2) Implied by the decomposition $V = V^G \oplus V'$.

- (3)

$$\begin{array}{ccc} V & \xrightarrow{b} & W \\ R_A \downarrow & & \downarrow R_A \\ V^G & \xrightarrow{b} & W^G \end{array}$$

commutes. □

Theorem 4.7 (Hilbert, Nagata). *If G is a linearly reductive group acting on an affine variety X , then $\mathcal{O}(X)^G$ is finitely generated.*

Proof. Done in class, uses R_A and graded Nakayama lemma. □

$\mathcal{O}(X)^G \subset \mathcal{O}(X)$ yields $\pi : X \rightarrow Y$, a G -invariant morphism of affine varieties, giving a good quotient.

Proposition 4.8. (1) π is regular.

(2) If $Z \subset X$ is closed and G -stable, then $\pi|_Z$ is the good quotient of Z .

(3) If $Z, Z' \subset X$ are closed G -stable and disjoint, then $\pi(Z), \pi(Z')$ are disjoint.

(4) Every fiber of π contains a unique closed G -orbit.

But if $U \subset X$ is open affine G -stable, then $\pi|_U$ is not the good quotient.

Example 4.9. $G = \mathbb{G}_n$ acts on \mathbb{A}^n by scalar, i.e. $g(x_1, \dots, x_n) = (gx_1, \dots, gx_n)$. Then $\mathcal{O}(\mathbb{A}^n)^G = k$, $\pi : \mathbb{A}^n \rightarrow \bullet$, but $f_n U = D(f)$, where f is a nonzero linear form,

$$\mathcal{O}(U)^{\mathbb{G}_m} = \bigcup_{d \geq 0} \frac{k[x_1, \dots, x_n]}{f^d}$$

Notation: $Y = X//G$, the space of closed orbits.

Definition 4.10. A point $x \in X$ is called *stable* if its orbit $G \cdot x \subset X$ is closed and the stabilizer G_x is finite.

This definition is different from that in Mumford's book on GIT (our definition there is called *properly* stable point.)

Proposition 4.11. *The set of stable points X^s is open, G -stable and $\pi(X^s) = Y^s$ is open. Moreover, $X^s = \pi^{-1}(Y^s)$ and $X^s \rightarrow Y^s$ is a geometric quotient.*

Warning: there might not be any stable points at all! (But it looks like there are some ways to get around this.)

Proof. The first trick very elementary; just playing with orbits. Let $x_0 \in X$ be stable. Then

$$G \cdot x_0 \subset \pi^{-1}(\pi(x_0)) = \{x \in X : \overline{G \cdot x} \supset G \cdot x\} = G \cdot x_0$$

since $\dim(G \cdot x_0) = \dim G$.

Then, roughly, remove higher dimensional orbits, and since the remaining orbits have all the same dimension, then they are all closed.

The final assertion $X^s \rightarrow Y^s$ is similar to what we did. No deep arguments. □

Example 4.12. $G = \mathrm{SL}_n$ acts on $X = k[x_1, \dots, x_n]_d$ by linear change of variables. Let's consider some values. If $d = 1$, then the quotient is $X//G = \mathrm{pt}$. For $d = 2$, the quadratic forms, we have the discriminant Δ as an invariant, and in fact $\mathcal{O}(\Delta) = k[\Delta]$ (**A nice exercise!**). Actually it looks like $\pi = \Delta : X \rightarrow \mathbb{A}^1$, and the stabilizer of f is $\mathrm{SO}(f)$, no stable points.

For $d \geq 3$, and $n \geq 5$, then $\mathcal{O}(X)^G$ is not known! But we also have the discriminant $\Delta \in \mathcal{O}(X)^G$ and in fact (Jordan) if $\Delta(f) \neq 0$ then f is stable. Let us prove this fact (we don't know how did Jordan prove this): we show that G_f is finite and

stable, i.e. $\text{Lie}(G_f) = 0$. The Lie algebra of SL_2 , the traceless matrices, acts on X by

$$A \cdot f = \sum_{i=1}^n \ell_i \frac{\partial f}{\partial x_i}$$

where $\ell_i = \sum_{j=1}^n a_{ij} x_j$. Then $A \cdot f = 0$. Now $\Delta(f) \neq 0$ iff $\frac{\partial f}{\partial x_j}$ have no nontrivial common zero, iff, $\frac{\partial f}{\partial x_j}$ form a regular sequence. This says that the ring Cohen-Macaulay. Then

$$\ell_i \frac{\partial f}{\partial x_i} = \sum_{j \neq i} \ell_j \frac{\partial f}{\partial x_j}$$

which in turn implies that

$$\ell_i \in \left(\frac{\partial f}{\partial x_j}, j \neq i \right) \iff \ell_i = 0 \forall i$$

... which says *what?*

Let X be a projective G -variety, G linearly reductive, $L \rightarrow X$ an ample line bundle

$$R(X, L) = \bigoplus_{n=0}^{\infty} \Gamma(X, L^{\otimes n})$$

which is a graded, finitely generated algebra, then $X = \text{Proj} R(X, L)$. If G acts on $R(X, L)$ “compatibly”, then $R(X, L)^G$ is finitely generated:

$$\text{Proj} R(X, L)^G = X // G$$

and we get a birational map

$$\begin{array}{ccc} X & \dashrightarrow & X // G \\ \uparrow & \nearrow \pi & \\ U & & \end{array}$$

where U is the domain of definition.

If G is finite, then $X = \mathbb{P}(V)$, V finite dimensional G -module. $L = \mathcal{O}_X(1)$: G acts on L and hence on $R(X, L)$ (“I’ll explain details soon”)

We have a finite ring extension $R(X, L)^G = R(X, L)$, and we get

$$X = \text{Proj} R(X, L) \rightarrow \text{Proj} R(X, L)^G = Y$$

Now let G be an algebraic group, X a G -variety, $f : L \rightarrow X$ a line bundle.

Definition 4.13. A G -linearization of L is a G -action on L such that f is equivariant and $g : L_x \rightarrow L_{g \cdot x}$ is linear for all $g \in G, x \in X$. That is, the G -action on L commutes with the \mathbb{G}_m -action by multiplication on fibers.

Then we get an action on the section ring: G acts on $L^\times = L \setminus \{\text{zero section}\}$ and hence $\mathcal{O}(L^\times)$ is a G -module. As we said before, $\mathcal{O}(L^\times) = \bigoplus_{n \in \mathbb{Z}} \Gamma(X, L^{\otimes n})$ is a graded module, and the grading corresponds to the G -action! That is, this is a G -module structure on $\Gamma(X, L^{\otimes n})$. And $R(X, L)$ is a graded G -algebra.

Definition 4.14. X a G -variety, L a G -linearized line bundle on X and $x \in X$. We say x is *semi-stable* w.r.t. L if there exists $n \geq 1$ and $\sigma \in \Gamma(X, L^{\otimes n})^G$ and that $x \in X_\sigma$ and X_σ is affine.

x is *stable* if in addition $G \cdot x$ is closed in X_σ and G_x is finite.

Theorem 4.15 (Mumford). *The set of semis-stable points $X^{ss}(L) \subset X$ is open, G -stable and has a good quotient $\pi : X^{ss}(L) \rightarrow Y(L)$. (Notice everything depends on the line bundle.) The set of stable points $X^s(L)$ is open G -stable and equal to $\pi^{-1}\pi(X^s(L))$ and $\pi|_{X^s(L)}$ is a geometric quotient.*

Proposition 4.16. *Let $x \in \mathbb{P}(V)$ and $x \in [v]$.*

- (1) $x \in \mathbb{P}(V)^{ss} \iff 0 \notin \overline{G \cdot v}$.
- (2) $x \in \mathbb{P}(V)^s \iff v \in V^s$ i.e. $G \cdot v$ is closed in V and G_v is finite.

Proof. Done in class. □

The following is our first example of moduli space.

Example 4.17 (Smooth hypersurfaces of degree d in \mathbb{P}^n). The isomorphism classes of such surfaces are given by: take degree d polynomials, projectivize, take away diagonal (this gives an open set) and finally quotient by SL_{n+1} :

$$\begin{aligned} \{\text{isom. classes}\} &= \mathbb{P}(k[x_0, \dots, x_n]_d)_\Delta / \mathrm{SL}_{n+1} \xrightarrow{\text{open}} \\ &\mathbb{P}(\cdot)^1 / \mathrm{SL}_{n+1} \hookrightarrow \mathbb{P}(\cdot)^n // \mathrm{SL}_{n+1} \end{aligned}$$

a projective variety, normal, very singular.

Remark 4.18. $G \curvearrowright X$ categorical quotient $\pi : X \rightarrow Y$, then π yields a coarse moduli space for

$$\begin{array}{ccc} X & \longrightarrow & [X/G] \\ & \searrow & \vdots \\ & & Y \end{array}$$

Proof. $\mathrm{Hom}(S, [X/G])$ is a category, not a set as usual. Its objects are

$$\begin{array}{ccc} & & G\text{-equivariant} \\ & & \text{morphism } b \\ P & \xrightarrow{\quad} & X \\ \text{principal} \downarrow & & \\ \mathbb{G}\text{-bundle} & & \\ S & & \end{array}$$

and morphisms are

$$\begin{array}{ccccc} & & X & & \\ & \nearrow b' & & \nwarrow b & \\ P & & & & P \\ & \xrightarrow{\varphi} & & & \\ & \searrow & & \swarrow & \\ & & S & & \end{array}$$

(continues...)

□

5. HILBERT-MUMFORD CRITERION

Definition 5.1. A *one-parameter subgroup* of G is a homomorphism $\lambda : \mathbb{G}_m \rightarrow G$. If $G \curvearrowright X$ and $x \in X$, we say that

$$\lim_{t \rightarrow 0} \lambda(t)$$

exists and equals $y \in X$ if $\mathbb{A}^1 \setminus \{0\} = \mathbb{G}_m \rightarrow X$, $t \mapsto \lambda(t) \cdot x$ extends to $\mathbb{A}^1 \rightarrow X$, $0 \mapsto y$.

Theorem 5.2 (Hilbert-Mumford criterion). *V finite-dimensional G -module, G linearly reductive, $v \in V$, $\overline{G \cdot v} \supset G \cdot v$ closed orbit. There exists a one-parameter subgroup λ such that $\lim_{t \rightarrow 0} \lambda(t) \cdot v \in G \cdot v$.*

We will not prove this theorem. Here are some corollaries.

Lemma 5.3. $v \notin V^{ss} \iff \exists \lambda$ such that $\lim_{t \rightarrow 0} \lambda(t) \cdot v = 0$. Then v is called unstable and

$$\{\text{unstable points}\} = \text{nilcone } \mathcal{N}(V) = V.$$

Lemma 5.4. $G \cdot v$ is closed $\iff \lim_{t \rightarrow 0} \lambda(t) \cdot v$ exists, then it lies in $G \cdot v$.

Proposition 5.5. $G \cdot v$ is affine $\iff G_v$ is linearly reductive.

Proof. (\implies) $G_v = H \iff G \cdot vG/H = G//H$.

(\impliedby) We show that the functor

$$\begin{aligned} (\text{left } H\text{-modules}) &\longrightarrow (\text{f.d. vector spaces}) \\ V &\longmapsto V^H \end{aligned}$$

(continues...)

□

We can characterize stable points using this proposition.

Lemma 5.6. $v \in V^s \iff G \cdot v$ is closed and v is fixed by no nontrivial one-parameter subgroup.

Proof. The reason is that if $G_v = H$ is linearly reductive then H^0 is generated by images of a one-parameter subgroup. □

Example 5.7 (Stable points for SL_2 acting on degree d polynomials). If $G = SL_2$ in characteristic zero, then

$$\lambda(t) = \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}$$

every nontrivial one-parameter subgroup of G is conjugate to $\lambda(t)^n$ for a unique $n \geq 1$. $V = V_d = k[x, y]_d \ni x^e y^{d-e}$.

$$\lambda(t)x^e y^{d-e} = t^{d-2e} x^e y^{d-e}$$

$f \in V$ is unstable $\iff \exists g \in G$ such that $gb \in \langle x^{d/2-t} y^{d/2+1}, \dots, y^d \rangle$ if dimension is even, and $\langle x^{(d-1)/2} y^{(d+1)/2}, \dots, y^d \rangle$ if dimension is odd, $\iff f$ has no root of multiplicity $d/2$. $f \notin V^s \iff f$ has no root of multiplicity $\geq d/2$.

If d is odd then $V^s = V^{ss}$. If d is even then $x^{d/2} y^{d/2}$ has a closed orbit and is fixed by $\text{textIm}(\lambda)$.

6. HYPERELLIPTIC CURVES

Let $C \rightarrow \mathbb{P}^1$ be a smooth projective curve of degree 2, genus g , branched in $2g+2$ distinct points, uniquely determined by the branch divisor.

{isomorphism classes of hyperelliptic curves of genus g }

$$= \mathbb{P}(V_{2g+2})_{\Delta}/SL_2 \hookrightarrow \mathbb{P}(V_{2g+2})^{ss}/SL_2$$

projective normal of dimension $2g-3$.

7. LOCAL STRUCTURE OF QUOTIENTS

X affine G -variety, G reductive, $x \in X$ such that $G \cdot x$ is closed, $H = G_x$ (reductive).

Theorem 7.1. *There exists an affine H -stable (locally ?) subvariety $S \subset X$ such that the sequence*

$$(g, s)H \longmapsto g \cdot s$$

$$\begin{array}{ccc} (G \times S)/H & \longrightarrow & X \\ \downarrow & & \downarrow \\ S//H & \longrightarrow & X//G \end{array}$$

$$x \longmapsto \bar{x}$$

$h(g, s) = (gh^{-1}, hs)$ is cartesian with étale horizontal arrows. If $x \in X$ is smooth, then there is an H -equivariant morphism.

$$\begin{aligned} S &\longrightarrow T_x(X)/T_x(G \cdot x) = N_{G \cdot x/X, x} \\ x &\longmapsto 0 \end{aligned}$$

This finishes the statement of the theorem. The last map may be interpreted as the inverse of the exponential map from differential geometry.

A corollary:

Lemma 7.2. *If $G_x = e$ then $X \rightarrow X//G$ is a principal G -bundle in a G -stable neighbourhood of x .*

Exercise 7.3. G finite, works with $S \subset X$ open.

Exercise 7.4. $G = \mathrm{SL}_2$, $V = V_G$, $x^3y^3 \in V_G$, stabilizer $= \mathbb{G}_m = \mathrm{Im}(\lambda)$. Prove that $V_G//\mathrm{SL}_2$ is singular at the image of x^3y^3 . ($S = \mathbb{G}_m$ -module with weights $-6, -4, 0, 4, 6$.) **Hint.** To check for non-smoothness you can check that $\mathcal{O}(S)^{\mathbb{G}_m}$ is not a polynomial ring; this is equivalent to non-smoothness for the quotient, and it should be easy to find: you just find that there are too many generators.

REFERENCES

- [Mum70] David Mumford, *Abelian varieties*, Tata Institute of Fundamental Research Studies in Mathematics, vol. 5, Oxford University Press, 1970.