

Lista 3

Problema 1 Seja $f : X \rightarrow Y$ un difeomorfismo entre duas variedades orientadas conexas. Prove que df_x preserva orientação para um ponto $x \in X$ se, e somente se, df_x preserva orientação para todo ponto $x \in X$.

Demonstração. Considere

$$\begin{aligned} D : X &\longrightarrow \mathbb{R} \\ x &\longmapsto \det d_x f \end{aligned}$$

É uma função contínua que nunca pode ser zero. Como é positiva em x , deve ser positiva sempre. \square

Problem 2 Seja X uma variedade orientável. Prove que a orientação induzida em $X \times X$ é independente da orientação de X .

Demonstração. A orientação de $X \times X$ está dada como segue: uma base (β_1, β_2) do espaço tangente $T_{(x,y)}X \times X$ é orientada se β_1 e β_2 são bases orientadas de X .

Agora considere a mesma construção usando $-X$. A base $(\tilde{\beta}_1, \tilde{\beta}_2)$ de $-X \times -X$ é orientada se $\tilde{\beta}_1$ e $\tilde{\beta}_2$ são bases orientadas de $-X$.

Porém, é equivalente que (β_1, β_2) seja orientada em $X \times X$ e que $(\tilde{\beta}_1, \tilde{\beta}_2)$ seja orientada em $-X \times -X$: tanto a transformação que manda $\beta_1 \mapsto \tilde{\beta}_1$ quanto a transformação que manda $\beta_2 \mapsto \tilde{\beta}_2$ tem determinante negativo, de modo que a transformação que manda $(\beta_1, \beta_2) \mapsto (\tilde{\beta}_1, \tilde{\beta}_2)$ tem determinante positivo! \square

Problem 3 Prove que $SO(n)$ é uma variedade orientável e calcule a sua dimensão. Usando teoria da interseção prove que $\chi(SO(n)) = 0$.

Demonstração. First notice that $SO(n)$ is one of the connected components of $O(n)$. Indeed, $SO(n) = \det^{-1}(1)$ for the submersion $\det : O(n) \rightarrow \{\pm 1\}$, making into a codimension-0 submanifold of $O(n)$ since $\dim\{\pm 1\} = 0$. This means that computing the dimension of $SO(n)$ is the same as computing the dimension of $O(n)$.

Now observe that a matrix in $O(n)$ is the same as an orthonormal frame of \mathbb{R}^n : the column vectors of any $A \in O(n)$ unitary and mutually orthogonal since $AA^T = \text{Id}$ says $\sum_k a_{ik}a_{jk} = \delta_{ij}$ for every i, j .

We can compute the dimension of $O(n)$ as follows. Take a vector v_1 in S^{n-1} , then a unitary vector in the orthogonal complement of v_1 , i.e. a vector in S^{n-2} , and so on until we choose either of the two vectors in S^0 . This means that we are choosing points in

$S^{n-1} \times S^{n-2} \times \dots \times S^0$, which gives $\dim O(n) = \sum_{i=0}^{n-1} i$. Gauss could tell at very early age that this number is $\frac{n(n-1)}{2}$.

To orient $SO(n)$ just notice that it acts on itself homogeneously (by orientation-preserving diffeomorphisms). Taking a basis at the identity matrix and moving it around our manifold using this action generates a smooth global choice of local orientations; i.e. a global orientation.

The fact that $\chi(SO(n)) = 0$ is immediate from the fact that its tangent bundle is trivial: there is a nowhere vanishing vector field (the orbit of any nonzero vector), giving the result by Poincaré-Hopf theorem. \square

Problem 4 Seja Σ uma superfície de gênero g . Construa um campo vetorial em Σ com um único zero de índice $2 - 2g$.

Solution. Primeiro note que é suficiente construir um campo vetorial com um único zero, já que pelo teorema do índice de Hopf ele necessariamente vai ter índice $2 - 2g = \chi(\Sigma)$. (Um modo de comprovar que essa é a característica de Euler de Σ é usando o campo gradiente da função altura, que tem dois zeros de índice 2 e $2g$ zeros de índice -1 .)

Para construir um campo vetorial com um único zero usarei ideias de [StackExchange](#).

Sabemos que Σ pode ser vista como um polígono com certas identificações no bordo. Podemos dar uma decomposição simplicial dele simplesmente botando um vértice no centro do polígono e as arestas que ligam os vértices do polígono com o centro. Usando o campo vetorial para triangulações discutido em aula,




Figure 3-22

obtemos um campo vetorial em Σ com uma quantidade finita de zeros.

A ideia é construir um grafo conexo em Σ cujos vértices sejam os zeros do campo, e que não tenha ciclos i.e. uma árvore. Desse jeito, uma vizinhança tubular do grafo é homeomorfa a um disco, e a restrição do campo a esse disco não tem zeros no bordo. Daí conseguirmos perturbar o campo usando à “Adaptação do truque de Alexander”:

Lemma Se v é um campo vetorial suave em D^2 sem zeros no bordo, existe outro campo vetorial suave com um único zero que coincide com v no bordo.

Prova do lema. Como v não tem zeros no bordo (e o nosso v na verdade tem uma quantidade finita de zeros) podemos pegar um $\varepsilon < 1$ tal que os zeros de v estão na bola B_ε . Daí pegamos uma bump function ρ que tenha um único zero em zero e seja constante 1 fora de B_ε .

Então defina

$$w(r, \theta) = \begin{cases} v(r, \theta) & r \geq \varepsilon \\ \rho(r)v(\varepsilon, \theta) & r \leq \varepsilon \end{cases}$$

Para concluir é só dar uma perturbada a $w : D^n \rightarrow TD^2 = \mathbb{R}^2$ para

□

□

Problem 5 Seja A uma matriz de $n \times n$ com coeficientes inteiros e seja $f : \mathbb{R}^2/\mathbb{Z}^2 \rightarrow \mathbb{R}^2/\mathbb{Z}^2$ tal que $f(x) = Ax$. Calcule o grau de f .

Demonstração. To compute the degree of f it's enough to compute the number of preimages of $[0] \in \mathbb{R}^n/\mathbb{Z}^n$. To do this consider the fundamental domain $[0, 1)^n$ and map it with A to \mathbb{R}^n . When we take the quotient, all the points with integer coordinates will be glued together in the class $[0]$. I.e. we are looking for the number of points with integer coordinates in $P := A([0, 1)^n)$.

I claim that this number is the volume of P , i.e. the determinant of A . To see why consider the case of dimension 2. We may assume that the image of e_1 is a vector in the x axis using a rotation (which preserves the volume of P and the amount of integer points inside P). The number of integer points inside P after rotating is simply the product of the lengths of Ae_1 and Ae_2 :

This product is $\det A$

the case of two dimensions. First use an isometry to assume that one of the sides of P is in the x axis. Then consider the map that projects the other side of P to the y axis, mapping P to a rectangle

the area of P is the same as the area of the rectangle we obtain when projecting the side

□

Problem 6 Prove que $\mathbb{R}P^{2n+1}$ é orientável e que $\mathbb{R}P^{2n}$ não é orientável.

Demonstração. First notice that $-\text{Id}$ preserves orientation iff n is odd. This map is a composition of n reflections, one about every axis of $\mathbb{R}^{n+1} \supset S^n$. Each of these reflections is orientation-reversing, and composing a map with an orientation-reversing map reverses orientation by the chain rule.

Now recall that $\mathbb{R}P^n = S^n / -\text{Id}$. Suppose $\mathbb{R}P^n$ is orientable, so that the quotient map is orientation-preserving since it is a submersion: the determinant of its differential is a nowhere-zero continuous function on a connected manifold, so it cannot be positive somewhere and negative elsewhere.

Choose an oriented basis of the tangent space of \mathbb{RP}^n at $[e_1]$. Pull back the basis using quotient map, this produces a basis at each of the preimages, namely e_1 and $-e_1$. These two bases must be in the same orientation of S^n since the quotient map is orientation-preserving and they are mapped to the same basis in the quotient. However, this only happens when $-\text{Id}$ is orientation-preserving. \square