

# Practice exercises on smooth manifolds

A pdf file with the questions may also be found [here](#).

## 1 Remedial topology

### 1.1 Topological spaces

**Definition 1.1** A set of all subsets of  $M$  is denoted  $2^M$ . **Topology** on  $M$  is a collection of subsets  $S \subset 2^M$  called *open subsets*, and satisfying the following conditions:

1. Empty set and  $M$  are open
2. A union of any number of open sets is open
3. An intersection of a finite number of open subsets is open.

A complement of an open set is called *closed*. A set with topology on it is called a *topological space*. An *open neighbourhood* of a point is an open set containing this point.

**Definition 1.2** A map  $\phi : M \rightarrow M'$  of topological spaces is called *continuous* if a preimage of each open set  $U \subset M'$  is open in  $M$ . A bijective continuous map is called a *homeomorphism* if its inverse is also continuous.

**Exercise 1.1** Let  $M$  be a set, and  $S$  a set of all subsets of  $M$ . Prove that  $S$  defines a topology on  $M$ . This topology is called *discrete*. Describe the set of all continuous maps from  $M$  to a given topological space.

*Solution.* Since all sets are open, topology axioms are satisfied by  $S$ . All maps from  $M$  to a given topological space are continuous.  $\square$

**Exercise 1.2** Let  $M$  be a set, and  $S \subset 2^M$  a set of two subsets: empty set and  $M$ . Prove that  $S$  defines a topology on  $M$ . This topology is called *codiscrete*. Describe the set of all continuous maps from  $M$  to a space with discrete topology.

*Solution.* It's trivial that  $S$  satisfies the axioms of topology. A map from  $M$  to a space  $N$  with discrete topology is always continuous: the preimage of  $N$  is always  $M$  by definition of function: every point of  $M$  corresponds to some point of  $N$ . Obviously the preimage of  $\emptyset$  is  $\emptyset$ .  $\square$

**Definition 1.3** Let  $M$  be a topological space, and  $Z \subset M$  its subset. *Open subsets* of  $Z$  are subsets obtained as  $Z \cap U$ , where  $U$  is open in  $M$ . This topology is called *induced topology*.

**Definition 1.4** A *metric space* is a set  $M$  equipped with a *distance function*  $d : M \times M \rightarrow \mathbb{R}^{\geq 0}$  satisfying the following axioms.

1.  $d(x, y) = 0$  iff  $x = y$ .
2.  $d(x, y) = d(y, x)$ .
3. (triangle inequality)  $d(x, y) + d(y, z) \geq d(x, z)$ .

An *open ball* of radius  $r$  with center in  $x$  is  $\{y \in M : d(x, y) < r\}$ .

**Definition 1.5** Let  $M$  be a metric space. A subset  $U \subset M$  is called *open* if it is obtained as a union of open balls. This topology is called *induced by the metric*.

**Definition 1.6** A topological space is called *metrizable* if its topology can be induced by a metric.

**Exercise 1.3** Show that discrete topology can be induced by a metric, and codiscrete cannot.

*Solution.* To induce the discrete metric define the distance between any two distinct points to be 1. This clearly satisfies the three axioms of metric, and the ball of radius  $1/2$  is an open set that contains only its center, making any point and thus any subset an open set.

If a metric space contains at least two points at distance  $d$ . The ball with radius  $d/2$  at any of these points is an open set distinct from the empty set and the total, so the topology induced by the metric cannot be discrete.  $\square$

**Exercise 1.4** Prove that an intersection of any collection of closed subsets of a topological space is closed.

*Solution.* As I recall this is due to de Morgan laws stating that for any collection  $F_\alpha$  of subsets

$$\left( \bigcap_{\alpha} F_{\alpha} \right)^c = \bigcup_{\alpha} F_{\alpha}^c \quad (1)$$

where superscript  $c$  means set complement. If this is true 'then' we are done because if  $F_\alpha$  are closed, we see that the intersection is also closed as its complement is open.

Let's try to show eq. (1) ...  $\square$

**Definition 1.7** An intersection of all closed supersets  $Z \subset M$  is called *closure* of  $Z$

**Definition 1.8** A *limit point* of a set  $Z \subset M$  is a point  $x \in M$  such that any neighbourhood of  $M$  contains a point of  $Z$  other than  $x$ . A *limit* of a sequence  $\{x_i\}$  of points in  $M$  is a point  $x \in M$  such that any neighbourhood of  $x \in M$  contains all  $x_i$  for all  $i$  except a finite number. A sequence which has a limit is called *convergent*.

**Exercise 1.5** Show that a closure of a set  $Z \subset M$  is a union of  $Z$  and all its limit points.

*Solution.* It's enough to show that the union of  $Z$  and all its limit points  $W$  is a closed set and that it is contained in any closed set containing  $Z$ .

To see  $W$  is closed choose a point in its complement  $p \in W^c$ . Since  $p$  is not a limit point of  $Z$  nor a point of  $Z$ , there is a neighbourhood of  $p$  not intersecting  $Z$ . This means that such neighbourhood is contained in  $W^c$ . We can do this for all points in  $W^c$ , thus obtaining a  $W^c$  as a union of open sets, which is open, and then  $W$  is closed.

To see  $W$  is contained in any closed set containing  $Z$ , suppose  $F$  contains  $Z$  but not  $W$ . Then there must be a limit point of  $Z$  that is not in  $F$ . But then  $F$  cannot be closed because there is no neighbourhood of such a limit point contained in  $F^c$ , which should be open. **Indeed, if  $F^c$  is open then every point contains a neighbourhood contained in  $F^c$ .**  $\square$

**Exercise 1.7** Let  $f : M \rightarrow M'$  be a map of metrizable topological spaces, such that  $f(\lim_i x_i) = \lim_i f(x_i)$  for any convergent sequence  $\{x_i \in M\}$ . Prove that  $f$  is continuous.

*Solution.* It is equivalent that the preimage of every open set is open (definition of  $f$  being continuous) with the preimage of every closed subset is closed: for any closed set  $M' \setminus U$  with  $U$  open,  $f^{-1}(M' \setminus U) = f^{-1}(M') \setminus f^{-1}(U)$  is closed.

Consider the closed set  $F \subset M'$  and let's check that its preimage is also closed. By the same reasoning as in Exercise 1.5, to show closedness it's enough to show the set contains all its limit points. Take a limit point  $p \in f^{-1}(F)$ . We construct a convergent sequence  $\{x_n\}$  taking balls of radius  $\frac{1}{n}$  around  $p$ , each of which must contain a point in  $f^{-1}(F)$ . This gives a sequence in  $F$ , which by hypothesis must converge to a limit point  $\lim_i f(x_i) = f(\lim_i x_i) \in F$ . This means  $p = \lim_i x_i$  is in the inverse image of  $F$ .  $\square$

**Exercise 1.8\*** Find a counterexample to the previous problem for non-metrizable, Hausdorff topological spaces (see the next subsection of a definition of Hausdorff).

*Sketch of solution.* Probably Sorgenfrey line is a counter-example? I should look for its definition to make sure it is Hausdorff (and how is it defined exactly—I think open sets are positive rays).  $\square$

**Exercise 1.9\*\*** Let  $f : M \rightarrow M'$  be a map of countable topological spaces, such that  $f(\lim_i x_i) = \lim_i f(x_i)$  for any convergent sequence  $\{x_i \in M\}$ . Prove that  $f$  is continuous, or find a counterexample.

*Sketch of solution.* Is a **countable space** a space whose cardinality is  $\mathbb{N}$ ? What are the possible topologies on  $\mathbb{N}$ ? Discrete topology gives that every map is continuous. Other topologies are maybe, again, rays.  $\square$

**Exercise 1.10\*** Let  $f : M \rightarrow N$  be a bijective map inducing homeomorphisms on all countable subsets of  $M$ . Show that it is a homeomorphism, or find a counterexample.

*Sketch of solution.* If we suppose that  $M$  and  $M'$  are metrizable, we can use Exercise 1.7 as follows. Choose any convergent sequence  $\{x_i \in M\}$ . Then the countable set  $\{x_i\} \cup \{\lim_i f(x_i)\}$  is mapped homeomorphically to  $\{f(x_i)\} \cup \{f(\lim_i x_i)\}$ . This implies that  $f(\lim_i f(x_i)) = f(\lim_i x_i)$ , so  $f$  is continuous. The same holds for  $f^{-1}$ , so  $f$  is a homeomorphism.

Probably the statement isn't true in general, so let's look for a counter-example.  $\square$

## 1.2 Hausdorff spaces

**Definition 1.9** Let  $M$  be a topological space. It is called *Hausdorff* or *separable*, if any two distinct points  $x \neq y \in M$  can be *separated* by open subsets, that is, there exist open neighbourhoods  $U \ni x$  and  $V \ni y$  such that  $U \cap V = \emptyset$ .

**Remark 1.1** In topology, the Hausdorff axiom is usually assumed by default. In subsequent handouts, it will be always assumed (unless stated otherwise).

**Exercise 1.11** Let  $M$  be a Hausdorff topological space. Prove that all points in  $M$  are closed subsets.

*Solution.* Fix a point  $x \in M$ . For every  $y \in M$  distinct from  $x$  we have the neighbourhoods  $U_y \ni x$  and  $V_y \ni y$  with  $U_y \cap V_y = \emptyset$ . Then  $X \setminus \{x\} = \bigcup_{y \neq x} V_y$ , which is open.  $\square$

**Exercise 1.13** Let  $M$  be a topological space, with all points of  $M$  closed. Prove that  $M$  is Hausdorff, or find a counterexample.

*Solution.* No solution yet.  $\square$

**Exercise 1.14** Count the number of non-isomorphic topologies on a finite set of 4 elements. How many of these topologies are Hausdorff.

*Solution.* For any set  $S$  of subsets of  $\{1, 2, 3, 4\}$  we can consider the *topology generated by*  $S$ , which consists of all unions and intersections of elements in  $S$ , along with the total space and the empty set.

For the following choices of  $S$  we get non-isomorphic topologies:

- |   |                                    |
|---|------------------------------------|
| 1. $S = \emptyset$ (codiscrete topology).                       | 6. $S = \{\{1, 2\}\}$ .            |
| 2. $S = \{\{1\}\}$  | 7. $S = \{\{1, 2\}, \{3\}\}$       |
| 3. $S = \{\{1\}, \{2\}\}$ .                                     | 8. $S = \{\{1, 2\}, \{3, 4\}\}$ .  |
| 4. $S = \{\{1\}, \{2\}, \{3\}\}$ .                              | 9. $S = \{\{1, 2, 3\}\}$ .         |
| 5. $S = \{\{1\}, \{2\}, \{3\}, \{4\}\}$<br>(discrete topology). | 10. $S = \{\{1, 2, 3\}, \{4\}\}$ . |

1-5 are Hausdorff while 6-10 are not Hausdorff.  $\square$

**Exercise 1.5 (!)** Let  $Z_1, Z_2$  be nonintersecting closed subsets of a metrizable space  $M$ . Find open subsets  $U \supset Z_1, V \supset Z_2$  which do not intersect.

*Solution.* Consider the distance between  $Z_1$  and  $Z_2$ :

$$d(Z_1, Z_2) := \inf\{d(z_1, z_2) : z_1 \in Z_1, z_2 \in Z_2\}.$$

We must argue that  $d(Z_1, Z_2) \neq 0$ . Suppose by contradiction that  $d(Z_1, Z_2) = 0$ . Then for every  $n \in \mathbb{N}$  there is a pair of points  $z_1^n$  and  $z_2^n$  such that  $d(z_1^n, z_2^n) < 1/n$ .  $\square$

**Definition 1.10** Let  $M, N$  be topological spaces. **Product topology** is a topology on  $M \times N$ , with open sets obtained as unions  $\bigcup_{\alpha} U_{\alpha} \times V_{\alpha}$ , where  $U_{\alpha}$  is open in  $M$  and  $V_{\alpha}$  is open in  $N$ .

**Exercise 1.16** Prove that a topology on  $X$  is Hausdorff if and only if the diagonal  $\Delta := \{(x, y) \in X \times X \mid x = y\}$  is closed in the product topology.

*Solution.* ( $\implies$ ) Suppose that  $X$  is Hausdorff. To check that  $\Delta$  is closed suppose there is a sequence  $(x_n, x_n)$  that converges to some point  $(a, b) \in X \times X$ . We need to show that  $(a, b) \in \Delta$ , i.e. that  $a = b$ . For every neighbourhood  $W = U \times V$  of  $(a, b)$  we know that all but a finite number of  $(x_n, x_n)$  belong to  $U$ . This means that  $x_n$  converges to  $a$  and also to  $b$ . But limits are unique in Hausdorff spaces: if  $a \neq b$  we can separate  $a$  and  $b$  by disjoint open subsets  $A$  and  $B$ , and then it cannot hold simultaneously that all but a finite number of points in the sequence  $x_n$  are in  $A$  and in  $B$ .

( $\impliedby$ ) Suppose  $\Delta$  is closed in the product topology.  $\square$

**Definition 1.11** Let  $\sim$  be an equivalence relation on a topological space  $M$ . **Factor-topology** (or **quotient topology**) is a topology on the set  $M/\sim$  of equivalence classes such that a subset  $U \subset M/\sim$  is open whenever its preimage in  $M$  is open.

**Exercise 1.17** Let  $G$  be a finite group acting (continuously) on a Hausdorff topological space  $M$ . Prove that the quotient map is closed (i.e. puts closed subsets to closed subsets).

*Solution.* The quotient map is  $\pi : M \rightarrow M/\sim$  where  $x \sim y$  if  $y = gx$  for some  $g \in G$ . To show  $\pi$  is closed pick  $F \subset M$  closed. We need to show that  $\pi(F)$  is closed, so we may show its complement is open. According to the definition of factor topology we want to show that

$$\pi^{-1}\left((M/\sim) \setminus \pi(F)\right) = M \setminus \pi^{-1}(\pi(F))$$

is open. Now  $\pi^{-1}(\pi(F))$  is the set of points that are  $G$ -related to points in  $F$ , namely  $\bigcup_{g \in G} gF$ . Since  $G$  is finite and acts by homeomorphisms, this set is a finite union of closed sets, which is closed. Looks like I didn't use the Hausdorff hypothesis.  $\square$

**Exercise 1.18\*** Let  $\sim$  be an equivalence relation on a topological space  $M$ , and  $\Gamma \subset M \times M$  its *graph*, that is, the set  $\{(x, y) \in M \times M \mid x \sim y\}$ . Suppose that the map  $M \longrightarrow M/\sim$  is open, and that  $\Gamma$  is closed in  $M \times M$ . Show that  $M/\sim$  is Hausdorff.

**Hint .** Prove that diagonal is closed in  $M \times M$ .

*Solution.* If we show that the diagonal is closed in  $M \times M$ , then its complement would be an open set, and when projecting to  $(M/\sim) \times (M/\sim)$  we obtain that the diagonal is closed in the latter space.  $\square$