

k3

1 Class 1

The most important invariant of a k3 surface is [intersection form](#).

There are three classes of manifolds

1. Smooth manifolds

$$\text{smooth manifolds} \xrightarrow{\text{forgetful functor}} \text{PL manifold} \longrightarrow \text{Topological manifolds}$$

Donaldson: countably many non-equivalent smooth structures on \mathbb{R}^4 . K3 surfaces has countably many smooth structures and only one of them is compatible with complex structure.

Definition. Intersection form. Given a quadratic form on a lattice $V_{\mathbb{Z}} = \mathbb{Z}^n$, so

$$q : V_{\mathbb{Z}} \times V_{\mathbb{Z}} \rightarrow \mathbb{Z}$$

is *unimodular* if

$$V_{\mathbb{Z}} \xrightarrow{q} \text{Hom}(V_{\mathbb{Z}}, \mathbb{Z})$$

is an isomorphism.

Theorem (Universal coefficients formula).

$$H_{n-1}(M, \mathbb{Z}) = \mathbb{Z}^{b_{n-1}(M)} \oplus T_{n-1}(M)$$

$$h^n(M, \mathbb{Z}) = \mathbb{Z}^{b_n(M)} \oplus T_{n-1}(M)$$

Corollary. $H^2(X, \mathbb{Z})$ is torsion free if $\pi_1(X) = 0$ because

Definition. *Signature* is $m - n$ if q has signature (m, n) .

Theorem (Rokhlin-Wu?). Signature is divisible by 16 for simply-connected (something else).

Remark. The methods used in surgery break down in smooth case because strange topological objects like infinite sums of spheres arise.

Theorem (Freedman, 1982). There are as many 4-manifolds as there are intersection forms. A simply connected 4 manifold homotopy class is uniquely determined by intersection form. Moreover, for every unimodular form there exists a unique M with this intersection form.

Theorem (Donaldson, 1986). M smooth compact manifold with positive definite odd intersection form q . Then

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Definition. Bilinear symmetric form is *indefinite* if it is not positive definite nor negative definite.

Theorem (Classification of unimodular symmetric bilinear forms). Odd are diagonalizable, while even are related to special Lie group E_8 .

Definition. A **K3 surface** is a Kähler complex surface M with $b_1 = 0$ (simply connected) and $c_1(M, \mathbb{Z}) = 0$.

Kodaira did what André Weil couldn't classify.

Theorem. K3 surfaces have trivial canonical bundle $K_M = \Lambda^2(\Omega^1 M)$.

2 Class 2

G topological group. **Principal G bundle** is a space with free G -action such that the quotient E/G is Hausdorff. There are several conditions that make this work. And then you have $\text{Homotopy}(X, BG) = \text{equivalence classes of } G\text{-bundles}$. Vector bundles of a manifold are the same as maps from X to $BU(n)$.

Vector bundles up to stable equivalence are classified basically by Chern classes, so by the cohomology in $H^*(BU) = \mathbb{Q}[c_1, c_2, \dots, c_n]$.

Now look at the loop space of X . Then $H^*(\Omega X)$ is a free graded commutative algebra. Loop space has the interesting property that $\Omega U = BU$ and $\Omega BU = U$.

2.1 Bialgebras

Let A be a superalgebra (graded with antisymmetric product). Then we ask the axiom of coassociativity and that .

Example. G group, and $C(G)$ the ring of k -valued functions $C(G \times G) = C(G) \times C(G)$ so

$$\begin{aligned} G \times G &\longrightarrow G \\ C(G) &\longmapsto C(G) \otimes C(G) \end{aligned}$$

2.2 H-spaces

Definition. H-space is a space M with a map $\mu : M \times M \rightarrow M$ that is homotopy associative,

$$\begin{array}{ccc} M \times M \times M & \xrightarrow{\mu \times \text{id}} & M \times M \\ \downarrow \text{id} \times \mu & & \downarrow \mu \\ M \times M & \xrightarrow{\mu} & M \end{array}$$

which is homotopy commutative. And with homotopy unit.

So it's like a homotopy algebra?

Example. The loop space.

2.3 Bialgebras of finite type

Definition. A bialgebra A is of *finite type* if it is the direct sum of $A = \bigoplus_{i \geq 0} A^i$ supercommutative and each A^1 is finite dimensional.

Remark. Free commutative algebra is polynomial algebra

Definition. $A = \mathbb{C}[x_1, \dots, x_n, \dots] \otimes \Lambda^\bullet(a_1, \dots, a_n, \dots)$ is a graded commutative free algebra. In the slides: it is $\text{Sym}_{\text{gr}} V^*$ where V^* is a graded vector space.

Theorem (Hopf). A graded commutative bialgebra of finite type over k of 0 characteristic is free graded commutative as a k algebra.

2.4 The cohomology algebra of $U(n)$

Claim. The cohomology algebra $H^*(U(n), \mathbb{Q})$ is a free graded commutative algebra with generators in degrees $1, 3, 5, \dots, 2n-1$.

Demonstração. Induction. $U(1)$ is clear because it is a circle. Then do Serre spectral sequence. Differentials vanish on the second page because there's only nonzero groups on even degrees! And we get that $E_2^{p,1} = H^p(S^{2n-1}) \otimes H^q(U(n-1))$. And then the sequence converges to that of the total space which is $U(n)$. \square

2.5 Grassman manifolds

Definition. The *fundamental bundle* B_{fun} is a rank n vector bundle over $\text{Gr}(n, m)$.

Claim. B, B' vector bundles of rank $n, m-n, B \oplus B'$

$$\varphi : X \rightarrow \text{Gr}(m, n)$$

$$\varphi(x) = B_x \subset B_x \oplus B'_x = \mathbb{K}^m$$

then $B = \varphi^* B_{\text{fun}}$.

Theorem. If you have B as a bundle on a manifold X then $B \oplus B'$ is trivial for some bundle B' .

Demonstração. Embed the total space in a large enough euclidean space. \square

Definition. $\text{Gr}(n, \infty) = \text{Gr}(n)$ is $\bigcup_{m=n_1}^{\infty} \text{Gr}(n, m) = \text{Gr}(n)$

Corollary. For every bundle B of rank n there is a function $\varphi : X \rightarrow \text{Gr}(n)$ such that $B = \varphi^* B_{\text{fun}}$.

Take a bundle $E \rightarrow X$ and G acts freely on E so E principal G bundle. Classifying space BG

Theorem (Atiyah-Bott). Classifying space is unique up to homotopy equivalence.

2.5.1 The fundamental bundle

In class 4 I finally understood that

Definition. The *fundamental bundle* on the Grassmanian $\text{Gr}(n)$ (the Grassmanian is this space where points are linear spaces) is the vector bundle such that the fiber of one point (which is a vector space) is the vector space that is the point. It's very tautological.

Theorem (Did we prove this?). Let B be a vector bundle of rank n on a cellular space X . Then there exists a continuous map $\varphi : X \rightarrow \text{Gr}(n)$ such that B is isomorphic to the pullback $\varphi^* B_{\text{fun}}$ of the fundamental bundle.

Remark. In fact $\text{Gr}(n)$ is the classifying space of vector bundles of rank n , in the sense that isomorphism classes of vector bundles of maps $\varphi : X \rightarrow \text{Gr}(n)$.

2.6 Stiefel spaces

Definition. \mathbb{K}^{∞} is the direct limit of \mathbb{K}^n so its just the direct sum $\bigoplus_{i=1}^{\infty} \mathbb{K}$. Stiefel space is the space of orthonormal n -frames.

If we prove that Stiefel is contractible we obtain our classifying space so let's prove that. We have a fibration

$$U(n) \hookrightarrow \text{St}(n, \infty) \rightarrow \text{Gr}(n, \infty)$$

Theorem. $\text{St}(n)$ is contractible.

Demonstração **Step 1** Locally trivial fibration with contractible fiber and base $Y \rightarrow X$ then Y is contractible, this is so trivial.

Step 2 Fibration $\text{St}(n) \rightarrow \text{St}(n-1)$ with fiber S^{∞}

Step 3 Show that S^{∞} is contractible.

Step 4 And then some map \mathbb{R} that is not surjective, and construct homotopy of identity to a constant map.

□

Exercise. If $X_\infty = \bigcup X_i$ is the inductive limit of contractible cellular spaces then it is contractible. Use Whitehead theorem.

Theorem (Important). $\text{Gr}(\infty) = B U$.

2.7 Stable equivalence

Definition. Vector bundles V, W are stable equivalent if $V \oplus A \cong W \oplus B$ for trivial vector bundles A and B .

Homotopy classes of equivalent vector bundles are in coorespondance with...

Theorem. $B U$ is H-space.

Corollary. $H^*(B U, \mathbb{Q})$ is a free supercommutative algebra.

Claim. $H^*(B U)$ is a free polynomial algebra generated by classes c_1, c_2, \dots in all even degrees.

3 Class 3

3.1 Reminder

Definition. *Bialgebra* is an algebra that is equipped with comultiplication, counit...

Remark. It is when the dual space also has an algebra structure, but we prefer to use the tensor notation.

Let $\sum_{i \geq 0} A^i$ with $\dim A^i < \infty$. *Free commutative algebra* is a polynomial algebra. *Free graded commutative algebra* is

$$\widetilde{\text{Sym}}^\bullet(W^\bullet \oplus V^\bullet) := \text{Sym}^\bullet(W^\bullet) \otimes \Lambda^\bullet(V^\bullet)$$

where

$$W = \bigoplus_i W^{\text{even}} \quad V = \bigoplus_i V^{\text{odd}}.$$

3.2 Hopf algebra

Definition. A bialgebra is a *Hopf algebra* when it is also equipped with an antipode map (S) such that the following diagram commutes

$$\begin{array}{ccccc}
 & H \otimes H & \xrightarrow{S \otimes \text{id}} & H \otimes H & \\
 \Delta \nearrow & & & & \searrow m \\
 H & \xrightarrow{\eta} & \mathbb{C} & \xrightarrow{u} & H \\
 \Delta \searrow & & & & \nearrow \\
 & H \otimes H & \xrightarrow{\text{id} \otimes S} & H \otimes H &
 \end{array}$$

[diagram from quantum group minicourse notes]

Example. The cohomology of the loop space, $H^\bullet(\Omega X)$.

3.3 Primitive elements in a bialgebra

Definition. An element of a bialgebra $x \in A$ is *primitive* if $\Delta(x) = x \otimes 1 + 1 \otimes x$.

$$\begin{aligned}
 \Delta(xy) &= \Delta(x)\Delta(y) \\
 &= (1 \otimes x + x \otimes 1)(y \otimes 1 + y \otimes y) \\
 &= 1 \otimes xy + xy \otimes 1 + x \otimes y + y \otimes x.
 \end{aligned}$$

Remark. We trying to show that Hopf algebras? bialgebras? are generated by primitive elements?

Definition. A^\bullet bialgebra, $\mathcal{P}^\bullet \subset A^\bullet$ space of primitive, and the natural embedding

$$\text{Sym}_{\text{gr}}(\mathcal{P}^\bullet) \rightarrow A$$

We say that A is *free up to defree* k if

$$\bigoplus_{i \leq k} \text{Sym}_{\text{gr}}^i(\mathcal{P}) \xrightarrow{\psi} A$$

is an embedding.

Lemma. Let A^\bullet be a bialgebra which is free up to degree k . Then A^\bullet is free up to degree $k + 1$.

Proof.

Step 1 Choose a basis of \mathcal{P} , $\{x_i\}$. Chose a polynomial condition $Q(x_1, \dots, x_n) = 0$ of degree $k + 1$. Write this as

$$Q = Q_m x_1^m + Q_{m-1} x_1^{m-1} + \dots + Q_0.$$

that is

$$Q = \sum_{i=0}^m Q_i x_1^i$$

with Q_i invariant somehow. Then we apply comultiplication to obtain

$$\Delta(Q) = Q \otimes 1 + 1 \otimes Q + R$$

where R is some sort of reminder with bounded degree:

$$R \in \mathfrak{U} := \bigoplus_{i \leq k} \text{Sym}_{\text{gr}}^i(P) \otimes \bigoplus_{i \leq k} \text{Sym}_{\text{gr}}^i(P)$$

which follows from a similar computation of that which we did after defining primitive elements.

Step 2 Project to drop the terms that have $Q \otimes 1 + 1 \otimes Q$:

$$\Pi : \mathfrak{U} \rightarrow x_1 \otimes \bigoplus_{i \leq k} \text{Sym}_{\text{gr}}^i(P)$$

since the x_i are primitive, i.e. $\Delta(x_i) = x_i \otimes 1 + 1 \otimes x_i$, one has

$$\Delta(x_1^m) = (x_1 \otimes 1 + 1 \otimes x_1)^m$$

we get that

$$\Pi(\Delta(x_1^m)) = mx_1 \otimes x_1^{m-1}$$

while on the board it is written that

$$\Pi(\Delta(x_1^m)) = \Pi((x_1 \otimes 1 + 1 \otimes x_1)^m)$$

Step 3 Let $\Pi(R) := x_1 \otimes R_0$. Since $Q = 0$ in A , its component R_0 is also equal to 0. So $\Pi(\Delta(Q)) = 0$. Then

$$\begin{aligned} 0 &= \Pi \left(\Delta \left(\sum_m x_1^m \cdot Q_m \right) \right) \\ &= \sum_m x_1 \otimes x_1^{m-1} Q_m + \Pi(mx_1 \otimes x_1^{m-1} \cdot \Delta(Q_m)) \\ &= \sum_m x_1 \otimes x_1^{m-1} Q_m \end{aligned}$$

so that

$$\begin{aligned} x_1 \otimes x_1^{m-1} Q_m &= 0 \\ \implies x_1^{m-1} Q_m &= 0 \end{aligned}$$

So we conclude that

$$Q_m = 0$$

□

Remark. We just proved that for any subalgebra generated by finite elements, we didn't use that it is free.

3.4 Algebras with filtration

Definition. A *filtration on algebra* is

$$A^\bullet \supset F_1 A^\bullet \supset F_2 A^\bullet \supset \dots$$

such that

$$F_i A^\bullet F_j \subset F_{i+j} A^\bullet$$

Definition. *Associated graded* to a filtered algebra is

$$A_{\text{gr}}^\bullet = \bigoplus_{i=0}^{\infty} \frac{F^i A^\bullet}{F^{i+1} A^\bullet}$$

$$F^0 A^\bullet = A^\bullet$$

Definition. $I \subset A$ ideal then *I-adic filtration* is the filtration by the degrees of the ideal

$$A \supset I \supset I^2 \supset I^3 \dots$$

Lemma. Choose an I-adic filtration. Then A_{gr} is generated by its first and second graded components $A/I \oplus I/I^2$.

Demonstração. Indeed, I^k / I^{k+1} is generated by products of k elements in (I/I^2) . □

Definition. A *augmentation ideal* in a bialgebra is the kernel of the counit homomorphism $\varepsilon : A \rightarrow k$. We denote it by $Z = \ker A$

Remark.

$$\Delta(x) = 1 \otimes x + x \otimes 1 \mod Z \otimes Z$$

Why? Because

$$\begin{aligned} x &= \varepsilon \otimes \text{id}(\Delta(x)) && \text{up to } Z \otimes A \\ \Delta(x) &= 1 \otimes x && \text{up to } A \otimes X \\ \Delta(x) &= x \otimes 1 \end{aligned}$$

Ok, now we can prove Hopf theorem.

Theorem (Hopf theorem). A finite type bialgebra is generated by primitive elements.

In slides: Let A be a graded bialgebra of finite type over a field k of characteristic 0. Then A is a free graded commutative k -algebra.

Proof.

Step 1 I think this is the computation above.

Step 2 A_{gr} is a bialgebra.

Step 3 A_{gr} is multiplicative generated by Z^1/Z^2 . All elements Z^1/Z^2 are primitive, so this algebra A_{gr} is generated by primitive elements.

Step 4 Let $\{\chi_i\}$ be a basis of primitive elements of A_{gr} . Then lifts of A have no relations because A_{gr} is already generated by primitive elements. Then there are no relations also for elements in A^\bullet (I think).

□

3.5 Grassmanians (Reminder)

B vector bundle of rank n on X then there exists a map (essentially unique) $\varphi : X \rightarrow \text{Gr}(n)$ such that

$$\varphi^*(B_{\text{fun}}) = B$$

which makes the Grassmanian a classifying space, and $\text{Gr}(1) = BU(n)$.

The infinite Grassmanian is important.

3.6 BU as an H-space (Reminder)

Bott periodicity identifies the space of loops on U and BU .

Proposition. Embed $\mathbb{C}^\infty \times \mathbb{C}^\infty$ into \mathbb{C}^∞ taking the basis vectors of the first copy to the even basis vectors and the basis of the second copy to the odd. Then for $L_1 \subset \mathbb{C}^\infty$, $L_2 \subset \mathbb{C}^\infty$, the map

$$L, L' \mapsto S(L, L')$$

defines a structure of H-space on the infinite Grassmanian BU .

Proof. Just show that H-associativity up to homotopy.

□

Corollary. $H^\bullet(BU, \mathbb{Q})$ is a free supercommutative algebra.

Proof. Follows from Hopf theorem.

□

3.7 Cohomology of BU

Claim. $H^\bullet(BU, \mathbb{Q})$ is a free polynomial algebra generated by classes c_1, c_2, \dots in all even degrees.

Demonstração. Leray-Serre spectral sequence.

□

3.8 Chern classes: axiomatic definition

Definition. *Chern classes* are classes $c_i(B) \in H^{2i}(X)$ for $i = 0, 1, 2, \dots$

Chern classes are $c_i(B) \in H^{2i}(X)$ for a complex vector bundle B over X with axioms

a. $c_0(B) = 1$

b. Functoriality (commutes with pullbacks): for $\varphi : X \rightarrow Y$ with B bundle on Y ,

$$\varphi^*(c_i(B)) = c_i(\varphi^*(B))$$

c. Define *total Chern class* $c_* := \sum_i c_i(B)$ then

$$c_i(B) \cdot c_i(B') = c_*(B \oplus B') \quad (\text{Whitney})$$

d. $\mathcal{O}(1)$ on \mathbb{CP}^n ,

$$c_i(\mathcal{O}(1)) = 1 + [H]$$

where $[H]$ is the fundamental class of a hyperplane section.

Suppose we have a class $a \in H^*(B U)$. Then for all B on X

$$\varphi : X \rightarrow B U$$

so

$$B \cong \varphi^*(B_{\text{fun}})$$

and so

$$\varphi_B^*(c_*) = c_*(B).$$

4 Class 4

4.1 Reminder

For each rank n bundle B on X there exists $\varphi_B : X \rightarrow \text{Gr}(n, \infty) = B U(n)$ such that $\varphi_B^*(B_{\text{fun}}) = B$.

The infinite grassmanian is classifying space for (?) stable bundles.

Some more review about H-space structure, primitive elements, a comment on last exercise of homework 2.

Chern classes of $\mathcal{O}(1)$ are hyperplane sections: $c_i(\mathcal{O}(1)) = 1 + [H]$.

4.2 The splitting principle

Exercise. Prove that $B U(1) = \mathbb{CP}^\infty$.

Solution. Hopf fibration on S^∞ ? It's easier, take $n = 1$, it's just by definition. □

Definition. The *fundamental bundle* on $BU(1)^n$ has fiber

$$\ell_1 \oplus \ell_2 \oplus \dots \ell_n$$

where $\ell_i \in BU(1)$ are product $\ell_1 \times \ell_2 \times \dots \times \ell_n$.

Remark. Chern classes of B_{fun} are uniquely determined by axioms, because every factor has Chern classes, and fibers are just sums, and pullbacks preserve sums...

$$B_{\text{fun}} = \bigoplus_i \pi_i \mathcal{O}(1)$$

where

$$\pi_i : BU(1)^n \rightarrow BU(1)$$

is a projection.

Remark. $H^\bullet(BU(1))^n = \mathbb{Z}[z_1, \dots, z_n]$ Here at least I remember that the cohomology of \mathbb{CP}^∞ is just polynomials so it looks reasonable that the n -th power is polynomials in more variables.

Theorem (Splitting principle). Let $\varphi_{\text{fun}} : BU(1)^n \rightarrow BU$, the *fundamental map*, it induces embedding on cohomology up to degree $2n$. For all primer generator $\sigma_i \in H^2(BU)$, $\varphi_{\text{fun}}(\sigma_i) = \lambda \sum_i z_i^k$ with $\lambda \neq 0$.

So

$$\begin{array}{ccc} BU(1)^n & \longrightarrow & BU \\ \uparrow & \nearrow & \\ X & & \end{array}$$

Remark. [Wiki](#) Thus, the set of isomorphism classes of circle bundles over a manifold M are in one-to-one correspondence with the homotopy classes of maps from M to \mathbb{CP}^∞

Theorem. Chern classes are unique (uniquely determined by axioms).

Proof.

Step 1 Every bundle is obtained as pullback of the fundamental bundle. So for $A \in H^\bullet(BU)$ and B bundle on X , $A(B) = \varphi_B^*(A) \in H^\bullet(X)$ so $c_i(B)$ are obtained as pullbacks of c in the fundamental bundle.

Step 2

$$BU(1)^\infty \xrightarrow{\varphi_{\text{fun}}} BU$$

pullback of fundamental bundle is fundamental. (This map is defined from the former by induction).

$$\varphi_{\text{fun}}^*(c_i(B_{\text{fun}})) = c_i(B_{\text{fun}} \text{ on } BU)$$

The Chern classes of the fundamental bundle are already known. Since φ_{fun}^* is injective by the splitting principle we are done.

□

4.3 Primitive generators of $H^*(BU)$

Recall the H-space multiplication:

$$\begin{aligned} BU \times BU &\longrightarrow BU \\ L_1 \times L_2 &\longmapsto L_1 \oplus L_2 \end{aligned}$$

and the comultiplication

$$\Delta : H^*(BU) \rightarrow H^*(BU)$$

Generators of $H^*(BU)$ are c_{h_1}, c_{h_2}, \dots with $c_{h_i} \in H^{2i}(BU)$ and we have the comultiplication $\Delta(c_{h_i}) = c_{h_i} \otimes 1 + 1 \otimes c_{h_i}$.

Remark.

$$\varphi = (\varphi_1, \varphi_2) : X \rightarrow BU \times BU$$

and we can compose so we have

$$\varphi \circ \mu : X \rightarrow BU$$

what does this map do?

$$\begin{aligned} \varphi \circ \mu : X &\longrightarrow BU \\ \varphi^*(B_{\text{fun}}) &\longmapsto B_1 \\ (\varphi \circ \mu)^*(B_{\text{fun}}) &= B_1 \oplus B_2 \end{aligned}$$

So then we have

$$\begin{aligned} \varphi^* : H^*(BU) \otimes H^*(BU) &\rightarrow H^*(X) \\ \Delta : H^*(BU) &\rightarrow H^*(BU) \otimes H^*(BU) \\ \Delta \circ \varphi^* : H^*(BU) &\rightarrow H^*(X) \end{aligned}$$

Corollary. For every $x \in H^*(BU)$

$$X(B_1 \oplus B_2) = \Delta(x)(B_1, B_2)$$

Corollary. If $x \in H^*(BU)$ is primitive, then $x(B_1 \oplus B_2) = x(B_1) \oplus x(B_2)$.

Proof. $\Delta(x) = x \otimes 1 + 1 \otimes x$ so $\Delta(x)$ evaluated on (B_1, B_2) □

Remark. We will construct the full Chern class $c_*(B)$ as a pullback of a class $C \in H^*(BU)$.

Remark. Then take exponential. Let $\chi_i \in H^{2i}(BU)$ be a primitive generator. Use Hopf theorem to see that it is unique by a constant. Since $\chi_i(B_1 \oplus B_2) = \chi_i(B_1) + \chi_i(B_2)$, the class $C = e^{\sum_i a_i \chi_i} = 1 + \dots + \frac{\chi_n}{n!} + \dots$ satisfies the Whitney formula.

To construct Chern classes satisfying the axioms it remains to arrange the coefficients a_i in such a way that $C(\mathcal{O}(1)) = 1 + [H]$ I think this means hyperplane section.

Lemma. An embedding

$$\mathrm{BU}(1) \xhookrightarrow{\varphi} \mathrm{BU}$$

with $\chi_i \in H^{2i}(\mathrm{BU})$ primitive generator. Then $\varphi^*(\chi_i) \neq 0$

Proof. $H^\bullet(\mathrm{BU}) = \text{symmetric polynomials in } H^i(\mathrm{BU}(1))^n$, $\varphi_{\mathrm{fun}}(x_N) = x \sum_{i=1}^n z_i^k$ so $\varphi(x_k) = \lambda x_1^k$. \square

Remark. $\varphi^*(c_i(\mathrm{B}_{\mathrm{fun}})) = c_i(\mathcal{O}(1)) = 1 + [H]$

Theorem. Choose generators $\chi_i \in H^2(\mathrm{BU})$ primitive. Then $\varphi^*(\sum_i \chi_i) = \log(1 + [H])$ where the logarithm is a formal power series, namely $\sum_{i=1}^{\infty} \frac{H^i}{i!} (-1)^i$.

That means $c(\mathrm{B}_{\mathrm{fun}}) = \exp\left(\sum \chi_i\right)$.

5 Class 5