

# Exam

## Contents

Exercise 1.3 . . . . .	1
Exercise 1.7 . . . . .	1
Exercise 2.7 . . . . .	2
A possible counter-example . . . . .	2
Proof of the statement without taking modulus . . . . .	3
Exercise 2.8 . . . . .	3
Exercise 3.3 . . . . .	4
Exercise 3.7 . . . . .	5
Exercise 4.2 . . . . .	5
Exercise 4.6 . . . . .	5
Exercise 5.2 . . . . .	6
Exercise 5.8 . . . . .	6

**Exercise 1.3** Let  $M$  be an almost complex manifold, and  $f : M \rightarrow \mathbb{R}$  a non-constant function which satisfies  $dJd(f) = 0$ . Prove that  $M$  admits a non-zero holomorphic 1-form (this means closed form).

*Solution.* It is unclear what is the definition of a holomorphic form on an almost complex manifold. Here's some ideas:

The form  $\alpha := df - \sqrt{-1}Jdf$  is complex linear since for any vector  $v$

$$\begin{aligned}
 \alpha(Jv) &= (df - \sqrt{-1}Jdf)(Jv) \\
 &= df(Jv) - \sqrt{-1}df(J^2v) \\
 &= df(Jv) + \sqrt{-1}df(v) \\
 &= -\sqrt{-1}^2 df(Jv) + \sqrt{-1}df(v) \\
 &= \sqrt{-1}(df(v) - \sqrt{-1}dJf(v)) \\
 &= \sqrt{-1}(df - \sqrt{-1}Jdf)(v) \\
 &= \sqrt{-1}\alpha(v).
 \end{aligned}$$

But then again it may be more suitable to find a form whose *differential* is complex-linear. After consulting with Misha, he said to find a form that is closed. . . but  $dJd(f)$  is already closed!  $\square$

**Exercise 1.7** Let  $\eta$  be a closed 1-form on a 1-dimensional complex manifold  $(M, I)$  such that  $I(\eta)$  is also closed.

- [Not part of the exam] Prove that  $\eta = df$ , where  $f$  is a real part of a holomorphic function if  $M$  is simply connected.
- Find an example of  $(M, I, \eta)$  where  $\eta$  is exact, but such  $f$  does not exist.

*Solution.*

(See [StackExchange](#).) The idea is to define the form as the exterior derivative of the real part of a logarithm. The logarithm cannot be defined globally so that function is not the real part of a holomorphic function.

Summary: on any non-simply connected open subset  $\Omega$  of  $\mathbb{C}$  we can find a function, namely  $u(z) = \log|z - w|$  with  $w \notin \Omega$  such that there is a curve with nonzero winding number around it contained in  $\Omega$ . This function is harmonic but it is not the real part of a holomorphic function (which would be a complex logarithm).

**Step 1** Define  $u(z) = \log|z - w|$  on a non-simply connected subset  $\Omega \subset \mathbb{C}$  not containing  $w$ . Also let  $\gamma$  be a curve with nonzero winding number with respect to  $w$ .

**Step 2** Notice that  $u$  is harmonic because it is *locally* the real part of a holomorphic function, the logarithm.

**Step 3** Suppose  $v$  is such that  $f := u + iv$  is holomorphic. Recall that harmonic conjugates are unique up to adding a constant by Cauchy-Riemann equations. This means that the derivative of  $f$  is the same as the derivative of the logarithm, ie.,  $f'(z) = \frac{1}{z-w}$ .

**Step 4** Remember the definition of the winding number:

$$n(\gamma, w) = \frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z - w} = \frac{1}{2\pi i} \int_{\gamma} f'$$

so  $n(\gamma, w) \neq 0$  by the choices of  $\gamma$  and  $w$ , but its also 0 because it's the integral of a holomorphic function over a closed curve.

Now let's solve our exercise. Consider  $\eta = du$  for the same  $u$  as above. We already know that  $u$  is not the real part of a holomorphic function but perhaps there is another function  $\tilde{u}$  such that  $\eta = d\tilde{u}$  and  $\tilde{u}$  is the real part of a holomorphic function. Then  $d(u - \tilde{u}) = 0$  so  $u - \tilde{u}$  is constant so  $\tilde{u}(z) = \log|z - w| + \lambda$ . But the same argument shows that such a function cannot be the real part of a holomorphic function.  $\square$

**Exercise 2.7** Let  $f_i$  be a collection of holomorphic functions on a disk such that  $\sum_{i=1}^{\infty} |f_i(z)|$  converges uniformly on  $\Delta$ . Prove that  $\sum_{i=0}^{\infty} |f'_i(z)|$  converges uniformly on  $\Delta$ .

*Solution.* I could prove that  $\sum_{i=0}^{\infty} f'_i(z)$  (without the bars) converges (uniformly) on  $\Delta$ . However, looks like the answers might be that such a function cannot exist.

### A possible counter-example

(See [StackExchange](#)). The proposed counter example is

$$f_n(z) = \frac{1}{n^2} \sqrt{1+z}$$

The reason is that the series converges to  $\frac{\pi^2}{6} \sqrt{1+z}$ , but the series of derivatives is unbounded, so "it cannot be uniformly convergent". But of course unboundedness alone does not imply that the sequence of derivatives cannot converge uniformly—for any unbounded function on a domain, just subtracting a  $1/n$ -term will produce a uniformly convergent series of unbounded functions.

### Proof of the statement without taking modulus

Now I show how that if  $\sum_{i=1}^{\infty} |f_i(z)|$  converges uniformly on  $\Delta$ , then  $\sum_{i=0}^{\infty} |f'(z)|$  converges on  $\Delta$ .

Recall from Ahlfors *Complex Analysis* (p. 176) Weierstrass theorem that if a sequence of holomorphic functions on a region  $\Omega$  (a region is a nonempty connected open set) converges to a limit function  $f$  in  $\Omega$  and uniformly on every compact subset of  $\Omega$ , then  $f$  is analytic on  $\Omega$ . Moreover,  $f'_i$  converges uniformly to  $f'$  on every compact subset of  $\Omega$ .

This is almost exactly what we need. Following I sketch the Ahlfors's proof slightly adapted to the exercise.

Recall that an absolutely uniform convergent series is uniformly convergent, so that the sequence of partial sums  $F_n = \sum_{i=1}^n f_i$  without the bars is uniformly convergent. Choose any point  $z \in \Delta$  and a closed curve contained in  $\Delta$  around  $z$ . Then Cauchy formula for the first derivative says  $F'_n(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{F_n(\zeta)}{(\zeta-z)^2} d\zeta$  for every  $n \in \mathbb{N}$ . Now take limit and put it inside the integral using uniform convergence of  $F_n$  to get that

$$\sum_{i=1}^{\infty} f'_i = \frac{1}{2\pi i} \int_{\gamma} \frac{\sum_{i=1}^{\infty} f_i(\zeta)}{(z-\zeta)^2} d\zeta$$

which says that  $\sum_{i=1}^{\infty} f'_i$  converges to the derivative of  $\sum_{i=1}^{\infty} f_i$ . Then Ahlfors says a simple estimate shows convergence is uniform but I'm not sure how exactly.  $\square$

**Exercise 2.8** Construct a non-zero bounded holomorphic function on  $\{z \in \mathbb{C} : \operatorname{Re}(z) > 0\}$  such that  $f(n) = 0$  for all  $n \in \mathbb{Z}^{>0}$ .

*Solution.* (See [StackExchange](#)) Such a function doesn't exist. We will show it must be constant.

**Step 1** Pass to a function whose domain is the unit disk by composing  $f$  with a Möbius transformation that maps the disk to the right-hand plane, namely  $z \mapsto \frac{1}{1+z} - \frac{1}{2}$ . So define  $g(z) = f\left(\frac{1}{1+z} - \frac{1}{2}\right)$ .

**Step 2**  $g(z) = 0$  when  $\frac{1}{1+z} - \frac{1}{2}$  is an integer  $n$  so define  $z_n = -1 + \frac{1}{n+\frac{1}{2}}$ .

$\prod_n |z_n|$  diverges to zero. That means that it converges to zero but the infinite sum (series) the logarithms diverges.

**Step 3** Consider the inverse of a finite Blaschke product that vanishes on the  $z_n$  (this is why we put everything inside the unit disk). It is  $\tilde{B}_k(z) = \prod_{n=1}^k \frac{1-\bar{z}_n z}{z-z_n}$ . Also consider the product  $g \cdot \tilde{B}_k$ , which is analytic because  $g$  vanishes at the poles of  $\tilde{B}_k$ .




Figure 1: Domain coloring plot for  $\tilde{B}$

**Step 4** Recall that the modulus of such a function takes its maximum at the boundary of its domain. Then show that  $|\tilde{B}_k(z)|$  is bounded by 1 (I'm not sure how). We already know that  $g$  is bounded by some number  $M$  because  $f$  is bounded by hypothesis. We have  $|g(z)| \leq \frac{M}{|\tilde{B}_k(z)|}$  for all  $|z| < 1$  and all  $k$ .

**Step 5** Put  $z = 0$ , you get  $|g(z)| \leq M \cdot \prod_{n \leq k} |z_n|$  and take limit as  $k \rightarrow \infty$ . The product diverges to zero so  $g(0) = 0$ .

**Step 6** That means  $\frac{g(z)}{z}$  is analytic so do everything again with  $\frac{g(z)}{z}$  instead of  $g(z)$ . And again for  $\frac{g(z)}{z^2}$ . And so on.  $g$  has a zero of infinite order, which is not possible and thus it is constant zero.

□

**Exercise 3.3** Let  $V$  be an  $n$ -dimensional Hermitian complex space of signature  $(1, n-1)$ . Prove that the space  $B \subset \mathbb{P}_{\mathbb{C}} V$  of all positive complex lines in  $V$  biholomorphic to a ball in  $\mathbb{C}^{n-1}$ .

*Solution.* (See [Wiki](#).) A positive line is a 1-dimensional subspace  $\ell$  such that  $q(x, x) > 0$  for some (and hence all)  $x \in \ell$ . For a line to be positive there must be a representative with non-zero first coordinate, which we may choose (uniquely) to be 1. Then

$$q(x, x) > 0 \iff 1 > \sum_{i=2}^n |x_i|^2.$$

We have constructed a biholomorphism

$$\begin{aligned} B &\longrightarrow \{|x| < 1\} \subset \mathbb{C}^{n-1} \\ [1 : x_2 : \dots : x_n] &\longmapsto (x_2, \dots, x_n) \end{aligned}$$

□

**Definition 3.1** *Horocycle* on a Poincaré plane is an orbit of a parabolic subgroup

$$P_t = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \in \mathrm{PSL}(2, \mathbb{R}) = \mathrm{SO}^+(1, 2).$$

**Exercise 3.7** Prove that the group of isometries  $\mathrm{Iso}(\mathbb{H}^2) = \mathrm{SO}^+(1, 2)$  acts transitively on the set of horocycles.

*Solution.* Let's fix two parabolic subgroups. I say that there is an isometry in  $\mathrm{Iso}(\mathbb{H}^2)$  that maps one to the other via conjugation. This gives a transitive action on horocycles: two horocycles, each determined by a point and a parabolic subgroup, are paired via a single element of  $\mathrm{Iso}(\mathbb{H}^2)$  (the pairing is via conjugation on subgroups and usual mapping on points). □

**Exercise 4.2** Let  $M$  be a compact, Kobayashi-hyperbolic complex manifold. Prove that  $M$  does not admit non-zero holomorphic vector fields.

*Solution.*

Integrating a non-zero vector field on  $M$  gives a complex 1-parameter subgroup acting on  $M$ . Let's show that  $\mathbb{C}$  cannot act nontrivially on  $M$ . Suppose that  $f : \mathbb{C} \times M \rightarrow M$  is an holomorphic action of  $\mathbb{C}$  on  $M$ . Then for each  $p \in M$ , the map  $a \in \mathbb{C} \mapsto f(a, p) \in M$  is holomorphic, so it is distance decreasing with respect to Kobayashi distances in  $\mathbb{C}$  and  $M$  (i.e. it is 1-Lipschitz). But Kobayashi distance in  $\mathbb{C}$  is trivial, making the action trivial. □

**Exercise 4.6** Let  $a, b, c \in \mathrm{Abs}$  be three distinct points on the absolute, and  $A \in \mathrm{SO}^+(1, 2)$  an isometry of Poincaré plane which fixes  $a, b, c$ . Prove that  $A = \mathrm{Id}$ .

*Solution.* We have seen in our lectures that  $SO^+(1,2)$  is isomorphic to the group of transformations  $z \mapsto \frac{az+b}{cz+d}$  with  $a, b, c, d \in \mathbb{R}$  and  $ad - bc > 0$ . Such transformations act on the upper-half plane as hyperbolic isometries. A fixed point of any such transformation is solution of a quadratic equation, which has potentially only two distinct solutions. Thus, if a transformation fixes three distinct points, it must be the identity.  $\square$

**Exercise 5.2** Let  $M$  be a Riemann surface with infinite fundamental group. Prove that any continuous map  $S^2 \rightarrow M$  is homotopic to a trivial map (map to a point).

*Solution.* All higher homotopy groups of Riemann surfaces of nonzero genus are trivial by covering space argument. For  $S^2$  itself, the statement does not hold since it would imply that  $-\text{Id}$  is homotopic to  $\text{Id}$ , which is not true (but of course, the fundamental group of  $S^2$  is not infinite).  $\square$

**Exercise 5.8** Let  $TS^2 \oplus C^\infty S^2$  be a direct sum of a tangent bundle  $TS^2$  and a 1-dimensional bundle. Is the bundle  $TS^2 \oplus C^\infty S^2$  trivial?

*Solution.* [From Hatcher VBK p. 10.] The direct sum of  $TS^2$  and its normal bundle  $NS^2$  is trivial since the map  $(x, v, tx) \mapsto (x, v + tx)$ , with  $v$  tangent at  $x \in S^2$  and  $t \in \mathbb{R}$ , is an isomorphism  $TS^2 \oplus NS^2 \cong S^2 \times \mathbb{R}^3$ . The idea here is that while any global section will vanish at one point  $x \in S^2$ , so  $v_x = 0$ , the point  $x$  will never be zero.

Recall that a vector bundle isomorphism consists of a homeomorphism of the total spaces that restricts to a linear isomorphism between corresponding fibers. It is clear that the map above is an homeomorphism, and fixing  $x \in S^2$  we also find a trivial isomorphism between the fibers—there isn't much to say here.

For a line bundle  $C^\infty S^2$  that is always linearly independent with respect to  $TS^2$  (which is the case since we have a direct sum), the map  $(x, v, y) \mapsto (x, v + y)$  yields an isomorphism  $TS^2 \oplus C^\infty \cong S^2 \times \mathbb{R}^2$  or  $S^2 \times \mathbb{R}^3$  depending on the rank of  $TS^2$ .  $\square$