

VERTEX ALGEBRAS

github.com/danimalabares/vertex-algebras

CONTENTS

1. Cartan subalgebra, Cartan matrix and Serre relations	1
2. Some infinite dimensional Lie algebras	2
3. Kac-Moody algebras	5
4. Affine Kac-Moody algebras	8
5. Weyl group	9
6. Weyl character formula	10
7. Characters of integrable highest weight for affine Kac-Moody algebras	15
8. θ -functions	17
References	18

1. CARTAN SUBALGEBRA, CARTAN MATRIX AND SERRE RELATIONS

Kac-Moody algebras are Lie algebras, whose definition is motivated by the structure of finite-dimensional simple Lie algebras over $\mathbb{C}$.

Let $\mathfrak{g}$ be a finite-dimensional semisimple Lie algebra over $\mathbb{C}$. Then $\mathfrak{g}$ has a *Cartan subalgebra* $\mathfrak{h} \subset \mathfrak{g}$ (abelian + ...) (see [Kac10, Definition 8.2]). Fixing $\mathfrak{h} \subset \mathfrak{g}$ gives a *root space decomposition* (see [Kac10, Proposition 8.5])

$$\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_{\alpha}$$

where $\Delta \subset \mathfrak{h}^*$ linear dual, and, by definition

$$\mathfrak{g}_{\alpha} = \{X \in \mathfrak{g} \mid [H, X] = \alpha(H)X \ \forall H \in \mathfrak{h}\}$$

Turns out the $\mathfrak{g}_{\alpha}$ are all 1-dimensional, though this property is lost when we go to Kac-Moody algebras.

$$[\mathfrak{g}_{\alpha}, \mathfrak{g}_{\beta}] \subset \mathfrak{g}_{\alpha+\beta}$$

The Killing form $\kappa : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$, $\kappa(x, y) = \text{Tr}_{\mathfrak{g}} \text{ad}(x)\text{ad}(y)$ is nondegenerate. “This is kind of the definition of semisimple.” (Think of $\mathfrak{h}$ as $\mathfrak{g}_0$, btw.)

$\kappa|_{\mathfrak{g}_{\alpha} \times \mathfrak{g}_{\beta}} \neq 0$ only when $\beta = -\alpha$. $\kappa|_{\mathfrak{h} \times \mathfrak{h}}$ is non-degenerate. This gives a linear isomorphism $\mathfrak{h} \xrightarrow{\nu} \mathfrak{h}$ via $\nu(H)(H') = \kappa(H, H')$.

So, $\mathfrak{h}^*$ comes with a non-degenerate bilinear form.

The *reflection* $r_{\alpha} : \mathfrak{h}^* \rightarrow \mathfrak{h}^*$ in $\alpha \in \mathfrak{h}^*$ (usually a root) is $r_{\alpha}(\lambda) = \lambda - 2 \frac{(\lambda, \alpha)}{(\alpha, \alpha)} \cdot \alpha$.

“Classify root systems [...] classify semisimple Lie algebras” It is a fact that $r_{\alpha}(\Delta) = \Delta$ for all $\alpha \in \Delta$, which motivates the definition of *root system* (see [Kac10, Definition 15.1]) and permits classification. (See [Kac10, Lecture 17] for comments on correspondence of root systems and semisimple Lie algebras.)

Example 1.1. $\mathfrak{g} = \mathfrak{sl}_2$, $\mathfrak{h}$ = diagonal matrices

$$\begin{pmatrix} 1 & & \\ & -1 & \\ & & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & & \\ & 1 & \\ & & -1 \end{pmatrix}$$

is a basis of $\mathfrak{h}$. There are 6 roots vectors

$$E_{12} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

E_{23}, E_{13} , etc.

Exercise 1.2. $[H_1, E_{12}] = 2E_{12}$, $[H_2, E_{12}] = -E_{12}$, $\alpha_{12} = (2, -1)$.

[Drawing of roots]

Notions of *positive roots* and *simple roots* (set of rank $\mathfrak{g}$ simple roots has ℓ elements, where $\ell = \dim(\mathfrak{h}^*)$). (See [Kac10, Definition 17.1].) This will also fail for Kac-Moody algebras more generally). Next write the *Cartan matrix* (see [Kac10, Definition 17.2])

$$A = (a_{ij}), \quad a_{ij} = 2 \frac{(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)}$$

for $1 \leq i, j \leq \ell$.

Example 1.3. $\mathfrak{sl}_3$. [Picture, hexagonal pattern]. $(\alpha_1, \alpha_1) = (\alpha_2, \alpha_2) = 2$, $(\alpha_1, \alpha_2) = -1$, so

$$A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$$

Example 1.4. $\mathfrak{sl}_5$. [Picture, square pattern]. $|\alpha_2| = 1$, $|\alpha_1| = 2$, $(\alpha_1, \alpha_2) = -2$, so

$$A = \begin{pmatrix} 2 & -1 \\ -2 & 2 \end{pmatrix}$$

Remark 1.5. Since $\mathfrak{g}_\alpha$ is 1-dimensional, set $\mathfrak{g}_\alpha = \mathbb{C}E_\alpha$ and $E_i = E_{\alpha_i}$, $i = 1, 2, \dots, \ell$ (simple root vectors). It turns out that

$$\mathrm{ad}(E_i)^{1-a_{ij}} E_j = 0.$$

This is called a *Serre relation*.

2. SOME INFINITE DIMENSIONAL LIE ALGEBRAS

Let $\mathfrak{g}$ be a finite-dimensional semisimple Lie algebra, and define the *loop algebra*

$$\begin{aligned} L\mathfrak{g} &= \mathfrak{g}[t, t^{-1}], \text{ (with basis } at^m | a \in \text{a basis of } \mathfrak{g} \text{ } m \in \mathbb{Z} \text{)} \\ &= \mathfrak{g} \otimes_{\mathbb{C}} \mathbb{C}[t, t^{-1}] \end{aligned}$$

with the Lie bracket

$$[at^m, bt^n] = [a, b]t^{m+n}.$$

“This construction is absurdly general — we don’t need $\mathfrak{g}$ to be semisimple [...]”

Take $\mathfrak{g} = \mathfrak{sl}_2$. Recall that

$$E = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad F = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

[Picture with $F, H, E, Ft, Ht, Et, Et^2 \dots$] E was a root vector, corresponding to the unique root in $\mathfrak{sl}_2$, call it α_1 . We seem to have a second simple root α_0 , corresponding to Ft .

This looks like it wants to have a Cartan matrix

$$A = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$$

We will indeed recover (a variant of) $L\mathfrak{g}$ as a Lie algebra “built from” $A = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$, a Kac-Moody algebra. But note first, $\mathfrak{h} = \mathbb{C}H$ is too small. “Problem with α_0 and α_1 being linearly independent ...”

Exercise 2.1. Consider $L\mathfrak{g} \oplus \mathbb{C}d$, and set $[d, at^m] = mat^m$, $[d, d] = 0$. Check this defines a Lie algebra.

Proof. Skew-commutativity, i.e. for all $x \in L\mathfrak{g} \oplus \mathbb{C}d$,

$$(2.1.1) \quad [x, x] = 0,$$

is immediate from skew commutativity in $L\mathfrak{g}$ and the hypothesis that $[d, d] = 0$.

To confirm Jacobi identity, i.e. that for all $x, y, z \in L\mathfrak{g} \oplus \mathbb{C}d$

$$(2.1.2) \quad [x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0,$$

notice that since this is a cyclic sum on x, y, z we only need to consider three elements in $L\mathfrak{g} \oplus \mathbb{C}d$ up to cyclic permutation. The cases in which the three elements are either in $L\mathfrak{g}$ or in $\mathbb{C}d$ are obvious, so that there are only two interesting possibilities:

$$(2.1.3) \quad x = d, \quad y = at^m, \quad z = bt^n$$

$$(2.1.4) \quad x = d, \quad y = d, \quad z = at^n$$

Case 2.1.3 gives

$$\begin{aligned} & [d, [at^m, bt^n]] + [at^m, [bt^n, d]] + [bt^n, [d, at^m]] \\ &= [d, [a, b]t^{m+n}] + [at^m, -nbt^n] + [bt^n, mat^m] \\ &= (m+n)[a, b]t^{m+n} - n[a, b]t^{m+n} + m[b, a]t^{m+n} \\ &= (m+n)[a, b]t^{m+n} - n[a, b]t^{m+n} - m[a, b]t^{m+n} \\ &= (m+n)[a, b]t^{m+n} - (m+n)[a, b]t^{m+n} = 0. \end{aligned}$$

Case 2.1.4 gives

$$\begin{aligned} & [d, [d, at^m]] + [d, [at^m, d]] + [at^m, [d, d]] \\ &= [d, mat^m] + [d, -mat^m] = 0. \end{aligned}$$

□

The Kac-Moody algebra turns out to be, not quite this, but slightly larger still.

Recall that an *invariant bilinear form* $(\cdot, \cdot)$ on a Lie algebra $\mathfrak{g}$ is a bilinear form such that

$$(2.1.5) \quad ([a, b], c) = (a, [b, c]) \quad \forall a, b, c \in \mathfrak{g}.$$

Exercise 2.2. Prove that an invariant bilinear form on a simple Lie algebra must in fact be symmetric.

Proof. It's enough to show that $\mathfrak{g}$ is *perfect*, i.e. that $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$. In this case, let $a, b \in \mathfrak{g}$ and suppose that $b = [x, y]$. Then

$$\begin{aligned} (a, b) &= (a, [x, y]) = (a, -[y, x]) = (-[a, y], x) = ([y, a], x) \\ &= (y, [a, x]) = (y, -[x, a]) = (-[y, x], a) = ([x, y], a) = (b, a) \end{aligned}$$

To confirm that $\mathfrak{g}$ is perfect just observe that $[\mathfrak{g}, \mathfrak{g}]$ is a nontrivial ideal of $\mathfrak{g}$. $\square$

Definition 2.3. Given $\mathfrak{g}$ simple, with $(\cdot, \cdot) : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$ invariant bilinear form, the *affine Lie algebra* is

$$\hat{\mathfrak{g}} = L\mathfrak{g} \oplus \mathbb{C}K,$$

with $[K, \hat{\mathfrak{g}}] = 0$, and $[at^m, bt^n] = [a, b]t^{m+n} + m(a, b)\delta_{m, -n}K$.

“For the construction to work it doesn't actually have to be nondegenerate.”

Exercise 2.4. Check that the affine Lie algebra $\hat{\mathfrak{g}}$ is a Lie algebra.

Proof. (Skew-commutativity.) Since $[K, \hat{\mathfrak{g}}] = 0$ and $K \in \hat{\mathfrak{g}}$, it is immediate that $[K, K] = 0$. For the case of an element in $L\mathfrak{g}$, we see that $[at^m, at^m] = 0$ by skew-commutativity of the bracket in $\mathfrak{g}$ and the Kronecker delta.

(Jacobi identity.) As in Exercise 2.1, any choice of x, y, z involving K is immediate by $[K, \hat{\mathfrak{g}}] = 0$. Thus the only interesting case is for Jacobi identity consider the cases

$$\begin{aligned} &[at^m, [bt^n, ct^\ell]] + [bt^n, [ct^\ell, at^m]] + [ct^\ell, [at^m, bt^n]] \\ &= [at^m, [b, c]t^{n+\ell} + n(b, c)\delta_{n, -\ell}K] \\ &+ [bt^n, [c, a]t^{\ell+m} + \ell(c, a)\delta_{\ell, -m}K] \\ &+ [ct^\ell, [a, b]t^{m+n} + m(a, b)\delta_{m, -n}K] \\ &= [at^m, [b, c]t^{n+\ell}] + [at^m, n(b, c)\delta_{n, -\ell}K] \\ &+ [bt^n, [c, a]t^{\ell+m}] + [bt^n, \ell(c, a)\delta_{\ell, -m}K] \\ &+ [ct^\ell, [a, b]t^{m+n}] + [ct^\ell, m(a, b)\delta_{m, -n}K] \\ &= [a, [b, c]]t^{m+(n+\ell)} + m(a, [b, c])\delta_{m, -(n+\ell)}K \\ &+ [b, [c, a]]t^{n+(\ell+m)} + n(b, [c, a])\delta_{n, -(\ell+m)}K \\ &+ [c, [a, b]]t^{\ell+(m+n)} + \ell(c, [a, b])\delta_{\ell, -(m+n)}K = 0 \end{aligned}$$

It is clear that we obtain a Jacobi equation on $\mathfrak{g}$. To see that the remaining terms vanish, notice that the condition on the Kronecker delta in its three appearances is the same, namely, $m + n + \ell = 0$. In this case, we only need to check that $(a, [b, c]) = (b, [c, a]) = (c, [a, b])$ to conclude. This follows from the invariance of $(\cdot, \cdot)$ and the fact that $\mathfrak{g}$ simple using Exercise 2.2. $\square$

We also have

Definition 2.5. The *extended affine Lie algebra* is

$$\tilde{\mathfrak{g}} = L\mathfrak{g} \oplus \mathbb{C}K \oplus \mathbb{C}d,$$

with $[d, at^m] = mat^m$ as before, and $[K, d] = 0$.

The extended affine Lie algebra is an example of a Kac-Moody algebra.

Exercise 2.6 (For those who like geometry). Let $R = \mathbb{C}[t, t^{-1}]$. If $D \in \text{Der}(R)$, then $L\mathfrak{g} \oplus \mathbb{C}d$ is a Lie algebra with $[d, a \otimes r] = a \otimes D(r)$. Is $(\mathfrak{g} \otimes R) \oplus \text{Der}(R)$ a Lie algebra? (The Lie algebra $L\mathfrak{g} \oplus \mathbb{C}d$ from Exercise 2.1 is a particular case, for $D = t \frac{d}{dt}$.)

Proof. Checking that $L\mathfrak{g} \oplus \mathbb{C}d$ is a Lie algebra with $[d, a \otimes r] = a \otimes D(r)$ is similar to Exercise 2.1: skew-commutativity is immediate from skew-commutativity in each of the components, while Jacobi identity is verified in two cases. For $x = y = d$ and $z = a \otimes r$ we quickly obtain

$$\begin{aligned} & [x, [y, z]] + [y, [z, x]] + [z, [x, y]] \\ &= [d, [d, a \otimes r]] + [d, [a \otimes r, d]] + [a \otimes r, [d, d]] \\ &= [d, a \otimes D(r)] + [d, -a \otimes D(r)] = 0. \end{aligned}$$

And for $x = d$, $y = a \otimes r$ and $z = b \otimes s$, we get

$$\begin{aligned} & [x, [y, z]] + [y, [z, x]] + [z, [x, y]] \\ (2.6.1) \quad &= [d, [a \otimes r, b \otimes s]] + [a \otimes r, [b \otimes s, d]] + [b \otimes s, [d, a \otimes r]] \\ &= [d, [a, b] \otimes rs] + [a \otimes r, -b \otimes D(s)] + [b \otimes s, a \otimes D(r)] \\ &= [a, b] \otimes D(rs) - [a, b] \otimes rD(s) + [b, a] \otimes sD(r) = 0. \end{aligned}$$

To check whether $(\mathfrak{g} \otimes R) \oplus \text{Der}(R)$ is a Lie algebra first put the Lie bracket on $\text{Der}(R)$ as $[D, D_1] = DD_1 - D_1D$. It is clear that this bracket is skew-commutative. Jacobi identity reads

$$\begin{aligned} & [D, [D_1, D_2]] + [D_1, [D_2, D]] + [D_2, [D, D_1]] \\ &= [D, D_1D_2 - D_2D_1] + [D_1, D_2D - DD_2] + [D_2, DD_1 - D_1D] \\ &= D(D_1D_2 - D_2D_1) - (D_1D_2 - D_2D_1)D + D_1(D_2D - DD_2) \\ &\quad - (D_2D - DD_2)D_1 + D_2(DD_1 - D_1D) - (DD_1 - D_1D)D_2 \\ &= DD_1D_2 - DD_2D_1 - D_1D_2D + D_2D_1D + D_1D_2D - D_1DD_2 \\ &\quad - D_2DD_1 + DD_2D_1 + D_2DD_1 - D_2D_1D - DD_1D_2 + D_1DD_2 = 0. \end{aligned}$$

Now put the bracket on $(\mathfrak{g} \otimes R) \oplus \text{Der}(R)$ as $[D, a \otimes r] = a \otimes D(r)$. Skew-commutativity is immediate. Jacobi identity for $x = D, y = a \otimes r$ and $z = b \otimes s$ is identical to the computation 2.6.1. In the case $x = D, y = D_1$ and $z = a \otimes r$, we get

$$\begin{aligned} & [D, [D_1, a \otimes r]] + [D_1, [a \otimes r, D]] + [a \otimes r, [D, D_1]] \\ &= [D, a \otimes D_1(r)] + [D_1, -a \otimes D(r)] + [a \otimes r, [D, D_1]] \\ &= a \otimes DD_1(r) - a \otimes D_1D(r) - a \otimes [D, D_1](r) = 0 \end{aligned}$$

□

3. KAC-MOODY ALGEBRAS

Recall the notion of the free Lie algebra on a vector space V of generators (or a set X , think of V as a vector space with basis X):

Definition 3.1. The *free Lie algebra* on V is characterized by the universal property

$$\begin{array}{ccc} V & \xrightarrow{f} & \mathfrak{g} \\ & \searrow i & \nearrow \exists! \tilde{f} \\ & F(V) & \end{array}$$

That is, for any linear map $f : V \rightarrow \mathfrak{g}$ with $\mathfrak{g}$ Lie algebra, there exists a unique $\tilde{f}$ homomorphism of Lie algebras $F(V) \rightarrow \mathfrak{g}$ such that $\tilde{f} \circ i = f$.

$$\mathrm{Hom}_{\mathrm{Lie}}(F(V), \mathfrak{g}) = \mathrm{Hom}_{\mathrm{Vec}}(V, \mathfrak{g})$$

naturally.

That is, F and the forgetful functor $G : \underline{\mathrm{Lie}} \rightarrow \underline{\mathrm{Vec}}$ are adjoint:

$$\mathrm{Hom}_{\underline{\mathrm{Lie}}}(F(V), \mathfrak{g}) \xrightarrow{\sim} \mathrm{Hom}_{\underline{\mathrm{Vec}}}(V, G(\mathfrak{g}))$$

A realisation of $F(V)$. Let

$$T(V) = \mathbb{C} \oplus V \oplus V^{\otimes 2} \oplus V^{\otimes 3} \oplus \dots$$

be the tensor algebra of V .

Then inside $T(V)$ consider $F(V)$ the span of iterated commutators of elements of V .

Proposition 3.2. *This realises the free Lie algebra.*

Proof. In online notes. □

In the finite dimensional simple case, we had

$$a_{ij} = \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)},$$

which we think also as $\alpha_i, \alpha_j \in \mathfrak{h}^*$, and $\alpha_i^\vee = \frac{2}{(\alpha_i, \alpha_i)} \nu^{-1}(\alpha_i) \in \mathfrak{h}$.

Clearly, $\alpha_{ii} = 2$ for all i . a_{ij} might not equal a_{ji} , but certainly $a_{ij} = 0 \iff a_{ji} = 0$. And $\forall i \neq j$, $a_{ij} \leq 0$.

One further property. Set

$$\varepsilon_i = \frac{2}{(\alpha_i, \alpha_i)}, \quad \text{and} \quad D = \begin{array}{c} \text{diagonal matrix} \\ \text{with entries } \varepsilon_i \end{array}$$

Then $A = DB$, where $B = ((\alpha_i, \alpha_j))$ is symmetric. If a matrix A is equal to $(\mathrm{diag})(\mathrm{symm})$, we call it *symmetrizable*.

Definition 3.3. A *generalized Cartan matrix* is an integer matrix $A = (a_{ij})$ which is

- symmetrizable,
- $a_{ii} = 2$ for all i ,
- $a_{ij} = 0 \iff a_{ji} = 0$,
- $a_{ij} \leq 0$ for $i \neq j$.

Definition 3.4. A *realisation* of a generalized Cartan matrix is a complex vector space $\mathfrak{h}$, and two sets

$$\begin{aligned}\Pi^\vee &= \{\alpha_1^\vee, \alpha_2^\vee, \dots, \alpha_n^\vee\}, \quad \text{and,} \\ \Pi &= \{\alpha_1, \alpha_2, \dots, \alpha_n\}\end{aligned}$$

such that $\langle \alpha_i^\vee, \alpha_j \rangle = a_{ij}$, $1 \leq i, j \leq n$.

Exercise 3.5. $\dim(\mathfrak{h}) \geq 2n - \text{rank}(A)$.

Proof. For $A = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$, a realisation is given by

$$\Pi^\vee = \{H_1, H_0\}, \quad \Pi = \{\alpha_0, \alpha_1\}$$

$$\mathfrak{h} = \mathbb{C}H, \mathbb{C}d, \mathbb{C}K,$$

$$\mathfrak{h}^* = \mathbb{C}\alpha_1 + \mathbb{C}\delta + \mathbb{C}\Lambda_0$$

(Canonical dual, $\langle \alpha_1, H \rangle = 2$, $\langle \delta, d \rangle = 1 = \langle \Lambda_0, K \rangle$, every other pairing 0.)

Then

$$\begin{cases} \alpha_1 = \alpha_1 \\ \alpha_0 = \delta - \alpha_1 \end{cases} \quad \begin{cases} \alpha_1^\vee = H \\ \alpha_0^\vee = K - H \end{cases}$$

So we obtain

$$\begin{aligned}\langle \alpha_0^\vee, \alpha_1 \rangle &= \langle K - H, \alpha_1 \rangle = 2 \\ \langle \alpha_1^\vee, \alpha_0 \rangle &= \langle H, \delta - \alpha_1 \rangle = -2 \\ \langle \alpha_0^\vee, \alpha_0 \rangle &= \langle K - H, \delta - \alpha_1 \rangle = +2\end{aligned}$$

□

Finally let's define Kac-Moody algebras.

Let A be a generalized Cartan matrix. Let

$$\tilde{\mathfrak{n}}_+ = F(e_1, \dots, e_n),$$

the free Lie algebra on n generators, and similarly

$$\tilde{\mathfrak{n}}_- = F(f_1, \dots, f_n).$$

Let $\mathfrak{h}$ be a realisation of A . Set $\tilde{\mathfrak{g}}(A) = \tilde{\mathfrak{n}}_- \oplus \mathfrak{h} \oplus \tilde{\mathfrak{n}}_+$.

Make $\tilde{\mathfrak{g}}(A)$ a Lie algebra by defining

- $[\mathfrak{h}, \mathfrak{h}] = 0$,
- $\forall H \in \mathfrak{h}, [H, e_i] = \langle \alpha_i, H \rangle e_i = \alpha_i(H)e_i$. And similarly, $[H, f_i] = -\alpha_i(H)f_i$.
- $[e_i, f_j] = \delta_{ij}\alpha_i^\vee$.

Then $\tilde{\mathfrak{g}}(A)$ is a Lie algebra (though not yet the Kac-Moody algebra). See Kac, [Kac90, Theorem 1.2].

Remark 3.6. In $\mathfrak{h}$ we have a lattice

$$Q^\vee = \mathbb{Z}\alpha_1^\vee + \dots + \mathbb{Z}\alpha_n^\vee, \quad \text{and}$$

$$Q = \mathbb{Z}\alpha_1 + \dots + \mathbb{Z}\alpha_n \text{ in } \mathfrak{h}^*$$

(root and coroot lattices). $\tilde{\mathfrak{g}}(A)$ is naturally Q -graded, with

$$\tilde{\mathfrak{g}}(A)_\beta = \text{span}\{\text{commutators of } e_i \text{ with } \sum \alpha_i = \beta\}.$$

$$\tilde{\mathfrak{g}}(A) = \mathfrak{h}.$$

Theorem 3.7 (Gabber-Kac). *Denote by $I \subset \tilde{\mathfrak{g}}(A)$ the maximal Q -graded ideal, such that $I \cap \mathfrak{h} = \{0\}$. Then I is generated by the Serre relations*

$$\text{ad}(e_i)^{1-a_{ij}} e_j \quad \text{and} \quad \text{ad}(f_i)^{1-a_{ij}} f_j, \quad i \neq j.$$

Proof. [Kac90, Theorem 9.11]. $\square$

(The existence of the ideal I does not need the theorem; the importance of the theorem is providing an expression for the generators.)

Definition 3.8. The Kac-Moody algebra $\mathfrak{g}(A)$ is $\tilde{\mathfrak{g}}(A)/I$.

4. AFFINE KAC-MOODY ALGEBRAS

Let $\mathfrak{g}$ be a finite-dimensional simple Lie algebra, with $(\cdot, \cdot) : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$ invariant bilinear form,

$$([x, y], z) = (z, [y, z]) \quad \forall x, y, z \in \mathfrak{g}$$

(Eg. the Killing form $\kappa(x, y) = \text{Tr}_{\mathfrak{g}} \text{ad}(x) \text{ad}(y)$ is invariant.)

Typically we normalise $(\cdot, \cdot)$ so that $(\alpha, \alpha) = 2$ for the long roots of $\mathfrak{g}$.

Then $\hat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}] \oplus \mathbb{C}K$ (affine Lie algebra),

$$[at^m, bt^n] = [a, b]t^{m+n} + m\delta_{m, -n}(a, b)K, \quad [K, \hat{\mathfrak{g}}] = 0$$

and $\tilde{\mathfrak{g}} = \hat{\mathfrak{g}} \oplus \mathbb{C}d$, $[d, K] = 0$, $[d, at^m] = mat^m$, (affine Kac-Moody algebra or “extended affine Lie algebra”)

Theorem 4.1. $\tilde{\mathfrak{g}}$ is a Kac-Moody algebra.

Let $\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_{\alpha}$, ($\mathfrak{g}_{\alpha} = \mathbb{C}E_{\alpha}$.)

The simple roots and coroots. $\tilde{\mathfrak{h}} = \mathfrak{h} \oplus \mathbb{C}K \oplus \mathbb{C}d$. We identify $\tilde{\mathfrak{h}}^*$ with $\mathfrak{h}^* \oplus \mathbb{C}\Lambda_0 \oplus \mathbb{C}\delta$ where

$$\Lambda_0(\mathfrak{h}) = \delta(\mathfrak{h}) = 0$$

$$\Lambda_0(d) = \delta(K) = 0$$

$$\Lambda_0(K) = \delta(d) = 1$$

The *real coroots* are

$$\hat{\Delta}^{V, re} = \{E_{\alpha}t^m | \alpha \in \Delta, m \in \mathbb{Z}\}$$

and there are also imaginary roots and coroots

$$\hat{\Delta}^{V, im} = \{Ht^m | H \in \mathfrak{h}, m \in \mathbb{Z} \setminus \{0\}\}$$

Roots:

$$\hat{\Delta}^{re} = \{\alpha + m\delta | \alpha \in \Delta, m \in \mathbb{Z}\}$$

$$\hat{\Delta}^{im} = \{m\delta | m \neq 0\}$$

Xt^m :

$$[H, Xt^m] = [H, x]t^m, \quad H \in \mathfrak{h}$$

$$[K, xt^m] = 0$$

$$[d, xt^m] = mxt^m$$

so it $x \in \mathfrak{g}_{\alpha}$, $xt^m \in \tilde{\mathfrak{g}}_{\alpha+m\delta}$.

The invariant bilinear form $(\cdot, \cdot)$ from $\mathfrak{g} \times \mathfrak{g}$ extends uniquely to $(\cdot, \cdot) : \tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}} \rightarrow \mathbb{C}$.

$(d, d) = (K, K) = 0$, $(d, K) = 1$ and $(d, \mathfrak{h}) = (K, \mathfrak{h}) = 0$.

So, in $\tilde{\mathfrak{h}}^*$:

$$\begin{aligned} (\Lambda_0, \Lambda_0) &= (\delta, \delta) = 0 \\ (\Lambda_0, \mathfrak{h}^*) &= (\delta, \mathfrak{h}^*) = 0 \\ (\Lambda_0, \delta) &= 1. \end{aligned}$$

Hence, $|\alpha + m\delta|^2 = |\alpha|^2$, $|m\delta|^2 = 0$.

Example 4.2. $\widetilde{\mathfrak{sl}}_2, \tilde{\mathfrak{h}}^* = \text{span}\{\alpha, \Lambda_0, \delta\}$ with Gram matrix ...

We can make a choice of positive roots,

$$\hat{\Delta}_+ = \{\alpha + m\delta | \alpha \in \Delta, m > 0\} \cup \{m\delta | m > 0\} \cup \Delta_+$$

Obviously, if $\alpha \in \Delta_+$ is simple, $\alpha \in \hat{\Delta}_+$ is simple.

Notation. Let $\theta \in \Delta_+$ be the highest root. ($\nexists \alpha \in \Delta_+$ such that $\alpha - \theta \in \mathbb{Z}_+ \Delta_+$.) and $\alpha = \delta - \theta$.

Then $\alpha_0 \in \hat{\Delta}_+$ is simple and the set of simple roots is $\hat{\Pi} = \{\alpha_0, \underbrace{\alpha_1, \dots, \alpha_\ell}_{\text{the finite simple roots}}\}$.

where $\ell = \text{rank}(\mathfrak{g})$.

The coroot corresponding to α_0 is

$$\alpha_0^\vee = K - \theta^\vee, \quad \theta^\vee = \frac{2}{(\theta, \theta)} \nu^{-1}(\theta) \in \mathfrak{h}$$

$$\text{and} \quad E_{\alpha_0} = E_{-\theta}t.$$

5. WEYL GROUP

Upshot. The Weyl group is a semidirect product of pseudoreflections and translations.

In any Kac-Moody algebra, we have

$$\begin{aligned} \text{roots} \quad \Pi &= \{\alpha_1, \dots, \alpha_\ell\} \subset \mathfrak{h}^* \\ \text{coroots} \quad \Pi^\vee &= \{\alpha_1^\vee, \dots, \alpha_\ell^\vee\} \subset \mathfrak{h}, \end{aligned}$$

and *reflections* $r_i \in \text{GL}(\mathfrak{h}^*)$, defined by

$$r_i(\lambda) = \lambda - \langle \lambda, \alpha_i^\vee \rangle \alpha_i.$$

One can check that

$$(r_i \lambda, r_i \mu) = (\lambda, \mu) \quad \forall \lambda, \mu \in \mathfrak{h}^*$$

The *Weyl group* W is $\langle r_i | i = 1, \dots, \ell \rangle \subset \text{GL}(\mathfrak{h}^*)$.

Example 5.1. For $\widetilde{\mathfrak{sl}}_2$, r_1 is easy,

$$\begin{aligned} r_1(\alpha) &= -\alpha \quad (\text{as in } \mathfrak{sl}_2) \\ r_1(\delta) &= \delta, \quad r_1(\Lambda_0) = \Lambda_0. \end{aligned}$$

To compute r_0 take an arbitrary element $m\alpha_1 + k\Lambda_0 + f\delta$ and do:

$$\begin{aligned} r_0(m\alpha_1 + k\Lambda_0 + f\delta) &= m\alpha_1 + k\Lambda_0 + f\delta - \langle \alpha_0^\vee, m\alpha_1 + k\Lambda_0 + f\delta \rangle \alpha_0 \\ \alpha_0 &= \delta - \alpha_1, \quad \alpha_0^\vee = K - \alpha^\vee \end{aligned}$$

so we obtain

$$\begin{aligned} &= m\alpha_1 + k\Lambda_0 + f\delta - (k - 2m)(\delta - \alpha_1) \\ &\quad (k - m)\alpha_1 + k\Lambda_0 + (f - k + 2m)\delta. \end{aligned}$$

Relative to basis $\{\alpha_1, \Lambda_0, \delta\}$.

$$\begin{aligned} r_1 &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, r_0 = \begin{pmatrix} -1 & 1 & 0 \\ 0 & 1 & 0 \\ 2 & -1 & 1 \end{pmatrix} \begin{pmatrix} m \\ k \\ f \end{pmatrix} \\ t = r_1 r_0 &= \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 2 & -1 & 1 \end{pmatrix} \end{aligned}$$

Notice that δ is fixed by all r_i . Also $m\alpha + k\Lambda_0 + f\delta$, the *coefficient* of Λ_0 is fixed by all r_i .

Then

$$t(m\alpha_1 + k\Lambda_0 + f\delta) = (m - k)\alpha_1 + k\Lambda_0 + (f - k + 2m)\delta.$$

Think of t as a translation.

The number k in

$$\mathfrak{h}^* \ni \hat{\lambda} = \lambda + k\Lambda_0 + f\delta$$

is called the *level* of $\hat{\lambda}$.

$\hat{\mathfrak{h}}$ = union of (hyper)planes of constant level which are stable under W . The roots α are all of level 0.

[Picture] “ r_1 changes the sign of the finite path”. And $t = r_1 r_0$ is a sort of translation. Indeed, in general we can consider $t_{\alpha_i} = r_{\alpha_i} \circ r_0 \in W$,

$$t_{\alpha}(\beta + m\delta) = \beta + (m + (\beta, \alpha_i))\delta$$

One can describe the action of t_{α} on $\hat{\lambda}$ in general (e.g. see [Kac90, Chapter 6])

Proposition 5.2. *For the affine Kac-Moody algebra $\hat{\mathfrak{g}}$ with $\hat{W} = \langle r_0, r_1, \dots, r_{\ell} \rangle$ its Weyl group (and $W = \langle r_1, \dots, r_{\ell} \rangle \subset \hat{W}$ the Weyl group of $\mathfrak{g}$), then $\hat{W} \simeq W \times t_{Q^{\vee}}$ (where it should be semidirect product instead of $\times \dots$) where $Q^{\vee}$ is the coroot lattice of $\mathfrak{g}$.*

Remark 5.3. For general Kac-Moody algebras, the Weyl groups are much larger, hyperbolic reflection groups.

In the affine case, $\hat{W}$ fixes level k , and $|\hat{\lambda}|$. One gets, in the intersection, paraboloids [Picture of section of hyperboloid that is a parabola].

6. WEYL CHARACTER FORMULA

Highest weight representations of Kac-Moody algebras. Let $\lambda \in \mathfrak{h}^*$, where $\mathfrak{g}(A) = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ is a Kac-Moody algebra. We define a *Verma module*

$$M(\Lambda) = U(\mathfrak{g}) \otimes_{U(\mathfrak{h} + \mathfrak{n}_+)} \mathbb{C}v_{\Lambda}$$

where $\mathfrak{h} + \mathfrak{n}_+$ acts on V_{Λ} by:

$$Xv_{\Lambda} = 0, \quad \forall x \in \mathfrak{n}_+, Hv_{\Lambda} = \Lambda(H)v_{\Lambda}, \quad \forall H \in \mathfrak{h}$$

So $\mathbb{C}v_\Lambda$ is a $U(\mathfrak{h} + \mathfrak{n}_+)$ -module,

$$\begin{array}{c} U(\mathfrak{h} + \mathfrak{n}_+) \\ \downarrow \\ U(\mathfrak{g}) \end{array}$$

By the PBW theorem, $M(\Lambda)$ has a linear $\mathbb{C}$ -basis.

Let $\{F_{\alpha,i} : i = 1, \dots, \dim \mathfrak{g}_\alpha\}$ be a basis of $\mathfrak{g}_{-\alpha}$, $\forall \alpha \in \Delta_+$. Also choose a total order on Δ_+ . (Some sort of lexicographical order that takes longer to write than to say.)

$$F_{\alpha_1, i_1}, F_{\alpha_2, i_2}, \dots, F_{\alpha_s, i_s}, v_\Lambda$$

$$\alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_s \text{ and if } \alpha_p = \alpha_{p+1}, i_p \leq i_{p+1}$$

We have $M(\Lambda)_\lambda = \{m | Hm = \lambda(H)m\}$ weight spaces.

$$M(\Lambda) = \bigoplus_{\lambda \in \mathfrak{h}^*} M(\Lambda)_\lambda$$

The vector v_Λ is in $M(\Lambda)_\Lambda$ by definition,

$$F_{\alpha,i} v_\Lambda \in M(\Lambda)_{\Lambda - \alpha}$$

$$H(Fv_\Lambda) = \underbrace{[H, F]v_\Lambda}_{= -\alpha(H)Fv_\Lambda} + \underbrace{FHv_\Lambda}_{=\Lambda(H)Fv_\Lambda}$$

So $\chi_{M(\Lambda)} = \sum_{\lambda \in \mathfrak{h}^*} \dim M(\Lambda)_\lambda e^\lambda$ is computed by counting monomials y with fixed $\sum_i \alpha_i$.

$$(6.0.1) \quad \chi_{M(\Lambda)} = e^\Lambda \prod_{\alpha \in \Delta_+} \frac{1}{(1 - e^{-\alpha})^{\dim \mathfrak{g}_\alpha}}.$$

The product on Eq. 6.0.1 is called *Weyl denominator*.

Exercise 6.1. Convince yourself of this.

Example 6.2. $\mathfrak{g} = \mathfrak{sl}_2$, [Picture]

$$\begin{aligned} \chi_{M(\Lambda)} &= e^\Lambda + e^{\Lambda - \alpha} + e^{\Lambda - 2\alpha} + \dots \\ &= e^\Lambda (1 + e^{-\alpha} + e^{-2\alpha} + \dots) \\ &= e^\Lambda \frac{1}{1 - e^{-\alpha}}. \end{aligned}$$

For certain Λ , $M(\Lambda)$ is *reducible* (i.e. there exists a submodule $0 \neq N \subset M(\Lambda)$ (with proper contention).

Lemma 6.3. For any submodule N ,

$$N = \bigoplus_{\mu \in \mathfrak{h}^*} N \cap M(\Lambda)_\mu.$$

Corollary. The sum of all proper submodules of $M(\Lambda)$ is proper, in particular there is a maximal proper submodule.

Notation. $L(\Lambda) = M(\Lambda) / \left(\begin{smallmatrix} \text{max. proper} \\ \text{submodule} \end{smallmatrix} \right)$

Example 6.4. $\mathfrak{sl}_2$. $\Lambda = 3\omega$ (ω : fundamental weight, $\alpha = 2\omega$.) $L(3\omega) = \mathbb{C}\langle e^{3\omega}, e^{-\omega}, e^{-3\omega} \rangle$.
[Picture]

Definition 6.5. A $\mathfrak{g}$ -module is *integrable* if

- $V = \bigoplus_{\mu \in \mathfrak{h}^*} V_\mu$ (weight module).
- For all simple roots α_i ; e_i and f_i are locally nilpotent on V (i.e. for all $v \in V$ there exists N such that $e_i^N v = f_i^N v = 0$.)

Remark 6.6. • Vermas are not integrable.

- $\dim V < \infty \implies V$ integrable.
- $\mathfrak{g}$ itself (Kac-Moody) is integrable.

Dominant integrable weights. Let $\{\alpha_1^\vee, \dots, \alpha_\ell^\vee\} \subset \mathfrak{h}$ be the simple coroots.

Definition 6.7. The *dominant integral weights* are the weights that pair with the coroots to give integers:

$$P_+ = \{\lambda \in \mathfrak{h}^* : \langle \lambda, \alpha_i^\vee \rangle \in \mathbb{Z}_{\geq 0}, i = 1, \dots, \ell\}.$$

For $L(\Lambda)$ to be integrable, it is necessary that $\Lambda \in P_+$.

Indeed, suppose $L(\Lambda)$ is integrable. Then $f_i^N v_\Lambda = 0$ in $L(\Lambda)$, or rather

$$\underbrace{e_i f_i^{N+1} v_\Lambda}_{K f_i^N = 0} \in M(\Lambda),$$

and K can only be zero if $\langle \Lambda, \alpha_i^\vee \rangle \in \mathbb{Z}_{\geq 0}$. Applying for all i , we find $\Lambda \in P_+$ is necessary.

Proposition 6.8. $L(\Lambda)$ is integrable if and only if $\Lambda \in P_+$.

Proof. For the converse, use induction and Serre relations (we know the result for the highest weight, and want to prove for others). $\square$

Example 6.9. $\mathfrak{sl}_3$. [Picture]

Example 6.10. $\widehat{\mathfrak{sl}_2}$. [Picture, P_+ looks like diagonal lines.]

$$\alpha_0^\vee = K - H, \quad \alpha_1^\vee = H \in \mathfrak{sl}_2, \quad \langle \delta, \alpha_i^\vee \rangle = 0, \quad i = 0, 1$$

Remark 6.11. For affine Kac-Moody algebras, almost nothing about the structure of $M(\Lambda)$ depends on the coefficient of δ in Λ . So it's common to consider

$$M(\Lambda) = M_k(\lambda) = M(k\Lambda_0 + \lambda), \quad \lambda \in \mathfrak{h}^*$$

where k , the level of Λ , is super important.

Then

$$\underbrace{\hat{P}_+}_{\substack{\delta\text{-coef.} \\ = 0}} = \bigcup_{k \in \mathbb{Z}_{\geq 0}} \{k\Lambda_0 + \lambda \mid \lambda \in P_+^k\} \{k\Lambda_0 + \lambda \mid \lambda \in P_+^k\}.$$

$$P_+^k = \{\lambda \in P_+ \mid \langle \lambda, \theta \rangle \leq k\} \subset P_+ \text{ for } \mathfrak{g}.$$

Consider $V = \bigoplus_{\mu \in \mathfrak{h}^*} V_\mu$ integrable, and $V_\lambda \neq 0$. Let $i \in \{1, \dots, \ell\}$. Consider $U = \bigoplus_{n \in \mathbb{Z}} V_{\lambda + n\alpha_i} \subset V$, and the action of e_i, f_i and $h_i = [e_i, f_i]$.

$\mathfrak{sl}_2 \curvearrowright U$, locally integrable. By structure of $\mathfrak{sl}_2$ -representations, U must be finite-dimensional with “symmetrical” weight space multiplicities, i.e.,

$$\{\lambda + n\alpha_i \mid n \in \mathbb{Z}\} \cap \{\text{weights of } V\} = \{\lambda + n\alpha_i \mid -p \leq n \leq q\}.$$

and

$$\langle \lambda - p\alpha_i, h_i \rangle = -\langle \lambda + q\alpha_i, h_i \rangle.$$

Consequently, the reflection $r_i(\lambda) := \lambda - \langle \lambda, \alpha_i^\vee \rangle \alpha_i$ has the same multiplicity as λ .

[Picture]

Now consider $M(\lambda)$ and $L(\Lambda)$ for $\Lambda \in P_+$. Actually, for general Λ , $M(\Lambda)$, while not necessarily irreducible, has an (Ω) -composition series* by irreducibles $L(\lambda)$.

$$M(\Lambda) = V_0 \supset V_1 \supset V_2 \supset \dots \supset V_n = 0,$$

such that V_i/V_{i+1} is $\simeq L(\lambda_i)$ (i.e. an irreducible highest weight module) for some $\lambda_i = \Lambda - \beta_i$, $\beta_i \in Q$. For Kac-Moody algebras we also consider the case that $(V_i/V_{i+1}) = 0$ for all $\mu \in \Omega + Q_+$, (which is the Ω -composition series).

[Picture.]

For an Ω -composition series, as above, we have

$$\text{ch}_{M(\Lambda)} = \sum_{\lambda \geq \Omega} \underbrace{[M(\Lambda) : L(\lambda)]}_{\substack{\# \text{ of times } L(\lambda) \\ \text{appears in the compos. series}}} \text{ch}_{L(\lambda)} \in \langle e^\mu : \mu \not\geq \Omega \rangle$$

Sending “ $\Omega \rightarrow -\infty$ ”, the identity

$$\text{ch}_{M(\Lambda)} = \sum_{\lambda \leq \Lambda} [M(\Lambda) : L(\lambda)] \text{ch}_{L(\lambda)}$$

makes sense.

Notation. $b_{\Lambda, \lambda} = [M(\Lambda) : L(\lambda)]$.

Remark 6.12. Recall the partial order on weights that $\lambda \leq \Lambda$ if $\Lambda - \lambda \in Q = \sum \mathbb{Z}_+ \alpha_i$. $b_{\Lambda, \lambda} = 1$ if $\lambda = \Lambda$ and $b_{\Lambda, \lambda} = 0$ if not ($\lambda \leq \Lambda$).

If we choose a total order on $\mathfrak{h}^*$, compatible with $\leq$. Then $\{b_{\Lambda, \lambda}\}$ is a lower triangular matrix with 1 on the diagonal. We can define $\{m_{\Lambda, \lambda}\}$ the *inverse matrix*. It's again lower triangular, 1 on the diagonal, and all $m_{\Lambda, \lambda}$ are integers (maybe negative now). And we have

$$(6.12.1) \quad \text{ch}_{L(\Lambda)} = \sum_{\lambda \leq \Lambda} m_{\Lambda, \lambda} \text{ch}_{M(\lambda)}.$$

Example 6.13. $\mathfrak{sl}_2$. $M(3) \underset{L(3)}{\supset} M(-5) \underset{L(-5)}{\supset} 0$. Since $M(-5)$ is already irreducible.

[Missing...]

We want to discover $m_{\Lambda, \lambda}$. In general massively difficult. For $\Lambda \in P_+$, $m_{\Lambda, \lambda}$ easy. Multiply Eq. 6.12.1 by R

$$R \text{ch}_{L(\Lambda)} = \sum_{\lambda \geq \Lambda} m_{\Lambda, \lambda} e^\lambda \cdot e^\rho R \text{ch}_{L(\Lambda)} = \sum_{\lambda \leq \Lambda} m_{\Lambda, \lambda} e^{\lambda + \rho}.$$

What's ρ ? It's $\rho \in \mathfrak{h}^*$ chosen so that $\langle \rho, \alpha_i^\vee \rangle = 1$, and it's called the *Weyl vector*.

Remark 6.14. For $\mathfrak{g}$ finite-dimensional, $\rho = \sum_{i=1}^\ell \omega_i$ necessarily. (And equals $\frac{1}{2} \sum_{\alpha \in \Delta_+} \alpha$.)

For $\mathfrak{g}$ finite dimensional and the affine $\hat{\mathfrak{g}}$, $\hat{\rho} = h^\vee \Lambda_0 + \rho$ works.

For $w \in W = \langle r_i | i = 1, \dots, \ell \rangle$, define $w(\text{ch}_V) = \sum \dim V_\mu e^{W(\mu)}$. We saw $w(\text{ch}_V) = \text{ch}_V$ if V integrable. In particular $w(\text{ch}_{L(\Lambda)}) = \text{ch}_{L(\Lambda)}$, $\Lambda \in P_+$.

Lemma 6.15. $m_{\Lambda, \lambda} = 0$ unless $\lambda + \rho = w(\Lambda + \rho)$ for some $w \in W$

Claim. $r_i(e^\rho R) = -e^\rho R$. So $w(e^\rho R) = \det(w)e^\rho R$ for all $w \in W$.

Proof.

$$R = \prod_{\alpha \in \Delta_+} (1 - e^{-\alpha})^{\text{mult}(\alpha)}.$$

Note that

- (1) $\text{mult}(r_i(\alpha)) = \text{mult}(\alpha)$ for all $\alpha \in \Delta$. (Since $\mathfrak{g}$ is integrable!)
- (2) $r_i(\Delta_+) = \{-\alpha_i\} \cup (\Delta_+ \setminus \{\alpha_i\})$.

Any $\alpha \in \Delta_+$ is of the form $\alpha = \sum_{i=1}^\ell k_i \alpha_i$. If $\alpha \neq \alpha_i$, some $k_{j_0} \neq 0$, $j_0 \neq i$ and

$$\begin{aligned} r_i(\alpha) &= \sum_j k_j \alpha_j \langle \alpha, \alpha_i^\vee \rangle \alpha_i \\ &= \sum_j k'_j \alpha_j \\ &= e^\rho (e^{-\alpha_i} - 1) \left(\prod_{\alpha \in \Delta_+ \setminus \alpha_i} \right) \\ &= -e^\rho R. \end{aligned}$$

for $k'_{j_0} = k_{j_0} > 0$. Can't have a mixture of signs, so $r_i(\alpha) \in \Delta_\perp$.
So

$$r_i(e^\rho R) = \prod_{\alpha \in \Delta_+ \setminus \alpha_i} (1 - e^{-\alpha})^{\text{mult}(\alpha)} \cdot (1 - e^{t\alpha_i} \cdot e^{\rho - \alpha_i})$$

$$r_i = \rho - \langle \alpha_i^\vee, \rho \rangle \alpha_i = \rho - \alpha_i$$

Hence

$$\sum_{\lambda \leq \Lambda} m_{\Lambda, \lambda} e^{\lambda + \rho}$$

is W -skew-invariant. (Lemma 6.15.)

Hence

$$\sum_{\lambda \leq \Lambda} m_{\Lambda, \lambda} e^{\lambda + \rho} = \sum_{w \in W} \det(w) \cdot e^{w(\Lambda + \rho)}$$

In conclusion, the *Weyl character formula*

$$\text{ch}_{L(\Lambda)} = \sum_{w \in W} \det(w) \cdot \frac{e^{w(\Lambda + \rho) - \rho}}{R}$$

□

Corollary (Weyl denominator formula).

$$e^\rho R = \sum_{w \in W} \det(w) e^{w(\rho)}.$$

Next time: Affine case, θ -functions. Modular forms (Poisson summation.)

7. CHARACTERS OF INTEGRABLE HIGHEST WEIGHT FOR AFFINE KAC-MOODY ALGEBRAS

Can be calculated using the Weyl character formula.

Recall: $\hat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}] \oplus \mathbb{C}K \oplus \mathbb{C}d$.

Dual Cartan $\hat{\mathfrak{h}}^* = \mathfrak{h}^* \oplus \mathbb{C}\Lambda_0 \oplus \mathbb{C}\delta$.

We consider weights $\Lambda = k\Lambda_0 + \lambda$, $\lambda \in \mathfrak{h}^*$, $k \in \mathbb{Z}_+$, $\lambda \in P_+^k$.

Simple roots: $\alpha_0 = \delta - \theta$, $\{\alpha_1, \dots, \alpha_\ell\} \subset \mathfrak{h}^*$.

The affine Weyl group is (the *semidirect* product!)

$$\widehat{W} = \langle r_0, \dots, r_\ell \rangle \cong W \times T$$

where $T = \{t_\alpha | \alpha \in Q\}$, where t_α is the translation

$$t_\alpha(\Lambda) = \Lambda + k\alpha - \left((\lambda, a) + k \frac{|\alpha|^2}{2} \right) \delta$$

$$\Lambda = k\Lambda_0 + la$$

Let's compute $\chi_{L(\Lambda_0)}$ using the Weyl character formula

$$\chi_{L(\Lambda)} = \frac{\sum_{w \in \widehat{W}} \varepsilon(w) e^{W(\Lambda + \rho) - \rho}}{R}$$

Firstly $R = \prod_{\alpha \in \hat{\Delta}} (1 - e^{-\alpha})^{\text{mult}(\alpha)}$

Recall

$$\begin{aligned} \hat{\Delta}_+ &= \{m\delta | m \in \mathbb{Z}_{\geq 1}\} \\ &\cup \{\alpha_1 + m\delta | m \in \mathbb{Z}_{\geq 0}\} \\ &\cup \{-\alpha_1 + m\delta | m \in \mathbb{Z}_{\geq 1}\} \end{aligned}$$

So let's write $e^{m\delta} = q^m$ (i.e. $q = e^{-\delta}$, is a symbol) and $y = e^{\alpha_1}$.

So

$$R = \prod_{n=1}^{\infty} \underbrace{(1 - y^{-1}q^{n-1})}_{\alpha + (n-1)\delta} \underbrace{(1 - q^n)}_{n\delta} \underbrace{(1 - yq^n)}_{-\alpha + n\delta}$$

Now let's express numerator also in terms of q and y

$$\hat{\rho} = h^\vee \Lambda_0 + \rho$$

$h^\vee$ dual Coxeter number for $\mathfrak{g}$. For $\mathfrak{g} = \mathfrak{sl}_2$, $h^\vee = 2$. (For $\mathfrak{g} = \mathfrak{sl}_n$, $h^\vee = n$ and $\mathfrak{g} = E_8$, $h^\vee = 30$.)

For $\mathfrak{sl}_2$, $\rho = \frac{1}{2}\alpha_1 = \omega_1$.

$\Lambda = \Lambda_0$, $\Lambda + \rho = 3\Lambda_0 + \omega_1$.

Using the formula, we find

$$\begin{aligned} t_{m\alpha_1}(3\Lambda_0 + \omega_1) - (\Lambda_0 + \omega_1) = \\ \dots, \Lambda_0 - 3\alpha, -2\delta, \Lambda_0, \Lambda_0 + 3\alpha_1 - 4\delta, \dots \end{aligned}$$

[Picture]

Notation. $w(\Lambda + \rho) := w \circ \Lambda$.

One finds

$$\sum_{w \in \widehat{W}} \varepsilon(w) e^{w \circ \Lambda_0} = e^{\Lambda_0} \left(\underbrace{1 + y^3 q^4 + y^{-3} q^2 + \dots}_{+ \text{ signs because } w \in T} \underbrace{-y^{-1} - y^2 q^2 - \dots}_{- \text{ signs because } w \in T_\sigma} \right)$$

We find explicitly

$$\chi_{L(\Lambda_0)} = e^{\Lambda_0} \frac{\sum_{m \in \mathbb{Z}} y^{3m} q^{3m^2+m} - \sum_{m \in \mathbb{Z}} y^{3m-1} q^{3m^2-m}}{\prod_{n=1}^{\infty} (1 - y^{-1} q^{n-1} (1 - q^n) (1 - y q^n))}$$

Exercise 7.1. Put this formula in mathematica and confirm that [Picture]

$$\chi_{L(\Lambda_0)} = e^{\Lambda_0} (1 + q(y^{-1} + 1 + y) + q^2(y^{-1} + 2 + y) + (q^3(2y^{-1} + 3 + 2y) + \dots))$$

Appears that the central column here are *partitions*, i.e. $p(n) = \#$ of partitions of n . To see why recall the generating function $\sum_{n=0}^{\infty} q^n p(n) = \prod_{k=1}^{\infty} \frac{1}{1-q^k}$, and expand.

It would appear that

$$\chi_{L(\Lambda_0)} = \left(\prod_{k=1}^{\infty} \frac{1}{1-q^k} \cdot \sum_{m \in \mathbb{Z}} y^m q^{m^2} \right)$$

This identity is true, and we will see it has a vertex-algebra interpretation.

Remark 7.2. If we compute $L(0) = 1$ using the formula, we obtain

$$\begin{aligned} \prod_{n=1}^{\infty} &= (1 - y q^{n-1})(1 - q^n)(1 - y^{-1} q^n) \\ &= \sum (\text{exercise}) \end{aligned}$$

This identity is called the *Jacobi triple product identity*.

These functions

$$\sum_{n \in \mathbb{Z}} q^{n^2}, \sum_{n \in \mathbb{Z}} y^n q^{n^2}, \sum_{n \in \mathbb{Z}} y^{3n} q^{3n^2+n}, \text{ etc. } \dots$$

are all examples of θ -functions.

Remark 7.3. In the formula for $\chi_{L(\Lambda_0)}(y, q)$, we could put $y = 1$ to get

$$\begin{aligned} \chi_{L(\Lambda_0)}(q) &= 1 + 3q + 4q^2 + 7q^3 + \dots \\ &= \prod \frac{1}{1-q^k} \cdot \sum_{m \in \mathbb{Z}} q^{m^2} \end{aligned}$$

If one looks at $L(\Lambda_0 + \omega_1)$ (the other $\Lambda \in P_+^1$),

$$\chi_{L(\Lambda+\omega_1)} = \prod \frac{1}{1-q^k} \sum_{m \in \mathbb{Z}} q^{(m+1/2)^2 - 1/4}$$

8. θ -FUNCTIONS

Let's consider

$$\theta(\tau) = \sum_{n \in \mathbb{Z}} q^{n^2/2}, \quad q = e^{2\pi i \tau}$$

This converges (absolutely on compact regions in) the domain $\text{Im}(\tau) > 0$.

Consider the Fourier transform

$$\hat{g}(y) = \int_{-\infty}^{\infty} g(x) e^{2\pi i x y} dx$$

Theorem 8.1 (Poisson summation).

$$\sum_{n \in \mathbb{Z}} g(n) = \sum_{n \in \mathbb{Z}} \hat{g}(n)$$

Let's take $g(x, t) = e^{-\pi t x^2}$. Then $\theta(it) = \sum_{n \in \mathbb{Z}} g(n, t)$.

In this case

$$\hat{g}(y) = \sqrt{t} e^{-\pi y^2/t}$$

(integral of Gaussian).

So we conclude that

$$\theta\left(-\frac{1}{\tau}\right) = \sqrt{\frac{\tau}{i}} \theta(\tau)$$

Note also that

$$\theta(\tau + 2) = \theta(\tau)$$

because $q^{\frac{1}{2}} = e^{\pi i \tau}$.

So $\theta(\tau)$ is an example of a modular form.

What is a modular form? Let

$$\text{SL}_2(\mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z}, \det = 1 \right\},$$

which acts on $\mathbb{H} = \{\tau \in \mathbb{C} \mid \text{Im}(\tau) > 0\}$ by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \tau = \frac{a\tau + b}{c\tau + d}.$$

Definition 8.2. Let $\Gamma = \text{SL}_2(\mathbb{Z})$ or some subgroup. A (weak) modular form (of weight k) is $f : \mathbb{H} \rightarrow \mathbb{C}$ holomorphic such that

$$(8.2.1) \quad f\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^k f(\tau).$$

If we demand that Γ contains $\begin{pmatrix} 1 & N \\ 0 & 1 \end{pmatrix}$ for some $N \geq 1$, (so $f(\tau + N) = f(\tau)$), and so $f(\tau) = \sum_{n=-\infty}^{\infty} a(n) q^{2\pi i n/2}$, for $f : \mathbb{H} \rightarrow \mathbb{C}$ satisfying Eq. 8.2.1 and such that $f(\tau)$ is “meromorphic at cusps”, i.e. $f(\tau) = \sum_{n \geq N_0} a(n) q^{n/N}$, etc.

Finally we would add a factor

$$f\left(\frac{a\tau + b}{c\tau + d}\right) = \varepsilon(A) (c\tau + d)^k f(\tau).$$

So in particular a weight-0 modular form is a modular function.

Denote $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. So $S\tau = -\frac{1}{\tau}$, and denote $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, so $T\tau = \tau + 1$.

Then $\Gamma = \langle S, T^2 \rangle \subset \mathrm{SL}_2(\mathbb{Z})$.

We saw

$$\begin{aligned}\theta(T^2\tau) &= \theta(\tau) \\ \theta(S\tau) &= \sqrt{\frac{\tau}{i}}\theta(\tau) = i^{-1/2}(c\tau + d)^{1/2}\theta(\tau).\end{aligned}$$

So $\theta(\tau)$ is a modular form of weight $1/2$, for the group $\Gamma = \langle S, T^2 \rangle$ with multiplier system $\varepsilon : \Gamma \rightarrow \mathbb{C}^\times$ defined by

$$\begin{aligned}\varepsilon(S) &= i^{-1/2} \\ \varepsilon(T^2) &= 1\end{aligned}$$

Theorem 8.3. *The Dedekind eta function*

$$\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n), \quad q = e^{2\pi i \tau}$$

is also a modular form of weight $1/2$ for $SL_2(\mathbb{Z}) = \langle S, T \rangle$.

$$\begin{aligned}\eta\left(-\frac{1}{\tau}\right) &= \sqrt{\frac{\tau}{i}}\eta(\tau) \\ \eta(T\tau) &= e^{2\pi i/24}\eta(\tau).\end{aligned}$$

REFERENCES

- [Kac90] V.G. Kac, *Infinite-dimensional lie algebras*, Progress in mathematics, Cambridge University Press, 1990.
- [Kac10] Victor Kac, *Lecture notes of 18.745 – introduction to lie algebras (fall 2010)*, <https://math.mit.edu/classes/18.745/classnotes.html>, 2010, Lecture notes, MIT.