

Farka's Lemma

Convex Optimization

Dan Calderone

Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.

1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$
2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.

1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$
2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

Core Intuition: Square Case

“each row of H is orthogonal to the (other) columns of H inverse”

$$HH^{-1} = \begin{bmatrix} - & H_1^\top & - \\ - & H_2^\top & - \end{bmatrix} \begin{bmatrix} | & | \\ G_1 & G_2 \\ | & | \end{bmatrix} = \begin{bmatrix} H_1^\top G_1 & H_1^\top G_2 \\ H_2^\top G_1 & H_2^\top G_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$H_i^\top G_j = 0 \quad i \neq j$$

Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.


1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$
2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

Core Intuition: Square Case

“each row of H is orthogonal to the (other) columns of H inverse”

$$HH^{-1} = \begin{bmatrix} - & H_1^\top & - \\ - & H_2^\top & - \end{bmatrix} \begin{bmatrix} | & | \\ G_1 & G_2 \\ | & | \end{bmatrix} = \begin{bmatrix} H_1^\top G_1 & H_1^\top G_2 \\ H_2^\top G_1 & H_2^\top G_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$H_i^\top G_j = 0 \quad i \neq j$$





Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.

1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$
2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

$$HH^{-1} = \begin{bmatrix} -H_1^\top & - \\ -H_2^\top & - \end{bmatrix} \begin{bmatrix} | & | \\ G_1 & G_2 \\ | & | \end{bmatrix} = \begin{bmatrix} H_1^\top G_1 & H_1^\top G_2 \\ H_2^\top G_1 & H_2^\top G_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$H_i^\top G_j = 0 \quad i \neq j$$



Core Intuition: Square Case

“each row of H is orthogonal to the (other) columns of H inverse”

Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.



1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$
2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

Core Intuition: Square Case

“each row of H is orthogonal to the (other) columns of H inverse”

$$HH^{-1} = \begin{bmatrix} -H_1^\top & - \\ -H_2^\top & - \end{bmatrix} \begin{bmatrix} | & | \\ G_1 & G_2 \\ | & | \end{bmatrix} = \begin{bmatrix} H_1^\top G_1 & H_1^\top G_2 \\ H_2^\top G_1 & H_2^\top G_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$H_i^\top G_j = 0 \quad i \neq j$$





Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.

1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$
2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

$$HH^{-1} = \begin{bmatrix} -H_1^\top & - \\ -H_2^\top & - \end{bmatrix} \begin{bmatrix} | & | \\ G_1 & G_2 \\ | & | \end{bmatrix} = \begin{bmatrix} H_1^\top G_1 & H_1^\top G_2 \\ H_2^\top G_1 & H_2^\top G_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$H_i^\top G_j = 0 \quad i \neq j$$



Core Intuition: Square Case

“each row of H is orthogonal to the (other) columns of H inverse”

Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.



Then exactly one of the following two assertions is true.

1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$
2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

Core Intuition: Square Case

“each row of H is orthogonal to the (other) columns of H inverse”

For square, invertible $H \quad \exists d \in \mathbb{R}^n$ s.t. $Hd \leq 0 \quad \Longleftrightarrow \quad d = H^{-1}z$ for $z \leq 0$



Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.



1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$

2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

True

False

Case 1.



Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.

1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$

2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

True



False

not

\Rightarrow

For all $d \in \mathbb{R}^n$ s.t. $Hd \leq 0$, $r^\top d \leq 0$

Case 1.



Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.



1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$

2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

False

True

Case 2.



Farka's Lemma Let $H \in \mathbb{R}^{m \times n}$ and $r \in \mathbb{R}^n$.

Then exactly one of the following two assertions is true.

1. There exists $v \in \mathbb{R}^m$ such that $v^\top H = r^\top$ and $v \geq 0$

2. There exists $d \in \mathbb{R}^n$ such that $Hd \leq 0$ and $r^\top d > 0$

False

True

Case 2.

not

\Rightarrow

For all $v \in \mathbb{R}^m$ s.t. $v^\top H = r^\top$, $v \not\geq 0$

