

Quadratic Forms, Definite Matrices, Congruence Transformations

Linear Algebra:

Winter 2022 - Dan Calderone

Definite (Symmetric) Matrices

Quadratic Form: $f(x) = x^T Q x \quad Q \in \mathbb{R}^{n \times n} \quad Q = Q^T$

Definiteness:	Short	Notation	Definition	Analogy	Eigenvalues
Positive definite:	PD	$Q \succ 0$	$x^T Q x > 0 \quad \forall x \quad x \neq 0$...positive orthant	$\lambda_i > 0 \quad \lambda_i \in \text{eig}(Q)$
Positive semi-definite	PSD	$Q \succeq 0$	$x^T Q x \geq 0 \quad \forall x$...positive orthant w/ boundary	$\lambda_i \geq 0 \quad \lambda_i \in \text{eig}(Q)$
Negative-definite	ND	$Q \prec 0$	$x^T Q x < 0 \quad \forall x \quad x \neq 0$...negative orthant	$\lambda_i < 0 \quad \lambda_i \in \text{eig}(Q)$
Negative semi-definite	NSD	$Q \preceq 0$	$x^T Q x \leq 0 \quad \forall x$...negative orthant w/ boundary	$\lambda_i \leq 0 \quad \lambda_i \in \text{eig}(Q)$
Indefinite:			$x^T Q x > 0 \quad \text{some } x$...the rest of the space	
			$x^T Q x < 0 \quad \text{some } x$		

Eigenvalue condition proof:

...consider eigenvector coordinates

$$x = V x'$$

since V is invertible... $\forall x \iff \forall x'$

$$x^T Q x = x V D V^T x = x'^T D x' = \sum_i \lambda_i x_i'^2$$

$$\sum_i \lambda_i x_i'^2 > 0 \quad \forall x' \iff \lambda_i > 0 \quad \forall \lambda_i \in \text{eig}(Q)$$

$x \neq 0$

Note: not a useful definition for general matrices

... condition only says something about the symmetric part of Q

Symmetric/Skew-symmetric Decomposition

$$Q = \underbrace{\frac{1}{2} \left(Q + Q^T \right)}_{\text{symmetric}} + \underbrace{\frac{1}{2} \left(Q - Q^T \right)}_{\text{skew-sym}}$$

$$x^T Q x = \frac{1}{2} x^T \left(Q + Q^T \right) x + \frac{1}{2} x^T \left(Q - Q^T \right) x$$

$$= \frac{1}{2} x^T \left(Q + Q^T \right) x + \frac{1}{2} x^T Q x - \underbrace{\frac{1}{2} x^T Q^T x}_{\text{...transpose}}$$

$$= \frac{1}{2} x^T \left(Q + Q^T \right) x + \underbrace{\frac{1}{2} x^T Q x - \frac{1}{2} x^T Q x}_{=0}$$

$$= \frac{1}{2} x^T \left(Q + Q^T \right) x$$



...only the symmetric part matters


Definite (Symmetric) Matrices

Quadratic Form: $f(x) = x^T Q x \quad Q \in \mathbb{R}^{n \times n} \quad Q = Q^T$


Definiteness:	Short	Notation	Definition	Analogy
Positive definite:	PD	$Q \succ 0$	$x^T Q x > 0 \quad \forall x \quad x \neq 0$...positive orthant
Positive semi-definite	PSD	$Q \succeq 0$	$x^T Q x \geq 0 \quad \forall x$...positive orthant w/ boundary
Negative-definite	ND	$Q \prec 0$	$x^T Q x < 0 \quad \forall x \quad x \neq 0$...negative orthant
Negative semi-definite	NSD	$Q \preceq 0$	$x^T Q x \leq 0 \quad \forall x$...negative orthant w/ boundary
Indefinite:			$x^T Q x > 0 \quad \text{some } x$ $x^T Q x < 0 \quad \text{some } x$...the rest of the space

Surfaces: $Q \succ 0$

$Q = I$




level sets




surface

Q diagonal




level sets




surface

Q general



level sets



surface

Eigenvalue condition proof:

...consider eigenvector coordinates

$$x = V x'$$

since V is invertible... $\forall x \iff \forall x'$

$$x^T Q x = x V D V^T x = x'^T D x' = \sum_i \lambda_i x_i'^2$$

$$\sum_i \lambda_i x_i'^2 > 0 \quad \forall x' \iff \lambda_i > 0 \quad \forall \lambda_i \in \text{eig}(Q)$$

$x \neq 0$


Definite (Symmetric) Matrices

Quadratic Form: $f(x) = x^T Q x$ $Q \in \mathbb{R}^{n \times n}$ $Q = Q^T$


Definiteness:	Short	Notation	Definition	Analogy
Positive definite:	PD	$Q \succ 0$	$x^T Q x > 0 \quad \forall x \quad x \neq 0$...positive orthant
Positive semi-definite	PSD	$Q \succeq 0$	$x^T Q x \geq 0 \quad \forall x$...positive orthant w/ boundary
Negative-definite	ND	$Q \prec 0$	$x^T Q x < 0 \quad \forall x \quad x \neq 0$...negative orthant
Negative semi-definite	NSD	$Q \preceq 0$	$x^T Q x \leq 0 \quad \forall x$...negative orthant w/ boundary
Indefinite:			$x^T Q x > 0$ some x $x^T Q x < 0$ some x	...the rest of the space

Surfaces: $Q \prec 0$

$Q = I$




level sets




surface

Q diagonal




level sets




surface

Q general



level sets



surface

Eigenvalue condition proof:

...consider eigenvector coordinates

$$x = V x'$$

since V is invertible... $\forall x \iff \forall x'$

$$x^T Q x = x V D V^T x = x'^T D x' = \sum_i \lambda_i x_i'^2$$

$$\sum_i \lambda_i x_i'^2 > 0 \quad \forall x' \iff \lambda_i > 0 \quad \forall \lambda_i \in \text{eig}(Q)$$

$x \neq 0$


Definite (Symmetric) Matrices

Quadratic Form: $f(x) = x^T Q x$ $Q \in \mathbb{R}^{n \times n}$ $Q = Q^T$


Definiteness:	Short	Notation	Definition	Analogy
Positive definite:	PD	$Q \succ 0$	$x^T Q x > 0 \quad \forall x \quad x \neq 0$...positive orthant
Positive semi-definite	PSD	$Q \succeq 0$	$x^T Q x \geq 0 \quad \forall x$...positive orthant w/ boundary
Negative-definite	ND	$Q \prec 0$	$x^T Q x < 0 \quad \forall x \quad x \neq 0$...negative orthant
Negative semi-definite	NSD	$Q \preceq 0$	$x^T Q x \leq 0 \quad \forall x$...negative orthant w/ boundary
Indefinite:			$x^T Q x > 0$ some x $x^T Q x < 0$ some x	...the rest of the space

Surfaces: $Q \succeq 0$

$$Q = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$




level sets




surface

$$Q = \begin{bmatrix} \lambda_1 & 0 \\ 0 & 0 \end{bmatrix}$$




level sets




surface

diagonal

$$Q = V \begin{bmatrix} \lambda_1 & 0 \\ 0 & 0 \end{bmatrix} V^T = \lambda_1 v_1 v_1^T \quad \text{general}$$



level sets



surface

Eigenvalue condition proof:

...consider eigenvector coordinates

$$x = V x'$$

since V is invertible... $\forall x \iff \forall x'$

$$x^T Q x = x V D V^T x = x'^T D x' = \sum_i \lambda_i x_i'^2$$

$$\sum_i \lambda_i x_i'^2 > 0 \quad \forall x' \iff \lambda_i > 0 \quad \forall \lambda_i \in \text{eig}(Q)$$

$x \neq 0$

Definite (Symmetric) Matrices

Quadratic Form: $f(x) = x^T Q x \quad Q \in \mathbb{R}^{n \times n} \quad Q = Q^T$

Definiteness:	Short	Notation	Definition	Analogy
Positive definite:	PD	$Q \succ 0$	$x^T Q x > 0 \quad \forall x \quad x \neq 0$...positive orthant
Positive semi-definite	PSD	$Q \succeq 0$	$x^T Q x \geq 0 \quad \forall x$...positive orthant w/ boundary
Negative-definite	ND	$Q \prec 0$	$x^T Q x < 0 \quad \forall x \quad x \neq 0$...negative orthant
Negative semi-definite	NSD	$Q \preceq 0$	$x^T Q x \leq 0 \quad \forall x$...negative orthant w/ boundary
Indefinite:			$x^T Q x > 0 \quad \text{some } x$ $x^T Q x < 0 \quad \text{some } x$...the rest of the space

Eigenvalues


$\lambda_i > 0 \quad \lambda_i \in \text{eig}(Q)$
 $\lambda_i \geq 0 \quad \lambda_i \in \text{eig}(Q)$
 $\lambda_i < 0 \quad \lambda_i \in \text{eig}(Q)$
 $\lambda_i \leq 0 \quad \lambda_i \in \text{eig}(Q)$

Eigenvalue condition proof:


...consider eigenvector coordinates
 $x = V x'$
since V is invertible...
 $\forall x \iff \forall x'$
 $x^T Q x = x V D V^T x = x'^T D x' = \sum_i \lambda_i x_i'^2$
 $\sum_i \lambda_i x_i'^2 > 0 \quad \forall x' \iff \lambda_i > 0 \quad \forall \lambda_i \in \text{eig}(Q)$
 $x \neq 0$

Surfaces: $Q \preceq 0$

$Q = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$




level sets




surface

$Q = \begin{bmatrix} \lambda_1 & 0 \\ 0 & 0 \end{bmatrix}$




level sets




surface

$Q = V \begin{bmatrix} \lambda_1 & 0 \\ 0 & 0 \end{bmatrix} V^T = \lambda_1 v_1 v_1^T$ general




level sets



surface

Definite (Symmetric) Matrices



Quadratic Form: $f(x) = x^T Q x \quad Q \in \mathbb{R}^{n \times n} \quad Q = Q^T$

Definiteness:	Short	Notation	Definition	Analogy
Positive definite:	PD	$Q \succ 0$	$x^T Q x > 0 \quad \forall x \quad x \neq 0$...positive orthant
Positive semi-definite	PSD	$Q \succeq 0$	$x^T Q x \geq 0 \quad \forall x$...positive orthant w/ boundary
Negative-definite	ND	$Q \prec 0$	$x^T Q x < 0 \quad \forall x \quad x \neq 0$...negative orthant
Negative semi-definite	NSD	$Q \preceq 0$	$x^T Q x \leq 0 \quad \forall x$...negative orthant w/ boundary
Indefinite:			$x^T Q x > 0 \quad \text{some } x$ $x^T Q x < 0 \quad \text{some } x$...the rest of the space

Eigenvalues

$\lambda_i > 0 \quad \lambda_i \in \text{eig}(Q)$
 $\lambda_i \geq 0 \quad \lambda_i \in \text{eig}(Q)$
 $\lambda_i < 0 \quad \lambda_i \in \text{eig}(Q)$
 $\lambda_i \leq 0 \quad \lambda_i \in \text{eig}(Q)$

Eigenvalue condition proof:

...consider eigenvector coordinates

$$x = V x'$$


since V is invertible... $\forall x \iff \forall x'$

$$x^T Q x = x V D V^T x = x'^T D x' = \sum_i \lambda_i x_i'^2$$


$$\sum_i \lambda_i x_i'^2 > 0 \quad \forall x' \iff \lambda_i > 0 \quad \forall \lambda_i \in \text{eig}(Q) \quad x \neq 0$$

Surfaces: Q indefinite

$$Q = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$




level sets




surface

$$Q = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix}$$




level sets



surface


diagonal

$$Q = V \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} V^T$$



level sets

general




surface

Definite (Symmetric) Matrices


Quadratic Form: $f(x) = x^T Q x \quad Q \in \mathbb{R}^{n \times n} \quad Q = Q^T$

Definiteness:	Short	Notation	Definition	Analogy
Positive definite:	PD	$Q \succ 0$	$x^T Q x > 0 \quad \forall x \quad x \neq 0$...positive orthant
Positive semi-definite	PSD	$Q \succeq 0$	$x^T Q x \geq 0 \quad \forall x$...positive orthant w/ boundary
Negative-definite	ND	$Q \prec 0$	$x^T Q x < 0 \quad \forall x \quad x \neq 0$...negative orthant
Negative semi-definite	NSD	$Q \preceq 0$	$x^T Q x \leq 0 \quad \forall x$...negative orthant w/ boundary
Indefinite:			$x^T Q x > 0 \quad \text{some } x$ $x^T Q x < 0 \quad \text{some } x$...the rest of the space

Surfaces: $Q \succ 0$



surface



level sets

Eigenvalues

$\lambda_i > 0 \quad \lambda_i \in \text{eig}(Q)$

$\lambda_i \geq 0 \quad \lambda_i \in \text{eig}(Q)$

$\lambda_i < 0 \quad \lambda_i \in \text{eig}(Q)$

$\lambda_i \leq 0 \quad \lambda_i \in \text{eig}(Q)$

Eigenvalue condition proof:

...consider eigenvector coordinates

$$x = V x'$$

since V is invertible... $\forall x \iff \forall x'$

$$x^T Q x = x V D V^T x = x'^T D x' = \sum_i \lambda_i x_i'^2$$

$$\sum_i \lambda_i x_i'^2 > 0 \quad \forall x' \iff \lambda_i > 0 \quad \forall \lambda_i \in \text{eig}(Q)$$

$$Q = V D V^T = \begin{bmatrix} | & | \\ v_1 & v_2 \\ | & | \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \begin{bmatrix} - & v_1^T & - \\ - & v_2^T & - \end{bmatrix} \quad ||v_i||_2 = 1$$


$$\begin{aligned} f\left(\frac{1}{\sqrt{\lambda_1}} v_1\right) &= \frac{1}{\sqrt{\lambda_1}} v_1^T Q v_1 \frac{1}{\sqrt{\lambda_1}} \\ &= \frac{1}{\sqrt{\lambda_1}} \begin{bmatrix} | \\ v_1 \\ | \end{bmatrix}^T \begin{bmatrix} | & | \\ v_1 & v_2 \\ | & | \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \begin{bmatrix} - & v_1^T & - \\ - & v_2^T & - \end{bmatrix} \begin{bmatrix} | \\ v_1 \\ | \end{bmatrix} \frac{1}{\sqrt{\lambda_1}} \\ &= \frac{1}{\sqrt{\lambda_1}} \begin{bmatrix} 1 \\ 0 \end{bmatrix}^T \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \frac{1}{\sqrt{\lambda_1}} = \frac{\lambda_1}{(\sqrt{\lambda_1})^2} = 1 \end{aligned}$$

Definite (Symmetric) Matrices


Quadratic Form: $f(x) = x^T Q x \quad Q \in \mathbb{R}^{n \times n} \quad Q = Q^T$

Definiteness:	Short	Notation	Definition	Analogy
Positive definite:	PD	$Q \succ 0$	$x^T Q x > 0 \quad \forall x \quad x \neq 0$...positive orthant
Positive semi-definite	PSD	$Q \succeq 0$	$x^T Q x \geq 0 \quad \forall x$...positive orthant w/ boundary
Negative-definite	ND	$Q \prec 0$	$x^T Q x < 0 \quad \forall x \quad x \neq 0$...negative orthant
Negative semi-definite	NSD	$Q \preceq 0$	$x^T Q x \leq 0 \quad \forall x$...negative orthant w/ boundary
Indefinite:			$x^T Q x > 0 \quad \text{some } x$ $x^T Q x < 0 \quad \text{some } x$...the rest of the space

Surfaces: Q indefinite



surface



level sets

Eigenvalues

$$\lambda_i > 0 \quad \lambda_i \in \text{eig}(Q)$$

$$\lambda_i \geq 0 \quad \lambda_i \in \text{eig}(Q)$$

$$\lambda_i < 0 \quad \lambda_i \in \text{eig}(Q)$$

$$\lambda_i \leq 0 \quad \lambda_i \in \text{eig}(Q)$$

Eigenvalue condition proof:

...consider eigenvector coordinates

$$x = V x'$$

since V is invertible... $\forall x \iff \forall x'$

$$x^T Q x = x V D V^T x = x'^T D x' = \sum_i \lambda_i x_i'^2$$

$$\sum_i \lambda_i x_i'^2 > 0 \quad \forall x' \iff \lambda_i > 0 \quad \forall \lambda_i \in \text{eig}(Q)$$

$x \neq 0$

$$Q = V D V^T = \begin{bmatrix} | & | \\ v_1 & v_2 \\ | & | \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \begin{bmatrix} - & v_1^T & - \\ - & v_2^T & - \end{bmatrix} \quad ||v_i||_2 = 1$$

$$f\left(\frac{1}{\sqrt{\lambda_1}} v_1\right) = \frac{1}{\sqrt{\lambda_1}} v_1^T Q v_1 \frac{1}{\sqrt{\lambda_1}}$$


$$= \frac{1}{\sqrt{\lambda_1}} \begin{bmatrix} | \\ v_1 \\ | \end{bmatrix}^T \begin{bmatrix} | & | \\ v_1 & v_2 \\ | & | \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \begin{bmatrix} - & v_1^T & - \\ - & v_2^T & - \end{bmatrix} \begin{bmatrix} | \\ v_1 \\ | \end{bmatrix} \frac{1}{\sqrt{\lambda_1}}$$

$$= \frac{1}{\sqrt{\lambda_1}} \begin{bmatrix} 1 \\ 0 \end{bmatrix}^T \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \frac{1}{\sqrt{\lambda_1}} = \frac{\lambda_1}{(\sqrt{\lambda_1})^2} = 1$$


Quadratic Form - Level Sets

Quadratic Form: $f(x) = x^T Q x$ $Q \in \mathbb{R}^{n \times n}$ $Q = Q^T$

Ellipsoids



Hyperboloids



Cones

