

Perturbation Analysis: KKT Conditions

Original
**Stationarity
Condition**

$$\frac{\partial f}{\partial x} \left(x \right) + \lambda^\top \frac{\partial g}{\partial x} \left(x, \theta \right) = 0$$

Feasibility Condition
(original constraint)

$$g \left(x, \theta \right) = 0$$



variables

parameters

(defines shape
of constraints)

Perturbation Analysis: KKT Conditions

Original
**Stationarity
Condition**

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0$$

Perturb
Shape of
Constraints

$$\theta \rightarrow \theta + \Delta\theta$$

Question:

How does this affect
other problem variables
at optimum?

Feasibility Condition
(original constraint)

$$g(x, \theta) = 0$$



variables

parameters

(defines shape
of constraints)

Perturbation Analysis: KKT Conditions

Original
**Stationarity
Condition**

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0$$

Feasibility Condition
(original constraint)

$$g(x, \theta) = 0$$



variables

parameters

(defines shape
of constraints)

Perturb
Shape of
Constraints

$$\theta \rightarrow \theta + \Delta\theta$$

Question:

How does this affect
other problem variables
at optimum?

Perturbation Analysis

Perturbed
**Feasibility
Conditions**

$$g(x + \Delta x, \theta + \Delta\theta) = 0$$

Perturbation Analysis: KKT Conditions

Original
**Stationarity
Condition**

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0$$

Feasibility Condition
(original constraint)

$$g(x, \theta) = 0$$



variables parameters

(defines shape
of constraints)

Perturb
Shape of
Constraints

$$\theta \rightarrow \theta + \Delta\theta$$

Question:

How does this affect
other problem variables
at optimum?

Perturbation Analysis

Perturbed
**Feasibility
Conditions**

$$g(x + \Delta x, \theta + \Delta\theta) = 0 \quad \approx$$

$$g(x, \theta) + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta\theta = 0$$

Perturbation Analysis: KKT Conditions

Original
**Stationarity
Condition**

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0$$

Feasibility Condition
(original constraint)

$$g(x, \theta) = 0$$



variables

parameters

(defines shape
of constraints)

Perturb
Shape of
Constraints

$$\theta \rightarrow \theta + \Delta\theta$$

Question:

How does this affect
other problem variables
at optimum?

Perturbation Analysis

Perturbed
**Feasibility
Conditions**

$$g(x + \Delta x, \theta + \Delta\theta) = 0 \quad \approx$$

$$g(x, \theta) + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta\theta = 0$$

Perturbed
**Stationarity
Conditions**

$$\frac{\partial f}{\partial x}(x + \Delta x) + (\lambda + \Delta\lambda)^\top \frac{\partial g}{\partial x}(x + \Delta x, \theta + \Delta\theta) = 0$$

Perturbation Analysis: KKT Conditions

Original
**Stationarity
Condition**

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0$$

Feasibility Condition
(original constraint)

$$g(x, \theta) = 0$$



variables

parameters

(defines shape
of constraints)

Perturb
Shape of
Constraints

$$\theta \rightarrow \theta + \Delta\theta$$

Question:

How does this affect
other problem variables
at optimum?

Perturbation Analysis

Perturbed
**Feasibility
Conditions**

$$g(x + \Delta x, \theta + \Delta\theta) = 0 \quad \approx$$

$$g(x, \theta) + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta\theta = 0$$

Perturbed
**Stationarity
Conditions**

$$\frac{\partial f}{\partial x}(x + \Delta x) + (\lambda + \Delta\lambda)^\top \frac{\partial g}{\partial x}(x + \Delta x, \theta + \Delta\theta) = 0$$

$$\frac{\partial f}{\partial x}(x) + \Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + (\lambda + \Delta\lambda)^\top \left[\frac{\partial g}{\partial x} \right] + \sum_i (\lambda_i + \Delta\lambda_i) \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i (\lambda_i + \Delta\lambda_i) \Delta\theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

Perturbation Analysis: KKT Conditions

Original
**Stationarity
Condition**

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0$$

Feasibility Condition
(original constraint)

$$g(x, \theta) = 0$$



variables

parameters

(defines shape
of constraints)

Perturb
Shape of
Constraints

$$\theta \rightarrow \theta + \Delta\theta$$

Question:

How does this affect
other problem variables
at optimum?

Perturbation Analysis

Perturbed
**Feasibility
Conditions**

$$g(x + \Delta x, \theta + \Delta\theta) = 0 \quad \approx$$

$$g(x, \theta) + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta\theta = 0$$

Perturbed
**Stationarity
Conditions**

$$\frac{\partial f}{\partial x}(x + \Delta x) + (\lambda + \Delta\lambda)^\top \frac{\partial g}{\partial x}(x + \Delta x, \theta + \Delta\theta) = 0$$

$$\frac{\partial f}{\partial x}(x) + \Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + (\lambda + \Delta\lambda)^\top \left[\frac{\partial g}{\partial x} \right] + \sum_i (\lambda_i + \Delta\lambda_i) \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i (\lambda_i + \Delta\lambda_i) \Delta\theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

Procedure:

1. Cancel terms...
2. Solve for $\Delta x, \Delta\lambda$ as functions of $\Delta\theta$.

Perturbation Analysis: KKT Conditions

Original Conditions

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0 \quad g(x, \theta) = 0$$

Perturbed Conditions

$$\begin{aligned} \frac{\partial f}{\partial x}(x) + \Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + (\lambda + \Delta \lambda)^\top \left[\frac{\partial g}{\partial x} \right] \\ + \sum_i (\lambda_i + \Delta \lambda_i) \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i (\lambda_i + \Delta \lambda_i) \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0 \end{aligned}$$
$$g(x, \theta) + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

Perturbation Analysis: KKT Conditions

Original Conditions

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0 \quad g(x, \theta) = 0$$

Perturbed Conditions

$$\cancel{\frac{\partial f}{\partial x}(x)} + \Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \cancel{(\lambda + \Delta \lambda)^\top} \left[\frac{\partial g}{\partial x} \right] + \sum_i (\lambda_i + \Delta \lambda_i) \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i (\lambda_i + \Delta \lambda_i) \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$
$$\cancel{g(x, \theta)} + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

1. Cancel zero-order terms...

Perturbation Analysis: KKT Conditions

Original Conditions

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0 \quad g(x, \theta) = 0$$

Perturbed Conditions

$$\begin{aligned} & \cancel{\frac{\partial f}{\partial x}(x)} + \Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \cancel{(\lambda + \Delta \lambda)^\top} \left[\frac{\partial g}{\partial x} \right] \\ & + \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0 \end{aligned}$$

$$\cancel{g(x, \theta)} + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

1. Cancel zero-order terms...
2. Cancel second-order terms...

Perturbation Analysis: KKT Conditions

Original Conditions

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0 \quad g(x, \theta) = 0$$

Perturbed Conditions

$$\cancel{\frac{\partial f}{\partial x}(x)} + \Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \cancel{(\lambda + \Delta \lambda)^\top} \left[\frac{\partial g}{\partial x} \right] + \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

$$\cancel{g(x, \theta)} + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

1. Cancel zero-order terms... 2. Cancel second-order terms...

$$\Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \Delta \lambda^\top \left[\frac{\partial g}{\partial x} \right] + \sum_i \lambda_i \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i \lambda_i \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

$$\frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

Perturbation Analysis: KKT Conditions

Original Conditions

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0 \quad g(x, \theta) = 0$$

Perturbed Conditions

$$\cancel{\frac{\partial f}{\partial x}(x)} + \Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \cancel{(\lambda + \Delta \lambda)^\top} \left[\frac{\partial g}{\partial x} \right] + \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

$$\cancel{g(x, \theta)} + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

1. Cancel zero-order terms... 2. Cancel second-order terms...

$$\Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \Delta \lambda^\top \left[\frac{\partial g}{\partial x} \right] + \sum_i \lambda_i \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i \lambda_i \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

$$\frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

organize into a system of equations...

$$\begin{bmatrix} Q & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta \lambda \end{bmatrix} = \begin{bmatrix} \sum_i \lambda_i \frac{\partial^2 g_i}{\partial x \partial \theta} \\ \frac{\partial g}{\partial \theta} \end{bmatrix} \Delta \theta$$

$$Q = \frac{\partial^2 f}{\partial x^2} + \sum_i \lambda_i \frac{\partial^2 g_i}{\partial x^2}$$

$$A = \frac{\partial g}{\partial x}$$

Perturbation Analysis: KKT Conditions

$$\frac{\partial f}{\partial x}(x) + \Delta x^\top \frac{\partial^2 f}{\partial x^2} + \lambda^\top \left[\frac{\partial g}{\partial x}(x, \theta) + \frac{\partial^2 g}{\partial x^2} \Delta x + \frac{\partial^2 g}{\partial x \partial \theta} \Delta \theta \right]$$

Original
Conditions

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0$$

$$g(x, \theta) = 0$$

$$+ \Delta \lambda^\top \left[\frac{\partial g}{\partial x}(x, \theta) + \frac{\partial^2 g}{\partial x^2} \Delta x + \frac{\partial^2 g}{\partial x \partial \theta} \Delta \theta \right]$$

Perturbed Conditions

$$\cancel{\frac{\partial f}{\partial x}(x)} + \Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \cancel{(\lambda + \Delta \lambda)^\top} \left[\frac{\partial g}{\partial x} \right]$$

$$+ \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

$$\cancel{g(x, \theta)} + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

1. Cancel zero-order terms... 2. Cancel second-order terms...

$$\Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \Delta \lambda^\top \left[\frac{\partial g}{\partial x} \right] + \sum_i \lambda_i \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i \lambda_i \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

$$\frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

organize into a system of equations...

$$\begin{bmatrix} \Delta x \\ \Delta \lambda \end{bmatrix} = \begin{bmatrix} Q & A^\top \\ A & 0 \end{bmatrix}^{-1} \begin{bmatrix} \sum_i \lambda_i \frac{\partial^2 g_i}{\partial x \partial \theta} \\ \frac{\partial g}{\partial \theta} \end{bmatrix} \Delta \theta$$

$$Q = \frac{\partial^2 f}{\partial x^2} + \sum_i \lambda_i \frac{\partial^2 g_i}{\partial x^2}$$

$$A = \frac{\partial g}{\partial x}$$

Perturbation Analysis: KKT Conditions

$$\frac{\partial f}{\partial x}(x) + \Delta x^\top \frac{\partial^2 f}{\partial x^2} + \lambda^\top \left[\frac{\partial g}{\partial x}(x, \theta) + \frac{\partial^2 g}{\partial x^2} \Delta x + \frac{\partial^2 g}{\partial x \partial \theta} \Delta \theta \right]$$

Original
Conditions

$$\frac{\partial f}{\partial x}(x) + \lambda^\top \frac{\partial g}{\partial x}(x, \theta) = 0$$

$$g(x, \theta) = 0$$

$$+ \Delta \lambda^\top \left[\frac{\partial g}{\partial x}(x, \theta) + \frac{\partial^2 g}{\partial x^2} \Delta x + \frac{\partial^2 g}{\partial x \partial \theta} \Delta \theta \right]$$

Perturbed Conditions

$$\cancel{\frac{\partial f}{\partial x}(x)} + \Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \cancel{(\lambda + \Delta \lambda)^\top} \left[\frac{\partial g}{\partial x} \right]$$

$$\cancel{g(x, \theta)} + \frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

$$+ \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i (\lambda_i + \cancel{\Delta \lambda_i}) \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

1. Cancel zero-order terms... 2. Cancel second-order terms...

$$\Delta x^\top \left[\frac{\partial^2 f}{\partial x^2} \right] + \Delta \lambda^\top \left[\frac{\partial g}{\partial x} \right] + \sum_i \lambda_i \Delta x^\top \left[\frac{\partial^2 g_i}{\partial x^2} \right] + \sum_i \lambda_i \Delta \theta^\top \left[\frac{\partial^2 g_i}{\partial \theta \partial x} \right] = 0$$

$$\frac{\partial g}{\partial x} \Delta x + \frac{\partial g}{\partial \theta} \Delta \theta = 0$$

organize into a system of equations...

$$\begin{bmatrix} \Delta x \\ \Delta \lambda \end{bmatrix} = \begin{bmatrix} Q & A^\top \\ A & 0 \end{bmatrix}^{-1} \begin{bmatrix} \sum_i \lambda_i \frac{\partial^2 g_i}{\partial x \partial \theta} \\ \frac{\partial g}{\partial \theta} \end{bmatrix} \Delta \theta$$

$$Q = \frac{\partial^2 f}{\partial x^2} + \sum_i \lambda_i \frac{\partial^2 g_i}{\partial x^2}$$

$$A = \frac{\partial g}{\partial x}$$

$$= \begin{bmatrix} Q^{-1} - Q^{-1} A^\top (A Q^{-1} A^\top)^{-1} A Q^{-1} & Q^{-1} A^\top (A Q^{-1} A^\top)^{-1} \\ (A Q^{-1} A^\top)^{-1} A Q^{-1} & -(A Q^{-1} A^\top)^{-1} \end{bmatrix} \begin{bmatrix} \sum_i \lambda_i \frac{\partial^2 g_i}{\partial x \partial \theta} \\ \frac{\partial g}{\partial \theta} \end{bmatrix} \Delta \theta$$