

COVERING SPACES.

In our proof of $\pi_1(S^1) \cong \mathbb{Z}$ we used $\mathbb{R} \rightarrow S^1$.
Can similarly show $\pi_1(T^2) \cong \mathbb{Z}^2$ using $\mathbb{R}^2 \rightarrow T^2$
or $\pi_1(S^1 \vee S^1) \cong \mathbb{F}_2$ using $T_4 \rightarrow S^1 \vee S^1$.

In each case, $\pi_1(X)$ gives symmetries of the space lying above.

A covering space of X is an \tilde{X} with
 $p: \tilde{X} \rightarrow X$

satisfying: \exists open cover $\{U_\alpha\}$ of X so that
each $p^{-1}(U_\alpha)$ is a disjoint union
of open sets, each homeomorphic to U_α .

Examples. $\mathbb{R} \rightarrow S^1$ $\mathbb{R} \times I \rightarrow S^1 \times I$ $\mathbb{R}^2 \rightarrow T^2$ $S^2 \rightarrow \mathbb{RP}^2$
 $S^1 \xrightarrow{\times n} S^1$ $\mathbb{R} \times I \rightarrow \text{Möbius Strip}$ $\mathbb{R}^2 \rightarrow \text{Klein bottle}$

A universal covering space is a covering space
that is simply connected.

We will see: ① $\pi_1(X) \leftrightarrow$ symmetries of univ. cover \tilde{X}
② Subgroups of $\pi_1(X) \leftrightarrow$ covers of X .






e.g. $X = S^1$.

① via path lifting, ② via path projecting

EXAMPLE

$$X = \begin{array}{c} b \quad a \\ \circ \quad \circ \\ \text{---} \end{array}$$

\tilde{X}



$$p_*(\pi_1(\tilde{X}))$$

$$\langle a, b^2, bab^{-1} \rangle$$

$$\langle a^2, b^2, ab \rangle$$

$$\langle a^4, b^4, ab, ba, a^2b^2 \rangle$$

$$\langle b^n a b^{-n} \rangle$$

$$\langle a \rangle$$

$$1$$

FUNDAMENTAL THEOREM

$p: \tilde{X} \rightarrow X$ covering map
 $G(\tilde{X}) =$ deck transformation group
 $=$ p -equivariant symmetries of \tilde{X}

Thm: $1 \rightarrow \text{Normalizer}(p_*(\pi_1(\tilde{X}))) \rightarrow \pi_1(X) \rightarrow G(\tilde{X}) \rightarrow 1$

So: $p_* \pi_1(\tilde{X})$ normal $\iff G(\tilde{X})$ acts transitively
on $p^{-1}(x_0)$
Since change of basept for \tilde{X}
 \iff conjugation of $p_* \pi_1(\tilde{X})$ in $\pi_1(X)$.

$$p_*(\pi_1(\tilde{X})) = 1 \iff \tilde{X} = \text{universal cover} \\ \iff G(\tilde{X}) \cong \pi_1(X)$$

Converse: Any $\pi_1(X) \rightarrow G$ is realized by some \tilde{X}
Any $H \hookrightarrow \pi_1(X)$ is realized by some \tilde{X} .

In particular: every 2-generator group is
the symmetry group of some cover of
 $S^1 \vee S^1$

LIFTING PROPERTIES

$p: \tilde{X} \rightarrow X$ covering space

A lift of $f: Y \rightarrow X$ is $\tilde{f}: Y \rightarrow \tilde{X}$ with $p\tilde{f} = f$.

Proposition 1 (Homotopy lifting property) Given a homotopy $f_t: Y \rightarrow X$ and $\tilde{f}_0: Y \rightarrow \tilde{X}$ lifting f_0 , $\exists!$ \tilde{f}_t lifting f_t .

Proof: Same as S^1 case.

$Y = \text{point} \rightsquigarrow$ path lifting property

$Y = I \rightsquigarrow$ homotopy lifting for paths

Cor: $p_*: \pi_1(\tilde{X}) \rightarrow \pi_1(X)$ is injective.

Note: $p_*(\pi_1(\tilde{X}))$ is the subgroup of $\pi_1(X)$ consisting of loops that lift to loops.

Degree of \tilde{p} cover: $|p^{-1}(x)|$ is locally constant, hence constant

Cor: X, \tilde{X} path connected.

$$\text{degree of } p = [\pi_1(X) : p_*\pi_1(\tilde{X})]$$

Proof: Let $H = p_*\pi_1(\tilde{X})$.

Define $\{\text{cosets of } H\} \rightarrow p^{-1}(x_0)$

$$H[g] \mapsto \tilde{g}(1).$$

Surjective: path proj. Injective: path lifting \square

Proposition 2 (Lifting existence criterion) $Y =$ connected,
 locally path connected. We can lift $f: (Y, y_0) \rightarrow (X, x_0)$
 to $\tilde{f}: (Y, y_0) \rightarrow (\tilde{X}, \tilde{x}_0)$ iff

$$f_*(\pi_1(Y)) \leq p_*\pi_1(\tilde{X}).$$

Proof: \Rightarrow \tilde{f} exists $\Rightarrow f = p\tilde{f} \Rightarrow f_* = p_*\tilde{f}_*$
 $\Rightarrow \text{Im } f_* \subseteq \text{Im } p_*.$

\Leftarrow Suppose $\text{Im } f_* \subseteq \text{Im } p_*$. Want to build \tilde{f} .

Let $y \in Y$, f a path from y_0 to y .
 Prop 1 $\Rightarrow f$ has unique lift $\tilde{f}f: Y \rightarrow \tilde{X}$.
 Define

$$\tilde{f}(y) = \tilde{f}f(1).$$

Why is \tilde{f} well-defined?

Let $f' =$ another path from y_0 to y .
 $\Rightarrow (ff')(f\bar{f})$ is a loop h_0 at x_0 .
 $\Rightarrow h_0 = f(f\bar{f}) \in f_*(\pi_1(Y))$
 $\Rightarrow h_0 \in p_*(\pi_1(\tilde{X}))$ by assumption.
 \Rightarrow the lifted path \tilde{h}_0 is a loop.

Uniqueness of lifted paths $\Rightarrow \tilde{h}_0 = \tilde{f}f \tilde{f}f'$
 $\Rightarrow \tilde{f}f, \tilde{f}f'$ share common endpoint.

Exercise: \tilde{f} continuous.



Proposition 3 (Uniqueness of lifts) Let $f: Y \rightarrow X$, Y connected.
 If lifts \tilde{f}_1, \tilde{f}_2 agree at one point, then they are equal.

Proof: Will show

$$A = \{y \in Y : \tilde{f}_1(y) = \tilde{f}_2(y)\}$$

is open and closed in Y .

Let $y \in Y$. Let U be open nbhd of Y as in definition of covering space.

Let \tilde{U}_1, \tilde{U}_2 be the components of $p^{-1}(U)$ containing $\tilde{f}_1(y), \tilde{f}_2(y)$.

Continuity of $\tilde{f}_i \Rightarrow \exists$ nbhd N of y with
 $\tilde{f}_i(N) \subseteq \tilde{U}_i$

$$\bullet \tilde{f}_1(y) \neq \tilde{f}_2(y) \Rightarrow \tilde{U}_1 \neq \tilde{U}_2 \Rightarrow \tilde{f}_1(N) \cap \tilde{f}_2(N) = \emptyset$$

$\Rightarrow A$ closed.

$$\bullet \tilde{f}_1(y) = \tilde{f}_2(y) \Rightarrow \tilde{U}_1 = \tilde{U}_2 \Rightarrow \tilde{f}_1|_N = \tilde{f}_2|_N$$

Thus A open. ▣