

Maximum Contextuality Gap up to Eleven Vertices and Ramsey Structure in the Maximizers

Dan Muşetoiu
dan.musetoiu@gmail.com

September 2026

Abstract

In the exclusivity-graph approach to contextuality, a noncontextuality inequality is a graph G , its noncontextual bound is the independence number $\alpha(G)$ and its quantum bound is the Lovász number $\vartheta(G)$. The absolute gap $\Delta(G) = \vartheta(G) - \alpha(G)$ was recently maximized exhaustively over all connected graphs on eight vertices. We extend the exhaustive search to nine, ten and eleven vertices — up to 1,006,700,565 graphs — and find $\Delta_{\max} = 2/3$ ($\vartheta = 11/3$), $1/\sqrt{2}$ ($\vartheta = 3 + 1/\sqrt{2}$) and 0.774889. The billion-graph search runs in hours on a consumer GPU through a certified screen: a feasible point of the dual semidefinite program is an explicit upper bound on ϑ , and only graphs whose certified bound can beat the record are solved exactly. At eleven vertices the maximizers change character, from dense graphs with $\alpha = 3$ to sparse triangle-free graphs with $\alpha = 4$. We explain this by a Ramsey-theoretic tension: triangle-free graphs with $\alpha \leq k - 1$ are the $R(3, k)$ -graphs and exist only below $R(3, k)$ vertices, so Ramsey numbers hold α down while ϑ grows. At the critical sizes the record graphs are $R(3, k)$ -graphs; the triangle-free maximizer on thirteen vertices is the unique $R(3, 5)$ -critical graph $C_{13}(1, 5)$, with ϑ in closed form. The complete catalogues of $R(3, k)$ -graphs give a ladder of lower bounds on $\Delta_{\max}(n)$ up to $n = 35$, and we prove $\Delta(G) \leq \nu(G) \leq n/2$, that $c^* = \lim \Delta_{\max}(n)/n$ exists with $0.126 \leq c^* \leq 1/2$, and hence $\Delta_{\max}(R(3, k) - 1) = \Omega(k^2/\ln k)$. Code, graph identifiers and certificates are released.

Keywords: quantum contextuality, exclusivity graphs, Lovász theta number, independence number, Ramsey graphs, semidefinite programming, exhaustive graph search

1 Introduction

Which exclusivity graphs separate the quantum from the classical bound of a noncontextuality inequality by the largest amount? For eight vertices the answer was found recently by exhaustive search [9]. Here we push the search to eleven vertices, over a billion graphs, and find that the answer changes character on the way: the maximizers stop being dense extensions of the eight-vertex record and become sparse, triangle-free graphs of a very specific kind — Ramsey graphs. The extremal problem, we argue, is governed not only by the semidefinite geometry of the Lovász number but by a Ramsey-theoretic tension between two invariants: the independence number must be kept small, which Ramsey numbers forbid beyond a certain size unless triangles are admitted, while the theta number grows with the number of vertices. The largest gaps occur exactly at the last sizes where a triangle-free graph can still hold its independence number down.

Cabello, Severini and Winter (CSW) showed that any noncontextuality (NC) inequality of the form $\sum_i P(e_i) \leq \alpha$ can be encoded by an *exclusivity graph* G whose vertices are the events e_i and whose edges join mutually exclusive events; the maximum of the left-hand side over noncontextual

models is the independence number $\alpha(G)$, and its maximum over quantum models is the Lovász theta number $\vartheta(G)$ [1]. The gap between the two numbers is therefore a graph invariant that measures how contextual an inequality can be. Two natural figures of merit have been studied: the ratio ϑ/α , used to identify experimentally accessible witnesses [6, 7] and as a dimension witness [8], and the absolute gap $\Delta(G) = \vartheta(G) - \alpha(G)$, whose maximization over all connected graphs on eight vertices was recently carried out by Tamer et al. [9]. Their exhaustive search over the 11,117 graphs identified a sparse ten-edge graph, Quad- C_5 , with $\Delta = 0.46784$, beating the Wagner graph ($\Delta = \sqrt{2} - 1$).

Two questions follow. What are the maximizers for more vertices, and is there structure behind them? Here we answer the first question exhaustively for $n = 9, 10, 11$ and, for the triangle-free class, for $n = 12, 13$; and we answer the second by exhibiting a Ramsey-theoretic mechanism and by proving what can be proved about it. The central open statement, for which everything from 8 to 35 vertices below is evidence, is:

At $n = R(3, k) - 1$, the maximum of Δ over all graphs on n vertices is attained by an $R(3, k)$ -critical graph (Conjecture 1).

It is verified over all graphs at $n = 8$ ($k = 4$: Quad- C_5 is triangle-free with $\alpha = 3$ on $8 = R(3, 4) - 1$ vertices [9]) and, in the triangle-free class, at $n = 13$ ($k = 5$); for the other critical sizes (17, 22, 27, 35) we can only report the maximum over the critical family itself.

Contributions.

1. **Exhaustive maxima for $n = 9, 10, 11$.** $\Delta_{\max}(9) = 2/3$ exactly, $\Delta_{\max}(10) = 1/\sqrt{2}$ exactly, $\Delta_{\max}(11) = 0.774889$ (Section 3). The eleven-vertex search covers all 1,006,700,565 connected graphs and is, to our knowledge, the first exhaustive computation of a Lovász-number extremum at this scale (the existing table of theta numbers [4, 5] stops at ten vertices).
2. **A certified GPU screening method** (Section 5): dual-feasible points of the theta SDP, found by numerical optimization in large batches on a consumer GPU, are explicit upper bounds on ϑ that eliminate all but a hundred graphs out of 10^9 ; the survivors are solved exactly. The distinction between the search (floating point, heuristic) and the certificate (an explicit matrix whose largest eigenvalue bounds ϑ) is kept throughout, and the method is validated against every known answer ($n \leq 10$) with zero misses.
3. **Ramsey structure in the maximizers** (Section 4). At the Ramsey-critical sizes the record graphs are $R(3, k)$ -graphs, i.e. triangle-free graphs with $\alpha \leq k - 1$ on fewer than $R(3, k)$ vertices; in between ($n = 9, 10$) the maximizers keep the previous independence number by admitting triangles. The thirteen-vertex triangle-free maximizer is the unique $R(3, 5)$ -critical graph $C_{13}(1, 5)$, whose theta number has a closed form. Ramsey numbers thus pin α while ϑ keeps growing, and $\Delta_{\max}(n)$ jumps at $n = R(3, k) - 1$. Computing ϑ over the complete catalogues of $R(3, k)$ -graphs gives a ladder of lower bounds up to $n = 35$.
4. **Growth** (Section 4.4). $\Delta(G) \leq \nu(G) \leq n/2$ for every graph (Proposition 2); the gap is superadditive, so $c^* = \lim \Delta_{\max}(n)/n$ exists and every record graph bounds it from below, $0.126 \leq c^* \leq \frac{1}{2}$ (Proposition 3); hence $\Delta_{\max}(R(3, k) - 1) = \Omega(k^2/\ln k)$ (Corollary 1).

2 Preliminaries

Throughout, $G = (V, E)$ is a simple graph on n vertices, interpreted as an exclusivity graph. The independence number $\alpha(G)$ is the size of a largest set of pairwise non-adjacent vertices. The Lovász number [2] can be written as the semidefinite program

$$\vartheta(G) = \max\{\langle J, X \rangle : \text{tr } X = 1, X_{ij} = 0 \forall ij \in E, X \succeq 0\}, \quad (1)$$

where J is the all-ones matrix, with the dual

$$\vartheta(G) = \min\{\lambda_{\max}(J + Y) : Y = Y^T, Y_{ij} = 0 \forall ij \notin E, Y_{ii} = 0\}. \quad (2)$$

The sandwich theorem gives $\alpha(G) \leq \vartheta(G) \leq \alpha^*(G)$, where α^* is the fractional packing number [3]. In the CSW framework $\alpha(G)$ and $\vartheta(G)$ are respectively the noncontextual and the quantum maximum of $\sum_{i \in V} P(e_i)$ over all correlation experiments whose exclusivity graph is G [1]; quantum theory violates the NC inequality iff $\vartheta(G) > \alpha(G)$, which happens iff G contains an induced odd cycle of length at least five or the complement of one [4]. We study the *absolute contextuality gap*

$$\Delta(G) = \vartheta(G) - \alpha(G), \quad \Delta_{\max}(n) = \max\{\Delta(G) : G \text{ connected}, |V(G)| = n\}. \quad (3)$$

We also write $f(n)$ for the maximum of Δ over *all* graphs on n vertices, connected or not. The two quantities are tied together by the following elementary facts, used repeatedly below.

Lemma 1 (Universal vertices and connectivity). *Adding a vertex adjacent to all vertices of G changes neither ϑ nor α . Consequently $f(n-1) \leq \Delta_{\max}(n) \leq f(n)$ and $\Delta_{\max}(n+1) \geq \Delta_{\max}(n)$.*

Proof. Independent sets are unchanged by a universal vertex v , so α is unchanged. In (1) the row and column of v must vanish off the diagonal, so $X = \text{diag}(X', s)$ with $\text{tr } X' = 1 - s$ and $\langle J, X \rangle = (1 - s)\vartheta(G) + s \leq \vartheta(G)$; hence ϑ is unchanged. A universal vertex makes any graph connected, so a (possibly disconnected) maximizer for $f(n-1)$ becomes a connected graph on n vertices with the same gap; the other inequality is trivial, and monotonicity follows by adding a universal vertex to a connected maximizer. $\square$

The two sequences therefore have the same growth, and the connected maximizers of Section 3 are the interesting objects: a disconnected graph never beats its best component with universal vertices added, since ϑ and α are additive over disjoint unions [3].

Lemma 2 (Ramsey pins α). *Let $R(3, k)$ denote the Ramsey number, and call a triangle-free graph with $\alpha \leq k-1$ an $R(3, k)$ -graph. For $R(3, k-1) \leq n < R(3, k)$, every triangle-free graph on n vertices has $\alpha \geq k-1$, and those with $\alpha = k-1$ are exactly the $R(3, k)$ -graphs on n vertices; for $n \geq R(3, k)$ there are none.*

Proof. Immediate from the definition of $R(3, k)$ as the least n such that every graph on n vertices contains a triangle or an independent set of size k : on $n \geq R(3, k-1)$ vertices a triangle-free graph has $\alpha \geq k-1$, on $n \geq R(3, k)$ it has $\alpha \geq k$. $\square$

The relevant values are $R(3, 4) = 9$, $R(3, 5) = 14$, $R(3, 6) = 18$, $R(3, 7) = 23$, $R(3, 8) = 28$, $R(3, 9) = 36$ [13]. McKay's catalogues [15, 14] list all $R(3, k)$ -graphs for every n when $k \leq 7$, and at the critical sizes $n = R(3, k) - 1$ for $k = 8, 9$; the $(3, 5, 13)$ and $(3, 9, 35)$ graphs are unique [13]. Both ϑ and α are additive over disjoint unions: for α trivially, for ϑ through Lovász's formula $\vartheta(G) = \max \sum_i \langle c, u_i \rangle^2$ over orthonormal representations $\{u_i\}$ (orthogonal on edges) and unit handles c [2]: the sum splits over components, each part being at most the component's ϑ , and the direct sum of optimal representations with handle $(\sqrt{\vartheta_1} c_1 \oplus \sqrt{\vartheta_2} c_2) / \sqrt{\vartheta_1 + \vartheta_2}$ attains $\vartheta_1 + \vartheta_2$ [3]. This fact is used several times below.

Table 1: Maximum contextuality gap over all connected graphs on n vertices, all rows from our own runs (those for $n \leq 8$ agree with [9]). Graphs are identified by their graph6 string [16]. The maximizers for $n = 9, 10, 11$ are unique up to isomorphism (the runners-up have $\Delta = 0.618, 0.667, 0.755$). The optimal primal matrix X in (1) has rank $r = 4$ for every row with $n \geq 8$ (Appendix B).

n	#graphs	graph6	$ E $	α	ϑ	$\Delta_{\max}(n)$	remarks
5	21	C_5	5	2	$\sqrt{5}$	0.236068	KCBS pentagon
6	112	EUZO	8	2	$\sqrt{5}$	0.236068	no improvement over $n = 5$
7	853	C_7	7	3	3.317667	0.317667	[9]
8	11,117	GCQb`o	10	3	3.467844	0.467844	Quad- C_5 [9]; $R(3, 4)$ -graph
9	261,080	HCRbdO{	15	3	11/3	2/3	new; $ \text{Aut} = 12$, one triangle
10	11,716,571	ICRb`yiu?	20	3	$3 + 1/\sqrt{2}$	$1/\sqrt{2}$	new; 4-regular, $ \text{Aut} = 16$
11	1,006,700,565	J?`D@pgd?{?	17	4	4.774889	0.774889	new; triangle-free, $R(3, 5)$ -graph

3 Exhaustive results

3.1 All connected graphs, $n \leq 11$

Table 1 lists the maximizers of Δ over all connected graphs for $n = 5, \dots, 11$; the values for $n \leq 8$ reproduce [9] exactly (including the full top-ten ranking for $n = 8$, Appendix A). Figure 1 draws the record graphs.

$n = 9$. The maximizer HCRbdO{ has 15 edges, degree sequence $3^6 4^3$, automorphism group of order 12 with vertex orbits of sizes 3 and 6, twelve induced pentagons and a single triangle. Its theta number is $11/3$ to twelve digits (Clarabel with tolerances 10^{-12}), so $\Delta = 2/3$ exactly; the optimal X has rank 4. The runner-up (HCRb`qi, 16 edges) is the Wagner graph plus one vertex and has $\vartheta = 2 + \varphi = 3.618034$, φ the golden ratio; it is followed by a large plateau of graphs at exactly Quad- C_5 ’s value 0.467844 (Quad- C_5 plus a vertex that changes neither ϑ nor α). The maximizer contains neither Quad- C_5 nor the Wagner graph as an induced subgraph. Its adjacency spectrum is $\{2 + \sqrt{2}, 1^{(2)}, (\sqrt{2} - 1)^{(2)}, 2 - \sqrt{2}, -2, (-1 - \sqrt{2})^{(2)}\}$.

$n = 10$. The maximizer ICRb`yiu? is 4-regular with 20 edges,  $|\text{Aut}| = 16$  (orbits of sizes 2 and 8), sixteen induced pentagons and four triangles, and contains the Wagner graph as an induced subgraph.  $\vartheta = 3 + 1/\sqrt{2}$  to twelve digits. The runner-up (ICQeR`[Mg, $\Delta = 0.667155$) contains both Quad- C_5 and the $n = 9$ maximizer; the “ $n = 9$ maximizer plus a vertex” lineage therefore reaches only second place, as the “Wagner plus a vertex” lineage did at $n = 9$. The values for $n = 9$ and $n = 10$ can in principle be read off the table of Cabello, Danielsen, López-Tarrida and Portillo, who computed α , ϑ (to four decimals) and α^* for all graphs with $\alpha < \vartheta$ on up to ten vertices [4, 5]; our runs reproduce that table graph-for-graph (16,533 graphs at $n = 9$, 975,330 at $n = 10$, $|\vartheta - \vartheta_{\text{table}}| \leq 5.3 \times 10^{-5}$). The maxima were, however, never reported.

$n = 11$. Here the picture changes. The five best graphs (Table 2) are all triangle-free, have $\alpha = 4$ and only 17–19 edges, while the best graph of the dense family that dominated $n \leq 10$ — the $n = 10$ maximizer plus one vertex, $\Delta = 0.735505$, which contains Quad- C_5 , the Wagner graph and the $n = 9$ and $n = 10$ maximizers as induced subgraphs — is only sixth. The maximizer J?`D@pgd?{? has girth 4, degree sequence $2^1 3^8 4^2$, $|\text{Aut}| = 2$, eleven induced pentagons and four induced heptagons, and an optimal X of rank 4. Note that $\Delta_{\max}(11) - \Delta_{\max}(10) = 0.068$ is the smallest increment in the table; the sparse family is only just taking over. Figure 2 shows all 100 survivors of the screen:




Figure 1: Record graphs. Top: the exhaustive maximizers for $n = 8$ (Quad- C_5 , [9]), $n = 9$ and $n = 10$. Bottom: the exhaustive maximizer for $n = 11$, the triangle-free maximizer for $n = 12$, and the triangle-free maximizer for $n = 13$, which is the circulant Ramsey graph $C_{13}(1, 5)$.

the two families are separated cleanly by edge count, and every graph above the $n = 10$ record with fewer than 20 edges is triangle-free.

Gap versus ratio. The maximizer of the ratio ϑ/α is a different graph from the gap maximizer at $n = 8, 10, 11$ (at $n = 10$ it is IUZurzmoo with $\alpha = 2$, $\vartheta = 5/2$; at $n = 11$ the dense graph in row 6 of Table 2); only at $n = 9$ do the two coincide.

3.2 Triangle-free graphs, $n = 12, 13, 14$

Motivated by the $n = 11$ result we searched exhaustively over connected triangle-free graphs, generated with `geng -t` [16]: 90,842 graphs for $n = 11$ (recovering the global top six), 1,144,061 for $n = 12$, 19,425,052 for $n = 13$ and 445,781,050 for $n = 14$ (6.1 hours on the GPU). The maxima for $n = 12, 13$ are

$$\begin{aligned} \Delta_{\max}^{\text{tf}}(12) &= 0.930021 \quad (\vartheta = 4.930021, \alpha = 4, 20 \text{ edges, } K_3\text{-free, } |\text{Aut}| = 8, \text{ rank } X = 5), \\ \Delta_{\max}^{\text{tf}}(13) &= 1.181737 \quad (\vartheta = 5.181737, \alpha = 4, 26 \text{ edges, } L_3(2, 1) \text{ at } YJODg). \end{aligned}$$

At $n = 13$ the maximizer is isolated: a second exhaustive pass over the class with threshold 0.80 shows that the next best connected triangle-free graph on 13 vertices has $\Delta = 0.854480$ ($\vartheta = 5.854480$, $\alpha = 5$, 19 edges, $L_3(2, 1)$ at AAD?oFAgDOAw); it is also the best connected graph with $\alpha = 5$ in the $R(3, 6)$




Figure 2: The 100 graphs on eleven vertices whose certified bound survived the screen (threshold $1/\sqrt{2}$), with their exact gaps: the sparse triangle-free family ($\alpha = 4$) and the dense family ($\alpha = 3$, built around the ten-vertex maximizer).

Table 2: The ten largest gaps among all 1,006,700,565 connected graphs on eleven vertices. “tf” = triangle-free.

rank	graph6	$ E $	α	ϑ	Δ	remarks
1	J?`D@pgd?{?	17	4	4.774889	0.774889	tf, $R(3, 5)$ -graph
2	J?`@f?kUDG_	17	4	4.754686	0.754686	tf
3	J?`@f@WFTH?	18	4	4.753664	0.753664	tf, contains Wagner
4	J?`FBQoFCi?	18	4	4.752419	0.752419	tf
5	J?`@f@Wl?y?	18	4	4.747377	0.747377	tf, contains Quad- C_5
6	JCRbeo{ifI?	25	3	3.735505	0.735505	$n=10$ maximizer + vertex
7	J?`FAqSFf_?	19	4	4.734338	0.734338	tf
8	J?`FAosFdI?	19	4	4.725688	0.725688	tf
9	J?`DD`gD`s?	17	4	4.719179	0.719179	tf
10	JCRbeo{ifM?	26	3	3.712900	0.712900	dense family

catalogue on 13 vertices, all 275,086 members of which we solved exactly. The maximizer is 4-regular with $|\text{Aut}| = 52$ and is isomorphic to the circulant graph $C_{13}(1, 5)$, the unique $(3, 5)$ -Ramsey-critical graph [13]. At $n = 14$, the first size at which triangle-free graphs must have $\alpha \geq 5$, the exhaustive maximum drops to

$$\Delta_{\max}^{\text{tf}}(14) = 1.072373 \quad (\vartheta = 6.072373, \alpha = 5, 24 \text{ edges, } \text{M?AA@agwAg@WM_Dc?}),$$

attained by an $R(3, 6)$ -graph — the best connected graph of the $R(3, 6)$ catalogue on 14 vertices (Table 3) — and all 12,764 triangle-free graphs on 14 vertices with certified bound above 1.0 have $\alpha = 5$. This is the drop predicted in Section 4, now established exhaustively rather than only within the catalogue.

4 The Ramsey structure of the maximizers

4.1 The record graphs at the critical sizes are $R(3, k)$ -graphs

For readers less used to Ramsey numbers, the mechanism in one paragraph. $R(3, k)$ is the least n such that every graph on n vertices contains a triangle or k pairwise non-adjacent vertices. Read the other way round: a triangle-free graph on $n \geq R(3, k)$ vertices is forced to have $\alpha \geq k$, while on $n = R(3, k) - 1$ vertices there still exist triangle-free graphs with $\alpha = k - 1$, and these are by definition the $R(3, k)$ -graphs (“critical” when $n = R(3, k) - 1$). Since $R(3, 4) = 9$, $R(3, 5) = 14$, $R(3, 6) = 18$, a triangle-free graph can have $\alpha = 3$ only up to 8 vertices, $\alpha = 4$ only up to 13, $\alpha = 5$ only up to 17. For the gap this matters because $\Delta = \vartheta - \alpha$ rewards a small α at fixed n , and ϑ of a sparse graph keeps growing with n ; so the best place to be is the last n at which α can still be held at $k - 1$.

By Lemma 2, a triangle-free graph with $\alpha \leq k - 1$ is an $R(3, k)$ -graph and exists only for $n < R(3, k)$. Checking the record graphs against McKay’s catalogues by isomorphism gives:

- $n = 8$: Quad- C_5 and the Wagner graph are two of the three $R(3, 4)$ -graphs on 8 vertices ($R(3, 4) = 9$).
- $n = 9, 10$: no triangle-free graph has $\alpha = 3$ any more; the maximizers keep $\alpha = 3$ by admitting one, respectively four, triangles.
- $n = 11$: the maximizer is one of the 105 $R(3, 5)$ -graphs on 11 vertices; indeed the 105 connected triangle-free graphs with $\alpha \leq 4$ on 11 vertices are exactly this catalogue.
- $n = 12$: the triangle-free maximizer is one of the 12 $R(3, 5)$ -graphs on 12 vertices.
- $n = 13$: the triangle-free maximizer is the unique $R(3, 5)$ -graph on 13 vertices, $C_{13}(1, 5)$.

The mechanism is transparent: ϑ grows with n (along the record family at least linearly, Section 4.4), whereas α is bounded below by Ramsey theory and jumps by one at every $n = R(3, k)$. Records are therefore expected just before the jumps, at $n = R(3, k) - 1$, and the gap should *decrease* within the triangle-free class right after a jump. Both regime changes we observe sit exactly on Ramsey numbers: at $n = 9 = R(3, 4)$ the triangle-free graphs with $\alpha = 3$ disappear and the maximizers switch to graphs with triangles; at $n = 14 = R(3, 5)$ the triangle-free graphs with $\alpha = 4$ disappear and the triangle-free maximum drops from 1.18 to 1.07. In between, at $n = 11$, the growth of ϑ has made $\alpha = 4$ triangle-free graphs competitive again, and they overtake the dense $\alpha = 3$ family.

4.2 Circulant Ramsey-critical graphs and closed forms

Two of the Ramsey-critical vertex counts carry circulant critical graphs, and for those the theta number can be computed in closed form. For a vertex-transitive graph the optimal X in (1) can be taken invariant under the automorphism group [2]; for a circulant $C_n(S)$ this reduces the SDP to a linear program over the Fourier coefficients of a symmetric vector $x \in \mathbb{R}^n$: $\vartheta = n \sum_j x_j$ maximized subject to $x_0 = 1/n$, $x_s = 0$ for $s \in \pm S$, and $\hat{x}_k = \sum_j x_j \cos(2\pi jk/n) \geq 0$ for all k . Its optimum is therefore an algebraic number in the maximal real subfield of $\mathbb{Q}(\zeta_n)$. When the graph is moreover edge-transitive, the Lovász–Hoffman bound is attained and the value is explicit [2]:

$$\vartheta(G) = \frac{n(-\lambda_{\min})}{\lambda_{\max} - \lambda_{\min}} \quad (G \text{ edge-transitive}). \quad (4)$$

Proposition 1. (i) *The unique $R(3, 5)$ -critical graph is the circulant $C_{13}(1, 5)$; it is edge-transitive, since the multiplier group $\{\pm 1, \pm 5\} \subset \mathbb{Z}_{13}^\times$ acts transitively on the connection set $\{\pm 1, \pm 5\}$ ($5^2 \equiv -1$). Its eigenvalues are $\lambda_j = 2 \cos(2\pi j/13) + 2 \cos(10\pi j/13)$, so $\lambda_{\max} = 4$, $\lambda_{\min} = 2 \cos(8\pi/13) - 2 \cos(\pi/13)$ (multiplicity 4) and*

$$\vartheta(C_{13}(1, 5)) = \frac{13(-\lambda_{\min})}{4 - \lambda_{\min}} = 5.181736625\dots, \quad \Delta_{\max}^{\text{tf}}(13) = \vartheta - 4 = 1.181736625\dots \quad (5)$$

(ii) *The unique $R(3, 9)$ -critical graph, on 35 vertices, is Kalbfleisch's circulant $C_{35}(1, 7, 11, 16)$ [12]; it is vertex-transitive but not edge-transitive, and $\vartheta = 12.410586\dots$ is the optimum of the Fourier linear program (the bound (4) gives 12.4979 and is not attained). Hence $\Delta_{\max}(35) \geq 4.410586$.*

(iii) *There is no circulant $R(3, k)$ -critical graph on $R(3, k) - 1$ vertices for $k = 6, 7, 8$ ($n = 17, 22, 27$); the circulant $R(3, k)$ -graphs that exist on 16 and 21 vertices are not the gap maximizers of their families (1.1648 vs 1.4654; 1.9768 vs 2.1413).*

The scan behind (iii) is exhaustive over connection sets of connected circulants (all $2^{\lfloor n/2 \rfloor}$ symmetric sets, triangle-freeness, α by clique search on the complement); a disconnected circulant on these vertex counts has isomorphic components and independence number at least twice that of a component, so it cannot be critical. The identifications in (i) and (ii) are by isomorphism test against McKay's catalogues, whose uniqueness statements for $(3, 5, 13)$ and $(3, 9, 35)$ are those of [13].

We note explicitly what Section 4 does and does not establish. It establishes that α of the triangle-free record graphs is dictated by Ramsey numbers (Lemma 2), it identifies every computed triangle-free maximizer, and the global maximizers at $n = 8$ and $n = 11$, with Ramsey graphs, and for the two circulant cases it gives ϑ in closed form. It does not provide a theorem singling out the Ramsey-critical graphs: a lower bound on ϑ for triangle-free graphs of small independence number, strong enough to show that the critical graphs beat everything else at $n = R(3, k) - 1$, is the missing ingredient. The general lower bounds we are aware of (through $\vartheta(G)\vartheta(\bar{G}) \geq n$ and $\chi(G) = O(\sqrt{n})$) are of order $\sqrt{n}$, below $k - 1 \approx \sqrt{n \log n}$ at $n = R(3, k) - 1$, and do not even certify positivity of the gap; what *can* be proved about growth is in Section 4.4.

4.3 A ladder of lower bounds up to $n = 35$

Since the $R(3, k)$ -graphs are completely catalogued for $k \leq 9$ [15, 14], and $\alpha = k - 1$ on all of them for $n \geq R(3, k - 1)$ (Lemma 2), the gap over each catalogue requires only ϑ . Table 3 gives the maxima; together with Lemma 1 they are lower bounds on $\Delta_{\max}(n)$ for all larger n . Figure 3 summarizes the results.

The $R(3, 6)$ -graphs on 13 vertices (275,086 graphs, for which $\alpha \in \{4, 5\}$) confirm the picture from the other side: the single graph with $\alpha = 4$ is $C_{13}(1, 5)$ with $\Delta = 1.181737$, and the best of the 275,085 graphs with $\alpha = 5$ reaches only 0.930021 (it is the twelve-vertex triangle-free maximizer plus an isolated vertex).

Within the triangle-free class the maximum gap is *not* monotone: it falls from 1.1817 at $n = 13$ to 1.0724 at $n = 14$ (both exhaustive), exactly as predicted by the jump of α at $R(3, 5) = 14$, and recovers from $n = 15$ on. The global $\Delta_{\max}(n)$ is monotone by Lemma 1, but the graphs realizing the monotone extension (a record graph plus universal vertices) are degenerate as witnesses. We are led to:

Table 3: Maximum gap over all $R(3, k)$ -graphs on n vertices (McKay’s catalogues; $\alpha = k - 1$ analytically, ϑ by SDP). These are lower bounds on $\Delta_{\max}(n)$. For $n = 14$ the catalogue maximum is $C_{13}(1, 5)$ plus an isolated vertex; the best connected graph is shown.

k	n	#graphs	α	$\vartheta_{\max}$	Δ	best graph (graph6)
5	13	1	4	5.181737	1.181737	$C_{13}(1, 5)$
6	14	263,520	5	6.072373	1.072373	MqG00H@_gQ?oGFCJ? (connected)
6	15	64,732	5	6.250964	1.250964	NoCOWGACGQIKGePIDI?
6	16	2,576	5	6.465350	1.465350	OoC?Igi_?E_eKcRCE@iO[
6	17	7	5	6.606664	1.606664	P@oGQA?`ACcdCtDYC]Ah?goC
7	21	1,118,436	6	8.141315	2.141315	THoGG?JODHsOAdS_wg?`GgSCZ??ig@@D@?GT
7	22	191	6	8.398747	2.398747	UsaC?GGC?DccQDCpKEJAOKW`Bo?kD_[_okHCUR??
8	27	477,142	7	10.060569	3.060569	(Appendix A)
9	35	1	8	12.410586	4.410586	the unique $(3, 9)$ -critical graph




Figure 3: (a) Maximum contextuality gap versus number of vertices: exhaustive over all connected graphs ($n \leq 11$), exhaustive over triangle-free graphs ($n \leq 14$), and lower bounds from the complete catalogues of $R(3, k)$ -graphs. Vertical lines mark the Ramsey numbers $R(3, k)$, where α is forced up by one for triangle-free graphs. (b) Gap per vertex of the record graphs, with the bounds $0.126 \leq c^* \leq \frac{1}{2}$ of Proposition 3; the sequence is still increasing at $n = 35$.

Call a maximizer of $\Delta_{\max}(n)$ *new* if it is not obtained from a maximizer on fewer vertices by adding universal vertices (Lemma 1); the degenerate extensions are never triangle-free, and they carry no information beyond the smaller record.

Conjecture 1. (a) For $n = R(3, k) - 1$ with $k \geq 4$, $\Delta_{\max}(n)$ is attained by an $R(3, k)$ -critical graph. (b) For $n \geq 11$, every new maximizer is triangle-free, with the least independence number that Ramsey theory allows on n vertices.

We state this as a conjecture with the appropriate caution. The evidence is: the exhaustive result at $n = 11$ (the only value where the statement is verified over all graphs), the exhaustive triangle-free results at $n = 12, 13$, the fact that the mechanism is forced for the two graph invariants separately (α by Ramsey theory, ϑ by its growth with n for sparse graphs), and the failure of the “previous winner plus a vertex” lineages. The evidence against unrestricted extrapolation is equally clear: for $n = 9, 10$ the maximizers contain triangles, because no triangle-free graph on nine or more vertices has $\alpha = 3$, so an $R(3, 4)$ -type structure with one or four triangles wins. What the conjecture does *not* say is that Ramsey-critical graphs dominate between the critical vertex counts, nor that

every maximizer is triangle-free: at $n = 14$ the triangle-free maximum drops (Section 4.4) and the global maximum is attained only by the degenerate extension of $C_{13}(1, 5)$, which has triangles and is not a new maximizer. A brute-force test at $n = 12$ would require 1.64×10^{11} graphs; a local search over graphs with triangles at $n = 12$ (Section 6) found nothing above the triangle-free record.

4.4 What can be proved: a linear upper bound and linear growth

The Ramsey mechanism pins α from below; the following elementary bound pins Δ from above.

Proposition 2. *For every graph G on n vertices, $\Delta(G) \leq \nu(G) \leq \lfloor n/2 \rfloor$, where $\nu(G)$ is the matching number.*

Proof. By the sandwich theorem $\vartheta(G) \leq \bar{\chi}(G)$, the clique cover number [3]. Covering the 2ν vertices of a maximum matching by its ν edges and the remaining vertices by singletons gives $\bar{\chi}(G) \leq n - \nu$. By the Tutte–Berge formula there is a set S with $n - 2\nu = \text{odd}(G - S) - |S|$; one vertex from each odd component of $G - S$ forms an independent set, so $\alpha(G) \geq \text{odd}(G - S) \geq n - 2\nu$. Hence $\Delta = \vartheta - \alpha \leq (n - \nu) - (n - 2\nu) = \nu$. $\square$

The bound is far from tight for small n ($\nu = 5$ versus $\Delta_{\max} = 0.775$ at $n = 11$), but it has the right order, and the matching lower bound is elementary: the gap is superadditive, so every record graph is a certificate of linear growth.

Proposition 3. *Let $f(n)$ be the maximum of $\Delta(G)$ over all graphs (connected or not) on n vertices. Then $f(m + n) \geq f(m) + f(n)$, the limit*

$$c^* = \lim_{n \rightarrow \infty} \frac{\Delta_{\max}(n)}{n} = \lim_{n \rightarrow \infty} \frac{f(n)}{n} = \sup_G \frac{\Delta(G)}{|V(G)|}$$

exists, and for every graph H on h vertices and every $n \geq 1$,

$$\Delta_{\max}(n) \geq \left\lfloor \frac{n-1}{h} \right\rfloor \Delta(H).$$

In particular, with $H = C_{35}(1, 7, 11, 16)$,

$$0.1260 \leq \frac{\Delta(C_{35}(1, 7, 11, 16))}{35} \leq c^* \leq \frac{1}{2}, \quad \Delta_{\max}(n) \geq 0.126n - 4.6 \quad (n \geq 1). \quad (6)$$

Proof. Both ϑ and α are additive over disjoint unions [3], so Δ is additive and f is superadditive; $f(n) \leq n/2$ by Proposition 2, so Fekete’s lemma gives $\lim f(n)/n = \sup f(n)/n$. Adding a vertex adjacent to everything changes neither ϑ nor α (proof of Lemma 1) and makes the graph connected, hence $f(n-1) \leq \Delta_{\max}(n) \leq f(n)$ and the two limits coincide. For the explicit bound take $\lfloor (n-1)/h \rfloor$ disjoint copies of H and fill up with universal vertices. The numbers in (6) are those of Table 3. $\square$

Proposition 3 is what Figure 3 shows: the slope of the Ramsey ladder is a lower bound on c^* , and each new record graph with a larger ratio Δ/n raises it. Our data give $\Delta/n = 0.091, 0.095, 0.109, 0.113, 0.126$ for the record graphs on 13, 17, 22, 27, 35 vertices, so the best current bound comes from the largest Ramsey-critical graph, and the sequence is still increasing. On the Ramsey side, combining (6) with Lemma 1 and the lower bound $R(3, k) \geq (\frac{1}{4} - o(1))k^2 / \ln k$ [18, 19] gives

Corollary 1. $\Delta_{\max}(R(3, k) - 1) \geq 0.126 R(3, k) - 4.8 = \Omega(k^2 / \ln k)$.

This explains the growth of the ladder in Table 3 through the critical vertex counts; it does not explain why the maximizers at those counts are the Ramsey-critical graphs themselves, which remains Conjecture 1. Two heuristics point to a limit strictly between the bounds: for the binomial random graph $G(n, d/n)$, $\vartheta = \Theta(n/\sqrt{d})$ [20] while $\alpha \approx (2n/d) \ln d$ [21], so sparse random graphs form a second, non-Ramsey family with linear gap; and the upper bound $\nu \leq n/2$ is attained only when $\vartheta = \bar{\chi}$ and $\alpha = n - 2\nu$ simultaneously, which no graph in our tables approaches. Determining c^* , and whether the maximizers stay triangle-free, are the natural open problems.

5 Methods

5.1 Enumeration and exact computation

Connected graphs on up to ten vertices were taken from McKay’s lists and, for eleven vertices and for the triangle-free classes, generated with `geng` from `nauty` 2.8.9 [16], split into resumable `res/mod` parts. α was computed exactly by enumerating all 2^n vertex subsets (bit operations; on the GPU for $n \geq 11$). ϑ was computed with the interior-point solver Clarabel [17] on the primal (1) with tolerances 10^{-9} (and 10^{-12} for the record graphs), at about 8,000 graphs per second on 20 CPU processes; the full $n = 10$ run took 51 minutes.

5.2 Certified GPU screening for $n = 11$

Exact SDPs for 10^9 graphs would take several days. Instead we use the dual (2) as a source of certificates and a GPU as a way to search for them. We separate the two roles explicitly.

The certificate. For any graph G and *any* symmetric matrix Y with $Y_{ij} = 0$ whenever $ij \notin E$ (diagonal included), weak duality gives $\vartheta(G) \leq \lambda_{\max}(J+Y)$. Thus a pair (Y, λ) with $\lambda \geq \lambda_{\max}(J+Y)$ is a proof that $\vartheta(G) \leq \lambda$, and if moreover $\lambda - \alpha(G) < \Delta_{\text{thr}}$ then G cannot beat the threshold. Nothing about how Y was found enters this statement. In our implementation Y is a single-precision matrix whose support is enforced exactly by masking, so it is an exactly known rational matrix; the only numerical step in the certificate is the evaluation of $\lambda_{\max}(J+Y)$, done with the batched Jacobi eigensolver of `cuSOLVER` (through `torch.linalg.eigh`), which is backward stable. The computed largest eigenvalue of an $n \times n$ symmetric matrix differs from the exact one by at most $c n u \|J+Y\|_2$ ($u = 2^{-24}$ in single precision, c a modest constant); in our runs the entries of Y stayed below 7 in magnitude and $\|J+Y\|_2$ below 15, so the error is at most $c \cdot 10^{-5}$ and $\lambda := \lambda_{\max}^{\text{computed}} + 10^{-3}$ is a valid upper bound with a factor of about a hundred to spare. The survivors’ final certificates are recomputed in double precision, where the margin is 10^9 times the error, and their α is recomputed independently on the CPU in the exact stage. The run is split into 50 `res/mod` parts of `geng`, each producing its own survivor file (in the repository), so it is resumable and auditable part by part; the whole pipeline for $n = 11$, including the exact stage, took under three hours of wall time.

The search. For a batch of 2^{15} graphs we optimize Y per graph with 100 Adam steps on the softmax-smoothed largest eigenvalue $\tau \log \sum_i e^{\lambda_i/\tau}$ (τ annealed from 0.3 to 0.02; gradient $V \text{diag}(\text{softmax}(\lambda/\tau)) V^T$ masked to E), using batched single-precision eigendecompositions. Every ten steps the certificate test above is applied and graphs that fail it are removed from the batch; since every intermediate Y is a valid certificate, an early removal is never a false elimination, and a poor optimization can only cost time (a graph that should have been eliminated survives to the exact stage). Survivors are re-solved exactly with Clarabel. With $\Delta_{\text{thr}} = 1/\sqrt{2} - 10^{-6}$ (the $n = 10$ record, valid by Lemma 1), the screen processed all 1,006,700,565 graphs in 2.41 hours on




Figure 4: Validation of the GPU screen. (a) Certified upper bound minus exact ϑ for all connected graphs on eight vertices, without pruning. (b) At $n = 10$: number of graphs with $\Delta \geq x$ from the exact run, and the number of survivors of the screen at threshold 0.65, which coincides with the exact count.

an RTX 2000 Ada (8 GB) at $0.95\text{--}1.3 \times 10^5$ graphs per second, leaving 100 survivors, 51 of which indeed have $\Delta \geq 1/\sqrt{2}$.

Validation. With no threshold, the certified bounds exceed ϑ by a median of 0.006 (max 0.21) over the 11,117 graphs on eight vertices, and never fall below ϑ (Figure 4a). With thresholds 0.60 at $n = 9$ and 0.65 at $n = 10$, the survivors are exactly the two, respectively thirty, graphs known to exceed the threshold (from the exact runs and from [5]), with no misses and no false survivors (Figure 4b).

5.3 Ramsey catalogues

For the $R(3, k)$ -graphs we used McKay’s files `r35_13`, `r36_14`–`r36_17`, `r37_21`, `r37_22`, `r38_27`, `r39_35` [15]; $\alpha = k - 1$ on all of them by Lemma 2, and ϑ was computed with Clarabel (74 minutes for the 1,118,436 graphs on 21 vertices, 88 minutes for the 477,142 graphs on 27 vertices).

6 Discussion

Certificates. For every record graph we provide an explicit primal feasible matrix X and an explicit dual feasible matrix Y of (1)–(2), recomputed in double precision independently of the solver; they bracket ϑ in an interval of width below 10^{-9} (Appendix B, Table 5), and the values were reproduced with a second solver (SCS). The exact values $11/3$ and $3 + 1/\sqrt{2}$ for $n = 9, 10$ are thus certified to nine digits; closed forms for the eleven-vertex maximizer and for the triangle-free maximizers on 12 vertices remain to be identified (their minimal polynomials are not obvious from the spectrum, which for $n = 11$ contains $2 + \sqrt{2}$ but is otherwise irrational).

Implications for experiments. From the optimal $X = B^\top B$ one reads off an orthogonal representation of the exclusivity graph in the CSW sense: unit vectors $u_i = b_i/\|b_i\|$, orthogonal for exclusive events, and a state $\psi \propto \sum_i b_i$ with $\sum_i \langle \psi, u_i \rangle^2 = \vartheta(G)$ (Cauchy–Schwarz gives $\geq$, Lovász’s theorem gives $\leq$). The rank of the maximum-rank optimal X is 4 for all maximizers with $n \leq 11$

and for Quad- C_5 and Wagner, and 5 for the triangle-free maximizers on 12 and 13 vertices, so all witnesses are realizable with a ququart, respectively a five-level system. We stress that this is an *upper* bound on the minimal dimension: an interior-point solver returns the analytic centre of the optimal face, i.e. a maximum-rank optimum, and we make no claim that four (or five) is minimal; settling the minimal dimension requires a separate rank-constrained computation, as done for Quad- C_5 in [9]. Following [9], the critical visibility under white noise — the least v for which $v\vartheta + (1-v)n/d > \alpha$, i.e. $v^* = (\alpha - n/d)/(\vartheta - n/d)$, smaller being more robust — is 0.529 for the $n = 9$ maximizer and 0.414 for the $n = 10$ maximizer (with $d = 4$), against 0.681 for Quad- C_5 and 0.707 for the Wagner graph; for the eleven-vertex maximizer $v^* = 0.617$ and for $C_{13}(1, 5)$ (in $d = 5$) $v^* = 0.542$. Two conclusions for experiment design follow. First, the new nine- and ten-vertex witnesses are markedly more robust than the eight-vertex ones: the ten-vertex maximizer tolerates 59% white noise, the Wagner graph 29%. Second, *a larger gap does not imply a more robust witness*: the eleven-vertex maximizer has a larger gap than the ten-vertex one (0.775 vs 0.707) but needs visibility 0.617 instead of 0.414, and $C_{13}(1, 5)$, with gap 1.18, needs 0.542. The reason is the noise term n/d , which grows with n at fixed dimension; the sparse Ramsey-type witnesses pay for their larger gaps with more vertices per dimension. For a given experimental dimension the right figure of merit is v^* , and on that count the dense ten-vertex graph is the best witness in our tables. The quoted v^* refer to the realizations we exhibit (dimension r); should a lower-dimensional realization of the same graph exist, its noise term n/d would be larger and its v^* worse, so the values above are the best we can certify, not the best possible.

Beyond triangle-free at $n = 12$. A local search over all graphs on twelve vertices, with no triangle-free restriction — edge additions, deletions and moves with plateau acceptance and random restarts, started from the triangle-free maximizers and from the eleven-vertex records extended by one vertex, with exact ϑ and α at every step — evaluated 1,425,556 distinct graphs in 90 minutes. The best gap found was the triangle-free record 0.930021 itself; the best graph containing a triangle reached $\Delta = 0.866462$ (one triangle, 21 edges, $\alpha = 4$). This is evidence, not proof, for Conjecture 1 at $n = 12$.

Relation to previous work. The theta number was introduced to bound the Shannon capacity, $\alpha(G) \leq \Theta(G) \leq \vartheta(G)$, with $\vartheta(C_5) = \sqrt{5}$ as the founding example [2]; its interplay with independence numbers of graph powers, and through $\vartheta(G)\vartheta(\bar{G}) \geq n$ with Ramsey-type constructions, is surveyed in [22]. In that literature ϑ is a tool for upper-bounding α of large product graphs; here the object is the *difference* $\vartheta - \alpha$ of the Ramsey graphs themselves, as a measure of contextuality, and the Ramsey numbers enter through the vertex counts at which α is forced up, not through ϑ . The graph-enumeration programme of [4, 7] computed α , ϑ and α^* for all graphs with up to ten vertices and studied ratio maximizers and “fully contextual” graphs ($\vartheta = \alpha^*$); none of the gap maximizers found here is fully contextual (e.g. $\alpha^* = 13/2$ for $C_{13}(1, 5)$), and the two figures of merit select different graphs from $n = 8$ on.

7 Conclusion

We have pushed the exhaustive maximization of the absolute contextuality gap from eight to eleven vertices, over a billion graphs, with a GPU screening scheme whose every elimination is backed by a dual certificate, and we have found that the answer has a structure: at the Ramsey-critical sizes the record graphs are Ramsey $R(3, k)$ -graphs, triangle-free graphs whose independence number is held at $k - 1$ by $n < R(3, k)$ while their theta number keeps growing, and in between the maximizers

keep the previous independence number by admitting triangles. The thirteen-vertex triangle-free maximizer is the Ramsey graph $C_{13}(1, 5)$ itself, with a closed-form theta number, and the catalogues of $R(3, k)$ -graphs yield gap records up to 35 vertices. Two questions are left open: whether the maximizer at twelve vertices — beyond exhaustive reach at 1.6×10^{11} graphs, though within reach of a more aggressive screen — is still triangle-free (Conjecture 1), and the value of the linear growth constant $c^* = \lim \Delta_{\max}(n)/n$, which Proposition 3 confines to $[0.126, \frac{1}{2}]$. For experiments, the new witnesses on nine and ten vertices have exact theta numbers $11/3$ and $3 + 1/\sqrt{2}$, realizations in dimension four (from the rank of the SDP optimum; minimality is not claimed), and are given explicitly in the appendix and in the accompanying repository.

Code and data availability

All scripts (exact search, GPU screening, Ramsey catalogue processing), the result tables with graph6 identifiers, and the certificates for the record graphs are available at <https://github.com/danmusetoiu/contextuality-gap>.

A Reproduction of the eight-vertex ranking and graph data

Table 4: Top ten connected graphs on eight vertices by Δ (our run, Clarabel), matching Table II of [9] rank for rank.

rank	graph6	$ E $	α	ϑ	Δ	ϑ/α	
1	GCQb`o	10	3	3.467844	0.467844	1.155948	Quad- C_5
2	GCR`r_	11	3	3.438447	0.438447	1.146149	
3	GCr b`o	12	3	3.414214	0.414214	1.138071	Wagner graph
4	GCRb`w	12	3	3.372281	0.372281	1.124094	
5	GCQbd o	11	3	3.372281	0.372281	1.124094	CHSH-type graph, ratio maximizer
6	GCp`d o	10	3	3.372281	0.372281	1.124094	
7	GQyurg	16	2	2.343146	0.343146	1.171573	
8	GQyurw	17	2	2.338044	0.338044	1.169022	
9	GCpdag	11	3	3.338044	0.338044	1.112681	
10	GUZurw	18	2	2.333333	0.333333	1.166667	

Edge lists of the record graphs Vertices are $0, \dots, n-1$ in the labelling induced by the graph6 string.

- Quad- C_5 ($n = 8$), GCQb`o: 03, 05, 14, 16, 25, 26, 27, 36, 37, 47.
- $n = 9$ maximizer, HCRbd0{: 03, 05, 07, 14, 15, 16, 25, 26, 27, 28, 36, 38, 47, 48, 58.
- $n = 10$ maximizer, ICRb`yiu?: 03, 05, 08, 09, 14, 15, 16, 19, 25, 26, 27, 28, 36, 37, 39, 47, 48, 49, 57, 68.
- $n = 11$ maximizer, J?`D@pgd?{?: (0,4), (0,6), (0,9), (1,5), (1,8), (2,6), (2,7), (2,8), (3,7), (3,9), (3,10), (4,7), (4,8), (4,10), (5,9), (5,10), (6,10).
- $n = 12$ triangle-free maximizer, K?`DA`gd?{Dg: (0,4), (0,6), (0,9), (1,5), (1,7), (1,8), (2,6), (2,8), (2,11), (3,7), (3,9), (3,10), (4,8), (4,10), (4,11), (5,9), (5,10), (5,11), (6,10), (7,11).

- $C_{13}(1, 5)$, L?`DE`g1@YJODg: (0,4), (0,6), (0,7), (0,9), (1,5), (1,7), (1,8), (1,11), (2,6), (2,8), (2,9), (2,10), (3,7), (3,9), (3,11), (3,12), (4,8), (4,10), (4,11), (5,9), (5,10), (5,12), (6,11), (6,12), (7,10), (8,12).

The $R(3, 8)$ record on 27 vertices graph6 Z@0\MagCADc`e@0\_\_`KoHO_CCHN?_G@D0?d??oCwQ i??GTSPK?Dc?@YdI?c?, $|E| = 91$, $\vartheta = 10.060569$, $\Delta = 3.060569$.

B Certificates

Table 5: Certificates for the record graphs. $[\vartheta_{\text{lb}}, \vartheta_{\text{ub}}]$ is the interval certified in double precision by an explicit primal feasible X (trace one, zero on edges, PSD after eigenvalue clipping; the residual on edge entries is below 10^{-9}) and an explicit dual feasible Y (supported on the edges, $\lambda_{\max}(J + Y)$ recomputed with LAPACK). r is the rank of the maximum-rank optimal X returned by the interior-point solver, hence an upper bound on the minimal dimension of an orthogonal representation attaining ϑ ; $v^* = (\alpha - n/r)/(\vartheta - n/r)$ is the critical visibility under white noise in dimension r . All matrices and the orthogonal representations are in the repository (`certificates/*.json`).

graph	n	$ E $	α	ϑ_{lb}	ϑ_{ub}	r	v^*
Quad- C_5 ($n = 8$)	8	10	3	3.467843730	3.467843730	4	0.6813
Wagner ($n = 8$)	8	12	3	3.414213562	3.414213562	4	0.7071
$n = 9$ maximizer	9	15	3	3.666666667	3.666666667	4	0.5294
$n = 10$ maximizer	10	20	3	3.707106781	3.707106781	4	0.4142
$n = 11$ maximizer	11	17	4	4.774888533	4.774888533	4	0.6173
$n = 11$, rank 2	11	17	4	4.754686045	4.754686045	4	0.6235
$n = 12$ triangle-free max.	12	20	4	4.930021465	4.930021467	5	0.6324
$C_{13}(1, 5)$	13	26	4	5.181736625	5.181736625	5	0.5423
$R(3, 6)$ best, $n = 16$	16	35	5	6.465350297	6.465350300	5	0.5512
$R(3, 6)$ best, $n = 17$	17	40	5	6.606663834	6.606663840	6	0.5742
$R(3, 7)$ best, $n = 22$	22	66	6	8.398747394	8.398747399	6	0.4931

References

- [1] A. Cabello, S. Severini, and A. Winter, Graph-theoretic approach to quantum correlations, *Phys. Rev. Lett.* **112**, 040401 (2014).
- [2] L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inf. Theory* **25**, 1 (1979).
- [3] D. E. Knuth, The sandwich theorem, *Electron. J. Combin.* **1**, A1 (1994).
- [4] A. Cabello, L. E. Danielsen, A. J. López-Tarrida, and J. R. Portillo, Basic exclusivity graphs in quantum correlations, *Phys. Rev. A* **88**, 032104 (2013).
- [5] A. Cabello, L. E. Danielsen, A. J. López-Tarrida, and J. R. Portillo, Database of quantum graphs, https://www.codetables.de/larsed/quantum_graphs/.
- [6] E. Amselem, L. E. Danielsen, A. J. López-Tarrida, J. R. Portillo, M. Bourennane, and A. Cabello, Experimental fully contextual correlations, *Phys. Rev. Lett.* **108**, 200405 (2012).
- [7] G. Cañas, E. Acuña, J. Cariñe, J. F. Barra, E. S. Gómez, G. B. Xavier, G. Lima, and A. Cabello, Experimental demonstration of the connection between quantum contextuality and graph theory, *Phys. Rev. A* **94**, 012337 (2016).

- [8] M. Ray, N. G. Boddu, K. Bharti, L.-C. Kwek, and A. Cabello, Graph-theoretic approach to dimension witnessing, *New J. Phys.* **23**, 033006 (2021).
- [9] U. Tamer, Ö. E. Müstecaplıoğlu, A. Dizdar, and Z. Gedik, The Quad- C_5 graph: maximum contextuality gap on eight vertices, *arXiv:2605.12828* (2026).
- [10] A. Muller and M. Saniga, Automated search for highly contextual Kochen–Specker proofs, *arXiv:2609.19862* (2026).
- [11] A. A. Klyachko, M. A. Can, S. Binicioğlu, and A. S. Shumovsky, Simple test for hidden variables in spin-1 systems, *Phys. Rev. Lett.* **101**, 020403 (2008).
- [12] J. G. Kalbfleisch, Chromatic graphs and Ramsey’s theorem, Ph.D. thesis, University of Waterloo (1966).
- [13] S. P. Radziszowski, Small Ramsey numbers, *Electron. J. Combin.*, Dynamic Survey DS1 (revised regularly).
- [14] J. Goedgebeur and S. P. Radziszowski, New computational upper bounds for Ramsey numbers $R(3, k)$, *Electron. J. Combin.* **20**, P30 (2013).
- [15] B. D. McKay, Ramsey graphs, <https://users.cecs.anu.edu.au/~bdm/data/ramsey.html>.
- [16] B. D. McKay and A. Piperno, Practical graph isomorphism, II, *J. Symbolic Comput.* **60**, 94 (2014).
- [17] P. J. Goulart and Y. Chen, Clarabel: An interior-point solver for conic programs with quadratic objectives, *arXiv:2405.12762* (2024).
- [18] T. Bohman and P. Keevash, Dynamic concentration of the triangle-free process, *Random Structures Algorithms* **58**, 221 (2021).
- [19] G. Fiz Pontiveros, S. Griffiths, and R. Morris, The triangle-free process and the Ramsey number $R(3, k)$, *Mem. Amer. Math. Soc.* **263**, no. 1274 (2020).
- [20] A. Coja-Oghlan, The Lovász number of random graphs, *Combin. Probab. Comput.* **14**, 439 (2005).
- [21] A. M. Frieze, On the independence number of random graphs, *Discrete Math.* **81**, 171 (1990).
- [22] N. Alon, Lovász, vectors, graphs and codes, in *Building Bridges II: Mathematics of László Lovász*, edited by I. Bárány, G. O. H. Katona, and A. Sali, *Bolyai Soc. Math. Stud.* **28** (Springer, Berlin, 2019), pp. 1–16.