

Maximum Contextuality Gap up to Eleven Vertices and the Ramsey Structure of Its Maximizers

Dan Muşetoiu
dan.musetoiu@gmail.com

September 2026

Abstract

In the exclusivity-graph approach to contextuality, a noncontextuality inequality is a graph G , its noncontextual bound is the independence number $\alpha(G)$ and its quantum bound is the Lovász number $\vartheta(G)$. The absolute contextuality gap $\Delta(G) = \vartheta(G) - \alpha(G)$ was recently maximized exhaustively over the 11,117 connected graphs on eight vertices (Quad- C_5 , $\Delta = 0.46784$). We extend the exhaustive search to nine, ten and eleven vertices — 261,080, 11,716,571 and 1,006,700,565 connected graphs respectively — and find $\Delta_{\max} = 2/3$ (with $\vartheta = 11/3$), $\Delta_{\max} = 1/\sqrt{2}$ (with $\vartheta = 3 + 1/\sqrt{2}$) and $\Delta_{\max} = 0.774889$. The search over 10^9 graphs was made feasible on a single consumer GPU by a rigorous screening scheme: for each graph a feasible point of the dual semidefinite program certifies an upper bound on ϑ , and only graphs whose certified bound can beat the running record are solved exactly. The maximizers are not a nested family, and at eleven vertices a regime change occurs: the dense, Wagner-based maximizers with $\alpha = 3$ are overtaken by sparse triangle-free graphs with $\alpha = 4$. We explain this through Ramsey theory: triangle-free graphs with $\alpha \leq k-1$ are exactly the $R(3, k)$ -graphs, and every record graph we find is one — Quad- C_5 and the Wagner graph are $R(3, 4)$ -graphs, the eleven-vertex maximizer is an $R(3, 5)$ -graph, and the triangle-free maximizer on thirteen vertices is the unique $R(3, 5)$ -critical graph $C_{13}(1, 5)$, with $\vartheta = 13(-\lambda_{\min})/(\lambda_{\max} - \lambda_{\min}) = 5.181737$ in closed form. Computing ϑ over McKay’s complete catalogues of $R(3, k)$ -graphs yields a ladder of lower bounds on $\Delta_{\max}(n)$ up to $n = 35$ ($\Delta \geq 4.4106$ from the unique $R(3, 9)$ -critical graph). We release all code, graph6 identifiers and certificates.

Keywords: quantum contextuality, exclusivity graphs, Lovász theta number, independence number, Ramsey graphs, semidefinite programming, exhaustive graph search

1 Introduction

Cabello, Severini and Winter (CSW) showed that any noncontextuality (NC) inequality of the form $\sum_i P(e_i) \leq \alpha$ can be encoded by an *exclusivity graph* G whose vertices are the events e_i and whose edges join mutually exclusive events; the maximum of the left-hand side over noncontextual models is the independence number $\alpha(G)$, and its maximum over quantum models is the Lovász theta number $\vartheta(G)$ [1]. The gap between the two numbers is therefore a graph invariant that measures how contextual an inequality can be. Two natural figures of merit have been studied: the ratio ϑ/α , used to identify experimentally accessible witnesses [6, 7] and as a dimension witness [8], and the absolute gap $\Delta(G) = \vartheta(G) - \alpha(G)$, whose maximization over all connected graphs on eight vertices was recently carried out by Tamer et al. [9]. Their exhaustive search over the 11,117 graphs identified a sparse ten-edge graph, Quad- C_5 , with $\Delta = 0.46784$, beating the Wagner graph ($\Delta = \sqrt{2} - 1$).

Two questions follow. What are the maximizers for more vertices, and is there structure behind them? Here we answer the first question exhaustively for $n = 9, 10, 11$ and, for the triangle-free class, for $n = 12, 13$; and we answer the second by exhibiting a Ramsey-theoretic mechanism.

Contributions.

1. **Exhaustive maxima for $n = 9, 10, 11$.** $\Delta_{\max}(9) = 2/3$ exactly, $\Delta_{\max}(10) = 1/\sqrt{2}$ exactly, $\Delta_{\max}(11) = 0.774889$ (Section 3). The eleven-vertex search covers all 1,006,700,565 connected graphs and is, to our knowledge, the first exhaustive computation of a Lovász-number extremum at this scale.
2. **A rigorous GPU screening method** (Section 5): dual-feasible points of the theta SDP, optimized in large batches on a consumer GPU, certify upper bounds on ϑ that eliminate all but a hundred graphs out of 10^9 ; the survivors are solved exactly. The method is validated against every known answer ($n \leq 10$) with zero misses.
3. **The Ramsey structure of the maximizers** (Section 4). All record graphs are $R(3, k)$ -graphs, i.e. triangle-free graphs with $\alpha \leq k-1$ on fewer than $R(3, k)$ vertices; the thirteen-vertex triangle-free maximizer is the unique $R(3, 5)$ -critical graph $C_{13}(1, 5)$, whose theta number has a closed form. Ramsey numbers thus pin α while ϑ keeps growing, and $\Delta_{\max}(n)$ jumps at $n = R(3, k) - 1$. Computing ϑ over the complete catalogues of $R(3, k)$ -graphs gives a ladder of lower bounds up to $n = 35$.

2 Preliminaries

Throughout, $G = (V, E)$ is a simple graph on n vertices, interpreted as an exclusivity graph. The independence number $\alpha(G)$ is the size of a largest set of pairwise non-adjacent vertices. The Lovász number [2] can be written as the semidefinite program

$$\vartheta(G) = \max\{\langle J, X \rangle : \text{tr } X = 1, X_{ij} = 0 \ \forall ij \in E, X \succeq 0\}, \quad (1)$$

where J is the all-ones matrix, with the dual

$$\vartheta(G) = \min\{\lambda_{\max}(J + Y) : Y = Y^T, Y_{ij} = 0 \ \forall ij \notin E, Y_{ii} = 0\}. \quad (2)$$

The sandwich theorem gives $\alpha(G) \leq \vartheta(G) \leq \alpha^*(G)$, where α^* is the fractional packing number [3]. In the CSW framework $\alpha(G)$ and $\vartheta(G)$ are respectively the noncontextual and the quantum maximum of $\sum_{i \in V} P(e_i)$ over all correlation experiments whose exclusivity graph is G [1]; quantum theory violates the NC inequality iff $\vartheta(G) > \alpha(G)$, which happens iff G contains an induced odd cycle of length at least five or the complement of one [4]. We study the *absolute contextuality gap*

$$\Delta(G) = \vartheta(G) - \alpha(G), \quad \Delta_{\max}(n) = \max\{\Delta(G) : G \text{ connected}, |V(G)| = n\}. \quad (3)$$

Restricting to connected graphs loses nothing for the maximum: ϑ and α are both additive over disjoint unions, so a disconnected graph never beats its best component plus isolated vertices, and isolated vertices contribute equally to ϑ and α .

Two elementary facts will be used repeatedly.

Lemma 1 (Monotonicity). $\Delta_{\max}(n+1) \geq \Delta_{\max}(n)$.

Table 1: Maximum contextuality gap over all connected graphs on n vertices. Graphs are identified by their graph6 string [16]. d^* is the rank of the optimal primal matrix X in (1), i.e. the dimension of the Lovász orthonormal representation attaining ϑ .

n	#graphs	graph6	$ E $	α	ϑ	$\Delta_{\max}(n)$	remarks
5	21	C_5	5	2	$\sqrt{5}$	0.236068	KCBS pentagon
6	112	EUZO	8	2	$\sqrt{5}$	0.236068	no improvement over $n = 5$
7	853	C_7	7	3	3.317667	0.317667	[9]
8	11,117	GCQb`o	10	3	3.467844	0.467844	Quad- C_5 [9]; $R(3, 4)$ -graph
9	261,080	HCRbd0{	15	3	11/3	2/3	new; $ \text{Aut} = 12$, one triangle
10	11,716,571	ICRb`yiu?	20	3	$3 + 1/\sqrt{2}$	$1/\sqrt{2}$	new; 4-regular, $ \text{Aut} = 16$
11	1,006,700,565	J?`D@pgd?{?	17	4	4.774889	0.774889	new; triangle-free, $R(3, 5)$ -graph

Proof. Add a vertex adjacent to all vertices of a maximizer G . Independent sets are unchanged, so α is unchanged. In (1) the new row and column of X must vanish off the diagonal, so $X = \text{diag}(X', s)$ with $\text{tr } X' = 1 - s$ and $\langle J, X \rangle = (1 - s)\vartheta(G) + s \leq \vartheta(G)$; hence ϑ is unchanged as well. $\square$

Lemma 2 (Ramsey pins α). *Let $R(3, k)$ denote the Ramsey number. A triangle-free graph on $n \geq R(3, k - 1)$ vertices has $\alpha \geq k - 1$; a triangle-free graph on $n < R(3, k)$ vertices can have $\alpha = k - 1$, and the graphs achieving it are exactly the $R(3, k)$ -graphs on n vertices.*

Proof. Immediate from the definition of $R(3, k)$ as the least n such that every graph on n vertices contains a triangle or an independent set of size k . $\square$

The relevant values are $R(3, 4) = 9$, $R(3, 5) = 14$, $R(3, 6) = 18$, $R(3, 7) = 23$, $R(3, 8) = 28$, $R(3, 9) = 36$ [13]. Complete catalogues of $R(3, k)$ -graphs for $k \leq 9$ are available from McKay [15, 14].

3 Exhaustive results

3.1 All connected graphs, $n \leq 11$

Table 1 lists the maximizers of Δ over all connected graphs for $n = 5, \dots, 11$; the values for $n \leq 8$ reproduce [9] exactly (including the full top-ten ranking for $n = 8$, Appendix A). Figure 1 draws the record graphs.

$n = 9$. The maximizer HCRbd0{ has 15 edges, degree sequence $3^6 4^3$, automorphism group of order 12 with vertex orbits of sizes 3 and 6, twelve induced pentagons and a single triangle. Its theta number is 11/3 to twelve digits (Clarabel with tolerances 10^{-12}), so $\Delta = 2/3$ exactly; the optimal X has rank 4. The runner-up (HCrb`qi, 16 edges) is the Wagner graph plus one vertex and has $\vartheta = 2 + \varphi = 3.618034$, φ the golden ratio; it is followed by a large plateau of graphs at exactly Quad- C_5 's value 0.467844 (Quad- C_5 plus a vertex that changes neither ϑ nor α). The maximizer contains neither Quad- C_5 nor the Wagner graph as an induced subgraph. Its adjacency spectrum is $\{2 + \sqrt{2}, 1^{(2)}, (\sqrt{2} - 1)^{(2)}, 2 - \sqrt{2}, -2, (-1 - \sqrt{2})^{(2)}\}$.

$n = 10$. The maximizer ICRb`yiu? is 4-regular with 20 edges,  $|\text{Aut}| = 16$  (orbits of sizes 2 and 8), sixteen induced pentagons and four triangles, and contains the Wagner graph as an induced subgraph.  $\vartheta = 3 + 1/\sqrt{2}$  to twelve digits. The runner-up (ICQeR`[Mg, $\Delta = 0.667155$) contains both Quad- C_5 and the $n = 9$ maximizer; the “ $n = 9$ maximizer plus a vertex” lineage therefore reaches only second place, as the “Wagner plus a vertex” lineage did at $n = 9$. The values for




Figure 1: Record graphs. Top: the exhaustive maximizers for $n = 8$ (Quad- C_5 , [9]), $n = 9$ and $n = 10$. Bottom: the exhaustive maximizer for $n = 11$, the triangle-free maximizer for $n = 12$, and the triangle-free maximizer for $n = 13$, which is the circulant Ramsey graph $C_{13}(1, 5)$.

$n = 9$ and $n = 10$ can in principle be read off the table of Cabello, Danielsen, López-Tarrida and Portillo, who computed α , ϑ (to four decimals) and α^* for all graphs with $\alpha < \vartheta$ on up to ten vertices [4, 5]; our runs reproduce that table graph-for-graph (16,533 graphs at $n = 9$, 975,330 at $n = 10$, $|\vartheta - \vartheta_{\text{table}}| \leq 5.3 \times 10^{-5}$). The maxima were, however, never reported.

$n = 11$. Here the picture changes. The five best graphs (Table 2) are all triangle-free, have $\alpha = 4$ and only 17–19 edges, while the best graph of the dense family that dominated $n \leq 10$ — the $n = 10$ maximizer plus one vertex, $\Delta = 0.735505$, which contains Quad- C_5 , the Wagner graph and the $n = 9$ and $n = 10$ maximizers as induced subgraphs — is only sixth. The maximizer J7`D@pgd?{? has girth 4, degree sequence $2^1 3^8 4^2$, $|\text{Aut}| = 2$, eleven induced pentagons and four induced heptagons, and $d^* = 4$. Note that $\Delta_{\max}(11) - \Delta_{\max}(10) = 0.068$ is the smallest increment in the table; the sparse family is only just taking over. Figure 2 shows all 100 survivors of the screen: the two families are separated cleanly by edge count, and every graph above the $n = 10$ record with fewer than 20 edges is triangle-free.

Gap versus ratio. The maximizer of ϑ/α differs from the maximizer of Δ at $n = 8$ (a CHSH-type graph, [9]), $n = 10$ (IUZurzmmo, $\alpha = 2$, $\vartheta = 5/2$, ratio 1.25) and $n = 11$ (the dense graph of rank 6 in Table 2, ratio 1.2452, versus 1.1937 for the gap maximizer); at $n = 9$ the two coincide.




Figure 2: The 100 graphs on eleven vertices whose certified bound survived the screen (threshold $1/\sqrt{2}$), with their exact gaps: the sparse triangle-free family ($\alpha = 4$) and the dense family ($\alpha = 3$, containing the Wagner graph).

Table 2: The ten largest gaps among all 1,006,700,565 connected graphs on eleven vertices. “tf” = triangle-free.

rank	graph6	$ E $	α	ϑ	Δ	remarks
1	J??`D@pgd?{?	17	4	4.774889	0.774889	tf, $R(3,5)$ -graph
2	J??`@f?kUDG_	17	4	4.754686	0.754686	tf
3	J??`@f@WFTH?	18	4	4.753664	0.753664	tf, contains Wagner
4	J??`FBQoFCi?	18	4	4.752419	0.752419	tf
5	J??`@f@Wl?y?	18	4	4.747377	0.747377	tf, contains Quad- C_5
6	JCRbeo{ifI?	25	3	3.735505	0.735505	$n=10$ maximizer + vertex
7	J??`FAqSFf_?	19	4	4.734338	0.734338	tf
8	J??`FAosFdI?	19	4	4.725688	0.725688	tf
9	J??`DD`gD`s?	17	4	4.719179	0.719179	tf
10	JCRbeo{ifM?	26	3	3.712900	0.712900	dense family

3.2 Triangle-free graphs, $n = 12, 13$

Motivated by the $n = 11$ result we searched exhaustively over connected triangle-free graphs, generated with `geng -t` [16]: 90,842 graphs for $n = 11$ (recovering the global top six), 1,144,061 for $n = 12$ and about 1.9×10^7 for $n = 13$. The maxima are

$$\begin{aligned} \Delta_{\max}^{\text{tf}}(12) &= 0.930021 \quad (\vartheta = 4.930021, \alpha = 4, 20 \text{ edges, } K??`DA`gd?\{Dg, |Aut| = 8, d^* = 5), \\ \Delta_{\max}^{\text{tf}}(13) &= 1.181737 \quad (\vartheta = 5.181737, \alpha = 4, 26 \text{ edges, } L??`DE`gl@YJODg). \end{aligned}$$

At $n = 13$ the maximizer is isolated: the next best triangle-free graph has $\Delta = 0.854$ and $\alpha = 5$. The maximizer is 4-regular with $|Aut| = 52$ and is isomorphic to the circulant graph $C_{13}(1, 5)$, the unique $(3, 5)$ -Ramsey-critical graph.

4 The Ramsey structure of the maximizers

4.1 Every record graph is an $R(3, k)$ -graph

By Lemma 2, a triangle-free graph with $\alpha \leq k - 1$ is an $R(3, k)$ -graph and exists only for $n < R(3, k)$. Checking the record graphs against McKay's catalogues by isomorphism gives:

- $n = 8$: Quad- C_5 and the Wagner graph are two of the three $R(3, 4)$ -graphs on 8 vertices ($R(3, 4) = 9$).
- $n = 9, 10$: no triangle-free graph has $\alpha = 3$ any more; the maximizers keep $\alpha = 3$ by admitting one, respectively four, triangles.
- $n = 11$: the maximizer is one of the 105 $R(3, 5)$ -graphs on 11 vertices; indeed the 105 connected triangle-free graphs with $\alpha \leq 4$ on 11 vertices are exactly this catalogue.
- $n = 12$: the triangle-free maximizer is one of the 12 $R(3, 5)$ -graphs on 12 vertices.
- $n = 13$: the triangle-free maximizer is the unique $R(3, 5)$ -graph on 13 vertices, $C_{13}(1, 5)$.

The mechanism is transparent: ϑ grows with n (roughly like $\sqrt{n}$ for sparse graphs), whereas α is bounded below by Ramsey theory and jumps by one at every $n = R(3, k)$. Records are therefore expected just before the jumps, at $n = R(3, k) - 1$, and the gap should *decrease* within the triangle-free class right after a jump.

4.2 Circulant Ramsey-critical graphs and closed forms

Two of the Ramsey-critical vertex counts carry circulant critical graphs, and for those the theta number can be computed in closed form. For a vertex-transitive graph the optimal X in (1) can be taken invariant under the automorphism group [2]; for a circulant $C_n(S)$ this reduces the SDP to a linear program over the Fourier coefficients of a symmetric vector $x \in \mathbb{R}^n$: $\vartheta = n \sum_j x_j$ maximized subject to $x_0 = 1/n$, $x_s = 0$ for $s \in \pm S$, and $\hat{x}_k = \sum_j x_j \cos(2\pi jk/n) \geq 0$ for all k . Its optimum is therefore an algebraic number in the maximal real subfield of $\mathbb{Q}(\zeta_n)$. When the graph is moreover edge-transitive, the Lovász–Hoffman bound is attained and the value is explicit [2]:

$$\vartheta(G) = \frac{n(-\lambda_{\min})}{\lambda_{\max} - \lambda_{\min}} \quad (G \text{ edge-transitive}). \quad (4)$$

Proposition 1. (i) *The unique $R(3, 5)$ -critical graph is the circulant $C_{13}(1, 5)$; it is edge-transitive, since the multiplier group $\{\pm 1, \pm 5\} \subset \mathbb{Z}_{13}^\times$ acts transitively on the connection set $\{\pm 1, \pm 5\}$ ($5^2 \equiv -1$). Its eigenvalues are $\lambda_j = 2 \cos(2\pi j/13) + 2 \cos(10\pi j/13)$, so $\lambda_{\max} = 4$, $\lambda_{\min} = 2 \cos(8\pi/13) - 2 \cos(\pi/13)$ (multiplicity 4) and*

$$\vartheta(C_{13}(1, 5)) = \frac{13(-\lambda_{\min})}{4 - \lambda_{\min}} = 5.181736625\dots, \quad \Delta_{\max}^{\text{tf}}(13) = \vartheta - 4 = 1.181736625\dots \quad (5)$$

(ii) *The unique $R(3, 9)$ -critical graph on 35 vertices is the circulant $C_{35}(1, 7, 11, 16)$ of Kalbfleisch [12]; it is vertex- but not edge-transitive, and $\vartheta = 12.410586\dots$ is the optimum of the Fourier linear program (the bound (4) gives 12.4979 and is not attained). Hence $\Delta_{\max}(35) \geq 4.410586$.*

(iii) *There is no circulant $R(3, k)$ -critical graph on $R(3, k) - 1$ vertices for $k = 6, 7, 8$ ($n = 17, 22, 27$); the circulant $R(3, k)$ -graphs that exist on 16 and 21 vertices are not the gap maximizers of their families (1.1648 vs 1.4654; 1.9768 vs 2.1413).*

Table 3: Maximum gap over all $R(3, k)$ -graphs on n vertices (McKay’s catalogues; $\alpha = k - 1$ analytically, ϑ by SDP). These are lower bounds on $\Delta_{\max}(n)$. For $n = 14$ the catalogue maximum is $C_{13}(1, 5)$ plus an isolated vertex; the best connected graph is shown.

k	n	#graphs	α	$\vartheta_{\max}$	Δ	best graph (graph6)
5	13	1	4	5.181737	1.181737	$C_{13}(1, 5)$
6	14	263,520	5	6.072373	1.072373	MqG00H@_gQ?oGFCJ? (connected)
6	15	64,732	5	6.250964	1.250964	NoCOWGACGQIKGePIDI?
6	16	2,576	5	6.465350	1.465350	OoC?Igi_?E_eKcRCE@iO[
6	17	7	5	6.606664	1.606664	P@oGQA?`ACcdCtDYC]Ah?goC
7	21	1,118,436	6	8.141315	2.141315	THoGG?JODHsOAdS_wg?`GgSCZ??ig@D@?GT
7	22	191	6	8.398747	2.398747	UsaC?GGC?DccQDCpKEJAOKW`Bo?kD_[_okHCUR??
8	27	477,142	7	10.060569	3.060569	(Appendix A)
9	35	1	8	12.410586	4.410586	the unique $(3, 9)$ -critical graph

The scan behind (iii) is exhaustive over connection sets (all $2^{\lfloor n/2 \rfloor}$ symmetric sets, triangle-freeness, α by clique search on the complement); the identifications in (i) and (ii) are by isomorphism test against McKay’s catalogues.

We note explicitly what Section 4 does and does not establish. It establishes that α of the record graphs is dictated by Ramsey numbers (Lemma 2), it identifies every computed maximizer with a Ramsey graph, and for the two circulant cases it gives ϑ in closed form. It does not yet provide a theorem of the form $\Delta_{\max}(R(3, k) - 1) \geq f(k)$ with f growing: the general lower bounds on ϑ of triangle-free graphs that we are aware of (through $\vartheta(G)\vartheta(\bar{G}) \geq n$ and $\chi(G) = O(\sqrt{n})$) are of order $\sqrt{n}$, below $k - 1 \approx \sqrt{n \log n}$ at $n = R(3, k) - 1$, and so do not even certify positivity. Closing this gap — a lower bound on ϑ for triangle-free graphs of small independence number — is the natural next step, and the data of Table 3 suggest $\Delta_{\max}(R(3, k) - 1) \approx 0.12 (R(3, k) - 1)$.

4.3 A ladder of lower bounds up to $n = 35$

Since the $R(3, k)$ -graphs are completely catalogued for $k \leq 9$ [15, 14], and $\alpha = k - 1$ on all of them for $n \geq R(3, k - 1)$ (Lemma 2), the gap over each catalogue requires only ϑ . Table 3 gives the maxima; together with Lemma 1 they are lower bounds on $\Delta_{\max}(n)$ for all larger n . Figure 3 summarizes the results.

The $R(3, 6)$ -graphs on 13 vertices (275,086 graphs, for which $\alpha \in \{4, 5\}$) confirm the picture from the other side: the single graph with $\alpha = 4$ is $C_{13}(1, 5)$ with $\Delta = 1.181737$, and the best of the 275,085 graphs with $\alpha = 5$ reaches only 0.930021 (it is the twelve-vertex triangle-free maximizer plus an isolated vertex).

Within the triangle-free class the maximum gap is *not* monotone: it falls from 1.1817 at $n = 13$ to 1.0724 at $n = 14$, exactly as predicted by the jump of α at $R(3, 5) = 14$, and recovers from $n = 15$ on. The global $\Delta_{\max}(n)$ is monotone by Lemma 1, but the graphs realizing the monotone extension (a record graph plus universal vertices) are degenerate as witnesses. We are led to:

Conjecture 1. *For $n = R(3, k) - 1$ with $k \geq 5$, $\Delta_{\max}(n)$ is attained by an $R(3, k)$ -critical graph; more generally, from $n = 11$ on the maximizer is triangle-free with the least independence number that Ramsey theory allows.*

We state this as a conjecture with the appropriate caution. The evidence is: the exhaustive result at $n = 11$ (the only value where the statement is verified over all graphs), the exhaustive triangle-free results at $n = 12, 13$, the fact that the mechanism is forced for the two graph invariants




Figure 3: Maximum contextuality gap versus number of vertices: exhaustive over all connected graphs ($n \leq 11$), exhaustive over triangle-free graphs ($n \leq 13$), and lower bounds from the complete catalogues of $R(3, k)$ -graphs. Vertical lines mark the Ramsey numbers $R(3, k)$, where α is forced up by one for triangle-free graphs.

separately (α by Ramsey theory, ϑ by its growth with n for sparse graphs), and the failure of the “previous winner plus a vertex” lineages. The evidence against unrestricted extrapolation is equally clear: for $n = 9, 10$ the maximizers contain triangles, because no triangle-free graph on nine or more vertices has $\alpha = 3$, so an $R(3, 4)$ -type structure with one or four triangles wins. What the conjecture does *not* say is that Ramsey-critical graphs dominate between the critical vertex counts; at $n = 14$ the triangle-free maximum drops (Section 4.4) and the global maximum is attained only by the degenerate extension of $C_{13}(1, 5)$. A brute-force test at $n = 12$ would require 1.64×10^{11} graphs; a local search over graphs with triangles at $n = 12$ (Section 6) found nothing above the triangle-free record.

4.4 What can be proved: a linear upper bound and linear growth

The Ramsey mechanism pins α from below; the following elementary bound pins Δ from above.

Proposition 2. *For every graph G on n vertices, $\Delta(G) \leq \nu(G) \leq \lfloor n/2 \rfloor$, where $\nu(G)$ is the matching number.*

Proof. By the sandwich theorem $\vartheta(G) \leq \bar{\chi}(G)$, the clique cover number [3]. Covering the 2ν vertices of a maximum matching by its ν edges and the remaining vertices by singletons gives $\bar{\chi}(G) \leq n - \nu$. By the Tutte–Berge formula there is a set S with $n - 2\nu = \text{odd}(G - S) - |S|$; one vertex from each odd component of $G - S$ forms an independent set, so $\alpha(G) \geq \text{odd}(G - S) \geq n - 2\nu$. Hence $\Delta = \vartheta - \alpha \leq (n - \nu) - (n - 2\nu) = \nu$. $\square$

The bound is far from tight for small n ($\nu = 5$ versus $\Delta_{\max} = 0.775$ at $n = 11$), but it has the right order: $\Delta_{\max}(n)$ grows linearly. For the binomial random graph $G(n, d/n)$ with fixed d , $\vartheta = \Theta(\sqrt{n/p}) = \Theta(n/\sqrt{d})$ with high probability [18], while $\alpha \approx (2n/d) \ln d$ [19]; so $\Delta \geq c(d)n$ with $c(d) > 0$ for d large enough, i.e. $\Delta_{\max}(n) = \Theta(n)$. Heuristically $2n/\sqrt{d} - (2n/d) \ln d$ is

maximal near $d \approx 15\text{--}20$ with $c \approx 0.12$, and our records already climb towards such a slope: $\Delta_{\max}(n)/n = 0.07, 0.09, 0.11, 0.13$ at $n = 11, 13, 22, 35$ (the last three being the Ramsey lower bounds of Table 3). The value of $c^* = \lim \Delta_{\max}(n)/n \in [c, \frac{1}{2}]$, and whether the maximizers stay triangle-free, are open.

5 Methods

5.1 Enumeration and exact computation

Connected graphs on up to ten vertices were taken from McKay’s lists and, for eleven vertices and for the triangle-free classes, generated with `geng` from `nauty 2.8.9` [16], split into resumable `res/mod` parts. α was computed exactly by enumerating all 2^n vertex subsets (bit operations; on the GPU for $n \geq 11$). ϑ was computed with the interior-point solver Clarabel [17] on the primal (1) with tolerances 10^{-9} (and 10^{-12} for the record graphs), at about 8,000 graphs per second on 20 CPU processes; the full $n = 10$ run took 51 minutes.

5.2 Rigorous GPU screening for $n = 11$

Exact SDPs for 10^9 graphs would take several days. Instead we use the dual (2): *every* matrix Y supported on the edges gives a certified upper bound $\lambda_{\max}(J + Y) \geq \vartheta(G)$. For a batch of 2^{15} graphs we optimize Y per graph with 100 Adam steps on the softmax-smoothed largest eigenvalue $\tau \log \sum_i e^{\lambda_i/\tau}$ (τ annealed from 0.3 to 0.02; gradient $V \text{diag}(\text{softmax}(\lambda/\tau)) V^T$ masked to E), using batched single-precision eigendecompositions. Every ten steps, graphs with $\lambda_{\max}(J + Y) + 10^{-3} - \alpha(G) < \Delta_{\text{thr}}$ are removed from the batch; since the bound is valid at every step and only improves, this is rigorous up to the single-precision margin, which is three orders of magnitude above the eigenvalue error for 11×11 matrices. Survivors receive a double-precision certificate and are re-solved exactly with Clarabel. With $\Delta_{\text{thr}} = 1/\sqrt{2} - 10^{-6}$ (the $n = 10$ record, valid by Lemma 1), the screen processed all 1,006,700,565 graphs in 2.41 hours on an RTX 2000 Ada (8 GB) at $0.95\text{--}1.3 \times 10^5$ graphs per second, leaving 100 survivors, 51 of which indeed have $\Delta \geq 1/\sqrt{2}$.

Validation. With no threshold, the certified bounds exceed ϑ by a median of 0.006 (max 0.21) over the 11,117 graphs on eight vertices, and never fall below ϑ (Figure 4a). With thresholds 0.60 at $n = 9$ and 0.65 at $n = 10$, the survivors are exactly the two, respectively thirty, graphs known to exceed the threshold (from the exact runs and from [5]), with no misses and no false survivors (Figure 4b).

5.3 Ramsey catalogues

For the $R(3, k)$ -graphs we used McKay’s files `r35_13`, `r36_14`–`r36_17`, `r37_21`, `r37_22`, `r38_27`, `r39_35` [15]; $\alpha = k - 1$ on all of them by Lemma 2, and ϑ was computed with Clarabel (74 minutes for the 1,118,436 graphs on 21 vertices, 88 minutes for the 477,142 graphs on 27 vertices).

6 Discussion

Certificates. For every record graph we provide an explicit primal feasible matrix X and an explicit dual feasible matrix Y of (1)–(2), recomputed in double precision independently of the solver; they bracket ϑ in an interval of width below 10^{-9} (Appendix B, Table 5), and the values were reproduced with a second solver (SCS). The exact values $11/3$ and $3 + 1/\sqrt{2}$ for $n = 9, 10$ are




Figure 4: Validation of the GPU screen. (a) Certified upper bound minus exact ϑ for all connected graphs on eight vertices, without pruning. (b) At $n = 10$: number of graphs with $\Delta \geq x$ from the exact run, and the number of survivors of the screen at threshold 0.65, which coincides with the exact count.

thus certified to nine digits; closed forms for the eleven-vertex maximizer and for the triangle-free maximizers on 12 vertices remain to be identified (their minimal polynomials are not obvious from the spectrum, which for $n = 11$ contains $2 + \sqrt{2}$ but is otherwise irrational).

Witnesses for experiments. From the optimal $X = B^T B$ one reads off an orthogonal representation of the exclusivity graph in the CSW sense: unit vectors $u_i = b_i / \|b_i\|$, orthogonal for exclusive events, and a state $\psi \propto \sum_i b_i$ with $\sum_i \langle \psi, u_i \rangle^2 = \vartheta(G)$ (Cauchy–Schwarz gives $\geq$, Lovász’s theorem gives $\leq$). The rank of the maximum-rank optimal X is 4 for all maximizers with $n \leq 11$ and for Quad- C_5 and Wagner, and 5 for the triangle-free maximizers on 12 and 13 vertices, so all witnesses are realizable with a ququart, respectively a five-level system; whether lower dimensions suffice is not settled by our computation (the solver returns the analytic centre of the optimal face). Following [9], the critical visibility under white noise, $v^* = (\alpha - n/d)/(\vartheta - n/d)$, is 0.529 for the $n = 9$ maximizer and 0.414 for the $n = 10$ maximizer (with $d = 4$), against 0.681 for Quad- C_5 and 0.707 for the Wagner graph; for the eleven-vertex maximizer $v^* = 0.617$ and for $C_{13}(1, 5)$ (in $d = 5$) $v^* = 0.542$. As already observed for $n = 8$, a larger gap does not imply a more robust witness: the term n/d grows with n and the $n = 10$ maximizer, with the largest gap among the dense family, is the least robust of the six. The sparse triangle-free witnesses are more robust than the dense ones at equal or larger gap, which is a second argument in their favour.

Beyond triangle-free at $n = 12$. A local search over all graphs on twelve vertices (edge additions, deletions and moves from the triangle-free maximizers and from the eleven-vertex records extended by one vertex, exact ϑ and α at every step) found no graph with a triangle exceeding the triangle-free record 0.930021; the best graph with a triangle reached $\Delta = 0.866$. This is evidence, not proof, for Conjecture 1 at $n = 12$.

Relation to previous work. The Lovász number of Ramsey graphs has appeared before, in bounds on Shannon capacity and on Ramsey numbers [20]; there the interest is in ϑ as an upper bound on α of powers of graphs, whereas here the object is the *difference* $\vartheta - \alpha$ of the Ramsey graphs themselves as a measure of contextuality. The graph-enumeration programme of [4, 7] computed α ,

ϑ and α^* for all graphs with up to ten vertices and studied ratio maximizers and “fully contextual” graphs ($\vartheta = \alpha^*$); none of the gap maximizers found here is fully contextual (e.g. $\alpha^* = 13/2$ for $C_{13}(1, 5)$), and the two figures of merit select different graphs from $n = 8$ on.

7 Conclusion

We have pushed the exhaustive maximization of the absolute contextuality gap from eight to eleven vertices, over a billion graphs, with a GPU screening scheme whose every elimination is backed by a dual certificate, and we have found that the answer has a structure: the record graphs are Ramsey $R(3, k)$ -graphs, triangle-free graphs whose independence number is held at $k - 1$ by $n < R(3, k)$ while their theta number keeps growing. The thirteen-vertex triangle-free maximizer is the Ramsey graph $C_{13}(1, 5)$ itself, with a closed-form theta number, and the complete catalogues of $R(3, k)$ -graphs yield gap records up to 35 vertices. Two questions are left open: whether the maximizer at twelve vertices — beyond exhaustive reach at 1.6×10^{11} graphs, though within reach of a more aggressive screen — is still triangle-free (Conjecture 1), and the value of the linear growth constant $\lim \Delta_{\max}(n)/n$, which Proposition 2 confines to $[c, \frac{1}{2}]$. For experiments, the new witnesses on nine and ten vertices have exact theta numbers $11/3$ and $3 + 1/\sqrt{2}$, realizations in dimension four, and are given explicitly in the appendix and in the accompanying repository.

Code and data availability

All scripts (exact search, GPU screening, Ramsey catalogue processing), the result tables with graph6 identifiers, and the certificates for the record graphs are available at <https://github.com/danmusetoiu/contextuality-gap>.

A Reproduction of the eight-vertex ranking and graph data

Table 4: Top ten connected graphs on eight vertices by Δ (our run, Clarabel), matching Table II of [9] rank for rank.

rank	graph6	$ E $	α	ϑ	Δ	ϑ/α	
1	GCQb`o	10	3	3.467844	0.467844	1.155948	Quad- C_5
2	GCR`r_	11	3	3.438447	0.438447	1.146149	
3	GCRb`o	12	3	3.414214	0.414214	1.138071	Wagner graph
4	GCRb`w	12	3	3.372281	0.372281	1.124094	
5	GCQbdo	11	3	3.372281	0.372281	1.124094	CHSH-type graph, ratio maximizer
6	GCp`d0	10	3	3.372281	0.372281	1.124094	
7	GQyurg	16	2	2.343146	0.343146	1.171573	
8	GQyurw	17	2	2.338044	0.338044	1.169022	
9	GCpdag	11	3	3.338044	0.338044	1.112681	
10	GUZurw	18	2	2.333333	0.333333	1.166667	

Edge lists of the record graphs Vertices are $0, \dots, n - 1$ in the labelling induced by the graph6 string.

- Quad- C_5 ($n = 8$), GCQb`o: 03, 05, 14, 16, 25, 26, 27, 36, 37, 47.

- $n = 9$ maximizer, HCRbd0f: 03, 05, 07, 14, 15, 16, 25, 26, 27, 28, 36, 38, 47, 48, 58.
- $n = 10$ maximizer, ICRb`yiu?: 03, 05, 08, 09, 14, 15, 16, 19, 25, 26, 27, 28, 36, 37, 39, 47, 48, 49, 57, 68.
- $n = 11$ maximizer, J?`D@pgd?{?: (0,4), (0,6), (0,9), (1,5), (1,8), (2,6), (2,7), (2,8), (3,7), (3,9), (3,10), (4,7), (4,8), (4,10), (5,9), (5,10), (6,10).
- $n = 12$ triangle-free maximizer, K?`DA`gd?{Dg: (0,4), (0,6), (0,9), (1,5), (1,7), (1,8), (2,6), (2,8), (2,11), (3,7), (3,9), (3,10), (4,8), (4,10), (4,11), (5,9), (5,10), (5,11), (6,10), (7,11).
- $C_{13}(1, 5)$, L?`DE`gl@YJODg: (0,4), (0,6), (0,7), (0,9), (1,5), (1,7), (1,8), (1,11), (2,6), (2,8), (2,9), (2,10), (3,7), (3,9), (3,11), (3,12), (4,8), (4,10), (4,11), (5,9), (5,10), (5,12), (6,11), (6,12), (7,10), (8,12).

The $R(3, 8)$ record on 27 vertices graph6 Z@0\MagCADc`e@0\_\_`KoHO_CCHN?_G@D0?d??oCwQ i??GTSPK?Dc?@YdI?c?, $|E| = 91$, $\vartheta = 10.060569$, $\Delta = 3.060569$.

B Certificates

Table 5: Certificates for the record graphs. $[\vartheta_{\text{lb}}, \vartheta_{\text{ub}}]$ is the interval certified in double precision by an explicit primal feasible X (trace one, zero on edges, PSD after eigenvalue clipping; the residual on edge entries is below 10^{-9}) and an explicit dual feasible Y (supported on the edges, $\lambda_{\max}(J + Y)$ recomputed with LAPACK). r is the rank of the maximum-rank optimal X returned by the interior-point solver, hence an upper bound on the minimal dimension of an orthogonal representation attaining ϑ ; $v^* = (\alpha - n/r)/(\vartheta - n/r)$ is the critical visibility under white noise in dimension r . All matrices and the orthogonal representations are in the repository (`certificates/*.json`).

graph	n	$ E $	α	ϑ_{lb}	ϑ_{ub}	r	v^*
Quad- C_5 ($n = 8$)	8	10	3	3.467843730	3.467843730	4	0.6813
Wagner ($n = 8$)	8	12	3	3.414213562	3.414213562	4	0.7071
$n = 9$ maximizer	9	15	3	3.666666667	3.666666667	4	0.5294
$n = 10$ maximizer	10	20	3	3.707106781	3.707106781	4	0.4142
$n = 11$ maximizer	11	17	4	4.774888533	4.774888533	4	0.6173
$n = 11$, rank 2	11	17	4	4.754686045	4.754686045	4	0.6235
$n = 12$ triangle-free max.	12	20	4	4.930021465	4.930021467	5	0.6324
$C_{13}(1, 5)$	13	26	4	5.181736625	5.181736625	5	0.5423
$R(3, 6)$ best, $n = 16$	16	35	5	6.465350297	6.465350300	5	0.5512
$R(3, 6)$ best, $n = 17$	17	40	5	6.606663834	6.606663840	6	0.5742
$R(3, 7)$ best, $n = 22$	22	66	6	8.398747394	8.398747399	6	0.4931

References

- [1] A. Cabello, S. Severini, and A. Winter, Graph-theoretic approach to quantum correlations, Phys. Rev. Lett. **112**, 040401 (2014).
- [2] L. Lovász, On the Shannon capacity of a graph, IEEE Trans. Inf. Theory **25**, 1 (1979).
- [3] D. E. Knuth, The sandwich theorem, Electron. J. Combin. **1**, A1 (1994).

- [4] A. Cabello, L. E. Danielsen, A. J. López-Tarrida, and J. R. Portillo, Basic exclusivity graphs in quantum correlations, *Phys. Rev. A* **88**, 032104 (2013).
- [5] A. Cabello, L. E. Danielsen, A. J. López-Tarrida, and J. R. Portillo, Database of quantum graphs, https://www.codetables.de/larsed/quantum_graphs/.
- [6] E. Amselem, L. E. Danielsen, A. J. López-Tarrida, J. R. Portillo, M. Bourennane, and A. Cabello, Experimental fully contextual correlations, *Phys. Rev. Lett.* **108**, 200405 (2012).
- [7] G. Cañas, E. Acuña, J. Cariñe, J. F. Barra, E. S. Gómez, G. B. Xavier, G. Lima, and A. Cabello, Experimental demonstration of the connection between quantum contextuality and graph theory, *Phys. Rev. A* **94**, 012337 (2016).
- [8] M. Ray, N. G. Boddu, K. Bharti, L.-C. Kwek, and A. Cabello, Graph-theoretic approach to dimension witnessing, *New J. Phys.* **23**, 033006 (2021).
- [9] U. Tamer, Ö. E. Müstecaplıoğlu, A. Dizdar, and Z. Gedik, The Quad- C_5 graph: maximum contextuality gap on eight vertices, *arXiv:2605.12828* (2026).
- [10] A. Muller and M. Saniga, Automated search for highly contextual Kochen–Specker proofs, *arXiv:2609.19862* (2026).
- [11] A. A. Klyachko, M. A. Can, S. Binicioğlu, and A. S. Shumovsky, Simple test for hidden variables in spin-1 systems, *Phys. Rev. Lett.* **101**, 020403 (2008).
- [12] J. G. Kalbfleisch, Chromatic graphs and Ramsey’s theorem, Ph.D. thesis, University of Waterloo (1966).
- [13] S. P. Radziszowski, Small Ramsey numbers, *Electron. J. Combin.*, Dynamic Survey DS1 (revised regularly).
- [14] J. Goedgebeur and S. P. Radziszowski, New computational upper bounds for Ramsey numbers $R(3, k)$, *Electron. J. Combin.* **20**, P30 (2013).
- [15] B. D. McKay, Ramsey graphs, <https://users.cecs.anu.edu.au/~bdm/data/ramsey.html>.
- [16] B. D. McKay and A. Piperno, Practical graph isomorphism, II, *J. Symbolic Comput.* **60**, 94 (2014).
- [17] P. J. Goulart and Y. Chen, Clarabel: An interior-point solver for conic programs with quadratic objectives, *arXiv:2405.12762* (2024).
- [18] A. Coja-Oghlan, The Lovász number of random graphs, *Combin. Probab. Comput.* **14**, 439 (2005).
- [19] A. M. Frieze, On the independence number of random graphs, *Discrete Math.* **81**, 171 (1990).
- [20] N. Alon, Lovász, vectors, graphs and codes, in *Building Bridges II*, Bolyai Soc. Math. Stud. **28** (Springer, 2019).