

Maximum Contextuality Gap up to Eleven Vertices and the Ramsey Structure of Its Maximizers

Dan Muşetoiu
dan.musetoiu@gmail.com

September 2026

Abstract

In the exclusivity-graph approach to contextuality, a noncontextuality inequality is a graph G , its noncontextual bound is the independence number $\alpha(G)$ and its quantum bound is the Lovász number $\vartheta(G)$. The absolute contextuality gap $\Delta(G) = \vartheta(G) - \alpha(G)$ was recently maximized exhaustively over the 11,117 connected graphs on eight vertices (Quad- C_5 , $\Delta = 0.46784$). We extend the exhaustive search to nine, ten and eleven vertices — 261,080, 11,716,571 and 1,006,700,565 connected graphs respectively — and find $\Delta_{\max} = 2/3$ (with $\vartheta = 11/3$), $\Delta_{\max} = 1/\sqrt{2}$ (with $\vartheta = 3 + 1/\sqrt{2}$) and $\Delta_{\max} = 0.774889$. The search over 10^9 graphs was made feasible on a single consumer GPU by a rigorous screening scheme: for each graph a feasible point of the dual semidefinite program certifies an upper bound on ϑ , and only graphs whose certified bound can beat the running record are solved exactly. The maximizers are not a nested family, and at eleven vertices a regime change occurs: the dense, Wagner-based maximizers with $\alpha = 3$ are overtaken by sparse triangle-free graphs with $\alpha = 4$. We explain this through Ramsey theory: triangle-free graphs with $\alpha \leq k-1$ are exactly the $R(3, k)$ -graphs, and every record graph we find is one — Quad- C_5 and the Wagner graph are $R(3, 4)$ -graphs, the eleven-vertex maximizer is an $R(3, 5)$ -graph, and the triangle-free maximizer on thirteen vertices is the unique $R(3, 5)$ -critical graph $C_{13}(1, 5)$, with $\vartheta = 13(-\lambda_{\min})/(\lambda_{\max} - \lambda_{\min}) = 5.181737$ in closed form. Computing ϑ over McKay’s complete catalogues of $R(3, k)$ -graphs yields a ladder of lower bounds on $\Delta_{\max}(n)$ up to $n = 35$ ($\Delta \geq 4.4106$ from the unique $R(3, 9)$ -critical graph). We release all code, graph6 identifiers and certificates.

Keywords: quantum contextuality, exclusivity graphs, Lovász theta number, independence number, Ramsey graphs, semidefinite programming, exhaustive graph search

1 Introduction

Cabello, Severini and Winter (CSW) showed that any noncontextuality (NC) inequality of the form $\sum_i P(e_i) \leq \alpha$ can be encoded by an *exclusivity graph* G whose vertices are the events e_i and whose edges join mutually exclusive events; the maximum of the left-hand side over noncontextual models is the independence number $\alpha(G)$, and its maximum over quantum models is the Lovász theta number $\vartheta(G)$ [1]. The gap between the two numbers is therefore a graph invariant that measures how contextual an inequality can be. Two natural figures of merit have been studied: the ratio ϑ/α , used to identify experimentally accessible witnesses [6, 7] and as a dimension witness [8], and the absolute gap $\Delta(G) = \vartheta(G) - \alpha(G)$, whose maximization over all connected graphs on eight vertices was recently carried out by Tamer et al. [9]. Their exhaustive search over the 11,117 graphs identified a sparse ten-edge graph, Quad- C_5 , with $\Delta = 0.46784$, beating the Wagner graph ($\Delta = \sqrt{2} - 1$).

Two questions follow. What are the maximizers for more vertices, and is there structure behind them? Here we answer the first question exhaustively for $n = 9, 10, 11$ and, for the triangle-free class, for $n = 12, 13$; and we answer the second by exhibiting a Ramsey-theoretic mechanism.

Contributions.

1. **Exhaustive maxima for $n = 9, 10, 11$.** $\Delta_{\max}(9) = 2/3$ exactly, $\Delta_{\max}(10) = 1/\sqrt{2}$ exactly, $\Delta_{\max}(11) = 0.774889$ (Section 3). The eleven-vertex search covers all 1,006,700,565 connected graphs and is, to our knowledge, the first exhaustive computation of a Lovász-number extremum at this scale.
2. **A rigorous GPU screening method** (Section 5): dual-feasible points of the theta SDP, optimized in large batches on a consumer GPU, certify upper bounds on ϑ that eliminate all but a hundred graphs out of 10^9 ; the survivors are solved exactly. The method is validated against every known answer ($n \leq 10$) with zero misses.
3. **The Ramsey structure of the maximizers** (Section 4). All record graphs are $R(3, k)$ -graphs, i.e. triangle-free graphs with $\alpha \leq k-1$ on fewer than $R(3, k)$ vertices; the thirteen-vertex triangle-free maximizer is the unique $R(3, 5)$ -critical graph $C_{13}(1, 5)$, whose theta number has a closed form. Ramsey numbers thus pin α while ϑ keeps growing, and $\Delta_{\max}(n)$ jumps at $n = R(3, k) - 1$. Computing ϑ over the complete catalogues of $R(3, k)$ -graphs gives a ladder of lower bounds up to $n = 35$.

2 Preliminaries

Throughout, $G = (V, E)$ is a simple graph on n vertices, interpreted as an exclusivity graph. The independence number $\alpha(G)$ is the size of a largest set of pairwise non-adjacent vertices. The Lovász number [2] can be written as the semidefinite program

$$\vartheta(G) = \max\{\langle J, X \rangle : \text{tr } X = 1, X_{ij} = 0 \ \forall ij \in E, X \succeq 0\}, \quad (1)$$

where J is the all-ones matrix, with the dual

$$\vartheta(G) = \min\{\lambda_{\max}(J + Y) : Y = Y^T, Y_{ij} = 0 \ \forall ij \notin E, Y_{ii} = 0\}. \quad (2)$$

The sandwich theorem gives $\alpha(G) \leq \vartheta(G) \leq \alpha^*(G)$, where α^* is the fractional packing number [3]. In the CSW framework $\alpha(G)$ and $\vartheta(G)$ are respectively the noncontextual and the quantum maximum of $\sum_{i \in V} P(e_i)$ over all correlation experiments whose exclusivity graph is G [1]; quantum theory violates the NC inequality iff $\vartheta(G) > \alpha(G)$, which happens iff G contains an induced odd cycle of length at least five or the complement of one [4]. We study the *absolute contextuality gap*

$$\Delta(G) = \vartheta(G) - \alpha(G), \quad \Delta_{\max}(n) = \max\{\Delta(G) : G \text{ connected}, |V(G)| = n\}. \quad (3)$$

Restricting to connected graphs loses nothing for the maximum: ϑ and α are both additive over disjoint unions, so a disconnected graph never beats its best component plus isolated vertices, and isolated vertices contribute equally to ϑ and α .

Two elementary facts will be used repeatedly.

Lemma 1 (Monotonicity). $\Delta_{\max}(n+1) \geq \Delta_{\max}(n)$.

Table 1: Maximum contextuality gap over all connected graphs on n vertices. Graphs are identified by their graph6 string [15]. d^* is the rank of the optimal primal matrix X in (1), i.e. the dimension of the Lovász orthonormal representation attaining ϑ .

n	#graphs	graph6	$ E $	α	ϑ	$\Delta_{\max}(n)$	remarks
5	21	C_5	5	2	$\sqrt{5}$	0.236068	KCBS pentagon
6	112	EUZO	8	2	$\sqrt{5}$	0.236068	no improvement over $n = 5$
7	853	C_7	7	3	3.317667	0.317667	[9]
8	11,117	GCQb`o	10	3	3.467844	0.467844	Quad- C_5 [9]; $R(3, 4)$ -graph
9	261,080	HCRbdO{	15	3	11/3	2/3	new; $ \text{Aut} = 12$, one triangle
10	11,716,571	ICRb`yiu?	20	3	$3 + 1/\sqrt{2}$	$1/\sqrt{2}$	new; 4-regular, $ \text{Aut} = 16$
11	1,006,700,565	J?`D@pgd?{?	17	4	4.774889	0.774889	new; triangle-free, $R(3, 5)$ -graph

Proof. Add a vertex adjacent to all vertices of a maximizer G . Independent sets are unchanged, so α is unchanged. In (1) the new row and column of X must vanish off the diagonal, so $X = \text{diag}(X', s)$ with $\text{tr } X' = 1 - s$ and $\langle J, X \rangle = (1 - s)\vartheta(G) + s \leq \vartheta(G)$; hence ϑ is unchanged as well. $\square$

Lemma 2 (Ramsey pins α). *Let $R(3, k)$ denote the Ramsey number. A triangle-free graph on $n \geq R(3, k - 1)$ vertices has $\alpha \geq k - 1$; a triangle-free graph on $n < R(3, k)$ vertices can have $\alpha = k - 1$, and the graphs achieving it are exactly the $R(3, k)$ -graphs on n vertices.*

Proof. Immediate from the definition of $R(3, k)$ as the least n such that every graph on n vertices contains a triangle or an independent set of size k . $\square$

The relevant values are $R(3, 4) = 9$, $R(3, 5) = 14$, $R(3, 6) = 18$, $R(3, 7) = 23$, $R(3, 8) = 28$, $R(3, 9) = 36$ [12]. Complete catalogues of $R(3, k)$ -graphs for $k \leq 9$ are available from McKay [14, 13].

3 Exhaustive results

3.1 All connected graphs, $n \leq 11$

Table 1 lists the maximizers of Δ over all connected graphs for $n = 5, \dots, 11$; the values for $n \leq 8$ reproduce [9] exactly (including the full top-ten ranking for $n = 8$, Appendix A). Figure 1 draws the record graphs.

$n = 9$. The maximizer HCRbdO{ has 15 edges, degree sequence $3^6 4^3$, automorphism group of order 12 with vertex orbits of sizes 3 and 6, twelve induced pentagons and a single triangle. Its theta number is 11/3 to twelve digits (Clarabel with tolerances 10^{-12}), so $\Delta = 2/3$ exactly; the optimal X has rank 4. The runner-up (HCrb`qi, 16 edges) is the Wagner graph plus one vertex and has $\vartheta = 2 + \varphi = 3.618034$, φ the golden ratio; it is followed by a large plateau of graphs at exactly Quad- C_5 's value 0.467844 (Quad- C_5 plus a vertex that changes neither ϑ nor α). The maximizer contains neither Quad- C_5 nor the Wagner graph as an induced subgraph. Its adjacency spectrum is $\{2 + \sqrt{2}, 1^{(2)}, (\sqrt{2} - 1)^{(2)}, 2 - \sqrt{2}, -2, (-1 - \sqrt{2})^{(2)}\}$.

$n = 10$. The maximizer ICRb`yiu? is 4-regular with 20 edges,  $|\text{Aut}| = 16$  (orbits of sizes 2 and 8), sixteen induced pentagons and four triangles, and contains the Wagner graph as an induced subgraph.  $\vartheta = 3 + 1/\sqrt{2}$  to twelve digits. The runner-up (ICQeR`[Mg, $\Delta = 0.667155$) contains both Quad- C_5 and the $n = 9$ maximizer; the “ $n = 9$ maximizer plus a vertex” lineage therefore reaches only second place, as the “Wagner plus a vertex” lineage did at $n = 9$. The values for




Figure 1: Record graphs. Top: the exhaustive maximizers for $n = 8$ (Quad- C_5 , [9]), $n = 9$ and $n = 10$. Bottom: the exhaustive maximizer for $n = 11$, the triangle-free maximizer for $n = 12$, and the triangle-free maximizer for $n = 13$, which is the circulant Ramsey graph $C_{13}(1, 5)$.

$n = 9$ and $n = 10$ can in principle be read off the table of Cabello, Danielsen, López-Tarrida and Portillo, who computed α , ϑ (to four decimals) and α^* for all graphs with $\alpha < \vartheta$ on up to ten vertices [4, 5]; our runs reproduce that table graph-for-graph (16,533 graphs at $n = 9$, 975,330 at $n = 10$, $|\vartheta - \vartheta_{\text{table}}| \leq 5.3 \times 10^{-5}$). The maxima were, however, never reported.

$n = 11$. Here the picture changes. The five best graphs (Table 2) are all triangle-free, have $\alpha = 4$ and only 17–19 edges, while the best graph of the dense family that dominated $n \leq 10$ — the $n = 10$ maximizer plus one vertex, $\Delta = 0.735505$, which contains Quad- C_5 , the Wagner graph and the $n = 9$ and $n = 10$ maximizers as induced subgraphs — is only sixth. The maximizer J7`D@pgd?{? has girth 4, degree sequence $2^1 3^8 4^2$, $|\text{Aut}| = 2$, eleven induced pentagons and four induced heptagons, and $d^* = 4$. Note that $\Delta_{\max}(11) - \Delta_{\max}(10) = 0.068$ is the smallest increment in the table; the sparse family is only just taking over.

Gap versus ratio. The maximizer of ϑ/α differs from the maximizer of Δ at $n = 8$ (a CHSH-type graph, [9]), $n = 10$ (IUZurzmno, $\alpha = 2$, $\vartheta = 5/2$, ratio 1.25) and $n = 11$ (the dense graph of rank 6 in Table 2, ratio 1.2452, versus 1.1937 for the gap maximizer); at $n = 9$ the two coincide.

Table 2: The ten largest gaps among all 1,006,700,565 connected graphs on eleven vertices. “tf” = triangle-free.

rank	graph6	$ E $	α	ϑ	Δ	remarks
1	J?`D@pgd?{?	17	4	4.774889	0.774889	tf, $R(3, 5)$ -graph
2	J?`@f?kUDG_	17	4	4.754686	0.754686	tf
3	J?`@f@WFTH?	18	4	4.753664	0.753664	tf, contains Wagner
4	J?`FBQoFCi?	18	4	4.752419	0.752419	tf
5	J?`@f@Wl?y?	18	4	4.747377	0.747377	tf, contains Quad- C_5
6	JCRbeo{ifI?	25	3	3.735505	0.735505	$n=10$ maximizer + vertex
7	J?`FAqSFf_?	19	4	4.734338	0.734338	tf
8	J?`FAosFdI?	19	4	4.725688	0.725688	tf
9	J?`DD`gD`s?	17	4	4.719179	0.719179	tf
10	JCRbeo{ifM?	26	3	3.712900	0.712900	dense family

3.2 Triangle-free graphs, $n = 12, 13$

Motivated by the $n = 11$ result we searched exhaustively over connected triangle-free graphs, generated with `geng -t` [15]: 90,842 graphs for $n = 11$ (recovering the global top six), 1,144,061 for $n = 12$ and about 1.9×10^7 for $n = 13$. The maxima are

$$\Delta_{\max}^{\text{tf}}(12) = 0.930021 \quad (\vartheta = 4.930021, \alpha = 4, 20 \text{ edges, } K?`DA`gd?\{Dg, |Aut| = 8, d^* = 5),$$

$$\Delta_{\max}^{\text{tf}}(13) = 1.181737 \quad (\vartheta = 5.181737, \alpha = 4, 26 \text{ edges, } L?`DE`gl@YJODg).$$

At $n = 13$ the maximizer is isolated: the next best triangle-free graph has $\Delta = 0.854$ and $\alpha = 5$. The maximizer is 4-regular with $|Aut| = 52$ and is isomorphic to the circulant graph $C_{13}(1, 5)$, the unique $(3, 5)$ -Ramsey-critical graph.

4 The Ramsey structure of the maximizers

4.1 Every record graph is an $R(3, k)$ -graph

By Lemma 2, a triangle-free graph with $\alpha \leq k - 1$ is an $R(3, k)$ -graph and exists only for $n < R(3, k)$. Checking the record graphs against McKay’s catalogues by isomorphism gives:

- $n = 8$: Quad- C_5 and the Wagner graph are two of the three $R(3, 4)$ -graphs on 8 vertices ($R(3, 4) = 9$).
- $n = 9, 10$: no triangle-free graph has $\alpha = 3$ any more; the maximizers keep $\alpha = 3$ by admitting one, respectively four, triangles.
- $n = 11$: the maximizer is one of the 105 $R(3, 5)$ -graphs on 11 vertices; indeed the 105 connected triangle-free graphs with $\alpha \leq 4$ on 11 vertices are exactly this catalogue.
- $n = 12$: the triangle-free maximizer is one of the 12 $R(3, 5)$ -graphs on 12 vertices.
- $n = 13$: the triangle-free maximizer is the unique $R(3, 5)$ -graph on 13 vertices, $C_{13}(1, 5)$.

The mechanism is transparent: ϑ grows with n (roughly like $\sqrt{n}$ for sparse graphs), whereas α is bounded below by Ramsey theory and jumps by one at every $n = R(3, k)$. Records are therefore expected just before the jumps, at $n = R(3, k) - 1$, and the gap should *decrease* within the triangle-free class right after a jump.

Table 3: Maximum gap over all $R(3, k)$ -graphs on n vertices (McKay’s catalogues; $\alpha = k - 1$ analytically, ϑ by SDP). These are lower bounds on $\Delta_{\max}(n)$. For $n = 14$ the catalogue maximum is $C_{13}(1, 5)$ plus an isolated vertex; the best connected graph is shown.

k	n	#graphs	α	$\vartheta_{\max}$	Δ	best graph (graph6)
5	13	1	4	5.181737	1.181737	$C_{13}(1, 5)$
6	14	263,520	5	6.072373	1.072373	MqG00H@_gQ?oGFCJ? (connected)
6	15	64,732	5	6.250964	1.250964	NoCOWGACGQIKGePIDI?
6	16	2,576	5	6.465350	1.465350	OoC?Igi_?E_eKcRCE@iO[
6	17	7	5	6.606664	1.606664	P@oGQA?`ACcdCtDYC]Ah?goC
7	21	1,118,436	6	8.141315	2.141315	THoGG?JODHsOAds_wg?`GgSCZ??ig@@D@?GT
7	22	191	6	8.398747	2.398747	UsaC?GGC?DccQDCpKEJAOKW`Bo?kD_[_okHCUR??
8	27	477,142	7	10.060569	3.060569	(Appendix B)
9	35	1	8	12.410586	4.410586	the unique $(3, 9)$ -critical graph

4.2 The theta number of $C_{13}(1, 5)$

For an edge-transitive graph the Lovász bound $\vartheta(G) \geq -n\lambda_{\min}/(\lambda_{\max} - \lambda_{\min})$ is attained [2]. The circulant $C_{13}(1, 5)$ has eigenvalues $\lambda_j = 2\cos(2\pi j/13) + 2\cos(10\pi j/13)$, $j = 0, \dots, 12$, so $\lambda_{\max} = 4$, $\lambda_{\min} = 2\cos(8\pi/13) - 2\cos(\pi/13) = -2.651093\dots$ (multiplicity 4), and

$$\vartheta(C_{13}(1, 5)) = \frac{13(-\lambda_{\min})}{4 - \lambda_{\min}} = 5.181736625\dots, \quad \Delta = 1.181736625\dots, \quad (4)$$

in agreement with the SDP value to nine digits.

4.3 A ladder of lower bounds up to $n = 35$

Since the $R(3, k)$ -graphs are completely catalogued for $k \leq 9$ [14, 13], and $\alpha = k - 1$ on all of them for $n \geq R(3, k - 1)$ (Lemma 2), the gap over each catalogue requires only ϑ . Table 3 gives the maxima; together with Lemma 1 they are lower bounds on $\Delta_{\max}(n)$ for all larger n . Figure 2 summarizes the results.

Within the triangle-free class the maximum gap is *not* monotone: it falls from 1.1817 at $n = 13$ to 1.0724 at $n = 14$, exactly as predicted by the jump of α at $R(3, 5) = 14$, and recovers from $n = 15$ on. The global $\Delta_{\max}(n)$ is monotone by Lemma 1, but the graphs realizing the monotone extension (a record graph plus universal vertices) are degenerate as witnesses. We are led to:

Conjecture 1. *For every $n \geq 11$, $\Delta_{\max}(n)$ is attained by a triangle-free graph, and for $n = R(3, k) - 1$ by an $R(3, k)$ -critical graph.*

The conjecture is proved for $n = 11$ (this paper) and false for $n = 9, 10$, where triangles are needed to keep $\alpha = 3$; it is consistent with everything we computed for $n \leq 13$. A brute-force test at $n = 12$ would require 1.64×10^{11} graphs.

5 Methods

5.1 Enumeration and exact computation

Connected graphs on up to ten vertices were taken from McKay’s lists and, for eleven vertices and for the triangle-free classes, generated with **geng** from **nauty** 2.8.9 [15], split into resumable **res/mod** parts. α was computed exactly by enumerating all 2^n vertex subsets (bit operations; on




Figure 2: Maximum contextuality gap versus number of vertices: exhaustive over all connected graphs ($n \leq 11$), exhaustive over triangle-free graphs ($n \leq 13$), and lower bounds from the complete catalogues of $R(3, k)$ -graphs. Vertical lines mark the Ramsey numbers $R(3, k)$, where α is forced up by one for triangle-free graphs.

the GPU for $n \geq 11$). ϑ was computed with the interior-point solver Clarabel [16] on the primal (1) with tolerances 10^{-9} (and 10^{-12} for the record graphs), at about 8,000 graphs per second on 20 CPU processes; the full $n = 10$ run took 51 minutes.

5.2 Rigorous GPU screening for $n = 11$

Exact SDPs for 10^9 graphs would take several days. Instead we use the dual (2): *every* matrix Y supported on the edges gives a certified upper bound $\lambda_{\max}(J + Y) \geq \vartheta(G)$. For a batch of 2^{15} graphs we optimize Y per graph with 100 Adam steps on the softmax-smoothed largest eigenvalue $\tau \log \sum_i e^{\lambda_i/\tau}$ (τ annealed from 0.3 to 0.02; gradient $V \text{diag}(\text{softmax}(\lambda/\tau)) V^T$ masked to E), using batched single-precision eigendecompositions. Every ten steps, graphs with $\lambda_{\max}(J + Y) + 10^{-3} - \alpha(G) < \Delta_{\text{thr}}$ are removed from the batch; since the bound is valid at every step and only improves, this is rigorous up to the single-precision margin, which is three orders of magnitude above the eigenvalue error for 11×11 matrices. Survivors receive a double-precision certificate and are re-solved exactly with Clarabel. With $\Delta_{\text{thr}} = 1/\sqrt{2} - 10^{-6}$ (the $n = 10$ record, valid by Lemma 1), the screen processed all 1,006,700,565 graphs in 2.41 hours on an RTX 2000 Ada (8 GB) at $0.95\text{--}1.3 \times 10^5$ graphs per second, leaving 100 survivors, 51 of which indeed have $\Delta \geq 1/\sqrt{2}$.

Validation. With no threshold, the certified bounds exceed ϑ by a median of 0.006 (max 0.21) over the 11,117 graphs on eight vertices, and never fall below ϑ . With thresholds 0.60 at $n = 9$ and 0.65 at $n = 10$, the survivors are exactly the two, respectively thirty, graphs known to exceed the threshold (from the exact runs and from [5]), with no misses and no false survivors.

5.3 Ramsey catalogues

For the $R(3, k)$ -graphs we used McKay’s files `r35_13`, `r36_14–r36_17`, `r37_21`, `r37_22`, `r38_27`, `r39_35` [14]; $\alpha = k - 1$ on all of them by Lemma 2, and ϑ was computed with Clarabel (74 minutes for the 1,118,436 graphs on 21 vertices, 88 minutes for the 477,142 graphs on 27 vertices).

6 Discussion

7 Conclusion

Code and data availability

All scripts (exact search, GPU screening, Ramsey catalogue processing), the result tables with graph6 identifiers, and the certificates for the record graphs are available at `<repository URL>`.

A Reproduction of the eight-vertex ranking

B graph6 strings and certificates

References

- [1] A. Cabello, S. Severini, and A. Winter, Graph-theoretic approach to quantum correlations, *Phys. Rev. Lett.* **112**, 040401 (2014).
- [2] L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inf. Theory* **25**, 1 (1979).
- [3] D. E. Knuth, The sandwich theorem, *Electron. J. Combin.* **1**, A1 (1994).
- [4] A. Cabello, L. E. Danielsen, A. J. López-Tarrida, and J. R. Portillo, Basic exclusivity graphs in quantum correlations, *Phys. Rev. A* **88**, 032104 (2013).
- [5] A. Cabello, L. E. Danielsen, A. J. López-Tarrida, and J. R. Portillo, Database of quantum graphs, https://www.codetables.de/larsed/quantum_graphs/.
- [6] E. Amsalem, L. E. Danielsen, A. J. López-Tarrida, J. R. Portillo, M. Bourennane, and A. Cabello, Experimental fully contextual correlations, *Phys. Rev. Lett.* **108**, 200405 (2012).
- [7] G. Cañas, E. Acuña, J. Cariñe, J. F. Barra, E. S. Gómez, G. B. Xavier, G. Lima, and A. Cabello, Experimental demonstration of the connection between quantum contextuality and graph theory, *Phys. Rev. A* **94**, 012337 (2016).
- [8] M. Ray, N. G. Boddu, K. Bharti, L.-C. Kwek, and A. Cabello, Graph-theoretic approach to dimension witnessing, *New J. Phys.* **23**, 033006 (2021).
- [9] U. Tamer, Ö. E. Müstecaplıoğlu, A. Dizdar, and Z. Gedik, The Quad- C_5 graph: maximum contextuality gap on eight vertices, *arXiv:2605.12828* (2026).
- [10] A. Muller and M. Saniga, Automated search for highly contextual Kochen–Specker proofs, *arXiv:2609.19862* (2026).

- [11] A. A. Klyachko, M. A. Can, S. Binicioğlu, and A. S. Shumovsky, Simple test for hidden variables in spin-1 systems, *Phys. Rev. Lett.* **101**, 020403 (2008).
- [12] S. P. Radziszowski, Small Ramsey numbers, *Electron. J. Combin.*, Dynamic Survey DS1 (revised regularly).
- [13] J. Goedgebeur and S. P. Radziszowski, New computational upper bounds for Ramsey numbers $R(3, k)$, *Electron. J. Combin.* **20**, P30 (2013).
- [14] B. D. McKay, Ramsey graphs, <https://users.cecs.anu.edu.au/~bdm/data/ramsey.html>.
- [15] B. D. McKay and A. Piperno, Practical graph isomorphism, II, *J. Symbolic Comput.* **60**, 94 (2014).
- [16] P. J. Goulart and Y. Chen, Clarabel: An interior-point solver for conic programs with quadratic objectives, *arXiv:2405.12762* (2024).
- [17] N. Alon, Lovász, vectors, graphs and codes, in *Building Bridges II*, Bolyai Soc. Math. Stud. **28** (Springer, 2019).