

Modeling light pollution using geospatial, meteorological, energy usage, and population data

Miles Leonard, Danny Nguyen
Group 1

Instructor: Dr. Charles Nicholson
DSA 5103– Fall 2025

December 9, 2025

Contents

1	Executive Summary	1
2	Problem Background	2
2.1	Problem Description	2
2.2	Data Description	3
2.3	Exploratory Data Analysis	3
3	Methodology	4
3.1	Feature Selection	4
3.2	Model Choice	6
3.3	Validation Approach	6
4	Results	6
4.1	Model Results and Performance	6
4.2	Key Findings	7
5	Conclusion	8

List of Figures

1	PCA Bi-Plot w/ Night-Sky Brightness Gradient	5
2	Predicted Values and Residuals	7

List of Tables

1	Summary of <code>caret</code> training results for <code>earth</code>	7
2	Performance of Post-Processed Predictions	7
3	Final <code>earth</code> Model Variable Importance	8

1 Executive Summary

Problem Statement Light pollution is a night-time phenomenon caused by the atmospheric scattering of excess light from things such as street lamps or other outdoor light fixtures that then bounces off of molecules in the air. Light pollution is commonly measured as night sky brightness (in magnitudes per square arcsecond), but is typically only directly measured at sites like observatories. Even then, data from these observatories is rarely accessible in real-time. The Globe at Night campaign created a monitoring network for observatories across the world to provide night sky brightness data for public use, with data going back as far as 2014. Light pollution, being an atmospheric phenomenon, is heavily impacted by weather conditions. The objective for this project is to predict light pollution (night sky brightness) at a given location using primarily meteorological, energy usage, and population data.

Major Concerns/Assumptions The training data for night sky brightness is limited to thirteen observation sites with relatively sparse readings is expected to be a significant training obstacle, likely to skew the models locally to those geographic locations. Additionally, the association of zip-codes to the nearest geographic electric-service provider is a rough approximation that may not accurately reflect the actual service-provider for a given area, most notably for high population density zones that are serviced by multiple providers.

Summary of Findings TO-DO

Recommendations TO-DO

2 Problem Background

2.1 Problem Description

The use of artificial light at night (ALAN) in outdoor spaces—e.g., through the use of light fixtures such as street lamps or flood lights—causes what is known as light pollution. More specifically, improper and excessive use of ALAN, which has risen dramatically alongside rapid urbanization and industrialization around the globe, is the primary source of light pollution. The definition of light pollution varies across different agencies, but the common denominator amongst them all is that it is the culmination of the unwanted, adverse aspects of ALAN. There are three main components that make up light pollution: glare, light trespass, and skyglow; for the purposes of this project, however, the focus is solely on skyglow[7]. Skyglow, the most broadly impactful aspect of light pollution, is the result of light being emitted upwards and then bouncing off of particles in the Earth’s lower atmosphere, scattering. It obscures the natural night sky, preventing those in the city from being able to marvel at the stars. Those with exceptional vision and access to pristine night skies are able to see up to 7,000 stars with the naked eye, however, that number plummets to fewer than 100 for the same observer in an urban area[5].

As it currently stands, light pollution-free skies are becoming more and more difficult to find. Over 99% of US and European populations are affected by some level of light pollution, and a recent analysis found that skyglow is rapidly worsening at a rate of nearly 10% per year[4]. Modeling the skyglow component of light pollution, typically referred to as night sky brightness, is a crucial aspect of many light pollution studies. Unfortunately, there is no definitive way to assess skyglow, and each method has its own pros and cons. Light pollution data for modeling often comes from a combination of ground-based measurements and satellite data. Ground-based measurements are most often taken using a sky quality meter (SQM), but have also been derived from citizen-science reports of naked-eye limiting magnitude (NELM). NELM is the magnitude (a logarithmic scale describing the apparent brightness of a star) of the faintest visible star to a naked-eye observer; the lower the magnitude, the brighter the object. The SQM is a handheld sensor that measures the brightness of the sky directly overhead in units of magnitudes per square arcsecond.

Due to the complexity of light emission behavior and limitations in computing power, research into different modeling methods is still ongoing [3]. Most light pollution models are numerical models based on a series of functions that describe upward light flux and atmospheric scattering. Many more radiative transfer-based numerical models exist, but are often tailored towards modeling meteorological conditions and therefore not suitable for modeling light pollution specifically [3]. Recently, it was found that most, if not all light pollution models are significantly, systematically biased through implicitly assuming that all particles in the lower atmosphere are homogenous and spherical (Mie theory), despite being a safe assumption for daytime light models [2]. A popular global map known as the New world atlas of artificial night sky brightness [1] is another numerical model utilizing both ground-based and satellite measurements, but assumed a perfectly clear, calm, atmosphere over the entire globe. This limitation, albeit understandable, greatly impacts the accuracy of its output. Atmospheric conditions vary widely across regions. Not only

that, but atmospheric conditions are always changing as well. Consecutive measurements of skyglow in the exact same location, even just hours apart, could produce different results.

While acknowledging that the extent of light pollution is influenced by a multivariate set of economic indicators [6], the current model specification is limited to assessing the predictive power of population metrics and energy consumption data.

2.2 Data Description

Night-Sky Brightness Globe at Night (GaN) The GaN-MN project, an extension of the original Globe at Night project, is a global night sky brightness monitoring network using a commercially available meter (SQM-LE by Unihedron) for long-term monitoring of the light pollution conditions in different places around the world. <https://globeatnight.org/gan-mn/>

Meteorological Data ERA5 provides hourly estimates for a large number of atmospheric, ocean-wave and land-surface quantities. An uncertainty estimate is sampled by an underlying 10-member ensemble at three-hourly intervals. Ensemble mean and spread have been pre-computed for convenience. Such uncertainty estimates are closely related to the information content of the available observing system which has evolved considerably over time. They also indicate flow-dependent sensitive areas. To facilitate many climate applications, monthly-mean averages have been pre-calculated too, though monthly means are not available for the ensemble mean and spread. <https://cds.climate.copernicus.eu/datasets/reanalysis-era5-single-levels?tab=overview>

Zip-Code Populations A tabulation of the total population per zip code per the 2020 U.S. Census. <https://data.census.gov/table>

Electric Retail Service Territories A table of electric power retail service territories in the U.S. These are areas serviced by electric power utilities responsible for the retail sale of electric power to local customers, whether residential, industrial, or commercial. <https://ornl.opendatasoft.com/explore/dataset/electric-retail-service-territories/information/>

Population-Weighted Centroids for Zip Code Tabulation Areas (ZCTAs) This dataset denotes ZIP Code centroid locations weighted by population. <https://hudgis-hud.opendata.arcgis.com/datasets/d032efff520b4bf0aa620a54a477c70e/about>

2.3 Exploratory Data Analysis

Night-Sky Brightness The Globe at Night dataset only contains thirteen sites and night sky brightness readings are relatively sparse (~63% missing); however most of the missing measurements are from the

daylight hours. Additionally the minimum time resolution of measurements is to the UTC hour instead of the local hour.

Meteorological Data Given that ERA5 is a reanalysis dataset, there are no missing values that must be accommodated.

Electric-Service Provider Coverage The service territories dataset is missing zip-code data for approximately 58% of electrical service providers.

Population 4320 zip-codes are missing between the intersection of the centroids and population datasets. This is due to zip-codes that did not have population data in the 2020 census data.

Principal Component Analysis Omitting the night-sky brightness (NSB) from the variables compiled via the sources above (26 total), when performing a PCA 43.4% of the data-set variance is contained within components 1 and 2. A random sample of 5% of the data was selected to plot against PC1 and PC2 for easier viewing, with each point colored by a gradient for the associated NSB measured value. This bi-plot is shown in Figure 1.

A cursory inspection of the figure implies there may be a skew of low NSB nights (the points of interest) in the direction of PC1.

3 Methodology

3.1 Feature Selection

Night-Sky Brightness Missing brightness readings are dropped from the training data entirely. The time of observation is assumed to be close enough to local time, so no time zone adjustments are made. The brightness readings are assumed to be correlated to the time as a cyclical feature, so the month and hour of observation are encoded as wave functions at an angular offset from typical brightness zeniths. Night sky brightness observations are recorded approximately every 5 minutes. However, the meteorological data is provided at an hourly resolution, so the mean night sky brightness of each hour is calculated and used instead. A local timestamp, as well as a UTC timestamp is recorded at the time of each observation as well, but only the UTC timestamp was kept, as the ERA5 timestamps are provided in UTC.

Meteorological Data ERA5 is currently one of the most popular and comprehensive meteorological reanalysis datasets publicly available, with a horizontal spatial resolution of 0.25 by 0.25 degrees and an hourly temporal resolution. While ERA5 provides a very large variety of atmospheric/meteorological variables, only the variables air temperature, dewpoint temperature, surface air pressure, and percentage of high cloud cover, medium cloud cover, and low cloud cover were chosen for use. With the python package




Figure 1: PCA Bi-Plot w/ Night-Sky Brightness Gradient

astropy, using a site's location and time of observation, the altitude and azimuth of the sun and moon were calculated to serve as feature variables as well.

Electric-Service Provider Rows with missing zip-code data for service providers are dropped from the training data due to the inability to map given providers to sensible zip-codes.

Population The zip-code centroids with missing population data are dropped from the training data. The values for population are not being estimated for time-adjusted values.

Expected Energy Usage The assumption is made that energy usage by nearby population centers is correlated to their nearest electric-service providers. Using the population-centroids of each zip-code,

the nearest electric-service provider is associated with each population center. The nominal energy usage of each population center is then encoded as a sine function of the month of observation to differentiate between the seasonal energy usage patterns. The `SUMMR_PEAK` and `WINTR_PEAK` values are used as the amplitudes of the wave functions for the respective seasons.

Distance from Population Centers Using the Haversine formula, the great circle distance between the nearest zip-code population centroid and the observation site is calculated and included as a feature. Since the radiance from light-sources are known to decay with distance, this distance is encoded as its inverse square.

3.2 Model Choice

The `earth` R package was used for its implementation of the MARS regression model. The `caret` package was used to train and tune the hyper-parameters `nprune` and `degree` of the model. Post-processing will be applied before the final prediction to clamp negative values up to the minimum magnitude of 1.

3.3 Validation Approach

The training control used for the model training was 10-fold CV with 5 repeats. The full dataset was split 75/25 between the training and testing holdout data.

4 Results

4.1 Model Results and Performance

Training Results The model training results are shown in Table 1. The hyper-parameters were tuned up to `degree`= 5 and `nprune`= 45, with marginal improvements in *RMSE* with each expansion of the tuning grid after `degree`= 4 and `nprune` > 30. More fine tuning of the hyper-parameters is possible, but the performance appears to be relatively plateaued with the current predictors.

Post-Processing After clamping predicted magnitudes < 1 up to 1, the resampled performance yielded marginal improvements across the categories as shown in Table 2.

Residual Analysis Figure ?? shows a plot demonstrating the post-processed prediction of the NSB against the actual NSB measured magnitude. Figure ?? shows a plot demonstrating the residual values for each predicted NSB magnitude. From these plots it is seen that the model has been relatively strongly impacted by the large cluster of bright nights in the dataset, and tends to over-estimate the night sky brightness.

degree	nprune	RMSE	R^2	MAE
3	35	0.8192	0.9213	0.5225
3	40	0.8141	0.9223	0.5199
3	45	0.8131	0.9225	0.5192
4	35	0.8011	0.9248	0.5200
4	40	0.7983	0.9253	0.5178
4	45	0.7977	0.9254	0.5172
5	35	0.8009	0.9248	0.5202
5	40	0.7981	0.9253	0.5182
5	45	0.7975	0.9254	0.5177

Table 1: Summary of `caret` training results for `earth`

RMSE	R^2	MAE
0.7957	0.9278	0.5268

Table 2: Performance of Post-Processed Predictions




Figure 2: Predicted Values and Residuals

Coefficients After the relatively aggressive pruning parameter of `nprune`= 45 in the final model, the original set of 26 predictors were trimmed down to a 5-way interaction between the 10 listed in Table 3.

4.2 Key Findings

TO-DO

variable	nsubsets	gcv	rss
sun_alt	44	100.00	100.00
moon_alt	42	79.415	79.42
elec_dist_sq	42	73.623	73.63
noon_cos	42	73.623	73.63
zip_pop	39	43.679	43.70
moon_phase	37	34.399	34.43
zip_dist_sq	32	18.826	18.89
summer_cos	26	12.786	12.86
es	22	10.295	10.38
elec_summer_peak	19	8.653	8.74

Table 3: Final `earth` Model Variable Importance

5 Conclusion

Summary TO-DO

Key Insights TO-DO

Final Recommendations and Impact TO-DO

References

- [1] Fabio Falchi et al. “The new world atlas of artificial night sky brightness”. In: *Science Advances* 2.6 (2016), e1600377. URL: <https://www.science.org/doi/abs/10.1126/sciadv.1600377>.
- [2] M. Kocifaj et al. “A systematic light pollution modelling bias in present night sky brightness predictions”. In: *Nature Astronomy* 7.269-279 (2023). URL: <https://doi.org/10.1038/s41550-023-01916-y>.
- [3] Z. Kolláth, D. Száz, and K. Kolláth. “Measurements and Modelling of Artificial Sky Brightness: Combining Remote Sensing from Satellites and Ground-Based Observations”. In: *Remote Sensing* 13.3653 (2021). URL: <https://doi.org/10.3390/rs13183653>.
- [4] C. C. M. Kyba et al. “Citizen scientists report global rapid reductions in the visibility of stars from 2011 to 2022”. In: *Science* 379.265-268 (2023). URL: <https://doi.org/10.1126/science.abq7781>.
- [5] B. Mizon. “Living with Light. Light Pollution: Responses and Remedies”. In: *The Patrick Moore Practical Astronomy Series, Springer* 3.31 (2012).
- [6] R. N. Olsen, T. Gallaway, and Mitchell D. “Modelling US light pollution”. In: *Journal of Environmental Planning and Management* 57.6 (2014). URL: <https://doi.org/10.1080/09640568.2013.774268>.
- [7] D. Welch and Coauthors. *The World At Night: Preserving natural darkness for heritage conservation and night sky appreciation*. 2024. URL: <https://portals.iucn.org/library/sites/library/files/documents/PAG-033-En.pdf>.