


ANNOUNCEMENTS

- Homework 10 (it's long; you started it, right?)
 - Due this Friday
- Final Homework is due Friday the 9th
 - Magnetic dipoles and some magnetic matter
- Final Exam (20%)
 - 12:45pm-2:45pm on Thursday the 15th in this room
- Detailed grade projections by Monday 12th
 - w/ clicker bonus, but not HW 11

The vector potential \mathbf{A} due to a long straight wire with current I along the z -axis is in the direction parallel to:

- A. \hat{z}
- B. $\hat{\phi}$ (azimuthal)
- C. \hat{s} (radial)

Assume the Coulomb Gauge



Consider a fat wire with radius a with uniform current I_0 that runs along the $+z$ -axis. We can compute the vector potential due to this wire directly. What is \mathbf{J} ?

- A. $I_0/(2\pi)$
- B. $I_0/(\pi a^2)$
- C. $I_0/(2\pi a)\hat{z}$
- D. $I_0/(\pi a^2)\hat{z}$
- E. Something else!?

Consider a fat wire with radius a with uniform current I_0 that runs along the $+z$ -axis. Given $\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{J}(\mathbf{r}')}{\mathfrak{R}} d\tau'$, which components of \mathbf{A} need to be computed?

- A. All of them
- B. Just A_x
- C. Just A_y
- D. Just A_z
- E. Some combination

Consider line of charge with uniform charge density, $\lambda = \rho/(\pi a^2)$. What is the magnitude of the electric field outside of the line charge (at a distance $s > a$)?

- A. $E = \lambda/(4\pi\epsilon_0 s^2)$
- B. $E = \lambda/(2\pi\epsilon_0 s^2)$
- C. $E = \lambda/(4\pi\epsilon_0 s)$
- D. $E = \lambda/(2\pi\epsilon_0 s)$
- E. Something else?!

Use Gauss' Law

Consider a shell of charge with surface charge σ that is rotating at angular frequency of ω . Which of the expressions below describe the surface current, \mathbf{K} , that is observed in the fixed frame.

- A. $\sigma \omega$
- B. $\sigma \dot{\mathbf{r}}$
- C. $\sigma \mathbf{r} \times \omega$
- D. $\sigma \omega \times \mathbf{r}$
- E. Something else?

What is the physical interpretation of $\oint \mathbf{A} \cdot d\mathbf{l}$?

- A. The current density \mathbf{J}
- B. The magnetic field \mathbf{B}
- C. The magnetic flux Φ_B
- D. It's none of the above, but is something simple and concrete
- E. It has no particular physical interpretation at all

Consider a square loop enclosing some amount of magnetic field lines with height H and length L . We intend to compute $\Phi_B = \oint \mathbf{A} \cdot d\mathbf{l}$? What happens to Φ_B as H becomes vanishingly small?

- A. Φ_B stays constant
- B. Φ_B gets smaller but doesn't vanish
- C. $\Phi_B \rightarrow 0$

Consider a square loop enclosing some amount of magnetic field lines with height H and length L . If $\Phi_B \rightarrow 0$ as $H \rightarrow 0$ (or $L \rightarrow 0$), what does that say about the continuity of \mathbf{A} ?

$$\Phi_B = \oint \mathbf{A} \cdot d\mathbf{l}$$

- A. \mathbf{A} is continuous at boundaries
- B. \mathbf{A} is discontinuous at boundaries
- C. ???