


I have two very long, parallel wires each carrying a current I_1 and I_2 , respectively. In which direction is the force on the wire with the current I_2 ?

- A. Up
- B. Down
- C. Right
- D. Left
- E. Into or out of the page



What is the magnitude of $\frac{d\mathbf{l} \times \hat{\mathbf{R}}}{R^2}$?


A. $\frac{dl \sin \phi}{z^2}$

B. $\frac{dl}{z^2}$

C. $\frac{dl \sin \phi}{z^2 + a^2}$


D. $\frac{dl}{z^2 + a^2}$

E. something else!




What is $d\mathbf{B}_z$ (the contribution to the vertical component of \mathbf{B} from this $d\mathbf{l}$ segment?)

- A. $\frac{dl}{z^2+a^2} \frac{a}{\sqrt{z^2+a^2}}$
- B. $\frac{dl}{z^2+a^2}$
- C. $\frac{dl}{z^2+a^2} \frac{z}{\sqrt{z^2+a^2}}$
- D. $\frac{dl \cos \phi}{\sqrt{z^2+a^2}}$
- E. Something else!



What is $\oint \mathbf{B} \cdot d\mathbf{l}$ around this purple (dashed) Amperian loop?




- A. $\mu_0(|I_2| + |I_1|)$
- B. $\mu_0(|I_2| - |I_1|)$
- C. $\mu_0(|I_2| + |I_1| \sin \theta)$
- D. $\mu_0(|I_2| - |I_1| \sin \theta)$
- E. $\mu_0(|I_2| + |I_1| \cos \theta)$

Stoke's Theorem says that for a surface S bounded by a perimeter L , any vector field \mathbf{B} obeys:


$$\int_S (\nabla \times \mathbf{B}) \cdot d\mathbf{A} = \oint_L \mathbf{B} \cdot d\mathbf{l}$$

Does Stoke's Theorem apply for any surface S bounded by a perimeter L , even this balloon-shaped surface S ?



- A. Yes
- B. No
- C. Sometimes

Rank order $\int \mathbf{J} \cdot d\mathbf{A}$ (over blue surfaces) where \mathbf{J} is uniform, going left to right:




- A. $\text{iii} > \text{iv} > \text{ii} > \text{i}$
- B. $\text{iii} > \text{i} > \text{ii} > \text{iv}$
- C. $\text{i} > \text{ii} > \text{iii} > \text{iv}$
- D. Something else!!
- E. Not enough info given!!

Much like Gauss's Law, Ampere's Law is always true (for magnetostatics), but only useful when there's sufficient symmetry to "pull \mathbf{B} out" of the integral.

So we need to build an argument for what \mathbf{B} looks like and what it can depend on.

For the case of an infinitely long wire, can \mathbf{B} point radially (i.e., in the \hat{s} direction)?



- A. Yes
- B. No
- C. ???


Continuing to build an argument for what **B** looks like and what it can depend on.

For the case of an infinitely long wire, can **B** depend on z or ϕ ?

A. Yes

B. No

C. ???




Finalizing the argument for what \mathbf{B} looks like
and what it can depend on.

For the case of an infinitely long wire, can \mathbf{B}
have a \hat{z} component?

A. Yes

B. No

C. ???



For the infinite wire, we argued that $\mathbf{B}(\mathbf{r}) = B(s)\hat{\phi}$. For the case of an infinitely long **thick** wire of radius a , is this functional form still correct? Inside and outside the wire?

- A. Yes
- B. Only inside the wire ($s < a$)
- C. Only outside the wire ($s > a$)
- D. No