



Advanced Time Series Analysis and Forecasting

Seminar 10: Long memory and rough volatility



Daniel Traian PELE

Bucharest University of Economic Studies

IDA Institute Digital Assets

Blockchain Research Center

AI4EFin Artificial Intelligence for Energy Finance

Romanian Academy, Institute for Economic Forecasting

MSCA Digital Finance

Acronyms used in this material (1/2)

- ▣ AI: Artificial Intelligence
- ▣ AR: AutoRegressive (model); in event studies: Abnormal Return
- ▣ ARCH: AutoRegressive Conditional Heteroskedasticity
- ▣ ARFIMA: AutoRegressive Fractionally Integrated Moving Average (model)
- ▣ BET: Bucharest Exchange Trading (index)
- ▣ BNR: Banca Națională a României (the National Bank of Romania)
- ▣ BUX: Budapești Értéktőzsde index (the Budapest Stock Exchange index)
- ▣ CAC: Cotation Assistée en Continu (CAC 40 index, Paris) — the CAC 40 index of the Paris bourse
- ▣ CEE: Central and Eastern Europe
- ▣ DM: Diebold–Mariano (test)
- ▣ ELW: Exact Local Whittle (estimator)
- ▣ EODHD: EOD Historical Data (market-data provider)
- ▣ FCVAR: Fractionally Cointegrated VAR
- ▣ FFT: Fast Fourier Transform
- ▣ FIGARCH: Fractionally Integrated GARCH
- ▣ FTSE: Financial Times Stock Exchange (FTSE Russell, index provider)
- ▣ GPH: Geweke–Porter–Hudak (estimator)
- ▣ HAC: Heteroskedasticity and Autocorrelation Consistent (standard errors)
- ▣ HAR: Heterogeneous AutoRegressive (model)
- ▣ LW: Local Whittle (estimator)
- ▣ MSE: Mean Squared Error
- ▣ OLS: Ordinary Least Squares
- ▣ PX: Prague Stock Exchange index (index Burzy cenných papírů Praha)
- ▣ QLIKE: Quasi-LIKelihood loss
- ▣ RFSV: Rough Fractional Stochastic Volatility (model)
- ▣ RV: Realised Variance

Acronyms used in this material (2/2)

- SD: Standard Deviation
- SE: Standard Error
- WIG20: Warszawski Indeks Giełdowy 20 (the Warsaw Stock Exchange index of the 20 largest companies)

Today's question and route

- **Question:** how do we measure, test and use long memory and roughness in volatility and inflation?
 - ▶ this seminar comes **before** Lecture 10: the section “What you need today” gives every definition the tasks use
- **Route**
 - ▶ Part A: ARFIMA autocorrelations, fractional weights, standard errors and bias of the estimators, fractional Gaussian noise, the RFSV kernel, circulant embedding, switching that mimics long memory
 - ▶ Part B: memory of FTSE 100 realised variance, BET and EUR/RON; roughness of the Euro Stoxx 50 and of Bitcoin; forecasting FTSE 100 and CAC 40 realised variance
 - ▶ Part C: co-memory of DAX and CAC 40 volatility; an AI answer to audit
- Notebook for today: [open the seminar notebook in Google Colab](#); each task names its notebook section

Exercise map

Task	Question	Type	Model
A1, A2	ARFIMA autocorrelations and the mean; fractional weights	Solved, Proposed	A1
A3, A4	standard errors of GPH and LW; bias and bandwidth	Solved, Proposed	A3
A5, A6	fractional Gaussian noise; the RFSV kernel	Solved, Proposed	A5
A7, A8	circulant embedding by hand; switching that mimics long memory	Solved, Proposed	A7, A1
B1, B2	memory of FTSE 100 log RV; BET and EUR/RON with the Qu test	Solved, Proposed	B1
B3, B4	roughness of the Euro Stoxx 50; Bitcoin by sampling interval	Solved, Proposed	B3
B5, B6	forecasting RV of the FTSE 100 and the CAC 40	Solved, Proposed	B5
C1, C2	fractional cointegration of two volatilities; what is wrong in an AI answer?	Proposed	B1, A1–A8

- **[Solved]**: full solution in the slides and in the notebook, a model to follow; **[Proposed]**: you solve it, following the model

Data used

Series	Source	Frequency	Period
RV of FTSE 100, Euro Stoxx 50, DAX, CAC 40	Heber et al. (2009)	daily, 5-minute returns	2000–2022
RV of Bitcoin	Binance one-minute prices	daily, 1–60 minutes	2018–2026
BET	EODHD	daily	2000–2026
EUR/RON	BNR reference rate	daily	2005–2026

- RV in $\%^2$, returns in %; in the notebook: `gph`, `local_whittle`, `elw`, `qu_stat`, `gjr_scaling`, `h_with_noise`, `rfsv_kernel`, `forecast_rv`, `fcvar_fit`; no account or key is needed

What you need today (1/4): long memory

- Long memory: $\rho(k) \sim c k^{2d-1}$ and $f(\lambda) \sim G\lambda^{-2d}$ near zero, $0 < d < 1/2$; Hurst exponent $H = d + 1/2$
 - ▶ ARFIMA(0, d , 0): $(1 - L)^d X_t = \varepsilon_t$; $(1 - L)^d = \sum_k \pi_k L^k$, $\pi_0 = 1$, $\pi_k = \pi_{k-1}(k - 1 - d)/k$ [Hosking \(1981\)](#)
 - ▶ $\rho(k) = \rho(k - 1)(k - 1 + d)/(k - d)$, $\rho(k) \sim \frac{\Gamma(1-d)}{\Gamma(d)} k^{2d-1}$; $\pi_k \sim k^{-d-1}/\Gamma(-d)$
- Variance of the mean: $\text{Var}(\bar{X}_n) = \frac{1}{n} \sum_{|k| < n} (1 - |k|/n) \gamma(k)$, of order n^{2d-1} , not n^{-1}
- Periodogram $I(\lambda_j) = \frac{1}{2\pi n} |\sum_t X_t e^{-i\lambda_j t}|^2$ at Fourier frequencies $\lambda_j = 2\pi j/n$; bandwidth m : the number of low frequencies used

What you need today (2/4): estimators of d

- ▣ **GPH** Geweke and Porter-Hudak (1983): OLS of $\log I(\lambda_j)$ on $-\log(4 \sin^2(\lambda_j/2))$, $j \leq m$; SE $\pi/\sqrt{24m}$
- ▣ **Local Whittle** Robinson (1995b): $\hat{d} = \arg \min_d \log \left(\frac{1}{m} \sum_j \lambda_j^{2d} I(\lambda_j) \right) - \frac{2d}{m} \sum_j \log \lambda_j$; SE $1/(2\sqrt{m})$
 - ▶ **exact local Whittle** Shimotsu and Phillips (2005), Shimotsu (2010): the periodogram of $(1-L)^d(X_t - \hat{\mu})$ instead of $\lambda_j^{2d} I(\lambda_j)$; valid also for $d \geq 1$
- ▣ Bias for ARFIMA(1, d , 0): $\mathbb{E}\hat{d} - d \approx C(m/n)^2$, $C = \frac{2\pi^2}{9} \cdot \frac{2\phi}{(1-\phi)^2}$; MSE-optimal $m^* = (n^4/(16C^2))^{1/5}$

What you need today (3/4): breaks and co-memory

- Level shifts, breaks and rare regime switches create a spectral pole: $\hat{d}$ is large although the process has short memory [Diebold and Inoue \(2001\)](#)
 - ▶ **Qu test** [Qu \(2011\)](#): $W = \sup_{r \in [\varepsilon, 1]} (\sum_j \nu_j^2)^{-1/2} |\sum_{j \leq mr} \nu_j (\lambda_j^{2\hat{d}} I(\lambda_j) / \hat{G} - 1)|$, $\nu_j = \log \lambda_j - \overline{\log \lambda}$; reject stationary long memory for large W
 - ▶ $m = n^{0.7}$, $\varepsilon = 0.02$; simulated critical values (10%, 5%, 1%): 1.09, 1.22, 1.56
- **FCVAR** [Johansen and Nielsen \(2012\)](#): $\Delta^d X_t = \alpha \beta' L_b \Delta^{d-b} X_t + \varepsilon_t$, $L_b = 1 - \Delta^b$; $\beta' X_t$ has memory $d - b$
 - ▶ for fixed (d, b) : Johansen reduced-rank regression; trace tests $\chi_{(p-r)^2}^2$ when $b < 1/2$

What you need today (4/4): rough volatility

- fBm W^H : $\mathbb{E}|W_{t+\Delta}^H - W_t^H|^2 = \Delta^{2H}$; increments (fGn) $\rho(k) = \frac{1}{2}(|k+1|^{2H} - 2|k|^{2H} + |k-1|^{2H})$ Mandelbrot and Van Ness (1968)
 - circulant embedding Davies and Harte (1987): first row $(\gamma_0, \dots, \gamma_{n-1}, \gamma_{n-2}, \dots, \gamma_1)$, eigenvalues = FFT of the row; exact if all ≥ 0
- Gatheral, Jaisson and Rosenbaum (2018): $m(q, \Delta) = \text{mean} |\log \sigma_{t+\Delta} - \log \sigma_t|^q \propto \Delta^{\zeta_q}$, $\zeta_q = qH$, $\sigma_t = \sqrt{RV_t}$; with i.i.d. error: $m(2, \Delta) = \nu^2 \Delta^{2H} + 2s^2$
- RFSV predictor: $\widehat{\log \sigma_{t+h}^2} = \sum_k w_k \log \sigma_{t-k}^2$, $w_k \propto \int_k^{k+1} \frac{du}{(u+h)u^{H+1/2}}$; variance: $\exp(\cdot + 2c\nu^2 h^{2H})$,
 $c = \frac{\Gamma(3/2-H)}{\Gamma(H+1/2)\Gamma(2-2H)}$
 - QLIKE loss $RV/F - \log(RV/F) - 1$ Patton (2011); DM test with Newey–West variance (Chapter 1)

A1: ARFIMA autocorrelations and the mean [Solved]

Task

- ▣ ARFIMA(0, 0.3, 0).
- ▣ 1. Show that $\rho(1) = d/(1 - d)$ and compute $\rho(1)$, $\rho(10)$, $\rho(100)$ with the recursion.
- ▣ 2. Compare $\rho(100)$ with $\frac{\Gamma(1-d)}{\Gamma(d)} 100^{2d-1}$.
- ▣ 3. For $n = 1000$, compute the ratio of the true SD of $\bar{X}$ to the i.i.d. formula $\sqrt{\gamma(0)/n}$ (notebook A1).
- ▣ Report: three numbers; two numbers; one number and one sentence.

Solution

- ▣ 1. $\rho(1) = \rho(0) \cdot \frac{0+d}{1-d}$; $\rho(1) = 0.429$, $\rho(10) = 0.173$, $\rho(100) = 0.069$
- ▣ 2. Asymptotic formula: 0.069, practically equal already at $k = 100$
- ▣ 3. Ratio 7.55: the i.i.d. standard error is 7.55 times too small
- ▣ Interpretation: with $d = 0.3$, 1000 observations carry the information of far fewer independent ones; HAC with a fixed bandwidth cannot repair this

A2: fractional differencing weights [Proposed]

Solution discussed in class

Task

- ▣ $d = 0.4$. Model: A1.
- ▣ 1. Compute $\pi_1, \dots, \pi_5$ of $(1 - L)^{0.4}$.
- ▣ 2. Compare π_{100} with $100^{-d-1}/\Gamma(-d)$.
- ▣ 3. Compute $\sum_{k=0}^{1000} \pi_k$; what is its limit, and what does the gap mean for a truncated filter?
- ▣ Report: five numbers; two numbers; one number and one sentence.

A3: standard errors of GPH and local Whittle [Solved]

Task

- ▣ $n = 5000$, $m = \lfloor n^{0.65} \rfloor$; a local Whittle estimate $\hat{d} = 0.45$.
- ▣ 1. Compute m and the asymptotic SE of GPH and of local Whittle.
- ▣ 2. Give the 95% interval for d and the relative efficiency of LW against GPH.
- ▣ 3. Which m gives a half-width of 0.05?
- ▣ Report: three numbers; an interval and one number; one number.

Solution

- ▣ 1. $m = 253$; GPH $\pi/\sqrt{24m} = 0.0403$; LW $1/(2\sqrt{m}) = 0.0314$
- ▣ 2. $[0.388, 0.512]$; efficiency $(\pi^2/24)/(1/4) = \pi^2/6 = 1.645$
- ▣ 3. $1.96/(2\sqrt{m}) \leq 0.05 \Rightarrow m \geq 385$
- ▣ Interpretation: the interval includes $1/2$, so stationarity is not established; precision is bought with bandwidth, and bandwidth costs bias (A4)

A4: bias and the optimal bandwidth [Proposed]

Solution discussed in class

Task

- ▣ ARFIMA(1, d , 0) with $\phi = 0.5$, $n = 5000$. Model: A3.
- ▣ 1. Compute the bias constant C .
- ▣ 2. For $m = \lfloor n^{0.8} \rfloor$, compute the bias and the SD of local Whittle.
- ▣ 3. Compute m^* , its exponent $\log m^* / \log n$, and the bias and SD at m^* .
- ▣ Report: one number; two numbers; four numbers and one sentence.

A5: fractional Gaussian noise and scaling [Solved]

Task

- ▣ fGn with $H = 0.1, 0.3$ and 0.7 .
- ▣ 1. Compute $\rho(1) = 2^{2H-1} - 1$ and $\rho(2)$ for each H .
- ▣ 2. Compute $m(2, 22)/m(2, 1) = 22^{2H}$: how much more does log volatility move over a month than over a day?
- ▣ 3. Show that $\zeta_q = qH$ for fBm.
- ▣ Report: six numbers; three numbers; one derivation.

Solution

- ▣ 1. $H = 0.1$: $-0.426, -0.026$; $H = 0.3$: $-0.242, -0.049$; $H = 0.7$: $0.320, 0.189$
- ▣ 2. $1.86, 6.39, 75.75$: with $H = 0.1$ the monthly variance of changes is less than twice the daily one
- ▣ 3. $W_{t+\Delta}^H - W_t^H \sim N(0, \Delta^{2H})$: $\mathbb{E}|\cdot|^q = \Delta^{qH} \mathbb{E}|Z|^q$; take logs
- ▣ Interpretation: rough log volatility moves a lot from day to day but drifts little: strong negative correlation of daily changes

A6: the RFSV kernel and the variance correction [Proposed]

Solution discussed in class

Task

- $H = 0.1$, $\nu = 0.3$, horizon $h = 1$; kernel truncated at 500 days. Model: A5.
- 1. Compute the share of the weight on the last day, the last 5 and the last 22 days (notebook A6).
- 2. Compute $c(H)$ and the correction $2c\nu^2 h^{2H}$ for $h = 1, 5, 22$.
- 3. By how much would ignoring the correction understate the variance forecast at $h = 22$?
- Report: three shares; four numbers; one number.

A7: circulant embedding by hand [Solved]

Task

- ▣ Three points of fGn with $H = 0.3$: $\gamma_0, \gamma_1, \gamma_2$.
- ▣ 1. Compute γ_1 and γ_2 .
- ▣ 2. Write the first row of the 4×4 circulant and its eigenvalues $\lambda_k = \sum_j c_j e^{-2\pi ijk/4}$.
- ▣ 3. Can the Davies–Harte method be used?
- ▣ Report: two numbers; a row and four numbers; one sentence.

Solution

- ▣ 1. $\gamma_1 = \frac{1}{2}(2^{0.6} - 2) = -0.2421$,
 $\gamma_2 = \frac{1}{2}(3^{0.6} - 2 \cdot 2^{0.6} + 1) = -0.0491$
- ▣ 2. $c = (\gamma_0, \gamma_1, \gamma_2, \gamma_1)$; $\lambda_0 = \gamma_0 + 2\gamma_1 + \gamma_2 = 0.4666$,
 $\lambda_1 = \lambda_3 = \gamma_0 - \gamma_2 = 1.0491$, $\lambda_2 = \gamma_0 - 2\gamma_1 + \gamma_2 = 1.4352$
- ▣ 3. Yes: all eigenvalues are positive, so the circulant is a valid covariance and its first 3 coordinates have the fGn covariance
- ▣ Interpretation: the embedding doubles the size but turns an $O(n^3)$ factorisation into two FFTs

A8: switching that mimics long memory [Proposed]

Solution discussed in class


Task

- ▣ $X_t = \mu S_t + \varepsilon_t$, $S_t \in \{-1, 1\}$ a Markov chain that stays with probability $q = 0.995$, $\mu = 1$, $\varepsilon_t \sim N(0, 1)$. Model: A1.
- ▣ 1. Show that $\rho(k) = \frac{\mu^2}{\mu^2 + 1} (2q - 1)^k$ and compute $\rho(1)$, $\rho(100)$, $\rho(500)$.
- ▣ 2. Compute the half-life of the autocorrelation.
- ▣ 3. Fit $\log \rho(k)$ on $\log k$ for $k = 5, \dots, 200$ and convert the slope to d .
- ▣ Report: three numbers; one number; one number and one sentence.

B1: memory of FTSE 100 realised variance [Solved]

- ▣ **Question:** how large is d for the log realised variance of the FTSE 100, and is it robust to the bandwidth and to level shifts?
- ▣ **Inputs:** Oxford-Man 5-minute RV of the FTSE 100, $n = 5\,586$ days to 25 February 2022
- ▣ **Tasks**
 1. Estimate d of log RV by GPH, local Whittle and exact local Whittle for $m = n^{0.5}, n^{0.6}, n^{0.7}$, with standard errors.
 2. Plot $\hat{d}(m)$ for $m = n^{0.40}, \dots, n^{0.85}$.
 3. Compute the Qu statistic with $m = n^{0.7}$ and $\varepsilon = 0.02$.
 4. Interpretation: is the memory of FTSE 100 volatility genuine, and is it stationary?
- ▣ **Report:** a table of nine estimates with SEs, one chart, one statistic, one sentence
- ▣ **Notebook:** section B1

B1: solution [Solved]



m	GPH (SE)	LW	ELW	SE (LW, ELW)
74	0.604 (0.075)	0.601	0.639	0.058
177	0.611 (0.048)	0.578	0.603	0.038
419	0.620 (0.031)	0.606	0.618	0.024

- Qu: $W = 0.33$ against the 10% value 1.09: no evidence of level shifts
- Interpretation: $\hat{d} \approx 0.6$ at every bandwidth with a flat $\hat{d}(m)$: genuine long memory, at or just beyond the stationarity border; report ELW


B2: BET and EUR/RON with the Qu test [Proposed]

- **Question:** is the memory of $|r_t|$ on the Bucharest market and for the leu genuine, or does it come from regime changes?
- **Inputs:** BET 2000–2026 and from 2010; EUR/RON (BNR) 2005–2026 and from 2014; model: B1
- **Tasks**
 1. Estimate d of $|r_t|$ by local Whittle with $m = n^{0.65}$ on the full sample and after the cut date.
 2. Compute the Qu statistic ($m = n^{0.7}$, $\varepsilon = 0.02$) on the same samples.
 3. Interpretation: which of the two markets has genuine long memory in volatility?
- **Report:** four estimates with SEs, four statistics, one sentence
- **Notebook:** section B2

B3: roughness of the Euro Stoxx 50 [Solved]

- **Question:** does the Gatheral–Jaisson–Rosenbaum scaling hold for the Euro Stoxx 50, and how rough is its volatility?
- **Inputs:** Oxford-Man 5-minute RV, $n = 5\,635$ days; $\sigma_t = \sqrt{RV_t}$; $q = 0.5, 1, 1.5, 2, 3$; $\Delta = 1, \dots, 50$
- **Tasks**
 1. Compute $m(q, \Delta)$ and the slopes ζ_q ; estimate H by regressing ζ_q on q through the origin.
 2. Estimate H with the measurement-error intercept $m(2, \Delta) = \nu^2 \Delta^{2H} + 2s^2$.
 3. Repeat the first estimate on the two halves of the sample.
 4. Interpretation: is ζ_q linear in q , and what does a departure mean?
- **Report:** five slopes, three values of H , ν , one sentence
- **Notebook:** section B3

B3: solution [Solved]



- $\zeta_q = 0.069, 0.132, 0.183, 0.217, 0.218$; $\hat{H} = 0.093$, $\hat{\nu} = 0.38$; with the error intercept 0.109; halves 0.091 and 0.095
- Interpretation: linear up to $q = 2$, flat between $q = 2$ and 3: large moments are driven by a few extreme days (heavy tails of the increments), so H from $q \leq 2$ is the safer reading; roughness is stable over time


B4: Bitcoin roughness by sampling interval [Proposed]

- **Question:** does the estimate of H for Bitcoin depend on the sampling interval of the realised variance?
- **Inputs:** Binance RV of Bitcoin from 1-, 5-, 15-, 30- and 60-minute returns, 2018–2026; model: B3
- **Tasks**
 1. Estimate H for each RV by the moment regression and with the error intercept.
 2. Estimate d of log RV by local Whittle with $m = n^{0.65}$.
 3. Interpretation: which direction does a coarser grid push H , and why?
- **Report:** ten values of H , five values of d , one sentence
- **Notebook:** section B4

B5: forecasting FTSE 100 realised variance [Solved]

- **Question:** does a rough or a long-memory model forecast FTSE 100 realised variance better than HAR?
- **Inputs:** rolling window of 1000 days, re-estimation every 20 days; out of sample 30 December 2003 – 26 January 2022 ($T = 4\,565$); horizons 1, 5, 22 days
- **Tasks**
 1. Forecast RV_{t+h} with HAR, ARFIMA(0, d , 0) and RFSV, with the lognormal corrections.
 2. Compute the average QLIKE of each model and the ratio to HAR.
 3. Compute the DM statistics against HAR.
 4. Interpretation: which model would you use at each horizon?
- **Report:** nine losses, six ratios, six DM statistics, one sentence
- **Notebook:** section B5

B5: solution [Solved]



h	QLIKE HAR	ARFIMA / HAR	DM	RFSV / HAR	DM
1	0.230	0.985	-0.98	1.038	1.49
5	0.324	0.968	-1.94	0.968	-1.33
22	0.561	0.895	-2.12	0.914	-1.93

- Interpretation: HAR at one day (RFSV is worse); at 22 days both ARFIMA ($p = 0.034$) and RFSV ($p = 0.053$) beat HAR by about a tenth; median rolling $\hat{H} = 0.11$

B6: forecasting CAC 40 realised variance [Proposed]

- **Question:** does the FTSE 100 ranking carry over to the CAC 40?
- **Inputs:** the same design, out of sample 22 December 2003 – 26 January 2022 ($T = 4\,631$); model: B5
- **Tasks**
 1. Repeat the three forecasts and the QLIKE ratios.
 2. Compute the DM statistics against HAR.
 3. Interpretation: is there a single best model across the two indices?
- **Report:** six ratios, six DM statistics, one sentence
- **Notebook:** section B6

C1: co-memory of DAX and CAC 40 volatility [Proposed]

- ▣ **Question:** do the volatilities of two neighbouring markets share one fractional trend?
- ▣ **Inputs:** log 5-minute RV of the DAX and the CAC 40 (Oxford-Man), $n = 5\,600$ common days; FCVAR with $k = 0$; model: B1
- ▣ **Tasks**
 1. Write a five-line pre-registration: hypothesis, model, grid of (d, b) , tests, what would change your mind.
 2. Estimate d of each series and of the spread $\beta'X_t$ by local Whittle, then the FCVAR by profile likelihood with trace tests.
 3. Interpretation: what do the data support, and what would you add to the model?
- ▣ **Report:** a five-line plan, five estimates, two tests, a project plan
- ▣ **Notebook:** section C1

C2: audit an AI answer [Proposed]

- ▣ A student asked an AI assistant to summarise long memory and rough volatility. The answer:
- ▣ (a) An ARFIMA with $d = 0.3$ has autocorrelations that decay exponentially, only more slowly than an AR(1).
- ▣ (b) The standard error of local Whittle is $\pi/\sqrt{24m}$.
- ▣ (c) A larger bandwidth m is always better, because the variance falls.
- ▣ (d) If the Qu test rejects, the series has structural breaks.
- ▣ (e) $H = 0.1$ means that volatility has short memory, because $H < 0.5$ is antipersistence.
- ▣ (f) FIGARCH with $0 < d < 1$ is covariance stationary with a finite unconditional variance.
- ▣ (g) Exact local Whittle remains consistent for $d = 1.2$, whereas local Whittle does not.
- ▣ Tasks
 - ▶ 1. For each statement, say whether it is correct; if not, give the correct statement and, where possible, the correct number (notebook section C2).
 - ▶ 2. Report: a list of seven verdicts with one line of justification each.

Key takeaways

- Long memory makes i.i.d. and HAC standard errors far too small
- Local Whittle is the default; exact local Whittle when d may exceed $1/2$; always vary m
- Test against level shifts before claiming long memory
- Log volatility is rough ($H \approx 0.1$) and persistent; the proxy affects H
- Forecast gains over HAR are small and horizon-dependent

After the seminar

- ▣ Lecture 10 develops each topic of today: spectral theory, asymptotics of the estimators, the Qu test, FCVAR, FIGARCH, rough volatility and its forecasts
- ▣ Try the [Proposed] tasks in the notebook
- ▣ C1 can grow into a team project: co-memory of Central and Eastern European volatilities
- ▣ Reading: [Robinson \(1995b\)](#); [Shimotsu and Phillips \(2005\)](#); [Qu \(2011\)](#); [Gatheral, Jaisson and Rosenbaum \(2018\)](#)
- ▣ **The seminar is for practice and is not graded; the solutions of [Proposed] tasks are discussed in class**

References

- ▣ Davies, R. B., & Harte, D. S. (1987). Tests for Hurst effect. *Biometrika*, 74(1), 95–101. doi:10.1093/biomet/74.1.95
- ▣ Diebold, F. X., & Inoue, A. (2001). Long memory and regime switching. *Journal of Econometrics*, 105(1), 131–159. doi:10.1016/S0304-4076(01)00073-2
- ▣ Gatheral, J., Jaisson, T., & Rosenbaum, M. (2018). Volatility is rough. *Quantitative Finance*, 18(6), 933–949. doi:10.1080/14697688.2017.1393551
- ▣ Geweke, J., & Porter-Hudak, S. (1983). The estimation and application of long memory time series models. *Journal of Time Series Analysis*, 4(4), 221–238. doi:10.1111/j.1467-9892.1983.tb00371.x
- ▣ Heber, G., Lunde, A., Shephard, N., & Sheppard, K. K. (2009). Oxford-Man Institute's realized library, version 0.3. Oxford-Man Institute, University of Oxford. web.archive.org
- ▣ Hosking, J. R. M. (1981). Fractional differencing. *Biometrika*, 68(1), 165–176. doi:10.1093/biomet/68.1.165
- ▣ Johansen, S., & Nielsen, M. Ø. (2012). Likelihood inference for a fractionally cointegrated vector autoregressive model. *Econometrica*, 80(6), 2667–2732. doi:10.3982/ECTA9299
- ▣ Mandelbrot, B. B., & Van Ness, J. W. (1968). Fractional Brownian motions, fractional noises and applications. *SIAM Review*, 10(4), 422–437. doi:10.1137/1010093
- ▣ Patton, A. J. (2011). Volatility forecast comparison using imperfect volatility proxies. *Journal of Econometrics*, 160(1), 246–256. doi:10.1016/j.jeconom.2010.03.034
- ▣ Qu, Z. (2011). A test against spurious long memory. *Journal of Business & Economic Statistics*, 29(3), 423–438. doi:10.1198/jbes.2010.09153
- ▣ Robinson, P. M. (1995b). Gaussian semiparametric estimation of long range dependence. *The Annals of Statistics*, 23(5), 1630–1661. doi:10.1214/aos/1176324317
- ▣ Shimotsu, K. (2010). Exact local Whittle estimation of fractional integration with unknown mean and time trend. *Econometric Theory*, 26(2), 501–540. doi:10.1017/S0266466609100075
- ▣ Shimotsu, K., & Phillips, P. C. B. (2005). Exact local Whittle estimation of fractional integration. *The Annals of Statistics*, 33(4), 1890–1933. doi:10.1214/009053605000000309