








Modelling Financial Markets

Seminar 0: Financial Markets in 2026 — Hands-on



Daniel Traian PELE
Drd. Antoaneta Amza, antoaneta.amza@csie.ase.ro

Bucharest University of Economic Studies
IDA Institute Digital Assets
Blockchain Research Center
AI4EFin Artificial Intelligence for Energy Finance
Romanian Academy, Institute for Economic Forecasting
MSCA Digital Finance

How Do Prices Become Growth and Risk?

- Question of the seminar
  - **how do we turn a price series into returns, growth and risk, measure the largest loss from a peak (drawdown) and check that the prices are right?**
- Outputs to produce
  - the answers to the A exercises
  - one figure or table per B exercise
  - one interpretation sentence per exercise
- Notebook for today: chapter0_seminar_notebook.ipynb in Colab
  - first step: Setup (save a copy, load the data); each task names its notebook section
- The exercises, their types and their order: the exercise map
- Seminar: Prof. Daniel Traian Pele, danpele@ase.ro; Antoaneta Amza, PhD candidate, antoaneta.amza@csie.ase.ro

Exercise Map

Exercise	Question	Type	Model
Core (in class)			
A1–A4	returns, compound annual growth, annualised volatility, drawdown on paper	Solved	–
B1	is one EUR/RON quote a real price?	Solved	–
B2	how did the SPY fund, gold and Bitcoin do since 2015?	Solved	–
A5, A6	your turn: the same calculations on new numbers	Proposed	A1–A4
Homework			
B3, B4	two more quotes; three other assets	Proposed	B1, B2
B5	how deep and long are drawdowns?	Proposed	A4, B2
C1	what is wrong in an artificial intelligence (AI) answer?	Proposed	A2–A3, B2
After the lecture (optional, at the end of this PDF)			
Packs 1–8	full data audit, calendars and correlations, precision of means and Sharpe ratios, extensions	Solved or Proposed (marked on each)	named on each task

- **[Solved]**: full solution in the slides (core) or in the notebook (after the lecture), a model to copy; **[Proposed]**: you solve it, the solution is discussed in class. For A9, A12, B4, B5, B6 extended and B10, the lecture gives benchmark values: reproduce them with your own calculations and interpret any differences

Data Used Today and in the Homework

- ETF (exchange-traded fund): a fund whose shares trade on an exchange like a stock; SPY holds the S&P 500 stocks, RSP the same stocks with equal weights, TLT US Treasury bonds with more than 20 years to maturity

Series	Price used	Currency	Calendar	Exercise
Standard & Poor's 500 (S&P 500)	SPY (ETF), adjusted close	USD	US trading days	B2, B4
Gold	spot gold, close	USD	weekdays only	B2
Bitcoin	BTC/USD, close	USD	daily, 24:00 Coordinated Universal Time (UTC)	B2
RSP, TLT	ETFs, adjusted close	USD	US trading days	B4
EUR/RON, official	National Bank of Romania (BNR), reference rate	RON per EUR	Romanian working days, 13:00	B1, B3
EUR/RON, market	series of EOD Historical Data (EODHD), the market-data provider, close	RON per EUR	every day with a quote	B1, B3

- Daily prices of the course data; sample 2 Jan 2015 – 18 Sep 2026
- Adjusted close: the closing price corrected for dividends and splits, so that its changes are total returns

Setup (1/2): Open the Notebook and Save a Copy

- Colab: Google's free notebook service; Python runs on Google's servers, nothing to install, a Google account is enough
- **Tasks**
  1. Open the seminar notebook in Colab.
  2. Choose *File* → *Save a copy in Drive*, or run the Drive-save cell under the banner: without a copy your changes are lost when the tab closes.
  3. Run the cells of the section *Setup* in order (Shift + Enter); they import the libraries and define the data loader.
  4. If a cell fails, choose *Runtime* → *Restart session* and run again from the top.
- **Report**: nothing; you are ready when the setup check of the next slide prints the expected numbers

Setup (2/2): Load the Data

```
# section "Setup" defines load(name, start):
# one daily series by its name,
# with the right price field
spy = load('SPY', start='2015-01-02')
bnr = load('EUR/RON', start='2015-01-01')
mkt = load('EUR/RON (market)',
           start='2015-01-01')
print(len(spy), spy.index[0].date(),
      spy.index[-1].date())
print(len(bnr), len(mkt))
```

- `load` calls `fetch_daily(symbol, source, field)`; `SERIES` in the notebook lists every name

- Expected: 2 945 SPY prices, 2 Jan 2015 to 18 Sep 2026; 2 939 BNR rates; 3 115 market quotes

date	close	adjusted
2015-01-02	205.43	169.26
2015-01-05	201.76	166.24
2015-01-06	199.82	164.64

- The two SPY columns differ: dividends are in the adjusted close only
- If your count or first value differs, stop and check the name and the start date

Prices and Returns: Definitions (1/2)

- P_t: the price on day t (here the close, or the adjusted close for ETFs)
- **Simple return** and **log return** of day t


$$R_t = \frac{P_t}{P_{t-1}} - 1, \qquad r_t = \ln \frac{P_t}{P_{t-1}} = \ln(1 + R_t)$$

  - R_t: the percentage gain of one day, what an investor earns
  - $r_t \approx R_t$ for small moves; $r_t < R_t$ for any $R_t \neq 0$, and the gap grows with the size of the move
- Over several days: simple returns **compound**, log returns **add**

$$\frac{P_n}{P_0} - 1 = \prod_{t=1}^{n}(1 + R_t) - 1, \qquad \ln \frac{P_n}{P_0} = \sum_{t=1}^{n} r_t$$

- Worked example, two days
  - $P_0 = 100$, $P_1 = 102$: $R_1 = 102/100 - 1 = 2.00\%$, $r_1 = \ln 1.02 = 1.98\%$
  - $P_2 = 96.9$: $R_2 = 96.9/102 - 1 = -5.00\%$, $r_2 = \ln 0.95 = -5.13\%$

Prices and Returns: Definitions (2/2) — Simple and Log Returns on a Chart




- Left: $r = \ln(1+R)$ against R; the gap is 9.5 percentage points (pp) at $R = +50\%$ and 19.3 pp at $R = -50\%$
- Right: a price that alternates $+20\%$ and -20% for 10 days (simulated)
  - the sum of the simple returns is 0, but the price ends at 81.5: $100 \times (1.2 \times 0.8)^{\mathbf{5}} = 81.5$
  - the sum of the log returns, $5(\ln 1.2 + \ln 0.8) = -20.4\%$, equals $\ln(81.5/100)$


A1 [Solved]: Adding Returns — Task

- **Question**: can two daily returns simply be added to get the two-day return?
- **Inputs**: prices $P_0 = 100$, $P_1 = 110$, $P_2 = 99$; no data file
- **Tasks**
  1. Compute the simple returns R_1, R_2 and the log returns r_1, r_2.
  2. Add the two simple returns and the two log returns.
  3. Compute the two-period returns $P_2/P_0 - 1$ and $\ln(P_2/P_0)$, and compare them with the sums of step 2.
- **Report**: a table with columns date, price, simple return, log return, and one sentence on which returns add up
- Notebook: section A1–A4

Back Exercise list

A1 [Solved]: The Price Path and Its Returns





MFM_ch0_seminar

A1 [Solved]: Adding Log Returns, Compounding Simple Returns


t	P_t	R_t	r_t
0	100	–	–
1	110	+10.00%	+9.53%
2	99	−10.00%	−10.54%
sum		0.00%	−1.005%
$0 \to 2$		−1.00%	−1.005%

- Interpretation: adding simple returns reports 0% for a path that lost 1%

- Log returns add up: $\ln P_2 - \ln P_1 + \ln P_1 - \ln P_0 = \ln(P_2/P_0)$, because the intermediate term $\ln P_1$ cancels
- Simple returns compound: $(1 + R_1)(1 + R_2) - 1 = 1.1 \times 0.9 - 1 = -1\%$

MFM_ch0_seminar

CAGR and Annualised Volatility: Definitions (1/3) — CAGR



- Y: the length of the sample in years

$$Y = \frac{\text{calendar days between the first and the last price}}{365.25}$$

- **Growth multiple** $P_{\text{end}}/P_{\text{start}}$ and **CAGR** (compound annual growth rate)

$$\text{CAGR} = \left(\frac{P_{\text{end}}}{P_{\text{start}}}\right)^{1/Y} - 1 = \exp\left(\frac{\ln(P_{\text{end}}/P_{\text{start}})}{Y}\right) - 1$$


  - the constant yearly growth that turns P_{start} into P_{end} (red line)
  - worked number: 100 becomes 200 in 10 years: $2^{1/10} - 1 = 7.18\%$, not $100\%/10 = 10\%$

- Chart: a simulated price over 10 years: CAGR 6.5%
  - growing at the mean simple return (8.3% a year, dashed) overshoots the final price: CAGR is geometric growth, not the average return

What Does Annualisation Mean?

- i.i.d. means independent and identically distributed: independent observations with the same distribution.
- The subscript A below means annualised; the same quantity has subscript a in the course and in exercise A7.
- The factor A is the number of returns per year: here it is estimated from each series calendar rather than assumed equal for all assets.
- For example, 252 daily changes and 365 daily changes cover one year under different calendars.

CAGR and Annualised Volatility: Definitions (2/3) — Volatility



- **Daily volatility**: the sample standard deviation of the n daily log returns

$$\hat{\sigma}_d = \sqrt{\frac{1}{n-1}\sum_{t=1}^{n}(r_t - \bar{r})^2}, \qquad \bar{r} = \frac{1}{n}\sum_{t=1}^{n} r_t$$


- **Return periods per year** $A = n/Y$: about 252 for US stocks, 260 for gold (weekdays), 365 for Bitcoin (every day)
- **Annualised volatility**

$$\hat{\sigma}_A = \hat{\sigma}_d\sqrt{A}$$

  - worked number: $\hat{\sigma}_d = 1.2\%$, $A = 252$: $1.2\% \times 15.87 = 19.05\%$

- $\sqrt{A}$, not A: for uncorrelated returns with equal variance, $\mathrm{Var}(r_1 + \cdots + r_h) = h\,\sigma_d^2$, so the s.d. of an h-day return is $\sigma_d\sqrt{h}$
  - chart: simulated sums of h daily returns lie on $\sigma_d\sqrt{h}$; one year: 15.9%, not 252%

CAGR and Annualised Volatility: Definitions (3/3) — Mean and Sharpe Ratio



- **Annualised mean** of the n daily simple returns

$$\bar{R} = \frac{1}{n}\sum_{t=1}^{n} R_t, \qquad \hat{\mu}_A = A\,\bar{R}$$

  - the mean scales with A, the standard deviation with $\sqrt{A}$
- **Sharpe ratio** (risk-free rate 0): the annualised mean per unit of annualised volatility

$$\mathrm{SR} = \frac{A\,\bar{R}}{\sqrt{A}\,s_R} = \sqrt{A}\,\frac{\bar{R}}{s_R}$$

  - s_R: the sample s.d. of the daily simple returns (divisor $n-1$)
  - reading: $\mathrm{SR} = 0.5$ means 0.5 pp of mean return per pp of volatility; a negative mean gives $\mathrm{SR} < 0$
  - chart: the slope of the line from the origin to each (hypothetical) asset; the steepest line has the best return per unit of risk

A2–A3 [Solved]: Annualisation and CAGR — Task

- **Question**: how do we turn daily numbers into annual ones?
- **A2 inputs**: daily volatility $\sigma_d = 1\%$; $A = 252$ return intervals a year for an equity index, $A = 365$ for Bitcoin
  1. Compute the annualised volatility $\sigma_d\sqrt{A}$ for both values of A.
  2. State the two assumptions under which $\sqrt{A}$ scaling is exact.
- **A3 inputs**: S&P 500 price index, multiplied by 3.72 between 2 Jan 2015 and 18 Sep 2026 (4 277 calendar days)
  3. Compute $Y = 4\,277/365.25$ and the CAGR $= 3.72^{1/Y} - 1$.
  4. Explain why this CAGR is lower than the 13.7% of SPY in B2.
- **Report**: two volatilities, one CAGR, two sentences

Back Exercise list

A2–A3 [Solved]: Annual Growth and Volatility

- A2
  1. $\sqrt{252} = 15.87$, so $1\% \times 15.87 = 15.87\%$; $\sqrt{365} = 19.10$, so 19.10%.
  2. Exact when daily returns are uncorrelated and have constant variance; A9 (Pack 3, after the lecture) shows what happens otherwise.
- A3
  3. $Y = 4\,277/365.25 = 11.71$;
     $\text{CAGR} = \exp(\ln 3.72/11.71) - 1 = \exp(1.3137/11.71) - 1 = e^{0.1122} - 1 = 11.87\%$.
  4. The S&P 500 index is a price index: it excludes dividends; SPY's adjusted close reinvests them (13.7% a year).
- Interpretation: CAGR is geometric growth per year, not the average annual return; for the same data the arithmetic mean is higher (A8, Pack 3)

Drawdown: Definitions (1/2)

- **Running peak**, **drawdown** and **maximum drawdown** (MDD)

$$M_t = \max_{s \le t} P_s, \qquad DD_t = \frac{P_t}{M_t} - 1 \le 0, \qquad \text{MDD} = \min_t DD_t$$

  - M_t: the highest price so far; DD_t: the loss of an investor who bought at that peak
  - MDD: the worst such loss in the sample; it comes with a peak date and a trough date (the day of the minimum)

- **Gain needed to recover**: after a loss $L = -DD_t$ the price is $M_t(1 - L)$


$$\frac{M_t}{M_t(1-L)} - 1 = \frac{L}{1-L}$$


  - worked number: peak 120, later price 90: $DD = -25\%$; back to the peak needs $120/90 - 1 = 0.25/0.75 = +33.3\%$

- **Underwater chart**: DD_t over time; 0 at every new peak, negative in between

- **Period under water**: from the day a peak is set to the first later close at or above it
  - a period still open at the end of the sample is **right-censored**: its true length is unknown, only a lower bound

Drawdown: Definitions (2/2) — Drawdown and Recovery on a Chart





- Left: a simulated price over 4 years; MDD -28.2% (green peak to red trough); the longest period under water lasts 321 trading days
- Right: the gain needed grows faster than the loss: -50% needs $+100\%$, -80% needs $+400\%$

A4 [Solved]: Drawdown — Task


- **Questions**
  - what is the worst loss from a previous peak?
  - when did it happen?
- **Inputs**: prices on days 0–5: $100, 120, 90, 110, 130, 100$
- **Tasks**: complete the worksheet
  1. Running peak $M_t = \max_{s \le t} P_s$.
  2. Drawdown $DD_t = P_t / M_t - 1$, reported as a negative number.
  3. Maximum drawdown $\min_t DD_t$ and its day.

Day	0	1	2	3	4	5
P_t	100	120	90	110	130	100
M_t	100	120	...	...	...	...
DD_t	0%	0%	...	...	...	...

- **Report**: the completed table and the maximum drawdown

Back Exercise list

A4 [Solved]: Drawdown from Each Running Peak



Day	0	1	2	3	4	5
P_t	100	120	90	110	130	100
M_t	100	120	120	120	130	130
DD_t	0%	0%	-25.00%	-8.33%	0%	-23.08%


- Maximum drawdown -25% on day 2; the later fall to 100 is smaller, -23.08%, because it is measured from the new peak of 130

MFM_ch0_seminar

Bad Ticks and the BNR Reference Rate (1/2)

- **Bad tick**: a recorded price at which nobody traded, for example a typing error, a wrong decimal or a quote from another instrument
- **BNR reference rate**: the official EUR/RON rate, published by the BNR every Romanian working day at about 13:00; an independent source to check a market series against
- The **EUR/RON series of the market-data provider** aggregates dealer quotes: it is not an official rate and is recorded at another time of day
- Three signs of a bad tick
  1. a one-day move far larger than usual for the series;
  2. the move reverses the next day;
  3. the quote is far from the independent source of the same day.
- A gap to BNR alone is not enough: on a day of real movement the two series differ because they are recorded at different times
- One bad tick creates two huge returns, the jump and its reversal, and inflates the volatility

Bad Ticks and the BNR Reference Rate (2/2): a Bad Tick on a Simulated Series




- Two checks of a quote P_t: against an independent reference rate B_t and against its own neighbours

$$|P_t - B_t| > 0.05 \text{ lei}, \qquad |P_t - \mathrm{med}(P_{t-\mathbf{5}}, \ldots, P_{t+\mathbf{5}})| > 0.05 \text{ lei}$$

  - med: the median of the 11 quotes centred on day t; one bad quote cannot move it
- Right: the bad price gives a jump of $+8.9\%$ and a reversal of -9.0%; annualised volatility 26.4% with it, 2.3% without it

B1 [Solved]: One Suspicious Quote — Task (1/2)



MFM_ch0_seminar

B1 [Solved]: One Suspicious Quote — Task (2/2)

- **Question**: is the market EUR/RON quote of 13 Aug 2025 a real price?
- **Data**: EUR/RON series of the market-data provider (close), 1 Jan 2015 – 18 Sep 2026, 3 115 rows; BNR reference rate
- **Tasks**
  1. Compute the daily log changes of the market series and find the largest one in absolute value.
  2. For 13 Aug 2025, report the previous quote, the quote and the next quote, and the two log changes.
  3. Compare the quote with the BNR rate of 13 Aug 2025, in RON and in percent.
  4. Compute the annualised volatility of the market series, of the same series without this quote and of the BNR rate, each with its own A.
- **Report**: the three quotes, the gap to BNR and the three volatilities
- Look first: spikes that reverse the next day are the suspects
- Notebook: section B1
- **Interpretation**: which of the three signs are present?


MFM_ch0_seminar

Back Exercise list

B1 [Solved]: Solution (1/2) — Step by Step

- Step 1: the largest daily log change is -15.41%, on 14 Aug 2025; the second largest, $+15.40\%$, is on 13 Aug 2025
- Step 2: 12 Aug 5.0636 $\rightarrow$ 13 Aug 5.9065 $\rightarrow$ 14 Aug 5.0629
  - $\ln(5.9065/5.0636) = +15.40\%$, then $\ln(5.0629/5.9065) = -15.41\%$: the move reverses the next day
- Step 3: BNR on 13 Aug 2025: 5.0633; gap $5.9065 - 5.0633 = 0.843$ RON, 16.7% above the official rate
- Step 4: annualised volatility
  - market series 12.4% ($A = 265.9$, it has some weekend quotes); without the quote of 13 Aug 2025: 10.6%
  - BNR rate 2.1% ($A = 251.1$)

B1 [Solved]: Solution (2/2) — Chart and Interpretation




- All three signs are present: a 15% move where the BNR rate moves about 0.13% a day ($2.1\%/\sqrt{251}$), reversed the next day, 0.84 RON away from BNR: a bad tick
- One quote adds 1.8 pp to the annualised volatility; the market series is still five times as volatile as the BNR rate: other bad ticks remain, the full audit is Pack 1
- Interpretation: check a market series against an official source before computing anything; for EUR/RON use the BNR rate

MFM_ch0_seminar

Growth of 1 USD and the Log Scale

- **Growth of 1 USD**: $G_t = P_t / P_0$, the value on day t of 1 USD invested on day 0; it puts assets with very different prices on one chart
- **Log scale**: the vertical axis shows $\ln G_t$ with the labels of G_t
  - equal distances are equal ratios: $1 \to 2$ and $100 \to 200$ have the same height
  - a constant growth rate is a straight line
- On a linear scale (left) Bitcoin's 257-fold growth flattens SPY into a line at the bottom



B2 [Solved]: Three Assets on One Chart — Task


- **Questions**
  - how much did 1 USD grow in SPY, gold and Bitcoin since 2015?
  - how volatile was each?
  - how deep was the worst fall?
- **Data**: SPY (adjusted close), gold (close, weekdays), Bitcoin (close, every day), 2 Jan 2015 – 18 Sep 2026, all in USD; each asset on its own calendar
- **Tasks**
  1. Plot the growth of 1 USD, P_t/P_0, on a log scale.
  2. For each asset compute the multiple, Y and the CAGR.
  3. Compute n, $A = n/Y$, the daily volatility $\hat{\sigma}_d$ and the annualised volatility $\hat{\sigma}_d\sqrt{A}$.
  4. Plot the drawdown curves and report each maximum drawdown with its peak and trough dates.
- **Report**: the two charts and a table (multiple, CAGR, A, volatility, MDD)
- **Interpretation**: does the asset with the highest CAGR also have the highest volatility and the deepest drawdown?
- Notebook: section B2

Back Exercise list

B2 [Solved]: Solution (1/4) — SPY Step by Step

- Prices: 169.26 on 2 Jan 2015, 761.69 on 18 Sep 2026; $Y = 4\,277/365.25 = 11.71$
- Step 2: multiple $761.69/169.26 = 4.50$; CAGR $= 4.50^{1/11.71} - 1 = e^{1.504/11.71} - 1 = 13.7\%$
- Step 3: $n = 2\,944$ returns, $A = 2\,944/11.71 = 251.4$; $\hat{\sigma}_d = 1.108\%$; volatility $1.108\% \times \sqrt{251.4} = 17.6\%$
- Step 4: running peak 307.63 on 19 Feb 2020, trough 203.91 on 23 Mar 2020:
  MDD $= 203.91/307.63 - 1 = -33.7\%$
- The same code runs for gold and Bitcoin: only the name changes; A follows each calendar (260.5 for gold, 365.25 for Bitcoin)


B2 [Solved]: Solution (2/4) — Growth of 1 USD



- Log scale: Bitcoin's falls of 2018 and 2022 are as visible as its rises; end labels are the growth multiples

MFM_ch0_seminar

B2 [Solved]: Solution (3/4) — Drawdowns



- Circles mark each maximum drawdown; on 18 Sep 2026 SPY is 1.8% below its peak, gold 20.6%, Bitcoin 35.2%

MFM_ch0_seminar

B2 [Solved]: Solution (4/4) — Table and Interpretation

	Multiple	CAGR	A	Volatility	MDD (peak → trough)
SPY	4.50	13.7%	251.4	17.6%	−33.7% (19 Feb 2020 → 23 Mar 2020)
Gold	3.68	11.8%	260.5	16.3%	−27.7% (28 Jan 2026 → 16 Jul 2026)
Bitcoin	256.8	60.6%	365.25	66.7%	−83.4% (16 Dec 2017 → 15 Dec 2018)

- Yes in this sample: Bitcoin has the highest CAGR, about 3.8 times the volatility of SPY and 4.1 times that of gold, and lost 83% from its 2017 peak
- SPY and gold have similar volatility; gold's worst fall is recent and not yet recovered
- One sample of 11.7 years: whether such differences are reliable is a question for the lecture and Pack 2

A5 [Proposed]: Returns, CAGR and Volatility — Task

- **Model**: A1 and A2–A3 [Solved]
- **Question**: do the rules of A1 and A2–A3 hold on new numbers?
- **Inputs**: a stock at $P_0 = 50$, $P_1 = 55$, $P_2 = 44$; a fund that grew from 100 to 250 in 2 922 calendar days; a daily volatility of 1.5%
- **Tasks**
  1. Compute R_1, R_2, r_1, r_2, the two sums and the two-day returns $P_2/P_0 - 1$ and $\ln(P_2/P_0)$.
  2. Compute Y and the CAGR of the fund.
  3. Annualise the daily volatility with $A = 252$ and with $A = 365$.
- **Report**: a small table and one sentence on which sum equals the two-day return

Back Exercise list

A6 [Proposed]: A Drawdown on Paper — Task

- **Model**: A4 [Solved]
- **Questions**
  - when was the investor furthest below the previous peak?
  - when did the price recover?
- **Inputs**: prices on days 0–5: $200, 180, 220, 165, 198, 230$
- **Tasks**
  1. Compute the running peak M_t and the drawdown DD_t for every day.
  2. Report the maximum drawdown and its day.
  3. Recovery after the trough:
     - on which day is the price back at or above the peak before the trough?
     - what percentage gain from the trough brings the price exactly back to the previous peak?
- **Report**: the completed table, the MDD and the recovery day with the gain needed

Back Exercise list

B3 [Proposed]: Two More Suspicious Quotes — Task

- **Model**: B1 [Solved]
- **Question**: the market EUR/RON quotes of 22 Apr 2022 and of 6 May 2025 are both more than 0.05 RON from the BNR rate; which one is a bad tick?
- **Data**: EUR/RON series of the market-data provider and BNR reference rate, as in B1
- **Tasks**
  1. For each date, report the previous quote, the quote and the next quote, and the two daily log changes.
  2. Compare each quote with the BNR rate of that day; if BNR published no rate that day (a public holiday), use its last earlier rate.
  3. Apply the two numeric checks: distance from BNR and from the centred 11-quote median, each above 0.05 RON. Relate them to the three signs of a bad tick, then decide: remove or keep.
  4. Plot both episodes, market series against BNR, over about three weeks.
- **Report**: one table row per date (quotes, changes, BNR, gap, signs present, decision) and the chart
- **Interpretation**: why is a gap to BNR alone not enough to remove a quote?

Back Exercise list

B4 [Proposed]: Three Other Assets — Task

- **Model**: B2 [Solved]
- **Question**: how did the S&P 500 with equal weights (RSP) and long US Treasuries (TLT) do against SPY since 2015?
- **Data**: SPY, RSP and TLT (adjusted close), 2 Jan 2015 – 18 Sep 2026, USD, US trading days
- **Tasks**
  1. Plot the growth of 1 USD of the three ETFs on a log scale.
  2. Compute the multiple, the CAGR, A and the annualised volatility of each.
  3. Plot the drawdown curves and report each MDD with its peak and trough dates and the drawdown on 18 Sep 2026.
- **Report**: the two charts and the table of B2 for the three ETFs
- **Interpretation**: are long Treasuries a safe asset in this sample?

B5 [Proposed]: Drawdowns and Recovery Time — Task

- **Model**: A4 and B2 [Solved]
- **Questions**
  - how deep are the drawdowns of the S&P 500 and of Bitcoin?
  - how long are they?
- **Data**: S&P 500 (close) from 3 Jan 2000 and Bitcoin (close) from 17 Sep 2014, both to 18 Sep 2026
- **Conventions**
  - a spell starts on the last day the running peak was set (equal peaks: the latest) and ends on the first close at or above that peak
  - a spell still open on 18 Sep 2026 is right-censored
- **Tasks**
  1. Compute DD_t for both assets and plot the two underwater curves.
  2. Report the maximum drawdown with its peak, trough and recovery dates and prices.
  3. Report the longest spell (start, end, calendar and trading days) and the current drawdown.
  4. Repeat 2–3 for the S&P 500 from 17 Sep 2014, the Bitcoin horizon.
- **Interpretation**: why must the two MDDs be compared on the same horizon?

Back Exercise list

C1 [Proposed]: Critique an AI Answer — the Answer

- **Prompt** sent to an AI assistant: "Using the daily prices of the course data, compute the annualised mean, the annualised volatility and the Sharpe ratio ($r_f = 0$) of SPY and Bitcoin from daily simple returns"
- **Answer**: the code below and the conclusion

```
import numpy as np, pandas as pd
URL = 'https://raw.githubusercontent.com/danpele/MFM/main/data/market/'
def load(sym, field='close'):
    return pd.read_csv(URL + sym + '.csv', index_col='date',
                       parse_dates=True)[field].dropna()
spy, btc = load('SPY.US'), load('BTC-USD.CC')
A = 252                                  # trading days per year
for name, p in [('SPY', spy), ('BTC', btc)]:
    r   = p.pct_change().dropna()        # daily simple returns
    mu  = r.mean() * A                   # annualised mean
    vol = r.std() * A                    # annualised volatility
    print(name, mu, vol, mu / vol)       # Sharpe ratio, rf = 0
```

- **Conclusion of the AI**: "Sharpe ratios: SPY 0.03, Bitcoin 0.07. Both assets barely reward their risk."

Back Exercise list

C1 [Proposed]: Critique an AI Answer — Design before Code

- **Model**: A2–A3 and B2 [Solved]
- **Sharpe ratio**: the definition in CAGR and Annualised Volatility: Definitions (3/3)
- **Data**: the full history, SPY from 29 Jan 1993 and Bitcoin from 13 Jul 2010, both to 18 Sep 2026
- **Step 1, research design** (write it down before you run any code)
  1. State the hypothesis behind the AI's conclusion as a testable sentence about the two Sharpe ratios.
  2. Fix the sample (dates, price field, calendar of each asset) and the comparison you will make.
  3. Name one rival explanation for any difference you will find (for example, the different sample periods).
  4. Say what the result cannot establish, whatever the numbers turn out to be.
- **Report**: four sentences, written before the code

C1 [Proposed]: Find and Log the Planted Errors

- **Step 2, the code**
  5. Find the three implementation errors planted in the code.
  6. For each, say how you would detect it: a check on the output, a published number, the lecture or the documentation.
  7. Check the corrected code on 2 Jan 2015–18 Sep 2026 against B2; then report the annualised mean, volatility and Sharpe ratio on each asset's full history. Interpret differences in light of the sample dates.
  8. Check the corrected result against Step 1:
     - is your rival explanation ruled out?
     - which limitation remains even with the code fixed?
  9. Log every error you actually found, in the format of `AI_ERRORS.md`: request, answer, error, how found, correction.
- **Interpretation**: does the conclusion "both assets barely reward their risk" survive?
- AI code usually runs without an error message: the mistakes are in the finance, not in the syntax

After the Lecture: Packs, Data and Notation

Pack	Exercises	Prerequisites
1. Data and calendars	B1 Full Audit, B2 Six Assets, B6 [all Solved]	core B1 and B2
2. Precision	A7, B2 Extended [Solved]	Pack 1, the Chapter 0 lecture
3. Return arithmetic	A8 [Solved], A9, A10	Pack 2
4. Correlation inference	B6 Extended, A11, A12, C2	Packs 1–2
5. Drawdown extremes	B5 Extended	B5, Pack 2
6. Romanian market	B7 [Solved], B8, B9, B9 Extended, B1 Extended	Pack 1, ordinary least squares (OLS)
7. Stablecoins	B10, B10 Extended	A2–A3, B7
8. Passive investing lab	C3	the lecture case study, Pack 2

- For after the Chapter 0 lecture, which introduces the standard errors, the bootstrap and the tests used here; one notebook section per pack, in this order; C2 and C3 are possible project starts
- Data (course data, daily, 2 Jan 2015 – 18 Sep 2026 unless stated)
  - SPY and TLT (adjusted close, total return)
  - S&P 500 index, gold (weekdays), Bitcoin (every day, 24:00 UTC), Euro Stoxx 50 and Nikkei 225 (close, prices converted to USD with Federal Reserve Economic Data (FRED), series `DEXUSEU`, `DEXJPUS`)
  - BNR EUR/RON and the EUR/RON series of the market-data provider
  - Bucharest Exchange Trading (BET), BET Total Return (BET-TR, dividends reinvested), Banca Transilvania, BRD
  - DefiLlama stablecoin supply
- Notation
  - P_t, R_t, r_t, n (also T), Y, $A = n/Y$, CAGR and $\hat{\sigma}_d\sqrt{A}$ as in the core
  - Sharpe ratio $\mathrm{SR} = \sqrt{A}\,\bar{R}/s_R$, defined in the core
  - iid: independent and identically distributed

Pack 1: B1 Full Audit [Solved]: All Bad Ticks

- **Task**
  - which quotes of the EUR/RON series of the market-data provider (close, 1 Jan 2015 – 18 Sep 2026, 3 115 rows) are errors?
  - how much do they distort volatility?
  - external check: the BNR reference rate

  1. Drop weekend rows.
  2. Flag weekday quotes more than 0.05 RON from the 11-day centred rolling median.
  3. Compare each flag with the BNR rate of that day (on a holiday, the last earlier fixing); remove only quotes more than 0.05 RON from BNR.
  4. Compute the annualised volatility of the raw market series, the cleaned series and the BNR rate, each with its own A.
- **Report**: an audit table (date, quote, median, BNR, gap, decision) and the three volatilities
- **Worked solution**: notebook section *B1 full audit*

Pack 1: B2 Six Assets [Solved]: A Cross-Asset Dashboard

- **Task**
  - how much did 1 USD grow since 2015 in SPY, Euro Stoxx 50, Nikkei 225, gold, TLT and Bitcoin?
  - how volatile was each?
  - period: 2 Jan 2015 – 18 Sep 2026, all in USD
  1. Convert Euro Stoxx 50 and Nikkei 225 to USD: $P^{USD} = P^{EUR} \times$ (USD per EUR), $P^{USD} = P^{JPY}/$(JPY per USD), with the FRED rate of the same day or, if missing, the last rate at most 5 calendar days earlier.
  2. Plot the growth of 1 USD, $P_t/P_{\mathbf{0}}$, on a log scale.
  3. For each asset compute n, Y, $A = n/Y$, the multiple, the CAGR and the volatility $\hat{\sigma}_d\sqrt{A}$.
  4. Recompute each volatility with $A = 252$.
- **Report**
  - the chart and a table (A, multiple, CAGR, volatility, volatility with 252)
  - for which assets does the choice of A matter?
  - why does it matter for them?
- **Worked solution**: notebook section *B2 six assets*

Pack 1: B6 [Solved]: Correlations and Calendars (1/2)

Task

- how much do the calendar and the estimator change a correlation?
- is the change of the stock–bond correlation larger than sampling error?
- data: S&P 500 and Bitcoin (close), TLT (adjusted close), daily log returns; Bitcoin – S&P 500 in 1 Jan 2022 – 18 Sep 2026; S&P 500 – TLT in 2010–2020 (n_1 days) and 2022–2026 (n_2 days)

Calendars

- join the *prices* on common days first, then compute returns (Friday → Monday for both)
- returns first and join later pairs the S&P 500 Friday → Monday return with Bitcoin's Sunday → Monday return and loses Bitcoin's weekend move
- forward-filling the S&P 500 creates zero returns paired with Bitcoin's weekend moves

1. Bitcoin – S&P 500 whole-period correlation on common days (prices joined first).
2. The same with the S&P 500 forward-filled on weekends, and with weekly returns: last common-day price of each week ending on Friday (Thursday if Friday is a holiday); an incomplete final week is dropped.
3. The mean of the 252-day rolling correlation over 2022–2026 (windows may start in 2021).
4. S&P 500 – TLT in both periods, with both estimators.

Back Exercise list

Pack 1: B6 [Solved]: Correlations and Calendars (2/2)

5. With SE denoting standard error, compute $\hat{\rho}_{\mathbf{1}}$, $\hat{\rho}_{\mathbf{2}}$ unrounded, $z_i = \operatorname{atanh} \hat{\rho}_i$, $\mathrm{SE}_z = \sqrt{1/(n_{\mathbf{1}} - 3) + 1/(n_{\mathbf{2}} - 3)}$ and $z = (z_{\mathbf{2}} - z_{\mathbf{1}})/\,\mathrm{SE}_z$ (Fisher (1915); iid pairs from a bivariate Normal distribution).
6. Compute the iid 95% interval of $\rho_{\mathbf{2}} - \rho_{\mathbf{1}}$: $(\hat{\rho}_{\mathbf{2}} - \hat{\rho}_{\mathbf{1}}) \pm 1.96\sqrt{(1 - \hat{\rho}_{\mathbf{1}}^{\mathbf{2}})^{\mathbf{2}}/(n_{\mathbf{1}} - 3) + (1 - \hat{\rho}_{\mathbf{2}}^{\mathbf{2}})^{\mathbf{2}}/(n_{\mathbf{2}} - 3)}$, where $(1 - \hat{\rho}^{\mathbf{2}})^{\mathbf{2}}/(n - 3)$ is the approximate variance of one $\hat{\rho}$.

- **Report**
  - a plot of the rolling correlations; a table method / sample size / correlation; z, its p-value, the change and its interval
  - which estimator answers "what was the correlation in 2022–2026"?
  - what can the interval not establish?
- **Worked solution**: notebook section *B6*

Pack 2: A7 [Solved]: How Precise Is a Mean Return?

- **Task**
  - SPY returned about 14% a year in 2015–2026
  - how precisely does this sample pin down the expected annual return?
  - SPY (adjusted close), daily simple returns R_t, 2 Jan 2015 – 18 Sep 2026
  - A and Y as in B2 Six Assets

  1. Compute the annualised mean $\hat{\mu}_a = A\,\bar{R}$ and the annualised volatility $\hat{\sigma}_a = s_R\sqrt{A}$.
  2. With iid returns, $\mathrm{SE}(\hat{\mu}_a) = A\,s_R/\sqrt{n} = \hat{\sigma}_a/\sqrt{Y}$ (since $n = AY$): compute it and the 95% interval $\hat{\mu}_a \pm 1.96\,\mathrm{SE}$.
  3. How many years of data would give a 95% interval of ± 2 pp with the same volatility?

- **Report**: $\hat{\mu}_a$, $\hat{\sigma}_a$, the SE, the interval and the number of years; why does sampling daily instead of monthly not make the mean more precise?

- **Worked solution**: notebook section *A7*

Pack 2: B2 Extended [Solved]: How Precise Is a Sharpe Ratio?

- **Task**
  - is SPY's Sharpe ratio precise enough to rank it above gold?
  - the six series of B2 Six Assets, daily simple returns R_t, risk-free rate 0
- **Stationary bootstrap** (Politis & Romano (1994))
  - blocks from random start days, continued with probability $1 - 1/b$ (mean block $b = 20$), wrapping around, until T days are drawn
  - each drawn day keeps the returns of all assets
  - A, Y fixed
  - `default_rng(42)`
  1. Compute $\mathrm{SR}_a = \sqrt{A}\,\bar{R}/s_R$ for SPY and the daily $\mathrm{SR}_d = \bar{R}/s_R$.
  2. Compute the asymptotic iid SE for returns following the Normal distribution (Lo (2002)), $\mathrm{SE} = \sqrt{A(1 + \mathrm{SR}_d^2/2)/T}$, and the 95% interval $\mathrm{SR}_a \pm 1.96$ SE.
  3. Bootstrap (block 20, 2 000 resamples): 95% percentile interval of SPY's SR and of the other five.
  4. On the common days of SPY and gold, bootstrap the paired difference of their Sharpe ratios (both with $A = 252$) and report its 95% interval.
- **Report**: the SPY calculation, a chart of the intervals, the difference with its interval; does B2 Six Assets rank SPY above gold?
- **Worked solution**: notebook section *B2 extended*

Pack 3: A8 [Solved]: Volatility Drag

- **Task**
  - can a positive average return go together with falling wealth?
  - daily $m = E[r]$, $a = E[R]$, $v_R = \mathrm{Var}(R)$, $v_r = \mathrm{Var}(r)$, $r = \ln(1 + R)$
  - Bitcoin close in USD, 17 Sep 2014 (457.33) to 18 Sep 2026 (80 901.46), 4 384 daily returns, 365 a year
  1. Expand $\ln(1 + R)$ to second order and show $m \approx a - (v_R + a^2)/2 \approx a - v_r/2$ for small daily returns.
  2. For Bitcoin, fill a daily-to-annual table: log mean $\times 365$, drag $v_r \times 365/2$, implied simple mean, CAGR $= e^{365m} - 1$; compare with the CAGR from the two endpoint prices.
  3. An asset has an annual arithmetic mean of 10% and a volatility of 60%: approximate its expected log growth and CAGR.
- **Report**: the expansion, the table, the two numbers of task 3; why can the typical path lose money when the mean return is positive?
- **Worked solution**: notebook section *A8*

Back Exercise list

Pack 3: A9 [Proposed]: When the $\sqrt{A}$ Rule Fails

Task

- daily log returns, stationary, variance σ^2, autocorrelations ρ_j
- data: BNR EUR/RON, 4 Jul 2005 – 18 Sep 2026, 5 352 daily log changes (models: A2–A3, A8)

1. For $k = 3$, write $\mathrm{Var}(r_1 + r_2 + r_3)$ as the sum of the nine covariances; then generalise to $\mathrm{Var}(\sum_{t=1}^{k} r_t) = k\sigma^2[1 + 2\sum_{j=1}^{k-1}(1 - j/k)\rho_j]$.
2. For a *hypothetical* AR(1), $\rho_j = \rho^j$, find the limit of the bracket as $k \to \infty$.
3. Compute $\hat{\rho}_1$, the $\sqrt{252}$ volatility and its hypothetical AR(1) correction.
4. Compute the autocorrelations at lags 1–10 and the volatility from month-end changes $\times\sqrt{12}$.
5. The EUR/RON series of the market-data provider (B1) has $\hat{\rho}_1 = -0.57$:
   - what would the AR(1) factor be?
   - why is it the wrong fix?

Report

- the two derivations, the volatilities and the autocorrelations
- does positive autocorrelation hide risk or invent it?
- is $\hat{\rho}_1$ enough?

Back Exercise list

Packs 3–4: A10 [Proposed]: The Standard Error of a Sharpe Ratio

- **Task**
  - T iid daily returns with mean μ, variance σ^2, finite central moments μ_3, μ_4, skewness $\gamma_3 = \mu_3/\sigma^3$, ordinary (not excess) kurtosis $\gamma_4 = \mu_4/\sigma^4$, daily $\mathrm{SR} = \mu/\sigma$
  - SPY: $\mathrm{SR}_a = 0.82$, $A = 251.4$, $T = 2\,944$, $Y = 11.71$, $\gamma_3 = -0.31$, $\gamma_4 = 17.1$ (models: A8, A7, B2 Extended)
  1. Gradient: for $g(\mu, \sigma^2) = \mu/\sigma$, compute ∇g.
  2. Quadratic form: $\sqrt{T}(\hat{\mu} - \mu, \hat{\sigma}^2 - \sigma^2) \to N(0, \Sigma)$, $\Sigma = \begin{pmatrix} \sigma^2 & \mu_3 \\ \mu_3 & \mu_4 - \sigma^4 \end{pmatrix}$; compute $\nabla g^\top \Sigma \nabla g$ and its Normal case (Lo (2002)).
  3. Normal case, annual scaling: with $\mathrm{SR}_a = \sqrt{A}\,\mathrm{SR}_d$ and $T = AY$, show $\mathrm{Var}(\widehat{\mathrm{SR}}_a) \approx (1 + \mathrm{SR}_a^2/(2A))/Y$ (large-T approximation, not an exact finite-sample result).
  4. SPY's asymptotic SE three ways: Normal daily, general daily (with γ_3, γ_4), and the Normal formula with $A = 1$, $\sqrt{(1 + \mathrm{SR}_a^2/2)/Y}$, as if the span were observed as annual returns; state the sampling assumption of each.
- **Report**: the derivations and the three SEs; why does sampling more often than monthly barely help?

Packs 3–4: B6 Extended [Proposed]: Does the Stock–Bond Change Survive Dependence?

- **Task**
  - S&P 500 – TLT daily log returns on common days, as in B6: 2010–2020 ($n_1 = 2\,769$) and 2022 – 18 Sep 2026 ($n_2 = 1\,182$)
  - bootstrap of B2 Extended

1. Stationary bootstrap each period separately (block 20, 2 000 resamples, `default_rng(42)`): recompute $\hat{\rho}_2^* - \hat{\rho}_1^*$ on each pair of resamples.
2. Report the 95% percentile interval of $\rho_2 - \rho_1$ and plot the histogram of the bootstrap changes with both intervals.
3. Compare its width with the width of the iid interval of B6 (task 6).

- **Report**
  - the bootstrap interval, the width ratio and the histogram
  - why is the bootstrap interval wider?
  - does the conclusion of B6 survive?

Pack 4: A11 [Proposed]: Why Is the Weekly Correlation Lower?

- **Task**
  - x_t = S&P 500, y_t = Bitcoin, daily log returns on common days, 2022 – 18 Sep 2026, $n = 1\,182$
  - $\gamma(j) = \mathrm{Cov}(x_t, y_{t-j})$
  - σ_x, σ_y: daily standard deviations
  - $\rho_{x,j}$: autocorrelation of x at lag j
  - $\rho_{\mathbf{1}}$, $\rho_{\mathbf{5}}$: correlation of daily and of 5-day returns (model: B6)

  1. For sums X, Y of 5 consecutive common-day returns, draw the 5×5 grid of covariances and show $\mathrm{Cov}(X, Y) = 5\gamma(0) + \sum_{j=1}^{4}(5-j)[\gamma(j) + \gamma(-j)]$ and $\mathrm{Var}(X) = 5\sigma_x^2 f_x$, $f_x = 1 + 2\sum_{j=1}^{4}(1 - j/5)\rho_{x,j}$; hence $\rho_{\mathbf{5}} = \{\rho_{\mathbf{1}} + \sum_{j=1}^{4}(5-j)[\gamma(j) + \gamma(-j)]/(5\sigma_x\sigma_y)\}/\sqrt{f_x f_y}$.
  2. Check the shortcut $f_x = f_y = 1$: own autocorrelations of x and y at lags 1–4, and f_x, f_y.
  3. Compute the lag term, the implied $\rho_{\mathbf{5}}$ with and without $\sqrt{f_x f_y}$, and the correlation of overlapping 5-day sums.
  4. Compute the weekly correlation for weeks anchored on Monday, . . . , Friday, each with a Fisher 95% interval.
  5. Report the joint week-block bootstrap interval of daily minus weekly (notebook section B6 extended).

- **Report**: the derivation and the numbers of tasks 2–5; is the gap 0.42 vs 0.30 a property of the markets or of the sampling? The Epps (1979) effect predicts the opposite sign

Back Exercise list

Pack 4: A12 [Proposed]: Asynchronous Closes

- **Task**
  - log prices $X(s) = W(s) + U(s)$ (Bitcoin), $Y(s) = W(s) + V(s)$ (S&P 500), W, U, V independent Brownian motions with variance rates $\sigma_W^2, \sigma_U^2, \sigma_V^2$, $\rho = \sigma_W^2 / \sqrt{(\sigma_W^2 + \sigma_U^2)(\sigma_W^2 + \sigma_V^2)}$
  - X closes at integer times t (24:00 UTC), Y a fraction δ of a day earlier (20:00 UTC in summer, 21:00 in winter)
  1. Draw the intervals covered by x_t, y_t and $y_{t+\mathbf{1}}$ on a time line and show $\mathrm{Corr}(x_t, y_t) = \rho(1-\delta)$, $\mathrm{Corr}(x_t, y_{t+\mathbf{1}}) = \rho\delta$, all other cross-correlations 0.
  2. Deduce that the lag -1, 0, $+1$ correlations sum to ρ (Scholes & Williams (1977)).
  3. Compute δ in summer and winter and the implied correlations for $\rho = 0.45$.
  4. Tabulate the observed $\mathrm{Corr}(x_t, y_{t+j})$, $j = -1, 0, 1$, for 2022–2026 and compare.
- **Report**: the derivation and the two tables; is the asynchronous-close bias important for this pair? Treat the sum as a diagnostic

Packs 4–5: C2 [Proposed]: Did Spot ETFs Change Bitcoin's Correlation with Equities?

- **Task**
  - US spot Bitcoin ETFs started trading on 11 Jan 2024
  - did the Bitcoin – S&P 500 correlation change at that date?
  - daily log returns on common days, 2018 – 18 Sep 2026 (models: B6, B2 Extended)

  1. Correlations in 2018–2019, 2020–2021, 1 Jan 2022 – 10 Jan 2024 (pre) and 11 Jan 2024 – 18 Sep 2026 (post), each with a stationary-bootstrap 95% interval (block 20).
  2. Test $H_0 : \rho_{post} - \rho_{pre} = 0$ with a bootstrap interval of the difference; plot the rolling correlation with the ETF date.
  3. Project extensions (optional): a counterfactual asset without an ETF event (state why it is comparable), an unknown-date break search, other calendars and windows.

- **Report**: a table period / n / $\hat{\rho}$ / interval, the difference, one figure, one limitation; a causal claim needs the extensions

Packs 4–5: B5 Extended [Proposed]: Is the 2009 Drawdown Extreme?

Task

- S&P 500 daily log returns r_t, 3 Jan 2000 – 18 Sep 2026, $n = 6\,717$
- your observed MDD and longest spell from B5, in trading days (open spells censored at the last day)

1. Build one path: resample n returns, set $\ln P_t = \ln P_{\mathbf{0}} + \sum_{s \le t} r_s^*$ (r_s^*: the resampled returns), compute its MDD and longest spell; plot it.
2. Repeat 5 000 times with the iid bootstrap and 5 000 times with the stationary bootstrap (block 20); seed 42.
3. Plot both distributions of the MDD and of the longest spell with the observed values; report the medians, the adverse-tail quantiles (5% for the MDD, 95% for the spell) and the tail shares $P(\mathrm{MDD}^* \le \mathrm{MDD}_{obs})$, $P(\text{spell}^* \ge \text{spell}_{obs})$.
4. Compute $\hat{\rho}_{\mathbf{1}}$ and the variance ratio (VR), $\widehat{\mathrm{VR}}(20) = \widehat{\mathrm{Var}}(r_t + \cdots + r_{t-\mathbf{19}})/(20\,\widehat{\mathrm{Var}}(r_t))$ from all overlapping 20-day sums (divisor: number of terms -1, no small-sample correction).

Report

- the charts and the numbers of task 3, $\hat{\rho}_{\mathbf{1}}$ and $\widehat{\mathrm{VR}}(20)$
- is the observed MDD extreme for this return distribution?
- The resampled returns include the crisis itself: a diagnostic, not a forecast

Abnormal Returns and Their Cumulative Sum

- **AR (event study)**: Abnormal Return.
- **CAR**: Cumulative Abnormal Return.
- If the observed return is 2% and the benchmark predicts 0.5%, the abnormal return is 1.5 percentage points.
- CAR sums these deviations over the event window. Deviations of 1.5 and -0.5 percentage points give a CAR of 1 percentage point.
- CAR measures the cumulative departure from the benchmark; its interpretation depends on the model and event window.

Pack 6: B7 [Solved]: Event Study, the Bank Tax of December 2018

- **Task**
  - how much did Banca Transilvania lose on the bank-tax news (announced on the evening of 18 Dec 2018
  - day 0 = 19 Dec 2018), beyond the market move?
  - TLV (adjusted close) and BET (close), daily log returns
  - estimation window $[-250, -11]$ (18 Dec 2017 – 4 Dec 2018, $T = 240$), event window $[0, +5]$ ($L = 6$)
  - MacKinlay (1997)
  - errors iid Normal, independent of the market return, with the same variance σ_ε^2 in both windows and independent across windows
  1. Estimate $r_{i,t} = a + b\, r_{m,t} + \varepsilon_t$ by OLS: $\hat{a}$, $\hat{b}$, $\hat{\sigma}_\varepsilon$, $\bar{r}_m$, $S_{mm} = \sum (r_{m,t} - \bar{r}_m)^2$.
  2. $AR_\tau = r_{i,\tau} - (\hat{a} + \hat{b}\, r_{m,\tau})$; table day, r_i, r_m, prediction, AR; $CAR[0,+5] = \sum_{\tau=0}^{5} AR_\tau$.
  3. Test $AR_0 = 0$ and $CAR = 0$ with the prediction-error variances $\hat{\sigma}_\varepsilon^2[1 + 1/T + (r_{m,0} - \bar{r}_m)^2 / S_{mm}]$ and $\hat{\sigma}_\varepsilon^2[L + L^2/T + (\sum_\tau r_{m,\tau} - L\bar{r}_m)^2 / S_{mm}]$; Student t with $T - 2$ degrees of freedom under the assumptions above.
- **Report**: the estimates, the AR table, CAR, t and p; BET contains both big banks: what does this do to $\hat{b}$ and to the AR?
- **Worked solution**: notebook section *B7*

Pack 6: B8 [Proposed]: The Same Event for BRD

- **Task**
  - the event of B7 for BRD (adjusted close) against the BET index, same windows
  - reuse the B7 function, only the symbol changes

  1. Report $\hat{b}$, $\hat{\sigma}_\varepsilon$, the AR table and $CAR[0,+5]$ with t and p.
  2. Build a two-bank table: $AR_{\mathbf{0}}$, its prediction-error t, CAR and its t.
  3. Plot the two CAR paths with ± 1.96 SE bands on one chart.
  4. Which bank was hit harder relative to its normal volatility? Use the prediction-error t of $AR_{\mathbf{0}}$.

- **Report**: the two tables and the chart; the reversal can mean overreaction or a partly withdrawn policy: what dated evidence would separate the two?

Back Exercise list

Pack 6: B9 [Proposed]: BET-TR in RON and in EUR

- **Data**: the BET-TR index (close) and BNR EUR/RON S_t (RON per EUR), prices joined on common days, 5 Jan 2015 – 18 Sep 2026 (model: the conversion of B2 Six Assets)
  1. Convert: $P_t^{EUR} = P_t^{RON}/S_t$; work one date step by step (13 Aug 2025).
  2. Show $r_t^{EUR} = r_t^{RON} - \Delta s_t$, with $s_t = \ln S_t$.
  3. Compute the CAGR and the volatility (observed A) in RON and in EUR, and the correlation of r^{RON} with Δs.
  4. Plot both wealth paths.
- **Report**: the two CAGRs, two volatilities, the correlation, the chart; for a euro investor, is the leu's depreciation a return drag, a risk, or both?

Back Exercise list

Pack 6: B9 Extended [Proposed]: Is the Currency Effect Significant?

- **Data**: the data of B9, $n = 2\,925$ common-day returns, $A = n/Y$ from the joined prices of B9 (models: B2 Extended, B7)
  1. Show $\mathrm{Var}(r^{EUR}) = \mathrm{Var}(r^{RON}) + \mathrm{Var}(\Delta s) - 2\,\mathrm{Cov}(r^{RON}, \Delta s)$; report each term annualised ($\times A$) in decimal units, then take square roots.
  2. Regress r_t^{RON} on a constant and Δs_t with Newey–West HAC (heteroskedasticity and autocorrelation consistent) standard errors (Newey & West (1987), $L = 8$ lags): slope, t, p; add a stationary-bootstrap 95% interval for the correlation.
  3. Estimate the annual log drift of the leu, $A\times$ mean of Δs_t, with its HAC SE, t and 95% interval.
  4. Show that $\ln(1 + \mathrm{CAGR}^{RON}) - \ln(1 + \mathrm{CAGR}^{EUR}) = \ln(S_{end}/S_{start})/Y$, the drift of task 3.
- **Report**: the variance terms, the regression, the drift with its interval; the hypothesis of task 3 is "zero log drift": is the gap between your two CAGRs of B9 distinguishable from zero?

Pack 6: B1 Extended [Proposed]: The Standard Hampel Filter

Task

- standard Hampel filter, Pearson, Neuvo, Astola & Gabbouj (2016), Section 2, eqs. (1)–(4): window $x_{k-K}, \ldots, x_{k+K}$ with $K = 5$ (11 quotes, as in their Sections 5–6)
- m_k = window median, $S_k = 1.4826 \times \text{median}_j\, |x_{k-j} - m_k|$
- x_k is flagged if $|x_k - m_k| > t\, S_k$
- the first and last 5 quotes are not tested
- data: BNR EUR/RON 2015–2026 (clean reference) and the weekday market series of B1 Full Audit

1. Work the window around 13 Aug 2025 step by step: m_k, median absolute deviation (MAD), S_k and $|x_k - m_k|/S_k$; for which t is the quote flagged?
2. On BNR: share of implosion windows ($S_k = 0$, their Section 3) and share of days flagged on their grid $t = 0, 0.5, \ldots, 21$ (Section 5.2), with a plot; compare the bound of their eq. (8) with the largest $|x_k - m_k|/S_k$.
3. On the market series, apply $t = 2$ (their Section 6) and $t = 5$ (their Section 5.2): flags, overlap with the 11 flags and 9 errors of B1 Full Audit, volatility after removing the flags.

Report

- the worked window, the two shares with the plot, the flags and the volatility
- why MAD, and why 1.4826?
- what happens in an implosion window?
- can a threshold replace the BNR check of B1 Full Audit?

Pack 7: B10 [Proposed]: Stablecoin Growth

- **Data**: DefiLlama, daily total circulating value of USD-pegged stablecoins, in bn USD (divided by 10^9), 31 Dec 2019 – 18 Sep 2026 (models: A2–A3, B2)
  1. Extract year-end values 2019–2025 and the calendar-year growth; report 2026 separately as year-to-date growth to 18 Sep.
  2. Find the 2022 peak and the following trough up to the end of 2023, with dates, and the peak-to-trough change.
  3. Compute the CAGR from 1 Jan 2020 to 18 Sep 2026 with the exact endpoints.
  4. Plot the series with the year-end points.
- **Report**: the year-end table, the peak, trough and change, the CAGR, the chart; supply is a USD value in circulation, not a price: can it fall without any token being burned?

Pack 7: B10 Extended [Proposed]: Did the Terra Collapse Change the Growth Regime?

Task

- daily log growth $g_t = \Delta \ln(\text{supply}_t)$, 2 Jan 2020 – 18 Sep 2026, $n = 2\,452$
- Terra collapse on 9 May 2022 (model: the OLS regression and t test of B7)

1. Known date (Chow (1960)): $g_t = a + b\,D_t + e_t$, $D_t = 1\{t \geq 9 \text{ May } 2022\}$, HAC SE (Newey & West (1987), $L = 8$ lags); a: daily growth before the date, b: its change after; report a, b (annualised $\times 365$) and t.
2. Unknown date (Andrews (1993)): the HAC Wald statistic $(\hat{b}/\,\mathrm{SE}_{\mathrm{HAC}}(\hat{b}))^2$ for every candidate date in the central 70% (3 Jan 2021 – 15 Sep 2025); compare its maximum with 8.85, the 5% critical value for this trimming.
3. Least-squares date: the date minimising the sum of squared residuals; plot the Wald and SSR profiles.
4. Check the residual variance before and after the date, and say whether the break-date interval of Bai (1997), which assumes the same error distribution in both regimes, can be reported.

Report

- a, b, t, the two dates and the profiles
- did the slowdown start with Terra?
- is the break in the level or in the drift of supply?

Pack 8: C3 [Proposed]: Passive Investing and Demand Elasticity (1/2)

Task

- is the one-third pass-through of Haddad, Huebner & Loualiche (2025) (the lecture case study) robust to who turns passive?
- stock k, active investor i
- w_{ik} = shares held by i as a fraction of all shares of k
- active share $\alpha_k = \sum_i w_{ik}$
- own elasticity $\bar{\mathcal{E}}_{ik}$
- χ: strength of the strategic response to the elasticity of the other investors in the stock (Table 2, row 1: $\chi = 2.97$)
- aggregate elasticity (footnote 53) $\mathcal{E}_{agg,k} = \sum_i w_{ik}\, \bar{\mathcal{E}}_{ik}/(1 + \chi\alpha_k)$

1. Two investors, $w_{\mathbf{1}} = 0.4$ with $\bar{\mathcal{E}}_{\mathbf{1}} = 1$, $w_{\mathbf{2}} = 0.3$ with $\bar{\mathcal{E}}_{\mathbf{2}} = 3$, $\chi = 2.97$; α falls from 0.70 to 0.60 (0.1 of all shares turns passive): compute $\mathcal{E}_{agg}$ before, and after the switch when both reduce in proportion, when investor 2 (elastic) switches, when investor 1 switches.
2. Compare each $\Delta \log \mathcal{E}_{agg}/\Delta \log \alpha$ with $1/(1 + \chi\alpha) = 0.325$ at the initial $\alpha = 0.70$ (a hypothetical example, distinct from the paper's time average 0.68).
3. FRED `BOGZ1LM563064100Q` (ETF corporate equities) over `BOGZ1LM893064105Q` (all sectors), 2001 Q1 – 2026 Q2: the ETF share in 2001 Q1, 2020 Q4, 2026 Q2 and the implied changes with 81% active and pass-through 0.33. Explain why the ETF share is not the paper's passive share: ETFs include active funds and omit index mutual funds.

Back Exercise list

Pack 8: C3 [Proposed]: Passive Investing and Demand Elasticity (2/2)

4. Simulation (starter code in the notebook, `numpy.random.default_rng(seed)`): 2 000 stocks, $\alpha_k \sim U(0.55, 0.85)$; 40 active investors per stock with $w_{ik} = \alpha_k \times$ weights drawn from the Dirichlet$(1, \ldots, 1)$ distribution (uniform on the 40-dimensional simplex); $\mathcal{E}_{ik}$ log-Normal with median 1.6 and log standard deviation 0.6; $\chi = 2.97$.
5. In each stock a fraction $f_k \sim U(0, 0.2)$ of α_k turns passive under three rules: all in proportion; most elastic first; least elastic first (whole holdings in order, the last one partly, until $f_k \alpha_k$ is reached).
6. Regress $\Delta \log \mathcal{E}_{agg,k}$ on a constant and $\Delta \log \alpha_k$ (Table 4, column 1).
7. Repeat for seeds 1–10 with the same draws for all rules, and for $\chi = 0$.

Report

- the ETF shares and implied changes; a table of slopes (mean, min, max over seeds; simulation variability, not a sampling confidence interval) next to the mean of $1/(1 + \chi\alpha_k)$
- why is the slope one-third only for representative switchers?
- how should the paper's 11% be read if the most elastic investors turned passive first?

References (1/2)

- Andrews, D.W.K. (1993). Tests for parameter instability and structural change with unknown change point. *Econometrica*, 61(4), 821–856.
- Bai, J. (1997). Estimation of a change point in multiple regression models. *Review of Economics and Statistics*, 79(4), 551–563.
- Boehmer, E., Musumeci, J., Poulsen, A.B. (1991). Event-study methodology under conditions of event-induced variance. *Journal of Financial Economics*, 30(2), 253–272.
- Campbell, J.Y., Lo, A.W., MacKinlay, A.C. (1997). *The Econometrics of Financial Markets*, Ch. 4 of the book, Event-Study Analysis. Princeton University Press.
- Chow, G.C. (1960). Tests of equality between sets of coefficients in two linear regressions. *Econometrica*, 28(3), 591–605.
- Epps, T.W. (1979). Comovements in stock prices in the very short run. *Journal of the American Statistical Association*, 74, 291–298.
- Fisher, R.A. (1915). Frequency distribution of the values of the correlation coefficient in samples from an indefinitely large population. *Biometrika*, 10(4), 507–521.
- Forbes, K.J., Rigobon, R. (2002). No contagion, only interdependence: measuring stock market comovements. *Journal of Finance*, 57(5), 2223–2261.
- Franke, J., Härdle, W.K., Hafner, C.M. (2019). *Statistics of Financial Markets: An Introduction* (5th ed.). Springer.
- Haddad, V., Huebner, P., Loualiche, E. (2025). How competitive is the stock market? Theory, evidence from portfolios, and implications for the rise of passive investing. *American Economic Review*, 115(3), 975–1018.

References (2/2)

- Härdle, W.K., Harvey, C.R., Reule, R.C.G. (2020). Understanding cryptocurrencies. *Journal of Financial Econometrics*, 18(2), 181–208.
- Koijen, R.S.J., Yogo, M. (2019). A demand system approach to asset pricing. *Journal of Political Economy*, 127(4), 1475–1515.
- Lo, A.W. (2002). The statistics of Sharpe ratios. *Financial Analysts Journal*, 58(4), 36–52.
- MacKinlay, A.C. (1997). Event studies in economics and finance. *Journal of Economic Literature*, 35(1), 13–39.
- Newey, W.K., West, K.D. (1987). A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica*, 55(3), 703–708.
- Opdyke, J.D. (2007). Comparing Sharpe ratios: so where are the p-values? *Journal of Asset Management*, 8(5), 308–336.
- Pearson, R.K., Neuvo, Y., Astola, J., Gabbouj, M. (2016). Generalized Hampel filters. *EURASIP Journal on Advances in Signal Processing*, 2016, article 87.
- Politis, D.N., Romano, J.P. (1994). The stationary bootstrap. *Journal of the American Statistical Association*, 89(428), 1303–1313.
- Scholes, M., Williams, J. (1977). Estimating betas from nonsynchronous data. *Journal of Financial Economics*, 5(3), 309–327.
- Public data: FRED, BNR reference rates, DefiLlama
- Code and data: github.com/danpele/MFM

Acronyms used in this material

- **ACF**: AutoCorrelation Function
- **AI**: Artificial Intelligence
- **AR**: AutoRegressive (model); in event studies: Abnormal Return
- **BET**: Bucharest Exchange Trading (index)
- **BET-TR**: BET Total Return (index)
- **BNR**: Banca Națională a României — *the National Bank of Romania*
- **CAGR**: Compound Annual Growth Rate
- **CAR**: Cumulative Abnormal Return
- **ETF**: Exchange-Traded Fund
- **FRED**: Federal Reserve Economic Data
- **FX**: Foreign eXchange
- **HAC**: Heteroskedasticity and Autocorrelation Consistent (standard errors)
- **MAD**: Median Absolute Deviation
- **MDD**: Maximum DrawDown
- **OLS**: Ordinary Least Squares
- **SE**: Standard Error
- **SR**: Sharpe Ratio
- **SSR**: Sum of Squared Residuals
- **US**: United States
- **UTC**: Coordinated Universal Time (Temps Universel Coordonné)
- **VR**: Variance Ratio