



Modelling Financial Markets

Seminar 1: Stylised Facts and Returns — Hands-on



Daniel Traian PELE
Antoaneta Amza

Bucharest University of Economic Studies
IDA Institute Digital Assets
Blockchain Research Center
AI4EFin Artificial Intelligence for Energy Finance
Romanian Academy, Institute for Economic Forecasting
MSCA Digital Finance

Acronyms used in this material (1/2)

- ▣ ACF: AutoCorrelation Function
- ▣ AI: Artificial Intelligence
- ▣ AR: AutoRegressive (model); in event studies: Abnormal Return
- ▣ ARCH-LM: ARCH Lagrange Multiplier (test)
- ▣ BET-TR: BET Total Return (index)
- ▣ BNR: Banca Națională a României (the National Bank of Romania)
- ▣ BVB: Bursa de Valori București (the Bucharest Stock Exchange)
- ▣ CAGR: Compound Annual Growth Rate
- ▣ CI: Confidence Interval
- ▣ COVID-19: COronaVirus Disease 2019
- ▣ CUSUM: CUmulative SUM (of squares)
- ▣ EGARCH: Exponential GARCH
- ▣ EODHD: EOD Historical Data (market-data provider)
- ▣ ETF: Exchange-Traded Fund
- ▣ FIGARCH: Fractionally Integrated GARCH
- ▣ FTSE: Financial Times Stock Exchange (FTSE Russell, index provider)
- ▣ GARCH: Generalised AutoRegressive Conditional Heteroskedasticity
- ▣ GJR-GARCH: Glosten–Jagannathan–Runkle GARCH
- ▣ HAC: Heteroskedasticity and Autocorrelation Consistent (standard errors)
- ▣ HAR: Heterogeneous AutoRegressive (model)
- ▣ JB: Jarque–Bera (test)
- ▣ MDD: Maximum DrawDown
- ▣ OHLC: Open, High, Low, Close
- ▣ OLS: Ordinary Least Squares
- ▣ QQ: Quantile–Quantile (plot)
- ▣ RMSE: Root Mean Squared Error

Acronyms used in this material (2/2)

- **SE: Standard Error**
- **US: United States**
- **UTC: Coordinated Universal Time (Temps Universel Coordonné)**
- **VIX: CBOE Volatility Index**

Today's Question and Required Outputs

- Question of the seminar: **which stylised facts of Lecture 1 survive inference that allows for heavy tails and volatility clustering?**
- Route
 - ▶ returns and moments: A1, B1, A3, B2
 - ▶ tails and dependence: A6, B3, A10, B9, A7, B11, B4
 - ▶ drawdowns and an AI answer to correct: A4(a), C2
- Required outputs (checklist on the last core slide): the A answers, one figure and one table per B exercise, the corrected C2 code, one interpretation sentence per exercise
- Notebook for today: [open the seminar notebook in Colab](#); each task names its notebook section
- Seminar: Antoaneta Amza, PhD candidate, antoaneta.amza@csie.ase.ro

Exercise Map

Exercise	Question	Type	Model
A1	how do returns add up over time?	Solved	–
B1	how heavy are the tails, and how fragile is the kurtosis?	Solved	–
A3	from a statistic to a p-value: JB and Ljung–Box	Solved	–
B2	how large is the VaR 1% (Value-at-Risk) of BET-TR?	Proposed	B1
A6	why does random volatility create heavy tails?	Solved	–
B3	are Bitcoin returns independent?	Solved	–
A10	why does clustering widen the autocorrelation band?	Proposed	A6
B9	which autocorrelations survive robust inference?	Solved	–
A7	the Hill estimator step by step	Solved	–
B11	are losses heavier than gains?	Solved	–
B4	do returns approach the Normal distribution with the horizon?	Proposed	B1, B3
A4(a)	how deep is a drawdown?	Proposed	A1
C2	what is wrong in an AI answer?	Proposed	A3, B3, B9
Extension packs	A1(c), A2, A4(b–e), A5, A8, A9, A11, B1(d), B5–B8, B10, B12–B15, C1, C3 (optional)	Solved or Proposed (marked on each)	named on each task

- ▣ **[Solved]**: full solution in the slides and in the notebook, a model to copy; **[Proposed]**: you solve it, the solution is discussed in class
- ▣ Part A: derivations on paper; Part B: verification and inference on real data, each ending with an interpretation question; Part C: open questions and AI critique

Data Used

Series	Price used	Period	Calendar	Returns
BET-TR	daily close	23 Sep 2014 – 18 Sep 2026	BVB trading days	2 992
Bitcoin	daily close (USD)	17 Sep 2014 – 18 Sep 2026	every day, 24:00 UTC	4 384
EUR/RON	BNR reference rate	4 Jul 2005 – 18 Sep 2026	Romanian working days	5 352
S&P 500	daily OHLC	2 Jan 1990 – 18 Sep 2026	US trading days	9 245
SPY ETF	OHLC, adjusted close (for dividends and splits)	2 Jan 2008 – 18 Sep 2026 (B8)	US trading days	–
VIX, 9-day VIX	daily close	C3 only	US trading days	–

- Daily prices of the course data, one series per asset (field used: close / adjusted close)
- EUR/RON is the official BNR reference rate, not the EUR/RON series from EODHD (market quotes)
- Units: returns in decimals in the code, reported in % on the slides; each series on its own calendar; multi-series analyses first align the *prices* on common days

Setup: Run and Verify

```
# notebook section "Data" defines load_data(name)
closes = {a: load_data(a)['Close'] for a in
           ['sp500', 'bettr', 'btc', 'eurron']}
rets = {a: np.log(c).diff().dropna()
        for a, c in closes.items()}
r_bet, r_btc = rets['bettr'], rets['btc']
r_eur, r_sp = rets['eurron'], rets['sp500']
for a, r in rets.items():
    print(a, r.index[0].date(), len(r))
```

- Expected output: first return date and count


Series	first return	T
S&P 500	1990-01-03	9 245
BET-TR	2014-09-24	2 992
Bitcoin	2014-09-18	4 384
EUR/RON	2005-07-05	5 352

- Moments: `scipy.stats.skew` and `kurtosis` (excess kurtosis, no small-sample correction); standard deviations with `ddof=1`; quantiles with `numpy.quantile` (linear)
- If a count differs, stop and check the series, the price type and the start date

Notation Used in Every Exercise

- ▣ P_t : price; simple return $R_t = P_t/P_{t-1} - 1$; log return $r_t = \ln(P_t/P_{t-1})$; T : number of returns
- ▣ $\bar{r}$, $\hat{\sigma}$: sample mean and standard deviation; $z_t = (r_t - \bar{r})/\hat{\sigma}$; $x_t = r_t - \bar{r}$
- ▣ **Skewness** $\hat{S} = \frac{1}{T} \sum_t z_t^3$; **kurtosis** $\hat{K} = \frac{1}{T} \sum_t z_t^4$; **excess kurtosis** $\hat{K} - 3$ (0 for the Normal distribution)
- ▣ **Autocorrelation** at lag k : $\hat{\rho}_k = \sum_{t=k+1}^T x_t x_{t-k} / \sum_{t=1}^T x_t^2$
- ▣ **Annualisation** with P return intervals a year: 252 (BET-TR, S&P 500, EUR/RON), 365 (Bitcoin); volatility $\sqrt{P} \hat{\sigma}$
- ▣ **p-value**: probability, under H_0 , of a statistic at least as extreme as the one observed; it is not the probability that H_0 is true
- ▣ One worked line: $P_0 = 100$, $P_1 = 102$: $R_1 = 2.00\%$, $r_1 = \ln 1.02 = 1.98\%$

First Look: Prices, Returns and Dated Extremes



- Every extreme day is a dated event: the bank-asset tax (19 Dec 2018), the COVID-19 crash (March 2020), the 2008 depreciation of the leu
- Calm and turbulent periods alternate in all three series
- EUR/RON is almost flat after 2013
- Look first, test later


A1 [Solved]: Returns and Annualisation — Task

- **Question:** which return adds up over time, and how do daily numbers become annual ones?
- **Inputs:** a share closes at 100, 110, 99, 104.5 on days 0–3 and pays no dividend; an index has a daily mean log return of 0.04% and a daily volatility of 1.2%; Bitcoin has a daily volatility of 3.5%; returns are serially uncorrelated with constant variance
- **Tasks**
 1. Compute the three simple returns R_t and the three log returns r_t , all in %.
 2. Compute the 3-day returns $P_3/P_0 - 1$ and $\ln(P_3/P_0)$ and compare them with $\sum_t R_t$ and $\sum_t r_t$.
 3. Annualise the index mean log return and volatility with $P = 252$.
 4. Annualise the Bitcoin volatility and justify the choice of P .
- **Report:** a table day, price, R_t , r_t and the three annual numbers
- **Notebook:** section A1

A1 [Solved]: Solution

t	P_t	R_t	r_t
0	100	—	—
1	110	+10.00%	+9.53%
2	99	-10.00%	-10.54%
3	104.5	+5.56%	+5.41%
sum		5.56%	4.40%
0 $\rightarrow$ 3		4.50%	4.40%

- Log returns telescope to $\ln(104.5/100)$; simple returns compound: $1.1 \times 0.9 \times 1.0556 - 1 = 4.5\%$



- Index: annualised mean log return $252 \times 0.04\% = 10.08\%$; volatility $\sqrt{252} \times 1.2\% = 15.87 \times 1.2\% = 19.05\%$
- Bitcoin trades every day: $\sqrt{365} \times 3.5\% = 66.9\%$; with $\sqrt{252}$ one reports 55.6%, 17% too low

B1 [Solved]: Descriptive Statistics — Task


- ▣ **Question:** how heavy are the tails of BET-TR, Bitcoin and EUR/RON, and how much does the kurtosis depend on a few days?
- ▣ **Data:** r_{bet} , r_{btc} , r_{eur} from the setup ($T = 2\,992, 4\,384, 5\,352$); $P = 252, 365, 252$
- ▣ **Tasks**
 1. Compute the mean, the standard deviation, the annualised volatility, the skewness, the excess kurtosis, the minimum and the maximum of each series.
 2. Compute $JB = T[\hat{S}^2/6 + (\hat{K} - 3)^2/24]$ and its two terms; compare JB with 5.99, the 5% critical value of χ^2_2 .
 3. Recompute each excess kurtosis after removing the day with the largest $|r_t|$, and after removing the five largest.
 4. Recompute the Crow–Siddiqui quantile kurtosis $\frac{q_{0.975} - q_{0.025}}{q_{0.75} - q_{0.25}} - 2.91$ on the same trimmed samples (0 for the Normal distribution).
- ▣ **Report:** one table for steps 1–2 and a chart of kurtosis against the number of days removed
- ▣ **Interpretation:** which measure of tail heaviness can you report with confidence?

B1 [Solved]: Solution (1/2) — BET-TR Step by Step

- **Step 1:** $\bar{r} = 0.077\%$, $\hat{\sigma} = 0.978\%$; volatility $\sqrt{252} \times 0.978\% = 15.5\%$;
 $\hat{S} = -1.40$, $\hat{K} - 3 = 17.79$
 - ▶ worst day -11.86% on 19 Dec 2018 (the bank-asset tax), best day $+6.71\%$ on 24 Dec 2018
- **Step 2:** skewness term $2992 \times 1.40^2/6 = 977$; kurtosis term $2992 \times 17.79^2/24 = 39455$
 - ▶ $JB = 40433$: 98% of the statistic comes from the kurtosis term;
 $p < 10^{-10}$
- **Step 3:** excess kurtosis 17.8 with all days, 11.8 without 19 Dec 2018, 5.95 without the five largest days (19 Dec 2018, 16 Mar 2020, 9 Mar 2020, 24 Dec 2018, 9 Mar 2022)
- **Step 4:** Crow–Siddiqui 1.06 with all days, 1.01 without the five largest days
- One day carries a third of the moment kurtosis; the quantile measure does not move

	BET-TR	Bitcoin	EUR/RON
Mean (% a day)	0.077	0.118	0.007
Std. dev. (% a day)	0.978	3.485	0.279
Volatility (% a year)	15.5	66.6	4.4
Skewness	-1.40	-0.70	0.73
Excess kurtosis	17.8	12.0	15.1
Min (%)	-11.9	-46.5	-2.5
Max (%)	6.7	22.5	2.9
JB	40433	26482	51575
Top 5 removed	5.95	4.24	11.17

B1 [Solved]: Solution (2/2) — Sensitivity



- All JB statistics exceed 5.99 by four orders of magnitude; Bitcoin without 12 Mar 2020 alone: kurtosis 12.0 $\rightarrow$ 5.3
- Interpretation: the tails are heavy by both measures; report the quantile kurtosis, whose value does not hinge on a few days


A3 [Solved]: Jarque–Bera and Ljung–Box, Step by Step — Task

- ▣ **Question:** how does a statistic become a decision, and which assumptions does the decision need?
- ▣ **Inputs:** two hypothetical samples of daily returns; tests at the 5% level
 - ▶ sample 1: $T = 2\,500$, skewness $\hat{S} = -0.5$, kurtosis $\hat{K} = 8$ (not excess)
 - ▶ sample 2: $T = 1\,000$, $\hat{\rho}_1 = 0.10$, $\hat{\rho}_2 = -0.05$, $\hat{\rho}_3 = 0.03$
- ▣ **Tasks**
 1. Compute the two terms of $JB = T[\hat{S}^2/6 + (\hat{K} - 3)^2/24]$ Jarque & Bera (1980), the statistic, and $p = P(\chi_2^2 \geq JB)$.
 2. Compute the three terms of $Q(3) = T(T+2) \sum_{k=1}^3 \hat{\rho}_k^2 / (T-k)$ Ljung & Box (1978), the statistic, and $p = P(\chi_3^2 \geq Q)$.
 3. State the H_0 of each test and the assumption of its χ^2 reference distribution that fails under volatility clustering.
- ▣ **Report:** the summands, the two statistics, the p-values and the decisions
- ▣ Notebook: section A3

A3 [Solved]: Solution (1/2) — Step by Step

1. $\hat{S}^2/6 = 0.25/6 = 0.042$; $(\hat{K} - 3)^2/24 = 25/24 = 1.042$
 - ▶ $JB = 2500 \times 1.084 = 2708.3$, $p < 10^{-10}$: reject; 96% of JB is the kurtosis term
2. $0.01/999 = 1.001 \times 10^{-5}$, $0.0025/998 = 0.251 \times 10^{-5}$, $0.0009/997 = 0.090 \times 10^{-5}$
 - ▶ $Q(3) = 1000 \times 1002 \times 1.342 \times 10^{-5} = 13.44$, $p = 0.004$: reject
3. JB: H_0 = i.i.d. and Normal distribution; Ljung-Box: H_0 = no autocorrelation, but the χ^2_3 reference also needs i.i.d. returns
 - ▶ under clustering $\text{Var}(\hat{\rho}_k) > 1/T$ and Q rejects too often (A10, B9)

A3 [Solved]: Solution (2/2) — Reference Distributions



- The p-value is the area of the χ^2 density beyond the observed statistic
- Interpretation: for an i.i.d. sample both rejections are sound; for clustered returns only the JB rejection of "i.i.d. and Normal" is sound, the $Q(3)$ p-value is too small

B2 [Proposed]: Heavy Tails and VaR 1% on BET-TR — Task

- ▣ **Model:** B1 [Solved]; notebook section B2
- ▣ **Question:** how large is the one-day loss exceeded on 1% of days, under three distributions?
- ▣ **Data:** r_{bet} , $T = 2\,992$; models fitted and checked on the **same** sample: a descriptive comparison, not a backtest
- ▣ **Tasks**
 1. Standardise $z_t = (r_t - \bar{r})/\hat{\sigma}$ and count the days with $|z_t| > 3$; compare their share with 0.27% under the Normal distribution.
 2. Fit a Student- t to z_t by maximum likelihood (`scipy.stats.t.fit`: degrees of freedom ν , location m , scale s).
 3. Compute the 1% quantile of z_t as $q_z = m + s t_{\nu}^{-1}(0.01)$, then $q_r = \bar{r} + \hat{\sigma} q_z$ and $\text{VaR}_{1\%} = -q_r$; repeat with $q_z = \Phi^{-1}(0.01) = -2.326$ and with the empirical 1% quantile of r_t .
 4. Count the breaches ($r_t < q_r$) of each VaR and compare them with 0.01 $T = 29.9$.
- ▣ **Report:** a table method / VaR / breaches and a lower-tail plot with the three quantiles
- ▣ **Interpretation:** why does the empirical VaR have about 30 breaches by construction?

A6 [Solved]: Kurtosis of a Volatility Mixture — Task


- ▣ **Question:** can random volatility alone create heavy tails?
- ▣ **Assumptions:** $r_t = \sigma_t \varepsilon_t$, ε_t i.i.d. $N(0, 1)$ and independent of σ_t ; $\sigma_t = 1\%$ with probability 0.8 and 2% with probability 0.2, drawn **independently every day**
- ▣ **Tasks**
 1. Using independence and $\mathbb{E}\varepsilon^4 = 3$, show that $K = \mathbb{E}r^4 / (\mathbb{E}r^2)^2 = 3\mathbb{E}\sigma^4 / (\mathbb{E}\sigma^2)^2$.
 2. Complete a two-state table with the probability, σ^2 (in $\%^2$) and σ^4 (in $\%^4$) and compute $\mathbb{E}\sigma^2$, $\mathbb{E}\sigma^4$ and K .
 3. Use Jensen's inequality, $\mathbb{E}[X^2] \geq (\mathbb{E}X)^2$ with $X = \sigma^2$, to show that $K \geq 3$.
 4. State whether this model produces volatility clustering.
- ▣ **Report:** the table, K and the excess kurtosis
- ▣ **Interpretation:** link the result to the mixture-of-distributions hypothesis of [Clark \(1973\)](#)
- ▣ Notebook: section A6

A6 [Solved]: Solution (1/2) — Moments

State	Prob.	σ^2	σ^4
calm	0.8	1	1
turbulent	0.2	4	16
mean		1.6	4.0

1. $\mathbb{E}r^4 = \mathbb{E}\sigma^4 \mathbb{E}\varepsilon^4 = 3\mathbb{E}\sigma^4$ and $\mathbb{E}r^2 = \mathbb{E}\sigma^2$
2. $K = 3 \times 4.0/1.6^2 = 12/2.56 = 4.69$; excess kurtosis 1.69
3. $\mathbb{E}[(\sigma^2)^2] \geq (\mathbb{E}\sigma^2)^2$, with equality only for a constant σ : $K \geq 3$
4. No clustering: the states are independent across days, so $|r_t|$ is uncorrelated; A8 makes them persistent

A6 [Solved]: Solution (2/2) — Densities




- $P(|r_t| > 4\%)$: 0.92% in the mixture, 0.16% for the Normal distribution with the same variance
- Interpretation: [Clark \(1973\)](#)'s random operational time (the pace of information arrival) is such a mixture: variation in volatility creates heavy tails

B3 [Solved]: Uncorrelated but Not Independent (Bitcoin) — Task

- ▣ **Question:** do past Bitcoin returns predict the direction or only the size of today's return?
- ▣ **Data:** `r_btc`, 18 Sep 2014 – 18 Sep 2026, $T = 4\,384$
- ▣ **Tasks**
 1. Plot the returns, then the ACF of r_t , $|r_t|$ and r_t^2 for lags 1–100 with the i.i.d. reference band $\pm 1.96/\sqrt{T} = \pm 0.030$.
 2. Compute the Ljung–Box $Q(10)$ on r_t and on r_t^2 (`acorr_ljungbox`), with p-values from χ_{10}^2 .
 3. Run the ARCH-LM test with 5 lags [Engle \(1982\)](#): regress x_t^2 on a constant and $x_{t-1}^2, \dots, x_{t-5}^2$ and compare $(T-5)R^2$ with χ_5^2 .
- ▣ **Report:** the chart, the lag-1 autocorrelations and the three statistics; label each with its series and its H_0
- ▣ **Interpretation:** the $Q(10)$ on r_t has $p \approx 3\%$; given step 3, is its i.i.d. reference distribution valid?
- ▣ **Notebook:** section B3

B3 [Solved]: Solution (1/2) — Returns and ACF



Direction: no structure in the ACF of r_t ; size: $|r_t|$ and r_t^2 autocorrelated for dozens of lags

B3 [Solved]: Solution (2/2) — Tests

1. $\hat{\rho}_1$: -0.022 for r_t (inside the band), 0.213 for $|r_t|$ and 0.137 for r_t^2
 2. $Q(10)$ on r_t : 20.2 , $p = 0.028$; on r_t^2 : 212.7 , $p \approx 4 \times 10^{-40}$
 3. ARCH-LM(5) = 114.7 , $p \approx 4 \times 10^{-23}$; its χ^2 reference needs $\mathbb{E}r^4 < \infty$, so the p-value is indicative
- ▣ The sizes of returns are dependent: returns are not i.i.d., so the 2.8% p-value of $Q(10)$ on r_t rests on a false assumption
 - ▣ The i.i.d. band is a reference, not a valid test for $|r_t|$ and r_t^2 ; A10 and B9 correct the inference for r_t


A10 [Proposed]: Why the Autocorrelation Band Widens under Clustering — Task

- ▣ **Model:** A6 [Solved]; notebook section A10
- ▣ **Setting:** $x_t = \sigma_t \varepsilon_t$ with ε_t i.i.d. $N(0, 1)$, independent of the stationary volatility process; then x_t is uncorrelated and $\text{Var}(\hat{\rho}_1) \approx v_1 / T$ with $v_1 = \mathbb{E}[x_t^2 x_{t-1}^2] / (\mathbb{E}x_t^2)^2$ [Lobato, Nankervis & Savin \(2001\)](#)
- ▣ **Inputs:** A6: $\mathbb{E}\sigma^2 = 1.6$, $\text{Var}(\sigma^2) = 4.0 - 1.6^2 = 1.44$ (in %⁴); persistent regimes (A8): $\text{Cov}(\sigma_t^2, \sigma_{t-1}^2) = \lambda \text{Var}(\sigma^2)$ with $\lambda = 0.875$; S&P 500: $T = 9\,245$, $\hat{\rho}_1 = -0.079$, $T\hat{\rho}_1 = 5.00$
- ▣ **Tasks**
 1. Show that $v_1 = 1$ for the A6 mixture, whose states are independent across days.
 2. Show that $v_1 = 1 + \lambda \text{Var}(\sigma^2) / (\mathbb{E}\sigma^2)^2$ with persistent regimes; compute it and the band-widening factor $\sqrt{v_1}$.
 3. Compute the S&P 500 lag-1 t -statistic with the i.i.d. SE $1/\sqrt{T}$ and with the robust SE $\sqrt{\hat{\tau}_1}$, and the two 95% band half-widths.
- ▣ **Interpretation:** which feature widens the band: heavy tails or persistent volatility?

B9 [Solved]: Heteroskedasticity-Robust Ljung–Box — Task

- ▣ **Question:** which autocorrelations of returns survive once the variance of $\hat{\rho}_k$ allows for clustering?
- ▣ **Data:** `r_sp`, `r_bet`, `r_eur`, `r_btc`; $x_t = r_t - \bar{r}$
- ▣ **Tasks**
 1. For $k = 1, \dots, 20$ compute $\hat{\rho}_k$ and $\hat{\tau}_k = \sum_{t=k+1}^T x_t^2 x_{t-k}^2 / (\sum_{t=1}^T x_t^2)^2$; plot $\hat{\rho}_k$ with the classical band $\pm 1.96/\sqrt{T}$ and the robust band $\pm 1.96\sqrt{\hat{\tau}_k}$ [Lobato, Nankervis & Savin \(2001\)](#).
 2. Compute the classical $Q(10)$ and the robust $\tilde{Q}(10) = \sum_{k=1}^{10} \hat{\rho}_k^2 / \hat{\tau}_k$; compare both with χ_{10}^2 (5% critical value 18.3).
 3. Count the lags among 1–20 outside each band: these are pointwise checks, distinct from the joint 10-lag test of step 2.
- ▣ **Assumptions of $\tilde{Q}$:** $\mathbb{E}r^4 < \infty$ and zero cross terms $\mathbb{E}[x_t^2 x_{t-k} x_{t-l}] = 0$ for $k \neq l$
- ▣ **Interpretation:** for which series does autocorrelation survive, and is the fourth-moment assumption credible?
- ▣ **Notebook:** section B9

B9 [Solved]: Solution (1/2) — ACF with Both Bands



- S&P 500, lag 1: $\hat{\rho}_1 = -0.079$, $T\hat{\tau}_1 = 5.00$; the robust band is $1.96\sqrt{5.00/9\,245} = 0.046$ instead of 0.020
- At lag 1 the robust band is $\sqrt{T\hat{\tau}_1} = 1.7$ (Bitcoin) to 2.6 (EUR/RON) times wider than the classical one

B9 [Solved]: Solution (2/2) — Joint Tests and Interpretation


	S&P 500	BET-TR	EUR/RON	Bitcoin
$Q(10)$ (p)	91.6 ($< 10^{-10}$)	70.0 ($< 10^{-10}$)	221.7 ($< 10^{-10}$)	20.2 (0.028)
$\tilde{Q}(10)$ (p)	19.0 (0.041)	16.9 (0.076)	37.4 (< 0.001)	11.9 (0.29)
Lags outside, classical / robust	8 / 3	5 / 2	6 / 2	3 / 0
Lags outside the robust band	1, 15, 16	2, 10	1, 3	none

- At 5% the classical test rejects for all four series; the robust test rejects for the S&P 500 (barely) and EUR/RON only
- Bitcoin (B3): the 2.8% p-value becomes 29%: no evidence of linear autocorrelation at lags 1–10
- EUR/RON keeps $\hat{\rho}_1 = 0.17$: the managed fixing smooths adjustments over several days
- Remaining assumptions: the thresholded test of [Dalla, Giraitis & Phillips \(2022\)](#) finds no significant cross terms for the S&P 500, BET-TR and Bitcoin; but with Hill $\hat{\alpha} \approx 3$ the fourth moment is doubtful, so the robust p-values are conditional on it

A7 [Solved]: The Hill Estimator, Step by Step — Task

- ▣ **Question:** how is a tail index estimated from the largest observations, and how precise is it?
- ▣ **Inputs:** the 11 largest absolute daily returns (%) of a hypothetical sample: 12.0; 9.5; 8.2; 7.4; 6.9; 6.1; 5.8; 5.5; 5.2; 5.0; 4.8
- ▣ **Assumption:** above the threshold $X_{(k+1)}$ the tail is Pareto, $P(X > x) \propto x^{-\alpha}$, and the observations are independent
- ▣ **Tasks**
 1. For $k = 10$ compute the log-ratios $\ln(X_{(i)}/X_{(11)})$, their mean and $\hat{\alpha} = 1/\text{mean}$ Hill (1975).
 2. Compute $\text{SE} = \hat{\alpha}/\sqrt{k}$ and the interval $\hat{\alpha} \pm 1.96 \text{SE}$.
 3. State which moments $\mathbb{E}|X|^p$ ($p < \alpha$) exist at the point estimate, and which claims hold at 95%.
 4. Repeat steps 1–2 on the 11 largest $|r_t|$ of BET-TR, using the worksheet printed by the notebook.
- ▣ **Report:** $\hat{\alpha}$, SE and interval for both samples
- ▣ Notebook: section A7

A7 [Solved]: Solution (1/2) — Hypothetical Sample



1. The labels are the log-ratios to 4.8; sum 3.616, mean 0.3616, $\hat{\alpha} = 1/0.3616 = 2.77$
2. $SE = 2.77/\sqrt{10} = 0.87$; interval [1.05; 4.48], a rough normal approximation for i.i.d. Pareto tails
3. At 2.77 the variance exists and the fourth moment does not, but the interval contains values below 2 and above 4: neither claim holds at 95%

A7 [Solved]: Solution (2/2) — BET-TR Worksheet


i	Date	r_t (%)	$X_{(i)}$ (%)	$\ln(X_{(i)}/X_{(11)})$
1	19 Dec 2018	-11.857	11.857	0.824
2	16 Mar 2020	-10.075	10.075	0.661
3	9 Mar 2020	-7.827	7.827	0.409
4	24 Dec 2018	+6.707	6.707	0.254
5	9 Mar 2022	+6.662	6.662	0.247
6	24 Aug 2015	-6.512	6.512	0.225
7	24 Mar 2020	+5.972	5.972	0.138
8	17 Mar 2020	+5.928	5.928	0.131
9	4 Mar 2022	-5.490	5.490	0.054
10	12 Mar 2020	-5.285	5.285	0.016
11	18 Jan 2016	-5.202	5.202	threshold

- Row 1 worked: $\ln(11.857/5.202) = \ln 2.279 = 0.824$
- Sum of the ten log-ratios 2.958;
 $\hat{\alpha}_{10} = 10/2.958 = 3.38$
- $SE = 3.38/\sqrt{10} = 1.07$; interval [1.28; 5.48]
- Six of the ten days fall in December 2018 or March 2020: the extremes cluster, so the i.i.d. SE understates the uncertainty
- Interpretation: ten observations cannot decide whether the fourth moment exists; B11 uses $k = 2\%$ of n

B11 [Solved]: Hill on Both Tails, BET-TR and Bitcoin — Task

- ▣ **Question:** are large losses more frequent than large gains of the same size?
- ▣ **Data:** r_{bet} ($n = 2\,992$), r_{btc} ($n = 4\,384$); losses $\text{loss} = -r[r < 0]$, gains $\text{gain} = r[r > 0]$, each sorted in decreasing order
- ▣ **Tasks**
 1. Plot the Hill estimate $\hat{\alpha}(k)$ of each tail for k from 1% to 10% of the full-sample n , with the band $\hat{\alpha}(1 \pm 1.96/\sqrt{k})$.
 2. Compute $\hat{\alpha}_-$ (losses) and $\hat{\alpha}_+$ (gains) at $k = \lfloor 0.02 n \rfloor$ (BET-TR 59, Bitcoin 87) with 95% intervals; the lecture uses $k = 2.5\%$ of n for $|r_t|$, which pools both tails.
 3. Compute the approximate independent-tail statistic $z = (\hat{\alpha}_- - \hat{\alpha}_+)/\sqrt{\hat{\alpha}_-^2/k + \hat{\alpha}_+^2/k}$ and its two-sided p-value.
 4. Compare with the Student- t $\hat{\nu}$ fitted to the standardised returns: 3.10 for BET-TR (B2), 2.23 for Bitcoin.
- ▣ **Interpretation:** is the gain/loss asymmetry statistically established, and does clustering make the true SE larger or smaller?
- ▣ Notebook: section B11

B11 [Solved]: Solution (1/2) — Hill Plots and Estimates



□ BET-TR: $\hat{\alpha}_- = 2.19$ [1.63; 2.75], $\hat{\alpha}_+ = 3.08$ [2.30; 3.87]; Bitcoin: $\hat{\alpha}_- = 2.89$ [2.28; 3.50], $\hat{\alpha}_+ = 3.62$ [2.86; 4.38]

□ E.g. BET-TR losses: $2.19 \times (1 \pm 1.96/\sqrt{59}) = 2.19 \times (1 \pm 0.255)$

B11 [Solved]: Solution (2/2) — Test and Interpretation

- ▣ BET-TR: $\sqrt{2.19^2/59 + 3.08^2/59} = \sqrt{0.0812 + 0.1612} = 0.492$; $z = (2.19 - 3.08)/0.492 = -1.82$, $p = 0.069$
- ▣ Bitcoin: $\sqrt{2.89^2/87 + 3.62^2/87} = 0.497$; $z = (2.89 - 3.62)/0.497 = -1.47$, $p = 0.14$
- ▣ Decision: losses are heavier in both markets, but $H_0: \alpha_- = \alpha_+$ is not rejected at 5%
 - ▶ the statistic treats the two tails as independent samples; extremes cluster in time, so the true SE is larger and the evidence weaker
 - ▶ a joint block bootstrap of the original returns keeps both features (extension)
- ▣ $\hat{\nu}$ is fitted to the whole distribution; Bitcoin's 2.23 lies below both tail indices: the Student- t does not fit centre and tails together, while $\hat{\alpha}$ uses only the tail

B4 [Proposed]: Aggregational Gaussianity (Bitcoin) — Task

- ▣ **Model:** B1 and B3 [Solved]; notebook section B4
- ▣ **Question:** do h -day Bitcoin returns approach the Normal distribution as h grows?
- ▣ **Data:** r_{btc} , $T = 4\,384$; non-overlapping blocks from the first day: the first 5-day block is 18–22 Sep 2014, the second 23–27 Sep 2014; the incomplete last block (4 days for $h = 5$ and $h = 20$) is dropped
- ▣ **Tasks**
 1. Build the h -day log returns (sums of h consecutive daily returns) for $h = 1, 5, 20$.
 2. Draw a QQ plot of each standardised series against the Normal distribution, with N , the number of blocks, in the title.
 3. Compute the excess kurtosis and the i.i.d.-Gaussian reference SE $\sqrt{24/N}$.
- ▣ **Report:** the three QQ plots and a table h , N , excess kurtosis, $\sqrt{24/N}$
- ▣ **Interpretation:** why is $\sqrt{24/N}$ too optimistic here, and why do non-overlapping blocks remove shared days but not volatility dependence?

A4(a) [Proposed]: How Deep Is a Drawdown? — Task

- ▣ **Model:** A1 [Solved]; notebook section A4
- ▣ **Question:** what is the worst loss from a previous peak, and how does it compare with the growth rate?
- ▣ **Inputs:** year-end portfolio values $V_0, \dots, V_5 = 100, 120, 90, 110, 130, 117$; 5-year CAGR 3.19%
- ▣ **Tasks:** complete the worksheet
 1. Running peak $M_t = \max_{s \leq t} V_s$.
 2. Drawdown $DD_t = V_t/M_t - 1$, as a negative number.
 3. Maximum drawdown $MDD = \min_t DD_t$, its year, and the Calmar ratio $\text{CAGR}/|MDD|$.

Year	0	1	2	3	4	5
V_t	100	120	90	110	130	117
M_t	100	120	...	...	...	...
DD_t	0%	0%	...	...	...	...

- ▣ **Interpretation:** why is the fall to 117 in year 5 not the maximum drawdown?

C2 [Proposed]: Critique an AI Answer — the Answer

- **Prompt** sent to an AI assistant: “Are the daily BET-TR log returns Normally distributed? Are they autocorrelated? Give Python code and a conclusion”
- **Answer:** the code below and the conclusion

```
from scipy import stats
from statsmodels.stats.diagnostic import acorr_ljungbox
r = np.log(bet["close"]).diff().dropna()      # daily log returns
k = stats.kurtosis(r)                          # sample kurtosis
print("excess_kurtosis:", k - 3)               # the Normal has kurtosis 3
jb, p = stats.jarque_bera(r)
lb = acorr_ljungbox(r**2, lags=[10])          # Ljung-Box, 10 lags
```

- **Conclusion of the AI**
 - ▶ “Excess kurtosis 14.8: heavy tails”
 - ▶ “JB p -value 0.000: the probability that BET-TR returns follow the Normal distribution is 0%”
 - ▶ “Ljung-Box $Q(10) = 713.5$, $p < 0.001$: returns are strongly autocorrelated, so BET-TR is predictable and not a random walk”

C2 [Proposed]: Critique an AI Answer — Task

- ▣ **Model:** A3, B3 and B9 [Solved]; notebook section C2
- ▣ **Data:** `r_bet`, the 2992 returns of the setup
- ▣ **Tasks**
 1. Find every substantive error in the code and in the three conclusions.
 2. For each error, say how you would detect it: a test on simulated data, a number from Lecture 1, a solved exercise or the library documentation.
 3. Write corrected, executable code and report the excess kurtosis, JB with its p-value, $Q(10)$ on r_t and on r_t^2 , and the robust $\tilde{Q}(10)$ of B9.
 4. State the assumptions behind the robust p-value.
 5. Write the corrected conclusion in three sentences.
- ▣ **Report:** the corrected code, a table AI value / correct value, a chart of the ACF of r_t and of r_t^2
- ▣ AI code usually runs without an error message: the mistakes are in the statistics, not in the syntax; log what you find in `AI_ERRORS.md`

Submission Checklist and Team Repository

Hand in before the next seminar

- ▣ A1, A3, A6, A7, A10, A4(a): tables and answers
- ▣ B1, B2, B3, B9, B11, B4: one figure and one table each, as in the solutions
- ▣ C2: corrected code, table, chart, corrected conclusion
- ▣ One interpretation sentence per exercise
- ▣ Homework choice: one extension pack

Team repository

- ▣ One Quantlet folder: B9 or B11 on an asset of your choice listed on the BVB
- ▣ Log AI use in AI_USE.md and caught errors in AI_ERRORS.md
- ▣ Project seeds: C1 (BET-TR after the 2020 reclassification), C3 (replication of [Guyon & Lekeufack \(2023\)](#))

Extension Packs: What Each One Needs

Pack	Exercises	Prerequisites
1. Return arithmetic and drawdowns	A1(c) [Solved], A2, A4(b–e)	A1, A4(a)
2. Mixtures, regimes and tails	A5 [Solved], A8, A9, A11	A6, A7, A10
3. Robust moments and model checks	B1(d) [Solved], B10, B14	B1, B9, B11
4. Dependence and asymmetry	B5, B12, B6, B15	B3, B9
5. Performance and range estimators	B7, B13, B8	A1, A5, B1
6. Projects	C1, C3	B9, the lecture case study

- Each task has its data, numbered steps and deliverable on one slide; notebook sections carry the same labels
- Solutions of [Proposed] exercises are discussed at the next seminar

A1(c) [Solved]: Portfolio Returns — Task and Solution

- ▣ **Question:** which return adds up across assets?
- ▣ **Inputs:** 60% in asset A and 40% in asset B for one day; $R_A = +5\%$, $R_B = -2\%$
- ▣ **Tasks**
 1. Compute the portfolio simple return $R_p = 0.6R_A + 0.4R_B$.
 2. Compute the weighted log return $0.6r_A + 0.4r_B$ and the true portfolio log return $\ln(1 + R_p)$.
- ▣ **Solution**
 1. $R_p = 0.6 \times 5\% + 0.4 \times (-2\%) = 2.20\%$
 2. $0.6 \times \ln 1.05 + 0.4 \times \ln 0.98 = 0.6 \times 4.879\% - 0.4 \times 2.020\% = 2.119\%$, while $\ln 1.022 = 2.176\%$
- ▣ **Interpretation:** simple returns add across assets, log returns add over time (A1); for daily moves the gap between the two rules is small, for large moves it is not

A2 [Proposed]: The $\sqrt{h}$ Rule under Autocorrelation — Task

- ▣ **Model:** A1 [Solved]; notebook section A2
- ▣ **Assumption:** stationary AR(1), $r_t = \rho r_{t-1} + e_t$, $|\rho| < 1$, $\text{Var}(r_t) = \sigma^2$, so $\gamma_j = \text{Cov}(r_t, r_{t-j}) = \sigma^2 \rho^j$
- ▣ **Scaffold**, $h = 3$: $\text{Var}(r_1 + r_2 + r_3) = 3\gamma_0 + 2(2\gamma_1 + \gamma_2)$: two pairs at lag 1, one pair at lag 2
- ▣ **Tasks**
 1. Generalise the pair count: show that $\text{Var}(\sum_{j=1}^h r_{t+j}) = h\sigma^2[1 + 2\sum_{j=1}^{h-1}(1 - j/h)\rho^j]$.
 2. Using $\sum_{j=1}^{h-1} \rho^j = \rho(1 - \rho^{h-1})/(1 - \rho)$ and $\sum_{j \geq 1} j|\rho|^j = |\rho|/(1 - |\rho|)^2$, show that the variance ratio $VR(h) = \text{Var}(\cdot)/(h\sigma^2) \rightarrow (1 + \rho)/(1 - \rho)$.
 3. Illustrative calculation: treat the sample lag-1 autocorrelations as AR(1) parameters, EUR/RON $\rho = 0.170$ (volatility 4.44%), S&P 500 $\rho = -0.079$ (17.99%); compute $\sqrt{VR(252)}$ and the corrected annual volatilities.
 4. Explain why volatility clustering alone does not change step 1.
- ▣ **Interpretation:** which ACF check must precede step 3?

A4(b–e) [Proposed]: A Benchmark for Drawdowns — Task

- ▣ **Model:** A4(a); notebook section A4
- ▣ **Assumption:** the log price is $X_t = \mu t + \sigma W_t$, with W_t a standard Brownian motion, σ the annual volatility, T the horizon in years; maximum log drawdown $D = \max_{s \leq t \leq T} (X_s - X_t) = -\ln(1 + MDD)$; for $\mu = 0$:
 $\mathbb{E}[D] = \sqrt{\pi/2} \sigma \sqrt{T}$ Magdon-Ismail et al. (2004)
- ▣ **Inputs:** gold spot, 1990–2026: $\sigma = 16.3\%$, $T = 36.7$ years, observed $MDD = -44.7\%$ ($D = 0.592$), mean log return 6.5% a year
- ▣ **Tasks**
 1. Use $W(cT) \stackrel{d}{=} \sqrt{c} W(T)$ to explain why $\mathbb{E}[D]$ is proportional to $\sqrt{T}$ when $\mu = 0$.
 2. Compute the ratio of $\mathbb{E}[D]$ for 40 and 10 years of the same process.
 3. Compute the driftless benchmark for gold and compare it with the observed D .
 4. Simulate 2000 daily paths (seed 42) with $\mu = 0$ and with $\mu = 6.5\%$ and plot the mean D against T .
- ▣ **Interpretation:** what do steps 1–4 imply for Calmar ratios computed on samples of different length?


A5 [Solved]: Parkinson Volatility and Its Efficiency — Task

- ▣ **Question:** how much information does the daily range carry about volatility?
- ▣ **Notation:** o_t, h_t, l_t, c_t = log open, high, low, close; $h_t - l_t = \ln(H_t/L_t)$
- ▣ **Inputs:** H/L of five days: 1.020, 1.015, 1.030, 1.010, 1.025; $P = 252$
- ▣ **Assumption:** within the day the log price is a driftless Brownian motion with volatility σ ; no overnight jumps
- ▣ **Tasks**
 1. Complete a five-row table: $\ln(H/L)$, its square, and the mean of the squares.
 2. Compute $\hat{\sigma}_P^2 = \overline{(h - l)^2} / (4 \ln 2)$ [Parkinson \(1980\)](#), the daily and the annual volatility.
 3. Name the information the estimator ignores and the effect of a strong drift.
 4. Using $\mathbb{E}[(h - l)^2] = 4 \ln 2 \sigma^2$ and $\mathbb{E}[(h - l)^4] = 9\zeta(3) \sigma^4$ ($\zeta(3) = 1.202$), compare the per-day variance of $(h - l)^2 / (4 \ln 2)$ with $\text{Var}(r_t^2) = 2\sigma^4$.
- ▣ **Notebook:** section A5

A5 [Solved]: Solution (1/2) — The Estimate

H/L	$\ln(H/L)$	square ($\times 10^{-4}$)
1.020	0.0198	3.92
1.015	0.0149	2.22
1.030	0.0296	8.74
1.010	0.0100	0.99
1.025	0.0247	6.10
mean		4.39

- $\hat{\sigma}_P^2 = 0.000439/2.7726 = 0.000158$; $\hat{\sigma}_P = 1.26\%$ a day, $1.26\% \times 15.87 = 20.0\%$ a year
- Ignored: the open and the overnight jump Yang & Zhang (2000); a drift widens the range and biases $\hat{\sigma}_P$ up Rogers & Satchell (1991)



- A simulated day: the range $h_t - l_t$ uses the whole path, the close-to-close return only its end points

A5 [Solved]: Solution (2/2) — Efficiency

4. $\text{Var}((h - l)^2) = (9\zeta(3) - 16(\ln 2)^2)\sigma^4 = (10.819 - 7.687)\sigma^4 = 3.131\sigma^4$

- ▶ divided by $(4 \ln 2)^2 = 7.687$: $0.407\sigma^4$ per day for the Parkinson term
- ▶ close-to-close: $\text{Var}(r_t^2) = 3\sigma^4 - \sigma^4 = 2\sigma^4$ per day

- For an n -day average of independent days both variances are divided by n : the ratio stays $2/0.407 = 4.9$
- One day of ranges carries about the information of five close-to-close days: the “about 5” of Lecture 1
- Interpretation: the gain assumes driftless continuous trading without overnight jumps; a stale open affects the estimators that use the open (Garman–Klass, Rogers–Satchell, Yang–Zhang), not Parkinson (B8)

A8 [Proposed]: Where the Autocorrelation of r_t^2 Comes From — Task

- ▣ **Model:** A6 [Solved] and A10; notebook section A8
- ▣ **Assumption:** as in A6, $\sigma_t^2 \in \{1, 4\}$ (in %²) with stationary probabilities (0.8; 0.2), but the state follows a two-state Markov chain [Hamilton \(1989\)](#) with transition matrix $\begin{pmatrix} 0.975 & 0.025 \\ 0.1 & 0.9 \end{pmatrix}$ (rows: today calm, turbulent)
- ▣ **Tasks**
 1. Check the balance condition $0.8 \times 0.025 = 0.2 \times 0.1$ and compute $\lambda = p_{LL} + p_{HH} - 1$.
 2. With the indicator $I_t = 1$ in the turbulent state, $\sigma_t^2 = 1 + 3I_t$; show that $\text{Cov}(I_t, I_{t-k}) = \pi_H(1 - \pi_H)\lambda^k$ and hence $\text{Cov}(r_t^2, r_{t-k}^2) = \text{Var}(\sigma^2)\lambda^k$ for $k \geq 1$.
 3. Using A6, compute $\text{Var}(\sigma^2)$, $\text{Var}(r^2) = 3\mathbb{E}\sigma^4 - (\mathbb{E}\sigma^2)^2$ and $\text{corr}(r_t^2, r_{t-k}^2)$ at lags 1 and 10.
 4. Plot the implied ACF against the S&P 500 ACF of r_t^2 (0.31 at lag 1, 0.04 at lag 100).
- ▣ **Interpretation:** which two features of the data can this model not reproduce?

A9 [Proposed]: Can a Gaussian GARCH Produce the S&P 500 Tails? — Task

- ▣ **Model:** A6 [Solved]; an advanced preview of Chapter 5; notebook section A9
- ▣ **Model equations** [Bollerslev \(1986\)](#): $r_t = \mu + \varepsilon_t$, $\varepsilon_t = \sigma_t z_t$, z_t i.i.d. $N(0, 1)$, $\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + \beta \sigma_{t-1}^2$, $\omega > 0$, $\alpha, \beta \geq 0$, $\alpha + \beta < 1$; here α is a GARCH coefficient, not a tail index
- ▣ **Result:** if $1 - (\alpha + \beta)^2 - 2\alpha^2 > 0$, then $K = 3[1 - (\alpha + \beta)^2]/[1 - (\alpha + \beta)^2 - 2\alpha^2]$
- ▣ **Tasks**
 1. Compute K for $(\alpha, \beta) = (0.10; 0.85)$ and $(0.10; 0.88)$.
 2. State the condition for a finite fourth moment and compare it with $\alpha + \beta < 1$.
 3. With $\alpha + \beta = 0.99$, solve for the α that gives the S&P 500 kurtosis $K = 13.9$.
 4. Fit the model to `r_sp` in % (`arch_model(100*r_sp, mean='Constant', vol='GARCH', p=1, q=1, dist='normal')`); compute the implied K and draw a QQ plot of $\hat{z}_t = \hat{\varepsilon}_t / \hat{\sigma}_t$.
- ▣ **Interpretation:** what does the QQ plot say about the Normal distribution of z_t ?

A11 [Proposed]: The Hill Estimator as a Maximum Likelihood Estimator — Task

- ▣ **Model:** A7 [Solved]; notebook section A11
- ▣ **Assumption:** given the threshold $u = X_{(k+1)}$, the k exceedances are the order statistics of an i.i.d. Pareto sample with density $f(x) = \alpha u^\alpha x^{-\alpha-1}$, $x > u$
- ▣ **Tasks**
 1. Write the log-likelihood $\ell(\alpha)$ of the k exceedances (the ordering adds only the constant $\ln k!$).
 2. Solve $\ell'(\alpha) = 0$ and show that the maximiser is the Hill estimator $\hat{\alpha} = k / \sum_{i=1}^k \ln(X_{(i)}/u)$ Hill (1975).
 3. Compute $\ell''(\alpha)$, the information k/α^2 and $\text{SE}(\hat{\alpha}) \approx \hat{\alpha}/\sqrt{k}$; check on the data of A7.
 4. Simulate $n = 2\,000\,000$ values $|X|$, X Student- t with $\nu = 3$ (`numpy.random.default_rng(1)`); plot $\hat{\alpha}_k$ against k/n and report $k = 200, 2\,000, 20\,000, 200\,000$.
- ▣ **Interpretation:** why does $\hat{\alpha}_k$ drift away from 3 when k is large?

B1(d) [Solved]: Quantile-Based Moments with Bootstrap Intervals

- ▣ **Question:** do negative skewness and excess kurtosis survive measures that need no moments?
- ▣ **Measures Kim & White (2004):** Hinkley skewness $\frac{q_{0.95} + q_{0.05} - 2q_{0.5}}{q_{0.95} - q_{0.05}}$ and Crow–Siddiqui kurtosis $\frac{q_{0.975} - q_{0.025}}{q_{0.75} - q_{0.25}} - 2.91$; both 0 for the Normal distribution
- ▣ **Bootstrap:** draw $\lceil T/20 \rceil$ starting days uniformly, take 20 consecutive returns from each, cut to T ; recompute both measures; 1 000 resamples (`default_rng(42)`); 95% percentile interval

	Moment skew.	Hinkley [95% CI]	Excess kurt.	Crow–Siddiqui [95% CI]
BET-TR	−1.40	−0.036 [−0.085, 0.025]	17.8	1.06 [0.73, 1.43]
Bitcoin	−0.70	−0.018 [−0.069, 0.032]	12.0	2.25 [1.88, 2.64]
EUR/RON	0.73	0.041 [−0.022, 0.093]	15.1	4.87 [3.74, 6.13]

- ▣ Every kurtosis interval lies above 0: heavy tails are established without moments
- ▣ Moment skewness and Hinkley skewness measure different things; every Hinkley interval contains 0, which does not establish symmetry

B10 [Proposed]: A Kurtosis That Does Not Converge — Task

- ▣ **Model:** B1 [Solved]; notebook section B10
- ▣ **Data:** `r_sp`; window n = the returns dated before 3 January of year $1990 + n$, $n = 1, \dots, 36$ (253 returns for $n = 1$, 2 527 for $n = 10$, 9 067 for $n = 36$)
- ▣ **Simulation:** 500 Student- t paths of length 9 245 for $\nu = 3$ and for $\nu = 6$ (`default_rng(42)`), cut at the same window counts
- ▣ **Tasks**
 1. Compute the excess kurtosis of the S&P 500 on each window and list the three largest jumps with the days that cause them.
 2. For each ν compute the 5%, 50% and 95% quantiles of the simulated kurtosis at each n .
 3. Plot the data curve over both bands on a symmetric log scale.
- ▣ **Interpretation:** why does the $\nu = 6$ band settle slowly although the kurtosis of t_6 is finite? What does “excess kurtosis = 10.9” then mean?

B14 [Proposed]: Which Stylised Facts Does GARCH- t Reproduce? — Task

- **Model:** B3 and B11 [Solved], A9; notebook section B14
- **Data:** r_{sp} converted to % ($T = 9\,245$); constant mean; ω in %² a day

```
from arch import arch_model
from arch.univariate import StudentsT
am = arch_model(100 * r_sp, mean='Constant', vol='GARCH', p=1, q=1, dist='t')
res = am.fit(dispatch='off')
am.distribution = StudentsT(seed=np.random.default_rng(42)) # reproducible paths
sim = am.simulate(res.params, nobs=len(r_sp), burn=1000)['data']
```

- **Tasks**
 1. Fit the model and list all five parameters with units.
 2. Pilot: simulate one path and compute its excess kurtosis; then run 200 paths.
 3. For each path compute the excess kurtosis, the ACF of $|r_t|$ at lags 1, 20, 100, 250, the Hill $\hat{\alpha}$ of $|r_t|$ at $k = 2\%$ of T and $L(1) = \text{corr}(r_t, |r_{t+1}|)$.
 4. Build the pointwise 5–95% intervals and mark each observed value inside or outside.
- **Interpretation:** which facts are not reproduced, and which model component would you change for each?

B5 [Proposed]: Leverage Effect, Romania vs Crypto — Task

- ▣ **Model:** B3 [Solved]; notebook section B5
- ▣ **Lag convention:** $L(k) = \text{corr}(r_t, |r_{t+k}|)$ [Christie \(1982\)](#); k counts observation days of each series; $k > 0$: today's return and a later size
 - ▶ example, $k = 1$: the BET-TR return of Monday 16 Mar 2020 is paired with $|r|$ of Tuesday 17 Mar 2020
- ▣ **Data:** `r_bet` and `r_btc`, each on its own calendar
- ▣ **Tasks**
 1. Compute $L(k)$ for $k = -20, \dots, 20$ and plot both series with the i.i.d. reference bands $\pm 1.96/\sqrt{T}$ (0.036 and 0.030).
 2. Report $L(1)$ and $L(5)$ for both markets and state whether they lie outside the reference bands.
 3. Give two competing explanations for a weaker Bitcoin effect.
- ▣ **Interpretation:** why is a reference band not a confidence interval under clustering, and what does B12 add?

B12 [Proposed]: The Leverage Effect as a Regression — Task

- ▣ **Model:** B5, B9 [Solved]; notebook section B12
- ▣ **Regression:** $|r_{t+1}| = c + \beta^+ r_t^+ + \beta^- r_t^- + \sum_{j=1}^5 \delta_j |r_{t-j}| + u_{t+1}$, $r_t^+ = \max(r_t, 0) \geq 0$, $r_t^- = \min(r_t, 0) \leq 0$
 - ▶ a gain $+a$ adds $\beta^+ a$, a loss $-a$ adds $-\beta^- a$: a symmetric response means $\beta^+ = -\beta^-$
 - ▶ design row for day t : $(1, r_t^+, r_t^-, |r_{t-1}|, \dots, |r_{t-5}|)$, response $|r_{t+1}|$
- ▣ **Data:** r_sp, r_bet, r_btc, r_eur
- ▣ **Tasks**
 1. Estimate by OLS with Newey–West standard errors [Newey & West \(1987\)](#), lag truncation $\lfloor 4(T/100)^{2/9} \rfloor$.
 2. Test $H_0: \beta^+ + \beta^- = 0$ with $W = (\hat{\beta}^+ + \hat{\beta}^-)^2 / [\text{Var}(\hat{\beta}^+) + \text{Var}(\hat{\beta}^-) + 2 \text{Cov}(\hat{\beta}^+, \hat{\beta}^-)]$, χ_1^2 at 5%.
 3. Plot the predicted $|r_{t+1}|$ after a gain $+a$ and a loss $-a$, lag controls at their means.
 4. Show why $|r_t|$ cannot be added as a control.

B6 [Proposed]: EUR/RON Across Regimes — Task

- ▣ **Model:** B1, B9 [Solved]; notebook section B6
- ▣ **Data:** `r_eur` (BNR reference rate), split into Jul 2005 – 2011, 2012–2019, 2020–2026 (1 656, 2 013, 1 683 returns); lags built inside each period
- ▣ **Tasks**
 1. In each period regress $r_t = a + \rho r_{t-1} + e_t$ by OLS with a HAC SE (lag truncation $\lfloor 4(T/100)^{2/9} \rfloor$); report $\hat{\rho}$, the SE, the volatility and the excess kurtosis.
 2. Test equal ρ across periods: weights $w_j = 1/\text{SE}_j^2$, $\bar{\rho} = \sum_j w_j \hat{\rho}_j / \sum_j w_j$, $W = \sum_j w_j (\hat{\rho}_j - \bar{\rho})^2 \sim \chi_2^2$.
 3. Separately, locate one variance break with the cumulative sum of squares [Inclán & Tiao \(1994\)](#): $D_k = C_k / C_T - k / T$, $C_k = \sum_{t \leq k} x_t^2$, break at $\arg \max_k |D_k|$.
- ▣ **Interpretation:** why is the 2020–2026 SE so large, and why is the CUSUM date descriptive rather than a test?

B15 [Proposed]: Long Memory or Breaks? — Task

- ▣ **Model:** B9, B11 [Solved]; notebook section B15 (helper `local_whittle(x, m)`)
- ▣ **Method Robinson (1995):** demean $x_t = |r_t|$, periodogram $I(\lambda_j)$ at $\lambda_j = 2\pi j/T$; $\hat{d}$ minimises $\ln\left(\frac{1}{m} \sum_{j=1}^m \lambda_j^{2d} I(\lambda_j)\right) - \frac{2d}{m} \sum_{j=1}^m \ln \lambda_j$ over $d \in [-0.49, 0.99]$; $m = \lfloor T^p \rfloor$; $SE = 1/(2\sqrt{m})$
- ▣ **Data:** $|r_t|$ of the S&P 500, BET-TR and Bitcoin
- ▣ **Tasks**
 1. Estimate d with $p = 0.65$ and give the 95% interval.
 2. Plot $\hat{d}$ against p from 0.45 to 0.80, with the lines $d = 0$ and $d = 0.5$.
 3. Locate one variance break of r_t (B6); re-estimate $\hat{d}$ on each side and plot the ACF of $|r_t|$ before and after.
- ▣ **Interpretation:** is $\hat{d} > 0$ robust to p and to the break, or consistent with regimes Diebold & Inoue (2001)?
What does $\hat{d} \geq 0.5$ imply?

B7 [Proposed]: BET-TR vs S&P 500, Same Period — Task

- ▣ **Model:** A1, B1 [Solved]; notebook section B7
- ▣ **Data:** BET-TR (total return, RON) and S&P 500 (price index, USD); align the *prices* on common dates (23 Sep 2014 – 18 Sep 2026), then compute 2 903 returns
 - ▶ interval lengths: 2 231 one-day, 538 of two or three days, 134 longer (at most 6 days)
- ▣ **Tasks**
 1. Plot the rebased wealth (log scale) and the two underwater curves.
 2. Compute the CAGR, the volatility of simple returns ($P = 252$), the Sharpe ratio $252\bar{R}/(\sqrt{252}\hat{\sigma}_R)$ with $r_f = 0$, the Sortino ratio (downside deviation over all days) [Sortino & van der Meer \(1991\)](#), the MDD and the Calmar ratio.
- ▣ **Report:** the chart and a two-column table
- ▣ **Interpretation:** list three reasons why the comparison is descriptive only (dividends, currency, risk-free rate in RON)

B13 [Proposed]: Is the Sharpe Difference Real? — Task

- ▣ **Model:** B7, B1(d); notebook section B13
- ▣ **Data:** the 2 903 paired returns of B7; simple returns for Sharpe ratios (1.36 and 0.75), log returns for correlations
- ▣ **Tasks**
 1. Convert to daily Sharpe ratios ($\widehat{SR}/\sqrt{252}$) and compute the Jobson–Korkie test [Jobson & Korkie \(1981\)](#) with the correction of [Memmle \(2003\)](#): $z = (\widehat{SR}_1 - \widehat{SR}_2)/\sqrt{\widehat{V}}$, $\widehat{V} = \frac{1}{T} [2(1 - \hat{\rho}) + \frac{1}{2}(\widehat{SR}_1^2 + \widehat{SR}_2^2 - 2\widehat{SR}_1\widehat{SR}_2\hat{\rho}^2)]$.
 2. Paired circular block bootstrap: draw $\lceil T/21 \rceil$ start days, take 21 consecutive days from each (wrapping at the end), cut to T , apply the **same** indices to both series; 2 000 resamples (`default_rng(42)`); 95% percentile interval of the annualised difference.
 3. Compute four correlations: same day, BET-TR with the S&P 500 of the previous day and of the next day, and weekly (Friday) returns.
- ▣ **Interpretation:** is the difference significant, what do both methods assume, and which correlation belongs in a portfolio?

B8 [Proposed]: Range-Based Estimators on the S&P 500 — Task

- ▣ **Model:** A5 [Solved]; notebook section B8 (tested helpers for all five estimators; formulas in Lecture 1)
- ▣ **Data:** S&P 500 index OHLC, 2 Jan 2008 – 18 Sep 2026; SPY ETF on one adjusted basis: C^* = the adjusted close (for dividends and splits), $O^*, H^*, L^* = (C^*/C) \times O, H, L$
 - ▶ example, 16 Mar 2020: factor $218.08/239.85 = 0.9092$; $O^* = 241.18 \times 0.9092 = 219.29$, $H^* = 233.58$, $L^* = 215.82$
- ▣ **Tasks**
 1. Compute 21-day rolling close-to-close, Parkinson, Garman–Klass [Garman & Klass \(1980\)](#), Rogers–Satchell [Rogers & Satchell \(1991\)](#) and Yang–Zhang [Yang & Zhang \(2000\)](#) volatilities ($P = 252$) and their time averages.
 2. For the index and for SPY compare the 21-day averages of close-to-close and Yang–Zhang with the full-sample overnight volatility $\text{sd}(o_t - c_{t-1})\sqrt{252}$ and open-to-close volatility.
- ▣ **Interpretation:** why does Yang–Zhang stay below close-to-close on the index but not on SPY?

C1 [Proposed]: Did BET-TR Autocorrelation Fall after the FTSE Reclassification?

- ▣ **Model:** B6, B9 [Solved]; a project seed
- ▣ **Context:** BET-TR has a positive lag-1 autocorrelation, typical of less liquid markets; Romania became a FTSE Russell Secondary Emerging market on 21 Sep 2020
- ▣ **Data:** r_{bet} ; liquidity: daily traded value (volume $\times$ raw close, in lei) of 11 BVB blue chips (TLV, SNP, BRD, TGN, FP, SNG, H2O, SNN, EL, DIGI, TEL), summed over the stocks listed that day; report how many are listed each year
- ▣ **Tasks**
 1. Compute $\hat{\rho}_1$ on rolling 250-day windows (step 5 days) with the B9 bands, labelled as reference bands under $\rho_1 = 0$.
 2. Estimate ρ_1 before and after 21 Sep 2020 as an AR(1) slope with HAC SEs (B6); report $z = (\hat{\rho}_{after} - \hat{\rho}_{before}) / \sqrt{SE_1^2 + SE_2^2}$ and the interaction e in $r_t = a + b r_{t-1} + c \text{post}_t + e r_{t-1} \text{post}_t$ (HAC, 10 lags).
 3. Correlate the rolling $\hat{\rho}_1$ with the log 250-day average traded value.
- ▣ **Deliverable:** a Quantlet folder and one slide with a descriptive conclusion and its main limitation, not a causal claim

C3 [Proposed]: Is the VIX Mostly Path-Dependent? — Task (1/2)

- ▣ **Model:** the lecture case study; a replication project, not an in-class task
- ▣ **Question:** can past S&P 500 returns alone explain the VIX, as [Guyon & Lekeufack \(2023\)](#) find (94.6% of its variance in 2000–2018, 85.5% in 2019 – May 2022)?
- ▣ **Data:** daily closes of the S&P 500, the VIX and the 9-day VIX; simple returns $r_{t_i} = S_{t_i}/S_{t_{i-1}} - 1$; volatility indices divided by 100
- ▣ **Model**, eq. (3.7) in the paper: $\sigma_t = \beta_0 + \beta_1 R_{1,t} + \beta_2 \Sigma_t$, $R_{1,t} = \sum_{i=0}^{999} K_1(i\Delta t) r_{t-i}$, $\Sigma_t = \left(\sum_{i=0}^{999} K_2(i\Delta t) r_{t-i}^2 \right)^{1/2}$, $\Delta t = 1/252$
- ▣ **Kernels**, eq. (3.9) in the paper: $K_j(\tau) = (\tau + \delta_j)^{-\alpha_j} / Z_j$, with Z_j such that $\sum_{i=0}^{999} K_j(i\Delta t) \Delta t = 1$
- ▣ **Parameters and units:** β_0 in volatility units ($0.03 = 3$ VIX points), β_1, β_2 unitless, $\alpha_j > 1$, $\delta_j > 0$ in years
- ▣ **Start** (Table 3 in the paper, VIX row): $\beta_0 = 0.057$, $\beta_1 = -0.095$, $\beta_2 = 0.82$, $\alpha_1 = 1.06$, $\delta_1 = 0.020$, $\alpha_2 = 1.60$, $\delta_2 = 0.052$; bounds $\alpha_j \in [1.0001, 20]$, $\delta_j \in [10^{-4}, 5]$

C3 [Proposed]: Is the VIX Mostly Path-Dependent? — Task (2/2)

- ▣ **Tasks**, in stages
 1. Fixed kernels: compute $R_{1,t}$ and Σ_t with the Table 3 kernels and estimate $\beta_0, \beta_1, \beta_2$ by OLS on 2000–2018.
 2. Full model: estimate all seven parameters by nonlinear least squares (`scipy.optimize.least_squares`) on 1 Jan 2000 – 31 Dec 2018, recomputing the features at every trial value.
 3. Report r^2 and RMSE on the training set and on 1 Jan 2019 – 15 May 2022, for the full model and the restrictions $\beta_1 = 0$ and $\beta_2 = 0$; repeat the full model for the 9-day VIX (training from 2011).
 4. Keep the parameters fixed and score 16 May 2022 – 18 Sep 2026.
- ▣ **Report**: parameter table, scores next to Table 2 of the paper, observed and fitted VIX, the ratio of observed to fitted VIX, the fitted kernels
- ▣ **Interpretation**: the model fits the VIX on the same day; it is not a next-day forecast; which feature explains more, $R_{1,t}$ or Σ_t ?

References (1/3)

- Bollerslev, T. (1986). Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics*, 31(3), 307–327.
- Bollerslev, T., Hood, B., Huss, J., Pedersen, L.H. (2018). Risk everywhere: modeling and managing volatility. *Review of Financial Studies*, 31(7), 2729–2773.
- Christie, A.A. (1982). The stochastic behavior of common stock variances: value, leverage and interest rate effects. *Journal of Financial Economics*, 10(4), 407–432.
- Clark, P.K. (1973). A subordinated stochastic process model with finite variance for speculative prices. *Econometrica*, 41(1), 135–155.
- Cont, R. (2001). Empirical properties of asset returns: stylized facts and statistical issues. *Quantitative Finance*, 1(2), 223–236.
- Dalla, V., Giraitis, L., Phillips, P.C.B. (2022). Robust tests for white noise and cross-correlation. *Econometric Theory*, 38(5), 913–941.
- Diebold, F.X., Inoue, A. (2001). Long memory and regime switching. *Journal of Econometrics*, 105(1), 131–159.
- Dimson, E. (1979). Risk measurement when shares are subject to infrequent trading. *Journal of Financial Economics*, 7(2), 197–226.
- Engle, R.F. (1982). Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica*, 50(4), 987–1007.
- Franke, J., Härdle, W.K., Hafner, C.M. (2019). *Statistics of Financial Markets: An Introduction* (5th ed.). Springer.
- Garman, M.B., Klass, M.J. (1980). On the estimation of security price volatilities from historical data. *Journal of Business*, 53(1), 67–78.
- Guyon, J., Lekeufack, J. (2023). Volatility is (mostly) path-dependent. *Quantitative Finance*, 23(9), 1221–1258.

References (2/3)

- ▣ Hamilton, J.D. (1989). A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica*, 57(2), 357–384.
- ▣ Hill, B.M. (1975). A simple general approach to inference about the tail of a distribution. *Annals of Statistics*, 3(5), 1163–1174.
- ▣ Inclán, C., Tiao, G.C. (1994). Use of cumulative sums of squares for retrospective detection of changes of variance. *Journal of the American Statistical Association*, 89(427), 913–923.
- ▣ Jarque, C.M., Bera, A.K. (1980). Efficient tests for normality, homoscedasticity and serial independence of regression residuals. *Economics Letters*, 6(3), 255–259.
- ▣ Jobson, J.D., Korkie, B.M. (1981). Performance hypothesis testing with the Sharpe and Treynor measures. *Journal of Finance*, 36(4), 889–908.
- ▣ Kim, T.-H., White, H. (2004). On more robust estimation of skewness and kurtosis. *Finance Research Letters*, 1(1), 56–73.
- ▣ Ledoit, O., Wolf, M. (2008). Robust performance hypothesis testing with the Sharpe ratio. *Journal of Empirical Finance*, 15(5), 850–859.
- ▣ Ljung, G.M., Box, G.E.P. (1978). On a measure of lack of fit in time series models. *Biometrika*, 65(2), 297–303.
- ▣ Lo, A.W. (2002). The statistics of Sharpe ratios. *Financial Analysts Journal*, 58(4), 36–52.
- ▣ Lobato, I., Nankervis, J.C., Savin, N.E. (2001). Testing for autocorrelation using a modified Box–Pierce Q test. *International Economic Review*, 42(1), 187–205.
- ▣ Magdon-Ismael, M., Atiya, A.F., Pratap, A., Abu-Mostafa, Y.S. (2004). On the maximum drawdown of a Brownian motion. *Journal of Applied Probability*, 41(1), 147–161.

References (3/3)

- ▣ Memmel, C. (2003). Performance hypothesis testing with the Sharpe ratio. *Finance Letters*, 1(1), 21–23.
- ▣ Newey, W.K., West, K.D. (1987). A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica*, 55(3), 703–708.
- ▣ Parkinson, M. (1980). The extreme value method for estimating the variance of the rate of return. *Journal of Business*, 53(1), 61–65.
- ▣ Robinson, P.M. (1995). Gaussian semiparametric estimation of long range dependence. *Annals of Statistics*, 23(5), 1630–1661.
- ▣ Rogers, L.C.G., Satchell, S.E. (1991). Estimating variance from high, low and closing prices. *Annals of Applied Probability*, 1(4), 504–512.
- ▣ Sortino, F.A., van der Meer, R. (1991). Downside risk. *Journal of Portfolio Management*, 17(4), 27–31.
- ▣ Yang, D., Zhang, Q. (2000). Drift-independent volatility estimation based on high, low, open, and close prices. *Journal of Business*, 73(3), 477–492.