








Modelling Financial Markets

Chapter 5: Models of Conditional Volatility



Daniel Traian PELE

Bucharest University of Economic Studies
IDA Institute Digital Assets
Blockchain Research Center
AI4EFin Artificial Intelligence for Energy Finance
Romanian Academy, Institute for Economic Forecasting
MSCA Digital Finance


Can We Forecast Tomorrow's Risk?

- Start with the evidence: large price changes tend to follow large changes, even when their signs are unpredictable
- Build a model of conditional variance, then interpret its reaction to news and its persistence
- Estimate the model and check whether its residuals still contain predictable patterns
- Compare forecasts on future observations, with a stated loss function and uncertainty measure
- ARCH means AutoRegressive Conditional Heteroskedasticity: variance depends on past shocks
- For further study: derivations and additional results in the Appendix

Core Path and Deeper Study

- **Core path**: the modelling and forecasting topics listed below; it is all you need for the in-class part of Seminar 5 (A1, B1–B3, C1)
  - Clustering and the ARCH tests; the GARCH recursion, persistence and half-life; estimation with Student-t innovations; forecasts; asymmetry (GJR, EGARCH); residual diagnostics; forecast evaluation with QLIKE, Mincer–Zarnowitz and Diebold–Mariano
- Inference beyond the basic model
  - QMLE asymptotics, the delta method near $\alpha + \beta = 1$, tests on the boundary, corrected residual tests, Giacomini–White and the Model Confidence Set
  - Practised at home in Seminar 5, A2 (delta method), B4 (boundary bootstrap) and B5 (forecast comparison)
- The two sections after the extensions (a quasi score-driven volatility model and an AI-assisted discovery loop) are optional research material for projects
- Proofs, full tables and settings are collected at the end of the chapter

Volatility Comes in Bursts



Amsterdam Stock Exchange, 21 October 1987, after the crash of 19 October


Photo: Bart Molendijk / Anefo (1987); CC0; Wikimedia Commons

- Crashes are not confined to isolated days: large moves are followed by large moves
  - "Large changes tend to be followed by large changes, of either sign" Mandelbrot (1963)
- This **volatility clustering** is stylised fact 6 of Chapter 1
- Volatility enters every risk number: Value at Risk (Chapter 7), portfolio weights (Chapter 4), option prices
- Questions
  - can we turn the clustering into a **forecast** of tomorrow's risk?
  - can we conduct valid inference on that forecast?

Reading the Market Charts

- S&P 500: Standard & Poor's index of large companies in the United States
- BET (Bucharest Exchange Trading) tracks Romanian stock prices
- BET-TR (Total Return) also includes reinvested dividends
- BNR: Banca Națională a României, the source of the reference exchange rate
- COVID-19: Coronavirus Disease 2019, used to label the pandemic shock in the charts
- EUR/RON: euros quoted in Romanian lei
- USD: United States dollar; p.a. in chart labels means per annum
- XAU: the code for one troy ounce of gold
- ACF (AutoCorrelation Function): correlation with past values at each lag
- i.i.d.: independent and identically distributed observations

Volatility Clusters in Every Market




- Daily log returns $r_t = 100\,(\ln P_t - \ln P_{t-\mathbf{1}})$ in %, closes up to 18 September 2026; S&P 500 and BET from 2000, Bitcoin (BTC) from 17 September 2014
  - P_t: closing price (index level) on day t; r_t: log return from close $t-1$ to close t, expressed in percent
- Equity indices: weekdays; days whose close equals the previous close (holidays filled with the last price) are dropped: 3 for the S&P 500, 0 for the BET; Bitcoin: 7 days a week
- Quiet years alternate with turbulent months in all three: 2008, March 2020, April 2025

MFM_ch5_arch_effects

Log Returns versus Simple Returns

- A simple return measures the relative change in price: $100(P_t/P_{t-1} - 1)$
- The log return is $100 \ln(P_t/P_{t-1})$: log returns add across consecutive dates
- Example: a price falling from 100 to 88 has a simple return of -12.0% and a log return of $100 \ln(0.88) \approx -12.8\%$
- For small changes they are close; during crashes the difference matters. The charts and models here use log returns

The Sign Is Unpredictable, the Size Is Not



- Sample autocorrelation at lag k: $\hat{\rho}_k = \widehat{\text{Corr}}(x_t, x_{t-k})$, computed for $x_t = r_t$ and $x_t = r_t^2$
  - Band $\pm 1.96/\sqrt{n}$ (n: number of returns): 95% range of $\hat{\rho}_k$ if the series were i.i.d.
- S&P 500, lag 1: -0.099 for r_t, 0.313 for r_t^2; all 50 autocorrelations of r_t^2 up to lag 50 lie outside ± 0.024
- Heavy tails too: excess kurtosis (kurtosis minus 3) 10.7 (S&P 500), 11.1 (BET), 12.0 (Bitcoin); 0 for the Normal distribution
- Object to model: $\sigma_t^2 = \text{Var}(r_t \mid \mathcal{F}_{t-1})$, with $\mathcal{F}_{t-1}$ the information at the close of day $t-1$

MFM_ch5_arch_effects

Testing for ARCH Effects (1/2): the Two Tests

- H_0: no clustering, $\mathrm{Var}(r_t \mid \mathcal{F}_{t-1})$ constant; both tests look for autocorrelation in squared returns
- **McLeod–Li** McLeod & Li (1983): the Ljung–Box statistic applied to r_t^2
  - $Q(m) = n(n+2)\sum_{k=1}^{m} \dfrac{\hat{\rho}_k^2}{n-k}$
  - $\hat{\rho}_k$: autocorrelation of r_t^2 at lag k; n: sample size; $m = 10$ lags
  - Measures the joint size of the first m autocorrelations; $Q \geq 0$, χ^2_{10} under H_0; reject if $Q(10) > 18.3$ (5%)
- **Engle: Lagrange Multiplier (LM) test** Engle (1982): regress $\hat{\varepsilon}_t^2$ on a constant and $\hat{\varepsilon}_{t-1}^2, \ldots, \hat{\varepsilon}_{t-5}^2$
  - $\hat{\varepsilon}_t = r_t - \bar{r}$: demeaned return, $\bar{r}$: sample mean
  - $LM = nR^2$, with R^2 the coefficient of determination of this regression; H_0: all five slopes are 0
  - χ^2_5 under H_0; reject if $LM > 11.1$ (5%); a large LM means past squared shocks predict today's

Testing for ARCH Effects (2/2): Results

- $\hat{\rho}_1(r^2)$: first-order autocorrelation of squared returns
- Every statistic is far above its critical value (all $p < 0.001$): clustering in equities, crypto, a managed exchange rate and gold; weakest for Bitcoin and gold
- One-sided score tests: Appendix

Series	$Q(10)$ on r^2	LM(5)	$\hat{\rho}_1(r^2)$
S&P 500	6 133	1 684	0.313
BET	2 440	982	0.330
BET-TR	714	371	0.171
Bitcoin	213	115	0.137
EUR/RON	2 420	871	0.342
Gold	744	329	0.110

MFM_ch5_arch_effects

The ARCH Model of Engle (1982)

- Setting: $r_t = \mu + \varepsilon_t$, $\varepsilon_t = \sigma_t z_t$, innovations z_t i.i.d. with mean 0 and variance 1
  - μ: mean return; ε_t: shock (unexpected part of the return); σ_t: conditional volatility
  - Conditional variance $\sigma_t^2 = \mathrm{Var}(r_t \mid \mathcal{F}_{t-1})$; unconditional variance $\sigma^2 = \mathbb{E}\sigma_t^2$
- **ARCH**(q): today's variance is a constant plus a weighted sum of the last q squared shocks Engle (1982)
  - $\sigma_t^2 = \omega + \alpha_1 \varepsilon_{t-1}^2 + \cdots + \alpha_q \varepsilon_{t-q}^2$
  - q: number of lags; ω: variance floor; α_i: weight of the shock i days ago
  - $\omega > 0$, $\alpha_i \geq 0$: the variance is always positive
- A large shock yesterday (ε_{t-1}^2 large) raises today's variance: clustering by construction
- ARCH(1): unconditional variance $\sigma^2 = \omega/(1-\alpha_1)$, finite if $\alpha_1 < 1$
- Engle received the 2003 Nobel Prize for ARCH (history: Appendix)

From ARCH to GARCH (Bollerslev, 1986)

- GARCH means Generalised AutoRegressive Conditional Heteroskedasticity: the variance also depends on its own past
- Problem with ARCH: clustering in real data lasts for months, so q must be large (many parameters)
- **GARCH**(1,1) Bollerslev (1986): $\sigma_t^2 = \omega + \alpha\,\varepsilon_{t-1}^2 + \beta\,\sigma_{t-1}^2$
  - Today's variance = constant + reaction to yesterday's squared shock + part of yesterday's variance
  - $\omega > 0$, $\alpha \geq 0$, $\beta \geq 0$; α: reaction to news; β: memory of past variance
- Substituting backwards: $\sigma_t^2 = \frac{\omega}{1-\beta} + \alpha \sum_{j \geq 1} \beta^{j-1} \varepsilon_{t-j}^2$
  - Valid for $\beta < 1$; j: age of the shock in days, weight $\alpha\beta^{j-1}$
  - An ARCH(∞) with geometrically declining weights, using only three parameters
- GARCH(p,q) adds more lags (p lags of σ^2, q lags of ε^2); in practice GARCH(1,1) is the workhorse Hansen & Lunde (2005)

Model Labels and Innovations

- The model name describes the variance dynamics; the suffix describes the law of the standardised innovation
- GARCH-N: GARCH with innovations from the Normal distribution
- GARCH-t: GARCH with Student-t innovations, which allow heavier tails
- GJR-GARCH (Glosten–Jagannathan–Runkle) allows a different variance response to negative shocks
- GJR-t and GJR-GARCH-t denote the same model here: asymmetric variance dynamics and Student-t innovations

GARCH(1,1) as an ARMA Model for Squared Shocks

- Define $\nu_t = \varepsilon_t^2 - \sigma_t^2$, the surprise in the squared shock ($\mathbb{E}[\nu_t \mid \mathcal{F}_{t-1}] = 0$)
- Then $\varepsilon_t^2 = \omega + (\alpha + \beta)\,\varepsilon_{t-1}^2 + \nu_t - \beta\,\nu_{t-1}$
  - Obtained by adding ν_t to both sides of the GARCH recursion and writing $\sigma_{t-1}^2 = \varepsilon_{t-1}^2 - \nu_{t-1}$
  - An AutoRegressive Moving Average model, ARMA(1,1), in ε_t^2 with autoregressive coefficient $\alpha + \beta$ and moving-average coefficient $-\beta$
- Consequence: the autocorrelations of ε_t^2 decay geometrically, $\rho_k = \rho_1(\alpha + \beta)^{k-1}$
  - $\rho_k = \mathrm{Corr}(\varepsilon_t^2, \varepsilon_{t-k}^2)$: autocorrelation of squared shocks at lag k
  - $\rho_1 = \alpha(1 - \beta^2 - \alpha\beta)/(1 - \beta^2 - 2\alpha\beta)$ if $\mathbb{E}\varepsilon_t^4 < \infty$ (Seminar 5, A6)
  - S&P 500, GARCH(1,1) with Normal innovations: $\rho_1 = 0.356$, $\rho_{10} = 0.298$; sample 0.313 and 0.248; the model values are check values for A6, which derives them
- $\alpha + \beta$ is therefore called the **persistence** of volatility


Stationarity and the Unconditional Variance

- **Covariance stationarity**: $\alpha + \beta < 1$; then
  - $\sigma^2 = \mathbb{E}[\sigma_t^2] = \dfrac{\omega}{1 - \alpha - \beta}$
- Derivation: take expectations in the recursion, $\mathbb{E}\sigma_t^2 = \omega + (\alpha + \beta)\,\mathbb{E}\sigma_{t-1}^2$, and set $\mathbb{E}\sigma_t^2 = \mathbb{E}\sigma_{t-1}^2$
- **IGARCH** (integrated GARCH): $\alpha + \beta = 1$; the unconditional variance is infinite
  - Yet, if $\alpha > 0$ and z_t^2 is non-degenerate, the process is still **strictly** stationary: the condition $\mathbb{E}\ln(\alpha z_t^2 + \beta) < 0$ Nelson (1990) holds by Jensen, since $\mathbb{E}\ln(\alpha z_t^2 + \beta) < \ln(\alpha + \beta) = 0$
- Notation: σ^2 is the model's long-run variance (Seminar 5 writes $\bar{\sigma}^2$), $\hat{\sigma}^2 = \hat{\omega}/(1 - \hat{\alpha} - \hat{\beta})$ its estimate; s^2 is the sample variance of r_t
  - They need not agree: S&P 500 long-run volatility 27.3% p.a. against a sample volatility of 19.2%
- Variance targeting: fix $\omega = s^2(1 - \alpha - \beta)$, so that the long-run variance equals s^2 and only α, β are estimated

Fourth Moment and Kurtosis

- With Normal z_t, the kurtosis of ε_t is $K = \dfrac{3\left[1-(\alpha+\beta)^2\right]}{1-(\alpha+\beta)^2-2\alpha^2}$
  - $K = \mathbb{E}\varepsilon_t^4/(\mathbb{E}\varepsilon_t^2)^2$: 3 for the Normal distribution, above 3 for heavy tails
  - Finite only if $(\alpha+\beta)^2 + 2\alpha^2 < 1$ (Seminar 5, A4)
  - ARCH(1) is the case $\beta = 0$: $K = 3(1-\alpha^2)/(1-3\alpha^2)$, finite only if $\alpha < 1/\sqrt{3}$: heavy tails even with Normal innovations
- S&P 500, GARCH(1,1) with Normal innovations: $\hat{\alpha} = 0.120604$, $\hat{\beta} = 0.859878$
  - $(\hat{\alpha}+\hat{\beta})^2 + 2\hat{\alpha}^2 = 0.990435$: just below 1, so $K = 12.1$ (a check value for A4); sample kurtosis of the returns: 13.7
- K is ill-conditioned near the moment boundary: report it from unrounded estimates (a rounding example: Appendix; Seminar 5, A4)
- Clustering alone generates much of the fat tails; the rest comes from heavy-tailed z_t

Simulation: GARCH Produces Clustering



- Top: i.i.d. Normal returns; bottom: GARCH(1,1) with $\omega = 0.02$, $\alpha = 0.10$, $\beta = 0.88$, the same unconditional variance (1), and the bands $\pm 2\sigma_t$
- Quiet stretches and bursts appear only in the GARCH path
- Normal innovations, yet fat-tailed returns: a mixture of Normal distributions with varying variances has heavy tails (kurtosis on a long path: Appendix)

MFM_ch5_garch_simulation

Persistence and Half-Life

- Multi-step forecast: $\mathbb{E}[\sigma^2_{t+h} \mid \mathcal{F}_t] = \sigma^2 + (\alpha+\beta)^{h-1}(\sigma^2_{t+1} - \sigma^2)$
  - h: forecast horizon in days; σ^2: unconditional (long-run) variance; σ^2_{t+1}: one-day-ahead variance, known at t
  - The gap to the long-run variance shrinks by the factor $\alpha+\beta$ each day
- **Half-life**: number of days after which half of a variance shock has gone
  - $(\alpha+\beta)^h = 0.5 \;\Rightarrow\; h_{1/2} = \dfrac{\ln 0.5}{\ln(\alpha+\beta)}$
  - $h_{1/2}$ grows without bound as $\alpha+\beta \to 1$
- Examples: $\alpha+\beta = 0.95$: 13.5 days; 0.99: 69 days; 0.995: 138 days
- Small changes in $\alpha+\beta$ near 1 imply very different half-lives: estimation uncertainty matters (next section; Seminar 5, A2 and B4)

A Toy Check: Multi-Day Risk After a Turbulent Day

- Toy GARCH(1,1), not fitted to any data: $\omega = 0.04$, $\alpha = 0.08$, $\beta = 0.90$
  - Persistence $p = \alpha + \beta = 0.98$; long-run variance $\sigma^2 = 0.04/0.02 = 2$; half-life $\ln 0.5 / \ln 0.98 \approx 34.3$ days
- A turbulent day: $\sigma^2_{t+1} = 4$, twice the long-run level
  - Day 2: $\mathbb{E}[\sigma^2_{t+2} \mid \mathcal{F}_t] = 2 + 0.98 \times (4 - 2) = 3.96$: the gap to σ^2 shrinks by 2% a day
- H-day variance: daily shocks are uncorrelated, so the variance of their sum is the sum of the daily variance forecasts, a geometric sum
  - $V_H = \sum_{h=1}^{H} \mathbb{E}[\sigma^2_{t+h} \mid \mathcal{F}_t] = H\sigma^2 + (\sigma^2_{t+1} - \sigma^2)\,\dfrac{1 - p^H}{1 - p}$
  - $H = 5$: $V_5 = 10 + 2 \times (1 - 0.98^5)/0.02 = 10 + 2 \times 4.804 = 19.61$; the square-root-of-time rule gives $5 \times 4 = 20$, too high
- Your turn: redo $H = 5$ after a calm day, $\sigma^2_{t+1} = 1$ (answer: $V_5 = 10 - 4.804 = 5.20 > 5$, so the rule is now too low, as in the S&P 500 worked example)

EWMA and IGARCH: No Mean Reversion

- **Exponentially Weighted Moving Average (EWMA)** J.P. Morgan/Reuters (1996): $\sigma_t^2 = \lambda \sigma_{t-1}^2 + (1-\lambda) r_{t-1}^2$, with $\lambda = 0.94$ for daily data and zero-mean returns
  - $\lambda \in (0,1)$: decay factor; a larger λ means a longer memory
  - Weights $(1-\lambda)\lambda^{j-1}$ on past squared returns: the most recent days count most
  - EWMA is an IGARCH(1,1) with $\omega = 0$, $\alpha = 1-\lambda$, $\beta = \lambda$: no parameter to estimate
- Multi-step forecasts at the close of day t differ
  - $\mathbb{E}_t[\cdot] = \mathbb{E}[\cdot \mid \mathcal{F}_t]$: expectation given the information at t
  - EWMA: flat, $\mathbb{E}_t \sigma_{t+h}^2 = \sigma_{t+1}^2$ for every h
  - IGARCH with $\omega > 0$: rising, $\mathbb{E}_t \sigma_{t+h}^2 = \sigma_{t+1}^2 + (h-1)\omega$
  - Stationary GARCH: pulled towards σ^2 at the rate $\alpha + \beta$
- EWMA serves as a benchmark in the forecast comparison

Maximum Likelihood

- Maximum Likelihood Estimation (MLE) chooses the parameters that maximise the likelihood
- Parameters $\theta = (\mu, \omega, \alpha, \beta)$; given θ, the recursion produces $\sigma_1^2(\theta), \ldots, \sigma_n^2(\theta)$
- Normal log-likelihood: $\ell(\theta) = -\frac{1}{2} \sum_{t=1}^{n} \left[\ln 2\pi + \ln \sigma_t^2 + \varepsilon_t^2 / \sigma_t^2 \right]$
  - $\ell(\theta)$: log of the joint density of the n observed returns, read as a function of θ; $\hat{\theta}$ maximises it; $\varepsilon_t = r_t - \mu$
  - Each day contributes the density of ε_t given the past: the likelihood factorises in time
- No closed form: numerical optimisation under $\omega > 0$, $\alpha, \beta \geq 0$, $\alpha + \beta \leq 1$
  - The `arch` package imposes $\alpha + \beta \leq 1$, a closed set: an estimate can sit on the bound (Bitcoin and EUR/RON: 1.0000 in the six-market table)
- Start-up: σ_1^2 is set to the sample variance (its effect fades at rate β)
- Scale: returns in %, so that the variance is of order 1
  - EUR/RON (daily standard deviation about 0.28%) is fitted on 10 r_t: unscaled, the optimiser stopped at a local optimum

Heavy-Tailed Innovations

- **Student-t** Bollerslev (1987): $z_t = \sqrt{(\nu-2)/\nu}\, t_\nu$, unit variance for $\nu > 2$
  - t_ν: a standard Student-t variable, variance $\nu/(\nu-2)$; the factor rescales it to variance 1
  - ν (degrees of freedom): the smaller ν, the heavier the tails; the fourth moment exists only for $\nu > 4$
- **Skewed-t** Hansen (1994): adds an asymmetry parameter $\lambda \in (-1, 1)$
  - $\lambda < 0$: the left tail is longer (large losses more likely than large gains)
- **Generalised Error Distribution (GED)** Nelson (1991): density $\propto \exp(-\frac{1}{2}|z/\lambda_\nu|^\nu)$
  - GED: generalised error distribution; $\nu > 0$: shape parameter; λ_ν: scale constant
  - $\nu = 2$: the Normal distribution; $\nu = 1$: Laplace; $\nu < 2$: heavier tails (λ_ν gives unit variance)

Gaussian QMLE (1/2): Definition and Consistency

- **Gaussian QMLE** (Quasi-Maximum Likelihood Estimator): maximise the Normal likelihood even if z_t is not Normal Bollerslev & Wooldridge (1992)
  - The Normal $\ell(\theta)$ is used as a criterion, not as a claim about the law of z_t
  - θ_0: true parameter value; $\hat{\theta}$: the estimate, the maximiser of ℓ
- **Consistency** (GARCH(1,1)): $\hat{\theta} \to \theta_0$ as $n \to \infty$ Berkes, Horváth & Kokoszka (2003), Francq & Zakoïan (2004)
  - Strict stationarity: $\mathbb{E} \ln(\alpha z_t^2 + \beta) < 0$
  - θ_0 in a compact (closed and bounded) parameter set; z_t^2 non-degenerate (not a constant)
  - No moment condition on ε_t: consistency holds even for IGARCH

Gaussian QMLE (2/2): Asymptotic Normality and Standard Errors

- **Asymptotic normality** of $\vartheta = (\omega, \alpha, \beta)$ (pure GARCH, known mean): add ϑ_0 interior and $\kappa_z = \mathbb{E}z_t^4 < \infty$
  - $\sqrt{n}(\hat{\vartheta} - \vartheta_0) \to N\left(0, (\kappa_z - 1)J^{-1}\right)$, $J = \mathbb{E}[g_t g_t']$, $g_t = \partial \ln \sigma_t^2 / \partial \vartheta$
  - In words: the estimation error shrinks like $1/\sqrt{n}$ and is approximately Normal
  - κ_z: kurtosis of z_t (3 if Normal); g_t: gradient of $\ln \sigma_t^2$; J: its mean outer product; interior: no constraint binds
- Sandwich $\hat{A}^{-1}\hat{B}\hat{A}^{-1}$ estimates $(\kappa_z - 1)J^{-1}$; the inverse Hessian $\hat{A}^{-1}$ estimates $2J^{-1}$
  - A: minus the expected Hessian (second derivatives) of ℓ/n; B: variance of the score (first derivatives) per observation
  - Robust / classical standard error (s.e.) $= \sqrt{(\kappa_z - 1)/2}$: 1 under Normality, above 1 with heavy tails
- Variants (estimated mean, IGARCH, explosive case): ▸ Appendix

QMLE in Our Data: When the Theory Holds and When It Fails

- S&P 500, GARCH(1,1) with Normal likelihood: residual kurtosis $\hat{\kappa}_z = 4.73$
  - Theory: robust / classical s.e. $= \sqrt{(4.73-1)/2} = 1.37$ (a check value for A5); observed: 1.37 for $\hat{\alpha}$, 1.37 for $\hat{\beta}$
- Bitcoin: GARCH-t (suffix -t: Student-t innovations, previous slides) gives $\hat{\nu} = 3.18 < 4$ degrees of freedom: under that law $\mathbb{E}z_t^4 = \infty$, and the condition for Gaussian-QMLE normality fails
  - Gaussian QMLE residual kurtosis does not settle: 19.8 on the first half, 14.6 on the full sample
  - Full sample, robust / classical s.e.: 2.51 for $\hat{\alpha}$, 2.01 for $\hat{\beta}$, against the formula value 2.61, which assumes $\kappa_z < \infty$
- Hall & Yao (2003): then the Gaussian QMLE converges slower than $\sqrt{n}$, to a non-Gaussian limit: its robust t-statistics are not valid
- Remedies: a heavy-tailed quasi-likelihood with its scale adjustment Fan, Qi & Xiu (2014), or the Student-t MLE (bounded score: no fourth moment needed if the law is right), with bootstrap inference near bounds

MFM_ch5_inference


S&P 500: GARCH(1,1)-t Estimates

Parameter	Estimate	Robust s.e.	Classical s.e.	Ratio
ω	0.0157			
α	0.123	0.0107	0.0101	1.06
β	0.872	0.0103	0.0096	1.08
ν	6.26	0.48		
$\alpha + \beta$	0.9947			

- Daily returns in %, 2000–2026; mean $\hat{\mu} = 0.079\%$ per day; ratios use the unrounded s.e.; full table with every standard error: Seminar 5, B1
- $\hat{\beta}$: most of today's variance is yesterday's variance; $\hat{\alpha}$: news adds a moderate update; $\hat{\nu} = 6.26$: tails clearly heavier than those of the Normal distribution
- Half-life 131 days; long-run volatility $\sqrt{251.4\,\hat{\omega}/(1-\hat{\alpha}-\hat{\beta})} = 27.3\%$ p.a. ($A = 251.4$ returns a year), above the sample 19.2%: dividing by $1-\hat{\alpha}-\hat{\beta}$ makes it imprecise (next slide)

MFM_ch5_garch_estimation

Near-Unit Persistence: What the Delta Method Can and Cannot Do (1/2)



MFM_ch5_seminar

Near-Unit Persistence: What the Delta Method Can and Cannot Do (2/2)

- $g(\theta)$: a smooth function of the parameters (long-run volatility, half-life); ∇g: its gradient; $p = \alpha + \beta$, $h = \ln 0.5 / \ln p$
- Delta method: $\widehat{\text{s.e.}}\, g(\hat{\theta}) \approx \sqrt{\nabla g' \hat{V} \nabla g}$, $\hat{V}$ the robust covariance of $(\hat{\omega}, \hat{\alpha}, \hat{\beta})$; long-run volatility 27.3% p.a., s.e. 10.8 points
- Half-life: $\partial h / \partial p = 24\,789$ at $\hat{p} = 0.994712$, $\widehat{\text{s.e.}}(\hat{p}) = 0.0047$, so $\widehat{\text{s.e.}}(\hat{h}) \approx 116$ days, and $\hat{h} \pm 1.96\,\widehat{\text{s.e.}}$ includes negative half-lives: the linearisation fails
- Transform the interval for p instead: $[0.9855; 1.0039]$ gives $h \in [48; \infty)$; whether $p = 1$ is plausible needs a boundary test (next slides)

MFM_ch5_seminar

Testing on the Boundary (1/2): a Decision Table

Situation under H_0	**Example**	**Likelihood Ratio (LR):** $LR = 2(\ell_1 - \ell_0)$	**5% critical value**
Interior restriction	GJR (asymmetric GARCH) vs GARCH, $\gamma = 0$ (with $\alpha > 0$)	χ^2_1	3.84
One parameter on the boundary, the others identified and interior	Normal vs Student-t ($1/\nu = 0$); IGARCH ($\alpha + \beta = 1$)	$\frac{1}{2}\chi^2_0 + \frac{1}{2}\chi^2_1$ Self & Liang (1987), Andrews (2001)	2.71
A nuisance parameter unidentified	No ARCH: $\alpha = 0$ leaves β unidentified	non-standard Andrews (2001)	simulate
Several bounds active at once	GJR with $\hat{\alpha} = 0$, test of $\gamma = 0$	non-standard	simulate

- ℓ_1, ℓ_0: maximised log-likelihoods without and with the restriction; $\frac{1}{2}\chi^2_0 + \frac{1}{2}\chi^2_1$: $LR = 0$ with probability $\frac{1}{2}$, otherwise χ^2_1
- GJR (full treatment in the asymmetry section): $\sigma^2_t = \omega + (\alpha + \gamma I_{t-1})\,\varepsilon^2_{t-1} + \beta\,\sigma^2_{t-1}$, $I_{t-1} = 1$ if $\varepsilon_{t-1} < 0$: γ is the extra reaction to bad news

Testing on the Boundary (2/2): Why IGARCH Is a Boundary Case

- IGARCH is on the boundary only because the fit restricts $\alpha + \beta \leq 1$, as `arch` does
  - Under H_0: $\alpha + \beta = 1$ the unrestricted estimate would exceed 1 about half of the time; the restriction sets it to 1 and then $LR = 0$: hence the mixture and 2.71
  - Without that restriction, $\alpha + \beta = 1$ lies inside the strictly stationary region, the QMLE is still asymptotically Normal there Lumsdaine (1996), and the LR would be compared with χ^2_1 (3.84)
- Non-nested models: **Akaike Information Criterion (AIC)** $= -2\ell + 2k$, **Bayesian Information Criterion (BIC)** $= -2\ell + k \ln n$ (k parameters); lower is better
- A Wald t for an estimate at the bound (e.g. $\hat{\alpha} = 0$) is not $N(0, 1)$; a parametric bootstrap from $\hat{\theta}$ is inconsistent when θ_0 is on the boundary Andrews (2000): simulate under H_0 instead

Boundary Tests on Our Data

- Normal vs Student-t innovations, GARCH(1,1): $LR = 311.0$ (S&P 500), 1 082.0 (Bitcoin; a check value for B6)
  - Far above 2.71: heavy tails are decisive; the mixture matters only in borderline cases
- GJR-t (the GJR recursion two slides back), S&P 500: $\hat{\alpha} = 0.000$; its t-test is not standard, and the χ^2_1 law of the LR for $\gamma = 0$ assumes $\alpha_0 > 0$
- IGARCH, $H_0 : \alpha + \beta = 1$ (GARCH-t): $LR = 1.11$ (S&P 500), 1.14 (BET)
  - S&P 500: $p = 0.29$ (χ^2_1), 0.15 (mixture), 0.17 (parametric bootstrap under H_0, $B = 999$ simulated samples)
  - BET: $p = 0.28$, 0.14 and 0.17 (check values for B8)
- Under H_0, 52% of the re-estimates have $\hat{\alpha} + \hat{\beta} \geq 0.999$ (at the bound $\alpha + \beta = 1$), as the mixture predicts: a unit root in variance cannot be rejected for either index

MFM_ch5_seminar

Worked Example (1/2): The One-Day Forecast

- S&P 500, GARCH(1,1)-t: $\hat{\omega} = 0.0157$, $\hat{\alpha} = 0.123$, $\hat{\beta} = 0.872$, $\hat{\mu} = 0.079$
  - All results below use the unrounded estimates, in particular $\hat{\rho} = \hat{\alpha} + \hat{\beta} = 0.994712$
  - With the rounded values shown, $1 - \hat{\alpha} - \hat{\beta} = 0.005$ and the long-run variance would be 3.14 instead of 2.97: near $\hat{\rho} = 1$, rounding matters
- Last day $T =$ 18.09.2026: $\hat{\varepsilon}_T = r_T - \hat{\mu} = 0.088$, filtered variance $\hat{\sigma}^2_T = 0.565$
  - Filtered variance: the recursion run through the whole sample with $\hat{\theta}$, read off on the last day T
  - $\hat{\sigma}^2_{T+1} = \hat{\omega} + \hat{\alpha}\hat{\varepsilon}^2_T + \hat{\beta}\hat{\sigma}^2_T = \hat{\omega} + 0.0009 + 0.493 = 0.509$, i.e. 11.3% p.a.
- A quiet last day: $\hat{\sigma}^2_{T+1}$ is far below the long-run level, so the forecasts will rise (next slide)


MFM_ch5_garch_estimation

Worked Example (2/2): Ten Days Ahead

- Long-run variance $\hat{\sigma}^2 = \hat{\omega}/(1-\hat{\alpha}-\hat{\beta}) = 2.97$; day 10: $\hat{\sigma}^2 + (\hat{\alpha}+\hat{\beta})^9(\hat{\sigma}^2_{T+1} - \hat{\sigma}^2)$
  - $= 2.97 + 0.953 \times (0.509 - 2.97) = 0.624$
- 10-day variance $\sum_{h=1}^{10} \hat{\sigma}^2_{T+h} = 5.67$, volatility 2.38%
  - The sum in closed form (toy check in the persistence section): $10\,\hat{\sigma}^2 + (\hat{\sigma}^2_{T+1} - \hat{\sigma}^2)\,(1-\hat{\rho}^{10})/(1-\hat{\rho})$
  - $= 29.72 + (0.509 - 2.972) \times 9.765 = 5.67$
  - The square-root-of-time rule gives $\sqrt{10} \times 0.714 = 2.26\%$: too low, because $\hat{\sigma}^2_{T+1} < \hat{\sigma}^2$ and volatility is forecast to rise

MFM_ch5_garch_estimation

Volatility Term Structure



- Average volatility over the next h days, forecast from 18.09.2026 (GARCH(1,1)-t); dotted: long-run level
- S&P 500 rises from 11.3% to 20.1% in a year (below its long run today); gold falls from 23.9% to 22.4%
- Bitcoin, EUR/RON: IGARCH with $\hat{\omega} > 0$, the variance forecast grows by $\hat{\omega}$ per day (Bitcoin: 108.3% in a year): an artefact of the unit-root estimate

MFM_ch5_persistence


GARCH(1,1)-t in Six Markets

Series	$\hat{\omega}$	$\hat{\alpha}$	$\hat{\beta}$	$\hat{\alpha}+\hat{\beta}$	$h_{1/2}$ (days)	$\hat{\nu}$	Long-run vol.
S&P 500	0.0157	0.123	0.872	0.9947	131	6.26	27.3
BET	0.0448	0.192	0.799	0.9907	74	5.01	34.7
BET-TR	0.0755	0.206	0.713	0.9188	8	5.02	15.2
Bitcoin	0.1937	0.109	0.891	1.0000	∞	3.18	–
EUR/RON	0.0000	0.123	0.877	1.0000	∞	4.20	–
Gold	0.0107	0.049	0.944	0.9935	106	4.58	20.7

- Long-run volatility in % p.a., each series annualised with its own frequency; – when $\hat{\alpha}+\hat{\beta} \geq 0.999$; EUR/RON: $\hat{\omega}$ is converted back from the fit on $10\,r_t$ (divided by 100) to the $\%^2$ units of the other rows, and is 0.0000 at four decimals
- Equities and gold: persistence 0.99–0.995; the BET reacts more to news ($\hat{\alpha} = 0.192$) and forgets faster than the S&P 500
- Different samples: BET-TR starts in 2014, so its lower persistence is not comparable with the BET 2000–2026; Bitcoin: $\hat{\nu} < 4$, an infinite fourth moment under the fitted law
- EUR/RON, a managed rate: on 6 May 2025 a move of $+1.20\%$ after days of $\pm 0.004\%$ gives a standardised residual of 154

MFM_ch5_garch_estimation


Persistence: How Fast Do Shocks Fade?



- $(\hat{\alpha}+\hat{\beta})^h$: the share of a variance shock left after h days; dashed line: one half
- Half-lives: column $h_{1/2}$ of the six-market table; in Bitcoin and EUR/RON shocks never fade in the model
- Level shifts in volatility can mimic high persistence Lamoureux & Lastrapes (1990) (regime switching: Hamilton & Susmel (1994)): report half-lives as intervals (Seminar 5, B4)

MFM_ch5_persistence


S&P 500: GARCH Volatility and the VIX



- $\hat{\sigma}_t\sqrt{A}$ ($A = 251.4$ returns a year) from GARCH(1,1)-t and the VIX (Volatility Index of the Chicago Board Options Exchange, 30-day implied volatility), 2000–2026: correlation 0.89
- The VIX is higher on average: 19.8% against 17.0%, above GARCH on 84% of the days
- Not a measured variance risk premium: the premium compares VIX^2 with a 30-day variance forecast under the real-world probability; $\hat{\sigma}_t$ is a one-day volatility

MFM_ch5_sp500_vix

BET, Bitcoin, EUR/RON and Gold



- GJR-GARCH-t volatility, each series annualised with its own frequency (Bitcoin: 365 days)
- BET: 2008 dominates, now 26.4%; Bitcoin: 50–100% is normal, the highest peak is March 2020
- EUR/RON: a managed calm with bursts in 2022 and 2025, poorly described by a daily GARCH; gold: a new high of 53% in 2026

MFM_ch5_markets


GJR-GARCH (Glosten, Jagannathan and Runkle, 1993)

- Bad news raises future volatility more than good news of the same size (stylised fact 9, Chapter 1); GARCH(1,1) cannot see the sign: ε_{t-1}^2 is the same for -3 and $+3$
- GJR: a negative shock gets the extra slope γ Glosten, Jagannathan & Runkle (1993)
  - $\sigma_t^2 = \omega + (\alpha + \gamma\, I_{t-1})\,\varepsilon_{t-1}^2 + \beta\,\sigma_{t-1}^2$, with $I_{t-1} = 1$ if $\varepsilon_{t-1} < 0$, else 0
  - Good news: slope α; bad news: slope $\alpha + \gamma$; positivity: $\omega > 0$, $\alpha \geq 0$, $\alpha + \gamma \geq 0$, $\beta \geq 0$ (γ may be negative)
  - Persistence $\alpha + \gamma/2 + \beta$ for symmetric z_t; for skewed z_t replace $1/2$ by $\mathbb{E}[z_t^2 1(z_t < 0)]$, $1(\cdot)$ the indicator (Seminar 5, A3)
- S&P 500: $\hat{\alpha} = 0.000$, $\hat{\gamma} = 0.205$ (t = 10.0), $\hat{\beta} = 0.881$; symmetry test: H_0: $\gamma = 0$ by t or by LR against GARCH(1,1)
- Candidate mechanisms, not tested here: leverage (a falling price raises the debt-to-equity ratio) and volatility feedback (higher expected volatility raises required returns, so prices fall now)

EGARCH (Nelson, 1991)

- Exponential GARCH (EGARCH) models the log of the variance, driven by the size and the sign of yesterday's standardised shock Nelson (1991)
  - $\ln \sigma_t^2 = \omega + \alpha\left(|z_{t-1}| - \mathbb{E}|z_{t-1}|\right) + \gamma\, z_{t-1} + \beta \ln \sigma_{t-1}^2$
  - $z_{t-1} = \varepsilon_{t-1}/\sigma_{t-1}$: standardised shock; $|z_{t-1}| - \mathbb{E}|z_{t-1}|$: its size above the average size
  - Size effect α, sign effect γ; $\gamma < 0$: negative shocks raise volatility more
- The log form keeps $\sigma_t^2 > 0$ without any sign restriction on the parameters
- Persistence: β (stationary if $|\beta| < 1$)
- S&P 500: $\hat{\alpha} = 0.138$, $\hat{\gamma} = -0.165$ (t $= -15.2$), $\hat{\beta} = 0.979$
- In `arch`, $\mathbb{E}|z|$ is set to $\sqrt{2/\pi}$, its value under the Normal distribution


The News Impact Curve for the S&P 500



- **News impact curve** Engle & Ng (1993): σ_t^2 as a function of ε_{t-1}, with σ_{t-1}^2 fixed at the sample variance $s^2 = 1.47$ (%^{2}), not at the long-run variance of the model
- Shock of -3%: GJR 3.16, EGARCH 2.76; $+3$%: GJR 1.31, EGARCH 1.22; GARCH: 2.40 for both
- After bad news the forecast is 2.4 times (GJR) and 2.3 times (EGARCH) that after good news: ratios of levels, not of increases

MFM_ch5_asymmetry

Zoom: The COVID-19 Crash, 2020



- COVID-19 (Coronavirus Disease 2019), January–July 2020: $|r_t|$ annualised, GARCH-t and GJR-GARCH-t volatility, VIX
- 24 February ($r_t = -3.41\%$): the VIX went from 17.1% to 25.0% the same day; GARCH-t from 12.2% to 22.5% only the next day: GARCH reacts, markets price at once
- After $r_t = -12.8\%$ on 16 March, peaks on 17 March: GARCH-t 111%, GJR-t 127%, higher because the shocks were negative; full-sample (2000–2026) parameters; a forecast with parameters estimated before 2020 (as in Seminar 5, B5) gives a different peak

MFM_ch5_sp500_vix

Asymmetry in Six Markets

Series	$\hat{\gamma}_{GJR}$	t	LR vs GARCH	p-value	$\hat{\gamma}_{EGARCH}$	t
S&P 500	0.205	10.0	235.7	< 0.001	-0.165	-15.2
BET	0.043	2.1	4.9	0.026	-0.028	-2.6
BET-TR	0.111	2.9	10.3	0.001	-0.063	-3.5
Bitcoin	-0.013	-0.7	0.7	0.416	0.015	1.2
EUR/RON	0.039	3.0	9.3	0.002	-0.040	-2.5
Gold	-0.043	-4.7	27.6	< 0.001	0.038	5.7

- Student-t innovations; robust t; $LR \sim \chi^2_1$ (3.84) if the Student-t density is correct; the robust t is $N(0,1)$ only if the other parameters are interior (not for $\hat{\alpha} = 0$)
- Bad news weighs more (GJR $\hat{\gamma} > 0$, EGARCH $\hat{\gamma} < 0$): S&P 500 strongly, BET and BET-TR weakly, EUR/RON slightly; Bitcoin: none detected; gold: inverted sign
- Leverage for equities and a safe-haven role for gold are interpretations consistent with these signs, not tested mechanisms
- The BET row is the check value for Seminar 5, B3, which re-estimates it and explains the LR, AIC and BIC decisions

MFM_ch5_asymmetry

S&P 500: Twelve Models Compared by BIC

ΔBIC	$N(0,1)$	Student-t	Skewed-t	GED
GARCH(1,1)	648.9	346.7	307.4	339.8
GJR-GARCH(1,1)	394.7	119.9	49.6	141.6
EGARCH(1,1)	372.9	74.8	0.0	111.4


- ΔBIC: BIC $= -2\ell + k \ln n$ minus the lowest BIC among the 12 models (0 = preferred model)
- Two choices matter: Student-t instead of Normal innovations lowers BIC by about 300; asymmetry lowers it by a further 226.8–271.9 (Student-t column, GJR and EGARCH against GARCH); winner: EGARCH with skewed-t innovations
- BIC ranks in-sample fit, not forecast accuracy: the forecast section tests whether these gains survive out of sample


MFM_ch5_garch_estimation

Standardised Residuals and the Sign-Bias Tests

- **Standardised residual**: $\hat{z}_t = \hat{\varepsilon}_t / \hat{\sigma}_t$; if the model is right, $\hat{z}_t$ is close to i.i.d. with variance 1
- Checks, each with H_0: the model captures that feature
  - Mean equation: Ljung–Box $Q(10)$ on $\hat{z}_t$; variance equation: $Q(10)$ on $\hat{z}_t^2$ and LM(5); distribution: Quantile–Quantile (QQ) plot against the assumed density
- Sign-bias regression Engle & Ng (1993): $\hat{z}_t^2 = b_0 + b_1 S_{t-1}^- + b_2 S_{t-1}^- \hat{\varepsilon}_{t-1} + b_3 S_{t-1}^+ \hat{\varepsilon}_{t-1} + u_t$
  - $S_{t-1}^- = 1$ if $\hat{\varepsilon}_{t-1} < 0$, else 0; $S_{t-1}^+ = 1 - S_{t-1}^-$; u_t: regression error; the size terms use the unstandardised residual $\hat{\varepsilon}_{t-1}$ (eq. 18 in Engle & Ng (1993))
  - b_1: sign bias; b_2, b_3: negative and positive size bias; joint test $nR^2 \sim \chi_3^2$ (7.81 at 5%), conservative because it omits the gradient of the fitted variance

S&P 500: Are the Innovations Normal? (1/2)





Standardised residuals $\hat{z}_t$ (kernel density)
Normal N(0,1)
Student-t, $\nu = 6.3$ (unit variance)

vs Normal quantiles
vs Student-t ($\nu = 6.3$) quantiles

MFM_ch5_innovations

S&P 500: Are the Innovations Normal? (2/2)

- Standardised residuals of GARCH(1,1) with Normal innovations: excess kurtosis falls from 10.7 (returns) to 1.73; skewness -0.53, Jarque–Bera 1 153
- Jarque–Bera $= n[S^2/6 + E^2/24]$, S: skewness, E: excess kurtosis; χ^2_2 under Normality, 5% critical value 5.99: Normality is rejected by a wide margin
- 15 residuals below -4: 0.21 expected under the Normal, 8.5 under a Student-t with $\nu = 6.3$; largest: $\hat{z} = -6.6$ (27 February 2007)
- A longer left tail: skewed-t with $\hat{\lambda} = -0.150$ (GJR, $\hat{\nu} = 7.7$ degrees of freedom), hence BIC prefers skewed-t innovations

MFM_ch5_innovations

S&P 500: Diagnostics of Five Models

Model	$Q(10)$ on $\hat{z}$	$Q(10)$ on $\hat{z}^2$	LM(5)	Sign bias t	Joint χ^2_3
GARCH-N	21.2	16.4	8.0	3.64	41.8
GARCH-t	21.9	13.6	5.8	3.56	40.7
GJR-t	22.3	10.2	4.5	3.61	28.1
EGARCH-t	27.6	10.5	8.2	2.04	9.9
GJR-skewed-t	19.7	10.7	5.0	3.91	31.8

- 5% critical values: 18.3 (χ^2_{10}), 11.1 (χ^2_5), 7.81 (χ^2_3), $|t| > 1.96$; naive laws, conservative with estimated parameters (next slide)
- Mean: $Q(10)$ on $\hat{z}$ of 20–28 in every model: a little autocorrelation that a constant mean ignores
- Variance: $Q(10)$ on $\hat{z}^2$ falls from 6 133 (on r_t^2) to 10–16 and LM(5) is not significant: these naive tests detect no clustering left
- Asymmetry: symmetric GARCH fails the joint test (40.7), GJR still fails (28.1), EGARCH comes closest (9.9, p = 0.020); with $n = 6\,714$, judge economic size, not only p-values


MFM_ch5_innovations

Residual Diagnostics Have Estimated Parameters

- $\hat{z}_t$ depends on $\hat{\theta}$: the Ljung–Box statistic on $\hat{z}_t^2$ is not exactly χ^2_m Li & Mak (1994); the fitted variance has already absorbed part of the dependence at small lags
- S&P 500, Gaussian QMLE of GARCH(1,1), $Q(10)$ on $\hat{z}_t^2$
  - Naive Ljung–Box: 16.4, $p = 0.089$: not rejected at 5%
  - Corrected for the estimated parameters: 19.1, $p = 0.038$: rejected at 5%, the verdict flips
- The naive test is conservative: it rejects a misspecified variance equation too rarely; the joint sign-bias test carries the same caveat
- The correction matrix and its entries: Appendix

MFM_ch5_inference

Diagnostics: Squared Returns versus Squared Residuals



- S&P 500, lags 1–30: autocorrelations of r_t^2 (before the model) and of $\hat{z}_t^2$ from GJR-GARCH-t
- After the model they sit inside the i.i.d. band: no remaining variance dynamics are detected at these lags
- The band is the naive one: with estimated parameters it is too wide at small lags (previous slide)

MFM_ch5_innovations

Diagnostics in Other Markets: a Decision Table

Market	Residual pattern	Possible failure	Next check
BET	$Q(10)$ on $\hat{z}$ = 116.6	autocorrelation in the mean (thin trading, Chapter 2)	AutoRegressive model AR(1) for the mean
Bitcoin	$Q(10)$ on $\hat{z}^2$ = 4.7, joint sign bias 1.3: no rejection	$\hat{\nu} = 3.18 < 4$: $\mathrm{Var}(z_t^2) = \infty$, so the χ^2 calibration is not justified	tests on $\lvert\hat{z}_t\rvert$
EUR/RON	excess kurtosis 3 190	one residual of 154 (May 2025) dominates	inspect the largest residuals; a bounded-score model
Gold	EGARCH: $Q(10)$ on $\hat{z}^2$ = 48.8	clustering left despite a good BIC	choose by diagnostics, not by BIC alone

- Always look at the largest residuals: one observation can dominate every diagnostic

MFM_ch5_innovations

The Proxy Problem and Robust Losses

- Volatility is never observed: we compare forecasts with a noisy **proxy**, e.g. r_t^2
- $r_t^2 = \sigma_t^2 z_t^2$ is unbiased ($\mathbb{E}[r_t^2 \mid \mathcal{F}_{t-1}] = \sigma_t^2$ for $\mu = 0$) but very noisy
  - Its R^2 against the true variance is small even for a perfect forecast Andersen & Bollerslev (1998)
  - S&P 500: we ignore the mean, since $\hat{\mu}^2 = 0.0062$ against an average daily variance of 1.47 ($\%^2$), a bias below 1%
- Patton (2011): only some loss functions rank forecasts correctly with a noisy proxy
  - Robust: Mean Squared Error (MSE) $(r_t^2 - h_t)^2$ and **Quasi-Likelihood (QLIKE)** $r_t^2/h_t + \ln h_t$, with h_t the variance forecast for day t; both are averaged over the forecast days, lower is better
  - Not robust: losses on volatility $|r_t| - \sqrt{h_t}$ or on logs
- QLIKE is (up to constants) minus the Normal log-likelihood; its expected value is lowest at $h_t = \sigma_t^2$; it penalises under-prediction more


Forecast Evaluation (1/2): Mincer–Zarnowitz

- **Mincer–Zarnowitz (MZ) regression** Mincer & Zarnowitz (1969): regress the proxy on the forecast
  - $r_t^2 = a + b\, h_t + u_t$
  - a: intercept; b: slope; u_t: regression error
- Unbiased forecast: $a = 0$, $b = 1$
  - Joint Wald test with Heteroskedasticity and Autocorrelation Consistent (HAC) errors; χ_2^2 under H_0
  - $b < 1$: forecasts too dispersed (too high when high, too low when low)
  - $R^2 \in [0, 1]$: share of the variation of r_t^2 explained by h_t; small even for good forecasts, because r_t^2 is noisy

Forecast Evaluation (2/2): Diebold–Mariano

- **Diebold–Mariano test** Diebold & Mariano (1995): do two forecasts have the same expected loss?
  - Loss differential $d_t = L(r_t^2, h_t^A) - L(r_t^2, h_t^B)$
  - L: loss function (QLIKE or MSE); h_t^A, h_t^B: forecasts of models A and B; $d_t < 0$: A did better on day t
- Diebold–Mariano (DM) statistic: $DM = \bar{d}/\widehat{\text{s.e.}}(\bar{d})$
  - $\bar{d}$: mean of d_t over the forecast days; HAC standard error, since d_t is autocorrelated
  - $\approx N(0,1)$ under equal accuracy; $DM < -1.96$: A significantly better; $DM > 1.96$: B significantly better
- The $N(0,1)$ law assumes given forecasts; for estimated, nested models (GARCH-t inside GJR-t) it can fail, so DM t is read descriptively there (Giacomini–White, GW, below)


Out-of-Sample Design and Forecast Paths, 2016–2026



- S&P 500: 2 692 one-day forecasts; refit each January on all data since 2000 (expanding window), daily updates with fixed parameters
- Timing: h_t uses data up to the close of day $t-1$ and is compared with r_t^2; changing r_t must leave h_t unchanged
- Models: GARCH-N, GARCH-t, GJR-t, EGARCH-t; benchmarks: EWMA and Hist-20 (20-day historical variance); shown: GJR-t, EWMA, $|r_t|\sqrt{A}$ ($A = 251.4$)

MFM_ch5_forecast_evaluation

S&P 500: Does Anything Beat GARCH(1,1)?



- Mean QLIKE minus that of GARCH-t, with the DM t (HAC, 5 lags); lowest QLIKE: GJR-t (0.732)
- GARCH family: DM t from -1.47 to -0.36, descriptive for these nested pairs
- EWMA and Hist-20 are clearly worse ($t = 3.48$, 4.10): mean reversion and estimated parameters pay off Hansen & Lunde (2005)
- MZ: unbiasedness not rejected for GARCH-type forecasts, rejected for Hist-20 (all statistics: Appendix)
- Not rejecting is weak evidence: s.e. of b is 0.25 for GARCH-t, and a few crisis days drive b (Seminar 5, B5)

MFM_ch5_forecast_evaluation

Comparing Forecasting Methods (1/2): Giacomini–White and the Model Confidence Set

- **The limit of the DM test** West (1996), Clark & McCracken (2001)
  - With estimated parameters, nested models and an expanding window, the loss differential degenerates under H_0
  - Consequence: DM is no longer $N(0,1)$
- **Giacomini–White test** Giacomini & White (2006)
  - Tests the forecasting **method**: the model plus a rolling window of $W = 1\,000$ days; valid for nested models, because the estimation noise does not vanish
  - $GW = n\,\bar{Z}^{\top}\hat{\Omega}^{-1}\bar{Z} \to \chi^2_2$ under H_0: $\mathbb{E}[d_{t+1} \mid \mathcal{F}_t] = 0$, with $Z_{t+1} = (1, d_t)^{\top} d_{t+1}$
  - d_{t+1}: loss differential (previous slides); $\bar{Z}$: mean of Z_{t+1}; $\hat{\Omega}$: the uncentred second moment $n^{-1}\sum Z_{t+1}Z_{t+1}^{\top}$ for these one-step forecasts under the conditional null; n: number of forecasts; reject at 5% if $GW > 5.99$
- **Model Confidence Set** Hansen, Lunde & Nason (2011)
  - Six models: five tests against GARCH-t (DM, GW), or all 15 pairs inside the MCS: a multiple-comparison problem
  - Drops the worst model until equal accuracy is no longer rejected
  - The final set contains the best model with probability $\geq 1 - \alpha$; α: the chosen error rate (here 10%)


Comparing Forecasting Methods (2/2): Results for the S&P 500

- **Giacomini–White**: rolling window, 2 692 forecasts, each model against GARCH-t
  - Rejects only EWMA and Hist-20
- **Model Confidence Set (MCS) 90%**: expanding window, QLIKE, stationary bootstrap
  - Stationary bootstrap: resamples blocks of random (geometric) length, mean 10 days, so that serial dependence is kept
  - Keeps the four GARCH models, eliminates EWMA and Hist-20
- The DM verdict, now with calibrated inference
  - Full table and the MCS statistic: Appendix

Model	GW p	MCS p
GARCH-N	0.054	0.586
GARCH-t	–	0.353
GJR-t	0.288	1.000
EGARCH-t	0.201	0.586
EWMA	0.003	0.008
Hist-20	< 0.001	0.001

MFM_ch5_inference

One-Month Forecasts: Mincer–Zarnowitz in Logs



- GJR-t forecast of the 22-day variance against the realised sum of 22 squared returns, 122 non-overlapping windows; regression $\ln Y = a + b \ln \hat{h} + u$, Y: realised sum, $\hat{h}$: forecast; R^2 in logs 0.38
- Slope in logs $b = 0.75$ (s.e. 0.07): descriptive only, since $\mathbb{E}[\ln Y \mid \mathcal{F}] \neq \ln \mathbb{E}[Y \mid \mathcal{F}]$; $b < 1$ does not prove over-dispersed forecasts
- In levels b is only 0.27: two COVID-19 windows dominate; compare forecasts with QLIKE Patton (2011)


MFM_ch5_forecast_evaluation

GARCH-in-Mean: Is Risk Rewarded?

- **GARCH-in-Mean (GARCH-M)** Engle, Lilien & Robins (1987): $r_t = c + \kappa\,\sigma_t + \varepsilon_t$ (or $\kappa\,\sigma_t^2$)
  - c: constant; κ: price of risk, the extra expected return per unit of volatility
  - $\kappa > 0$: investors demand a higher expected return when risk is higher (risk-return trade-off)
- S&P 500, GARCH(1,1)-t in mean: $\hat{\kappa} = 0.096$ (t = 3.02) with σ_t; 0.028 (t = 2.35) with σ_t^2
  - Positive and significant in this sample, but the literature is mixed: Glosten, Jagannathan & Runkle (1993) found a negative relation
- Caution: a daily mean effect is small and fragile; it is sensitive to the sample and to the asymmetry in variance

MFM_ch5_long_memory

Long Memory: Where GARCH Falls Short



- S&P 500, ACF of $|r_t|$ (log scale): data 0.289, 0.082, 0.009 at lags 1, 100, 250; GARCH(1,1)-t too high at lag 1 (0.416), then an exponential collapse
- **Fractionally Integrated GARCH (FIGARCH)** Baillie, Bollerslev & Mikkelsen (1996)
  - Filter $(1-L)^d$: hyperbolic decay; L: lag operator, $Lx_t = x_{t-1}$; $d \in (0,1)$: fractional order
  - $\hat{d} = 0.60$ (s.e. 0.06), BIC lower by 34.2
  - But the memory is now too long (0.144 at lag 250)
- No single-scale model fits all horizons: realised volatility and Heterogeneous AutoRegressive (HAR) models in Chapter 9

MFM_ch5_long_memory

Three Extensions, Three Different Questions

- MIDAS means Mixed Data Sampling; GARCH-MIDAS combines information observed at different frequencies
- Question: can slowly changing macroeconomic conditions explain a long-run variance component? Engle, Ghysels & Sohn (2013)
- GAS means Generalised Autoregressive Score; its update uses the score, the gradient of a log-density
- Question: can a bounded score limit the effect of one extreme observation? Creal, Koopman & Lucas (2013)
- DCC means Dynamic Conditional Correlation, a model of correlations that vary over time
- Question: how does the joint risk of several assets evolve? See Chapter 6

Limits of GARCH and the Extensions They Motivate

Limit	Evidence in this chapter	Extension
Level shifts inflate $\alpha + \beta$ Lamoureux & Lastrapes (1990)	BET-TR (2014–2026) 0.9188 vs BET (2000–2026) 0.9907	GARCH-MIDAS, a slow long-run component Engle, Ghysels & Sohn (2013); regime switching Hamilton & Susmel (1994)
Reaction lag of one day	COVID-19: the VIX moved the same day, GARCH the next	Realized GARCH, with a realised-variance measurement equation Hansen, Huang & Shek (2012) (Chapter 9)
One outlier drives the update	EUR/RON residual of 154 (May 2025)	GAS models with a bounded score Creal, Koopman & Lucas (2013); the case study below
Short memory	ACF of $\lvert r_t \rvert$ at lags 100–250	FIGARCH; HAR on realised variance (Chapter 9)
One asset at a time	portfolios need covariances	Multivariate GARCH and DCC (Chapter 6)
Managed prices	EUR/RON: long calm, then jumps	Model interventions as jumps or regimes

- Each extension is judged like GARCH: out of sample, robust loss, Model Confidence Set (model equations: Appendix)

MFM_ch5_garch_estimation

Two Research Extensions and What They Assume

- This section and the next are optional: the in-class part of Seminar 5 does not use them
- **Quasi score-driven volatility**: a replication of a 2023 journal article
  - Builds on: the Student-t density, LR tests and their boundary versions, the score-driven (GAS) update of the extensions section
  - Continued in Seminar 5, C3
- **AI for scientific discovery**: an assistant-driven research loop with checks
  - Builds on: the out-of-sample design, QLIKE, Diebold–Mariano and the Model Confidence Set
  - Its checklist supports the AI audit of Seminar 5, C1 and the team project

QSD: Should Shocks Be Downweighted?

- **Question** Blasques, Francq & Laurent (2023)
  - Must a score-driven volatility model update σ_t^2 with the score of the same density that describes the returns?
- Score-driven (GAS) update Creal, Koopman & Lucas (2013): tomorrow's parameter moves in the direction that would have made today's observation more likely
  - $f_{t+1} = \omega + \alpha\, S(f_t)\, \partial \log p(y_t \mid f_t)/\partial f_t + \beta f_t$
  - y_t: the return; f_t: the time-varying parameter (here the variance); $p(y_t \mid f_t)$: the density of y_t given f_t
  - $\partial \log p/\partial f_t$: the score; $S(f_t) > 0$: a scaling factor; ω, α, β: as in GARCH
- With a Student-t density, one parameter ν sets both
  - the tails of z_t
  - and how strongly a large shock is downweighted in the update

QSD: Separating Tail Thickness from Shock Downweighting

- **Quasi Score-Driven (QSD) GARCH(1,1)-t**, eq. (30) of the paper: $y_t = \sqrt{f_t}\,\epsilon_t$, ϵ_t standardised Student-t with $1/\xi$ degrees of freedom (f_t is σ_t^2, ϵ_t is z_t)
  - $f_{t+1} = \omega + \varpi X_t + \alpha \dfrac{1+\zeta}{1-2\zeta+\zeta\epsilon_t^2}\,\epsilon_t^2 f_t + \beta f_t$
  - X_t: an exogenous variable, the squared VIX at a daily horizon; ϖ: its coefficient
  - ξ: the tail thickness of ϵ_t ($\xi = 0$: Normal); ζ: the downweighting of large ϵ_t^2 in the update
  - $\zeta = 0$: GARCH(1,1)-t; $\zeta = \xi$: Beta-t-GARCH (score-driven)
  - Why: with $\zeta = \xi = 1/\nu$, multiplying by ν gives $\dfrac{1+\zeta}{1-2\zeta+\zeta\epsilon_t^2} = \dfrac{\nu+1}{\nu-2+\epsilon_t^2}$, the weight in the Student-t score
- Data: 400 S&P 500 stocks, 3 January 1995 (or later) – 28 February 2019; maximum likelihood with an AR(1) mean
  - LR tests of $\zeta = 0$ and $\xi = \zeta$ against χ_1^2 (full method: Appendix)
  - Probability Integral Transform (PIT) test: $u_t = F(y_t \mid \mathcal{F}_{t-1})$, the fitted distribution function at the observed return, must be i.i.d. uniform on $(0,1)$ if the density is right

QSD: Key Results (Paper, Tables 4 and 6)


Rejected at 5%, % of the 400 stocks	Student-t (Table 4)	Skewed t (Table 6)
LR, H_0: $\zeta = 0$ (GARCH update)	89.5	98.8
LR, H_0: $\xi = \zeta$ (score-driven; skewed: also $\kappa = \tau$)	53.5	100.0
PIT test: GARCH(1,1)	40.0	33.1
PIT test: Beta-t-GARCH	46.0	72.2
PIT test: QSD GARCH(1,1)	45.7	11.3

- κ: asymmetry of the density; τ: asymmetry of the update; $\kappa = \tau = 0.5$: symmetric (skewed-t version, Table 6)
- Downweighting large shocks matters (Table 4)
  - $\zeta = 0$ is rejected for most stocks (first row)
  - Every $\hat{\zeta}$ is positive

QSD: Reading the Results (Paper, Figs. 2–6)

- **Beta-t-GARCH downweights too much** (Figs. 2–3)
  - Where $\hat{\xi} - \hat{\zeta}$ is significant, $\hat{\xi} > \hat{\zeta}$ for all stocks but one
  - Median $1/\hat{\zeta}$: 9.48 in QSD, against 5.07 in Beta-t-GARCH, where $\zeta = \xi$
  - Tail thickness and shock downweighting are two different parameters
- **Skewed t** (Figs. 5–6)
  - $\hat{\tau}$ far below 0.5 (median 0.333), while $\hat{\kappa}$ stays near 0.5
  - Leverage enters through the update equation, not through the density
- **Calibration** (Table 6)
  - With skewed-t updates, the PIT test rejects the QSD model for only 11.3% of the stocks, against 72.2% for Beta-t-GARCH


Case Study: Replication on 11 Stocks (cf. Paper, Fig. 2)



- Course sample: 11 large stocks still listed (the paper: 400), 1995 (or later) – February 2019, 4986–6081 returns each; $X_t = \mathrm{VIX}_t^2/252$
- $\zeta = 0$ rejected for 9 of 11 (paper: 89.5%); $\xi = \zeta$ rejected for 8 (paper: 53.5%); $\hat{\xi} > \hat{\zeta}$ for 10 of 11
- Median $1/\hat{\zeta}$: 21.4 (QSD) against 6.32 (Beta-t-GARCH); paper: 9.48 against 5.07; 11 selected survivors: imprecise and not representative

MFM_ch5_quasi_score_driven

QSD: News Impact Curves (cf. Paper, Fig. 3)



- Weight of ϵ_t in f_{t+1} (units of αf_t) at quantiles of the 11 estimates: updating functions at a common αf_t, not fitted paths
- A shock of 5 standard deviations adds 25 (GARCH-t), 12.6 (median QSD), 6.2 (median Beta-t-GARCH); at 10 standard deviations: 100, 18.8, 7.0 (bound $(1+\zeta)/\zeta = 7.3$)
- QSD lies between the two
- Read the vertical axis in units of αf_t: GARCH-t adds $\epsilon_t^2\, \alpha f_t$, i.e. $25\, \alpha f_t$ at 5 standard deviations, not $25 f_t$

MFM_ch5_quasi_score_driven

Open Questions and the Discovery Loop (1/2)

- Questions
  - do machine-learning or realised-measure forecasts beat a well-specified GARCH for daily equity variance, out of sample?
  - does the answer survive once the number of models compared is accounted for?
- Literature
  - Christensen, Siggaard & Veliyev (2023): machine-learning methods with little tuning are competitive with HAR-type models
  - Hansen, Lunde & Nason (2011): the Model Confidence Set is the right object when many models compete
- Formal claim: $d_t = L(r_t^2, h_t^A) - L(r_t^2, h_t^B)$, $H_0 \colon \mathbb{E}d_t = 0$
  - Falsified if A leaves the Model Confidence Set
  - Or if the sign of $\bar{d}$ flips across pre-registered subperiods

Open Questions and the Discovery Loop (2/2)

- The loop
  - An Artificial Intelligence (AI) assistant drafts the literature map, competing hypotheses, code, a robustness plan and a referee-style critique
  - The researcher checks every Digital Object Identifier (DOI, a permanent publication identifier), reproduces every number and fixes the plan before the first run
- Checks specific to volatility forecasts
  - Look-ahead: h_t must not move when r_t changes
  - A proxy-robust loss (QLIKE, MSE)
  - Model mining: Model Confidence Set or a Bonferroni correction
  - Scale and convergence of the optimiser
  - Invented DOIs or `arch` arguments
- Reusable prompts and the full checklist: ▸ Appendix

Mini-Case and a Team-Project Seed

- A typical assistant claim: "GJR-t beats GARCH-t because of the leverage effect"
  - d_t = QLIKE(GJR-t) − QLIKE(GARCH-t), 2 692 days, 2016–2026: $\bar{d} = -0.025$, DM $t = -1.22$
- Testable implication of leverage: the gain sits after down days
  - After $r_{t-\mathbf{1}} < 0$ (1 223 days): $\bar{d} = -0.015$, $t = -0.51$; after $r_{t-\mathbf{1}} \geq 0$ (1 469): $\bar{d} = -0.034$, $t = -1.12$
  - Not supported; by period: 2016–2020 $t = -0.02$, 2021–2026 $t = -4.06$
  - Descriptive only (nested models, expanding window); calibrated follow-up: a fixed-window GW test with d_t regressed on $1(r_{t-\mathbf{1}} < 0)$
- **Project seed**: do machine-learning forecasts enter the Model Confidence Set together with GARCH, GJR and EGARCH for the S&P 500 and the BET, 2016–2026?
  - Replicate the QLIKE table first; pre-register models, losses, splits and error rate
  - Log AI use in `AI_USE.md` and every caught error in `AI_ERRORS.md` (full brief: Appendix)

Synthesis: a Decision Workflow

- **1. Detect**
  - $Q(10)$ on r_t^2 and the LM test
  - Clustering is universal; persistence $\alpha + \beta$ is close to 1 in all our markets
- **2. Fit**
  - GARCH, GJR, EGARCH with Normal, Student-t, skewed-t innovations; returns in %
  - Check convergence and parameters at a bound
- **3. Diagnose**
  - Level, squares and signs of $\hat{z}_t$, and its largest values
  - Naive tests are conservative
- **4. Forecast**
  - The recursion gives multi-step variances
  - The square-root-of-time rule ignores mean reversion
- **5. Compare**
  - Out of sample, QLIKE, GW with a rolling window and the Model Confidence Set
  - GARCH beats EWMA and Hist-20; the GARCH family is hard to separate
- **6. Report uncertainty**
  - Robust s.e. (valid if $\mathbb{E}z_t^4 < \infty$) and boundary-aware tests
  - Half-lives as intervals $[h_{lo}; \infty)$ when $\alpha + \beta \approx 1$

Check Yourself, and Next: Chapter 6

Retrieval questions (answers on the next slide)

- Why does the square-root-of-time rule overstate the 10-day risk after a turbulent day?
- Why is the LR test of Normal against Student-t innovations compared with 2.71 and not 3.84?
- Why can a symmetric interval for the half-life contain negative values?
- Why is a gap between the VIX and GARCH volatility not a measured variance risk premium?

Next chapters

- Chapter 6: multivariate volatility and dynamic correlations (DCC)
- Chapters 7–8: Value at Risk and expected shortfall with GARCH volatility, and their backtesting
- Chapter 9: realised volatility and long memory
- Seminar 5: the variance recursion step by step, GARCH-t and GJR on real data, an AI answer to audit, bootstrap tests and forecast comparisons

Check Yourself: Short Answers and Where to Revise

- **Square-root-of-time rule**
  - After a turbulent day $\sigma^2_{t+1} > \sigma^2$; the daily forecasts decay towards σ^2, so their sum is below $H\sigma^2_{t+1}$
  - Revise: the toy check after the half-life slide, the S&P 500 worked example (the opposite case), Seminar 5, A1
- **2.71 instead of 3.84**
  - Under H_0, $1/\nu = 0$ is on the boundary: in about half of the samples the estimate sits at the bound and $LR = 0$, so the law is $\frac{1}{2}\chi^2_0 + \frac{1}{2}\chi^2_1$, whose 5% critical value is 2.71
  - Revise: the decision table for tests on the boundary
- **Negative half-lives**
  - $h = \ln 0.5 / \ln p$ explodes as $p \to 1$; the delta method replaces it by a straight line, so $\hat{h} \pm 1.96\,\widehat{\text{s.e.}}$ ignores the curvature; map the interval for p instead
  - Revise: the delta-method slides, Seminar 5, A2
- **VIX against GARCH**
  - The premium compares VIX^2, a 30-day risk-neutral expectation, with a 30-day variance forecast under the real-world probability; $\hat{\sigma}_t$ is a one-day volatility, so horizon and probability both differ
  - Revise: the S&P 500 GARCH and VIX slide

Data Used in This Chapter

Series	Owner / source	Start	Obs.	Obs. per year	Vol. (% p.a.)	Excess kurt.
S&P 500	S&P Dow Jones Indices	2000	6 714	251.4	19.2	10.7
BET	Bucharest Stock Exchange	2000	6 670	249.8	22.1	11.1
BET-TR	Bucharest Stock Exchange	2014	2 992	249.7	15.4	17.8
Bitcoin (BTC/USD)	crypto exchanges (composite)	2014	4 384	365.3	66.6	12.0
EUR/RON reference rate	BNR	2005	5 352	252.4	4.4	15.1
Gold (XAU/USD)	spot market	2000	6 922	259.2	17.6	7.7

- Daily closes up to 18 September 2026; returns $r_t = 100\,(\ln P_t - \ln P_{t-1})$, in %
- Equity indices and gold: weekdays only
- Vol.: sample volatility annualised with each series' observations per year; excess kurtosis: kurtosis minus 3
- Bitcoin trades 7 days a week; the VIX index of the Chicago Board Options Exchange is used as a benchmark

MFM_ch5_arch_effects

Back

One-Sided Score Tests and the Davies Problem

- The alternative is one-sided ($\alpha_i \geq 0$): the two-sided χ^2 test wastes power
  - Lee & King (1993): one-sided score test built on the sum of the score components, $N(0,1)$ under H_0, rejects in the upper tail only
- Against GARCH(1,1) the null $\alpha = 0$ leaves β unidentified (Davies problem)
  - The ARCH(q) LM statistic is also the score test against GARCH(p,q) Lee & King (1993): the same statistic serves both alternatives

Back

Robert Engle and the 2003 Nobel Prize



Robert F. Engle (2022)

Photo: Jon Demske (2022); CC BY-SA 4.0; Wikimedia Commons



Clive W.J. Granger (2008)

Photo: Olaf Storbeck (2008); CC BY-SA 2.0; Wikimedia Commons

- Nobel Prize 2003: Engle "for methods of analyzing economic time series with time-varying volatility (ARCH)"
  - Shared with Clive Granger (cointegration)
- Nobel lecture: Engle (2004), on risk and volatility in financial practice
- Tim Bollerslev, Engle's doctoral student, generalised ARCH to GARCH in 1986

Back

Kurtosis: Rounding and a Long Simulated Path

- S&P 500, GARCH-N: the moment condition is 0.990435, so $K = 12.1$ has the denominator $1 - 0.990435$
  - Rounding to $\hat{\alpha} = 0.121$, $\hat{\beta} = 0.860$ gives 0.9916 and $K = 13.5$: report such functions from unrounded estimates
- Simulated GARCH(1,1) of the simulation slide, one million days
  - Excess kurtosis 2.47 (theory 3.06); lag-1 autocorrelation of ε_t^2: 0.26
  - The sample kurtosis converges slowly: its precision depends on the eighth moment, infinite for these parameters

Back

QMLE: Variants of the Theorem

- Estimated mean: the score gains the term ε_t/σ_t^2 and the covariance $(\kappa_z - 1)J^{-1}$ changes form
- IGARCH ($\alpha + \beta = 1$): strict stationarity still holds, and the QMLE remains consistent and asymptotically Normal Lumsdaine (1996)
- Explosive case, $\mathbb{E}\ln(\alpha z_t^2 + \beta) \geq 0$: Jensen & Rahbek (2004) cover (α, β) with ω fixed, since ω is not identified
- Heavy tails, $\kappa_z = \infty$: slower than $\sqrt{n}$, non-Gaussian limits Hall & Yao (2003); heavy-tailed quasi-likelihoods with a scale adjustment Fan, Qi & Xiu (2014)

Back

The Li–Mak Correction for Squared-Residual Autocorrelations

- Squared-residual autocorrelations $\hat{r}_k$: $\sqrt{n}\,\hat{r} \to N(0, C)$, $C = I - H' J^{-1} H/(\kappa_z - 1)$ Li & Mak (1994)
  - $\hat{r} = (\hat{r}_1, \ldots, \hat{r}_m)'$; I: identity matrix; C: the corrected covariance, smaller than I on the diagonal; g_t, J: as in the QMLE theorem
  - $H_k = \mathbb{E}[g_t(z_{t-k}^2 - 1)]$ (k-th column of H): how the fitted variance absorbs lag-k dependence; the ARMA rule χ^2_{m-p-q} does not apply
- S&P 500, Gaussian QMLE of GARCH(1,1): $C_{1,1} = 0.78$, $C_{5,5} = 0.93$, $C_{10,10} = 0.97$
  - The correction is largest at small lags, where $\hat{\alpha}$ has already fitted the dependence; corrected statistic $n\hat{r}' C^{-1} \hat{r} = 19.1$, χ^2_{10} under H_0

MFM_ch5_inference

Back

S&P 500 Forecast Evaluation, All Statistics

Model	QLIKE	MSE	DM t vs GARCH-t	MZ b	MZ R^2	Wald p
GARCH-N	0.751	22.3	−1.47	0.97	0.250	1.00
GARCH-t	0.757	22.8	–	0.88	0.238	0.80
GJR-t	0.732	21.8	−1.22	0.84	0.276	0.68
EGARCH-t	0.745	21.3	−0.36	1.24	0.293	0.71
EWMA	0.815	24.7	3.48	0.89	0.172	0.69
Hist-20	0.889	26.1	4.10	0.68	0.152	0.03

- Mean losses on 2 692 days; DM on QLIKE, HAC with 5 lags; negative t: better than GARCH-t
- MZ: $r_t^2 = a + b\,h_t$; Wald p: test of $a = 0$, $b = 1$

MFM_ch5_forecast_evaluation

Back

Giacomini–White and the MCS: Full Table and Settings

Model	QLIKE (W)	t uncond.	GW	p	MCS p
GARCH-N	0.751	1.87	5.85	0.054	0.586
GARCH-t	0.743	–	–	–	0.353
GJR-t	0.738	−0.18	2.49	0.288	1.000
EGARCH-t	0.746	0.07	3.21	0.201	0.586
EWMA	0.815	3.20	11.40	0.003	0.008
Hist-20	0.889	3.87	16.41	< 0.001	0.001

- GW: the statistic of the main text, χ^2_2 under H_0 (5.99 at 5%); t uncond.: the unconditional test (instrument 1 only), a Diebold–Mariano t on the rolling-window losses
- Rolling window $W = 1\,000$ days, re-estimated each January, 2 692 forecasts; tests against GARCH-t: unconditional t uses HAC with 5 lags; GW uses the uncentred second moment, without HAC
- MCS on the expanding-window forecasts: QLIKE, statistic T_R (the largest pairwise $|t|$ of loss differences), stationary bootstrap (mean block 10, 10 000 draws); 90% MCS: GARCH-N, GARCH-t, GJR-t, EGARCH-t

MFM_ch5_inference

Back

Score-Driven, Realized and Component Models

- **Score-driven (GAS)** Creal, Koopman & Lucas (2013): $\sigma^2_{t+1} = \omega + \alpha\, s_t + \beta\, \sigma^2_t$, s_t the scaled score of the Student-t density with respect to σ^2_t
  - The score is bounded in z_t: an outlier such as the EUR/RON residual of 154 moves the variance far less than in GARCH
- **Realized GARCH** Hansen, Huang & Shek (2012): adds a measurement equation linking σ^2_t to a realised variance
  - Uses intraday information and reacts faster; the bridge to Chapter 9
- **GARCH-MIDAS** Engle, Ghysels & Sohn (2013): $\sigma^2_t = \tau_t\, g_t$, a slow long-run component τ_t driven by macro data (mixed frequencies)
  - g_t: a unit-mean GARCH(1,1) component for the short-run moves
  - A moving long-run level absorbs the level shifts that push $\hat{\alpha} + \hat{\beta}$ towards 1

Back

Beyond Variance: Entropy and Tail Risk

- Pele, Lazar & Dufour (2017): entropy of the intraday return distribution and market risk (EUR/JPY)
  - Higher entropy goes with lower intraday Value at Risk and expected shortfall; entropy then forecasts daily Value at Risk
- Pele & Mazurencu-Marinescu-Pele (2019): the entropy of intraday Bitcoin returns proposed as an additional signal for daily Value at Risk
- Pele, Lazar & Mazurencu-Marinescu-Pele (2019): a nonparametric tail measure based on entropy, compared with standard expected shortfall models
- GARCH volatility feeds directly into Value at Risk and expected shortfall: Chapters 7–8

Back

Case Study, Full Method

- **Why it matters**: GARCH is driven by ε_t^2 whatever the density, so an isolated extreme shock can produce an excessive volatility response lasting days or weeks (Section 1 of the paper)
  - With a left-skewed density, a pure score-driven model raises volatility more after positive returns: the opposite of the leverage effect (Example 2)
  - QSD models, eq. (3): $f_{t+1} = \omega + \alpha\, \psi(y_t, X_t, f_t, \theta) + \beta f_t$, ψ the scaled score of an updating function ρ that need not equal $\log p$; theory: consistency, asymptotic normality, LR and LM tests with χ^2 limits (Section 3)
- **Estimation**: $\zeta = 0$ is kept inside the parameter space by eqs. (31)–(32): $\zeta \in (-1, 1/2)$ and $\Psi(x) = x\,(1 - e^{-cx})/(1 + e^{-cx})$ applied to the weight, $c = 1000$
  - PIT test of the case study Diebold, Gunther & Tay (1998), with the MLE and Pierce (1982): reject if $|\sum_t \hat{u}_t - T/2| / \sqrt{T(1/12 - B\,\mathcal{I}(\hat{\theta})^{-1} B^\top)} > 1.96$, $B = \lim_{T\to\infty} \mathbb{E}[\partial \sum_t (u_t - T/2)/\partial\theta^\top]$, $\mathcal{I}$ the Fisher information, T the sample length; it detects a shifted PIT mean, not every departure from uniformity
- **Data**: stocks with fewer than 4000 observations or with convergence problems are dropped: 400 stocks (Section 5)
  - Skewed variant, eq. (47): a skewed Student-t with asymmetry κ for ϵ_t and separate (ζ, τ) in the update; $\tau < 0.5$ gives a leverage effect whatever κ (Section 5.2)

Back

Case Study, an Example and the Limits Stated by the Authors

- Centene, 4330 days: $\hat{\xi} \approx 0.26$ (about 4 degrees of freedom) in all three models, but $\hat{\zeta} = 0.048$ (s.e. 0.027); LR 8.81 ($p = 0.003$) for $\xi = \zeta$ and 8.42 ($p = 0.004$) for $\zeta = 0$ (Table 5); after a crash day in March–April 2008, GARCH-t volatility stays well above Beta-t-GARCH and QSD lies between (Fig. 4)
- **Limits** stated by the authors
  - Beating GARCH-t and Beta-t-GARCH does not make the Student-t QSD model adequate: the three models fail the PIT test for 40.0–46.0% of the stocks (Table 4)
  - In the unmodified recursion (30), $\zeta < 0$ or $\zeta > 1/2$ does not define a proper volatility model, and $\zeta = 1/2$ with $\varpi = 0$ is degenerate (Section 4.1)
  - The $N(T/2, T/12)$ law of $\sum_t u_t$ holds only for known parameters; with estimated ones the PIT test needs the correction above (Section 5.1)
- **Open extensions** (Seminar 5, C3)
  - Does the bounded QSD update beat GJR-GARCH-t in QLIKE out of sample, especially after March 2020?
  - Does it tame the EUR/RON residual of 154?

Back

The Discovery Loop: Reusable Prompts (1/2)

- Tools are interchangeable: an AI assistant (Claude, ChatGPT, Gemini, GitHub Copilot) and literature engines (Semantic Scholar, Elicit, Google Scholar)
- **Literature**: *"List out-of-sample comparisons of GARCH, HAR and machine-learning variance forecasts since 2015, with DOI, asset, loss function, test and horizon."*
  - Resolve every DOI on Crossref; check that the loss is robust to a noisy proxy Patton (2011)
- **Hypotheses**: *"For the claim that asymmetry improves one-day forecasts, list competing mechanisms (leverage, volatility feedback, a few crisis days) and what each predicts for d_t given the sign of r_{t-1} and by subperiod."*
- **Data and code**: *"Write Python with `arch` that re-estimates GARCH-t and GJR-t each January on an expanding window from the course daily price files and stores one-day variance forecasts; no parameter may use data after the forecast origin."*
  - Replicate first: QLIKE 0.757 (GARCH-t) and 0.732 (GJR-t), DM $t = -1.22$

Back

The Discovery Loop: Reusable Prompts (2/2)

- **Robustness**: *"Write a pre-registered plan: splits by the sign of r_{t-1} and by subperiod, MSE besides QLIKE, the 22-day horizon, the HAC lag, and a Bonferroni or Model Confidence Set correction."*
  - A random-split placebo is calibrated only if it reassigns blocks of days, preserving volatility clustering
- **Critique**: *"Act as a referee: could the proxy r_t^2, one crisis year or the HAC lag create my result? Give the test that would expose each."*
- **Report**: all splits and losses tried, adjusted p-values, code, seeds and package versions; a specification chosen after seeing results is exploratory and reported as such
- **Influential periods**: a crisis year can dominate $\bar{d}$; report subperiods and state in advance whether any year may be excluded

Back

The Team-Project Seed in Full

- Question: the project seed of the mini-case slide, on daily data
- Realised-variance (HAR) branch: the course data set of 5-minute realised variances holds 40 large American stocks, January 2015 – January 2026; there are no intraday data for the S&P 500 index or the BET, so this branch is run stock by stock
- Replicate first: the QLIKE table (GARCH-t 0.757, EWMA DM $t = 3.48$) and the mini-case splits
- Deliverables
  - A one-page pre-registration: models, losses, splits, error rate and what would falsify the claim
  - Code on the course market data, the Model Confidence Set at 10%, a referee-style self-critique; `AI_USE.md` and `AI_ERRORS.md`
- Code of the mini-case: Quantlets/Ch_05/ai_discovery_case.py

Back

Further Resources

- Books
  - Franke, Härdle & Hafner (2019), chapters on AutoRegressive Integrated Moving Average (ARIMA) and GARCH models for financial time series
  - Francq & Zakoïan (2019): the mathematical theory of GARCH models
- Quantinar courses
  - Statistics of Financial Markets
  - Applied Time Series Analysis with Python
- Code
  - All charts and tables: Quantlets/Ch_05; Python package Sheppard, `arch`

Back

Acronyms used in this material (1/2)

- **ACF**: AutoCorrelation Function
- **AI**: Artificial Intelligence
- **AIC**: Akaike Information Criterion
- **AR**: AutoRegressive (model)
- **ARCH**: AutoRegressive Conditional Heteroskedasticity
- **ARIMA**: AutoRegressive Integrated Moving Average (model)
- **ARMA**: AutoRegressive Moving Average (model)
- **BET**: Bucharest Exchange Trading (index)
- **BET-TR**: BET Total Return (index)
- **BIC**: Bayesian Information Criterion
- **BTC**: Bitcoin
- **BNR**: Banca Națională a României — *the National Bank of Romania*
- **COVID-19**: COronaVIrus Disease 2019
- **DCC**: Dynamic Conditional Correlation
- **DOI**: Digital Object Identifier
- **DM**: Diebold–Mariano (test)
- **EGARCH**: Exponential GARCH
- **EWMA**: Exponentially Weighted Moving Average
- **FIGARCH**: Fractionally Integrated GARCH
- **GARCH**: Generalised AutoRegressive Conditional Heteroskedasticity
- **GARCH-M**: GARCH-in-Mean
- **GARCH-MIDAS**: GARCH with Mixed Data Sampling (long-run component)
- **GARCH-N**: GARCH with Normal innovations

Acronyms used in this material (2/2)

- **GAS**: Generalised Autoregressive Score (model)
- **GED**: Generalised Error Distribution
- **GJR**: Glosten–Jagannathan–Runkle (asymmetric GARCH)
- **GJR-GARCH**: Glosten–Jagannathan–Runkle GARCH
- **GW**: Giacomini–White (test)
- **HAC**: Heteroskedasticity and Autocorrelation Consistent (standard errors)
- **HAR**: Heterogeneous AutoRegressive (model)
- **IGARCH**: Integrated GARCH
- **LM**: Lagrange Multiplier
- **LR**: Likelihood Ratio (test)
- **MCS**: Model Confidence Set
- **MIDAS**: MIxed DAta Sampling (regression)
- **MLE**: Maximum Likelihood Estimation
- **MSE**: Mean Squared Error
- **MZ**: Mincer–Zarnowitz (regression)
- **PIT**: Probability Integral Transform
- **QLIKE**: Quasi-LIKElihood (loss function)
- **QMLE**: Quasi-Maximum Likelihood Estimator
- **QQ**: Quantile–Quantile (plot)
- **QSD**: Quasi Score-Driven (model)
- **VIX**: Chicago Board Options Exchange Volatility Index
- **XAU**: gold, international currency code (one troy ounce)

References (1/6)

- Andersen, T.G., Bollerslev, T. (1998). Answering the skeptics: yes, standard volatility models do provide accurate forecasts. *International Economic Review*, 39(4), 885–905.
- Andrews, D.W.K. (2000). Inconsistency of the bootstrap when a parameter is on the boundary of the parameter space. *Econometrica*, 68(2), 399–405.
- Andrews, D.W.K. (2001). Testing when a parameter is on the boundary of the maintained hypothesis. *Econometrica*, 69(3), 683–734.
- Baillie, R.T., Bollerslev, T., Mikkelsen, H.O. (1996). Fractionally integrated generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics*, 74(1), 3–30.
- Berkes, I., Horváth, L., Kokoszka, P. (2003). GARCH processes: structure and estimation. *Bernoulli*, 9(2), 201–227.
- Blasques, F., Francq, C., Laurent, S. (2023). Quasi score-driven models. *Journal of Econometrics*, 234(1), 251–275.
- Bollerslev, T. (1986). Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics*, 31(3), 307–327.
- Bollerslev, T. (1987). A conditionally heteroskedastic time series model for speculative prices and rates of return. *Review of Economics and Statistics*, 69(3), 542–547.
- Bollerslev, T., Wooldridge, J.M. (1992). Quasi-maximum likelihood estimation and inference in dynamic models with time-varying covariances. *Econometric Reviews*, 11(2), 143–172.

References (2/6)

- Christensen, K., Siggaard, M., Veliyev, B. (2023). A machine learning approach to volatility forecasting. *Journal of Financial Econometrics*, 21(5), 1680–1727.
- Clark, T.E., McCracken, M.W. (2001). Tests of equal forecast accuracy and encompassing for nested models. *Journal of Econometrics*, 105(1), 85–110.
- Creal, D., Koopman, S.J., Lucas, A. (2013). Generalized autoregressive score models with applications. *Journal of Applied Econometrics*, 28(5), 777–795.
- Diebold, F.X., Gunther, T.A., Tay, A.S. (1998). Evaluating density forecasts with applications to financial risk management. *International Economic Review*, 39(4), 863–883.
- Diebold, F.X., Mariano, R.S. (1995). Comparing predictive accuracy. *Journal of Business & Economic Statistics*, 13(3), 253–263.
- Engle, R.F. (1982). Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica*, 50(4), 987–1007.
- Engle, R.F. (2004). Risk and volatility: econometric models and financial practice. *American Economic Review*, 94(3), 405–420.
- Engle, R.F., Ghysels, E., Sohn, B. (2013). Stock market volatility and macroeconomic fundamentals. *Review of Economics and Statistics*, 95(3), 776–797.
- Engle, R.F., Lilien, D.M., Robins, R.P. (1987). Estimating time varying risk premia in the term structure: the ARCH-M model. *Econometrica*, 55(2), 391–407.

References (3/6)

- Engle, R.F., Ng, V.K. (1993). Measuring and testing the impact of news on volatility. *Journal of Finance*, 48(5), 1749–1778.
- Fan, J., Qi, L., Xiu, D. (2014). Quasi-maximum likelihood estimation of GARCH models with heavy-tailed likelihoods. *Journal of Business & Economic Statistics*, 32(2), 178–191.
- Francq, C., Zakoïan, J.-M. (2004). Maximum likelihood estimation of pure GARCH and ARMA-GARCH processes. *Bernoulli*, 10(4), 605–637.
- Francq, C., Zakoïan, J.-M. (2019). GARCH Models: Structure, Statistical Inference and Financial Applications, 2nd ed. Wiley.
- Franke, J., Härdle, W.K., Hafner, C.M. (2019). Statistics of Financial Markets: An Introduction, 5th ed. Springer.
- Giacomini, R., White, H. (2006). Tests of conditional predictive ability. *Econometrica*, 74(6), 1545–1578.
- Glosten, L.R., Jagannathan, R., Runkle, D.E. (1993). On the relation between the expected value and the volatility of the nominal excess return on stocks. *Journal of Finance*, 48(5), 1779–1801.
- Hall, P., Yao, Q. (2003). Inference in ARCH and GARCH models with heavy-tailed errors. *Econometrica*, 71(1), 285–317.
- Hamilton, J.D., Susmel, R. (1994). Autoregressive conditional heteroskedasticity and changes in regime. *Journal of Econometrics*, 64(1–2), 307–333.

References (4/6)

- Hansen, B.E. (1994). Autoregressive conditional density estimation. *International Economic Review*, 35(3), 705–730.
- Hansen, P.R., Huang, Z., Shek, H.H. (2012). Realized GARCH: a joint model for returns and realized measures of volatility. *Journal of Applied Econometrics*, 27(6), 877–906.
- Hansen, P.R., Lunde, A. (2005). A forecast comparison of volatility models: does anything beat a GARCH(1,1)? *Journal of Applied Econometrics*, 20(7), 873–889.
- Hansen, P.R., Lunde, A., Nason, J.M. (2011). The model confidence set. *Econometrica*, 79(2), 453–497.
- J.P. Morgan/Reuters (1996). RiskMetrics — Technical Document, 4th ed. New York.
- Jensen, S.T., Rahbek, A. (2004). Asymptotic inference for nonstationary GARCH. *Econometric Theory*, 20(6), 1203–1226.
- Lamoureux, C.G., Lastrapes, W.D. (1990). Persistence in variance, structural change, and the GARCH model. *Journal of Business & Economic Statistics*, 8(2), 225–234.
- Lee, J.H.H., King, M.L. (1993). A locally most mean powerful based score test for ARCH and GARCH regression disturbances. *Journal of Business & Economic Statistics*, 11(1), 17–27.
- Li, W.K., Mak, T.K. (1994). On the squared residual autocorrelations in non-linear time series with conditional heteroskedasticity. *Journal of Time Series Analysis*, 15(6), 627–636.

References (5/6)

- Lumsdaine, R.L. (1996). Consistency and asymptotic normality of the quasi-maximum likelihood estimator in IGARCH(1,1) and covariance stationary GARCH(1,1) models. *Econometrica*, 64(3), 575–596.
- Mandelbrot, B. (1963). The variation of certain speculative prices. *Journal of Business*, 36(4), 394–419.
- McLeod, A.I., Li, W.K. (1983). Diagnostic checking ARMA time series models using squared-residual autocorrelations. *Journal of Time Series Analysis*, 4(4), 269–273.
- Mincer, J., Zarnowitz, V. (1969). The evaluation of economic forecasts. In J. Mincer (ed.), *Economic Forecasts and Expectations: Analysis of Forecasting Behavior and Performance*, NBER.
- Nelson, D.B. (1990). Stationarity and persistence in the GARCH(1,1) model. *Econometric Theory*, 6(3), 318–334.
- Nelson, D.B. (1991). Conditional heteroskedasticity in asset returns: a new approach. *Econometrica*, 59(2), 347–370.
- Patton, A.J. (2011). Volatility forecast comparison using imperfect volatility proxies. *Journal of Econometrics*, 160(1), 246–256.
- Pele, D.T., Lazar, E., Dufour, A. (2017). Information entropy and measures of market risk. *Entropy*, 19(5), 226.
- Pele, D.T., Lazar, E., Mazurencu-Marinescu-Pele, M. (2019). Modeling expected shortfall using tail entropy. *Entropy*, 21(12), 1204.

References (6/6)

- Pele, D.T., Mazurencu-Marinescu-Pele, M. (2019). Using high-frequency entropy to forecast Bitcoin's daily Value at Risk. *Entropy*, 21(2), 102.
- Pierce, D.A. (1982). The asymptotic effect of substituting estimators for parameters in certain types of statistics. *The Annals of Statistics*, 10(2), 475–478.
- Self, S.G., Liang, K.-Y. (1987). Asymptotic properties of maximum likelihood estimators and likelihood ratio tests under nonstandard conditions. *Journal of the American Statistical Association*, 82(398), 605–610.
- Sheppard, K. arch: ARCH models and other tools for financial econometrics in Python (software documentation).
- West, K.D. (1996). Asymptotic inference about predictive ability. *Econometrica*, 64(5), 1067–1084.