



## Statistics of Financial Markets

Seminar 7: Efficient markets, random walk and variance-ratio tests



Daniel Traian PELE

Bucharest University of Economic Studies

IDA Institute Digital Assets

Blockchain Research Center

AI4EFin Artificial Intelligence for Energy Finance

Romanian Academy, Institute for Economic Forecasting

MSCA Digital Finance

## Acronyms used in this material

- ▣ ACF: Autocorrelation Function
- ▣ ADF: Augmented Dickey–Fuller (test)
- ▣ AI: Artificial Intelligence
- ▣ AIC: Akaike Information Criterion
- ▣ BET: Bucharest Exchange Trading (index)
- ▣ BIST: Borsa İstanbul (BIST 100 index) (the Istanbul Stock Exchange (BIST 100 index))
- ▣ BUX: Budapesti Értéktőzsde index (the Budapest Stock Exchange index)
- ▣ BVB: Bursa de Valori București (the Bucharest Stock Exchange)
- ▣ CD: Chow–Denning (multiple variance-ratio test)
- ▣ COVID-19: COronaVirus Disease 2019
- ▣ EODHD: EOD Historical Data (market-data provider)
- ▣ FTSE: Financial Times Stock Exchange (FTSE Russell, index provider)
- ▣ KPSS: Kwiatkowski–Phillips–Schmidt–Shin (stationarity test)
- ▣ LB: Ljung–Box (test)
- ▣ OMV: Österreichische Mineralölverwaltung (Austrian Mineral Oil Administration (OMV group))
- ▣ PP: Phillips–Perron (unit-root test)
- ▣ PX: Prague Stock Exchange index (index Burzy cenných papírů Praha)
- ▣ RW1: Random Walk 1 (i.i.d. increments)
- ▣ RW2: Random Walk 2 (independent increments)
- ▣ RW3: Random Walk 3 (uncorrelated increments)
- ▣ VR: Variance Ratio
- ▣ WIG20: Warszawski Indeks Giełdowy 20 (the Warsaw Stock Exchange index of the 20 largest companies)

## Today's Question and Route

- **Question:** how can we test whether past prices predict future returns without fooling ourselves?
  - ▶ this seminar comes **before** Lecture 7: the section “What You Need for Today” gives every definition the tasks use
- Route
  - ▶ Part A: variance ratios, Ljung–Box statistics, runs and Dickey–Fuller decisions, on paper
  - ▶ Part B: autocorrelation, unit-root and variance-ratio tests on the S&P 500, DAX, BET, Bitcoin, BVB stocks and emerging markets, each with an interpretation question
  - ▶ Part C: an open question for a project and an AI answer to audit
- Notebook for today: [open the seminar notebook in Google Colab](#); each task names its notebook section
- The seminar is for practice and is not graded; the solutions of [Proposed] tasks are discussed in class

## Exercise Map

| Task   | Question                                                                      | Type                | Model  |
|--------|-------------------------------------------------------------------------------|---------------------|--------|
| A1, A2 | the square-root-of-time rule and variance ratios from autocorrelations        | Solved,<br>Proposed | A1     |
| A3, A4 | Box–Pierce and Ljung–Box statistics, step by step                             | Solved,<br>Proposed | A3     |
| A5, A6 | the runs test on a sequence of signs                                          | Solved,<br>Proposed | A5     |
| A7, A8 | ADF and KPSS decisions from regression output                                 | Solved,<br>Proposed | A7     |
| B1, B2 | ACF, Ljung–Box and robust tests, runs test: S&P 500; BET, Bitcoin, BVB stocks | Solved,<br>Proposed | B1     |
| B3, B4 | ADF, Phillips–Perron and KPSS on prices and returns: DAX; BET, Bitcoin, TLV   | Solved,<br>Proposed | B3     |
| B5, B6 | variance-ratio tests: S&P 500; five emerging markets                          | Solved,<br>Proposed | B5     |
| C1, C2 | has the BET become more efficient? what is wrong in an AI answer?             | Proposed            | B1, B5 |

- **[Solved]**: full solution in the slides and in the notebook, a model to follow; **[Proposed]**: you solve it, following the model

## Data Used

| Series                                     | Source | Price used           | Period         |
|--------------------------------------------|--------|----------------------|----------------|
| S&P 500, DAX                               | EODHD  | close                | 1990–2026      |
| BET                                        | EODHD  | close                | 1997–2026      |
| Bitcoin                                    | EODHD  | close, 7 days a week | 2014–2026      |
| Banca Transilvania (TLV), OMV Petrom (SNP) | EODHD  | adjusted close       | 2010–2026      |
| WIG20, BUX, PX, BIST 100                   | EODHD  | close                | last ten years |

- Daily log returns in %:  $r_t = 100 (\ln P_t - \ln P_{t-1})$ ; weekends and repeated holiday closes are dropped (except Bitcoin); the last day is 18 September 2026
- TLV: the days 30 and 31 May 2016 are removed (a price adjustment applied one day late, [Chapter 2](#))
- In the notebook: `returns('sp500')`, `log_price('dax')`; no account or key is needed

## What You Need for Today (1/5): Efficiency and Random Walks

- **Weak-form efficiency** Fama (1970): past prices cannot be used to earn returns above the return that pays for risk
  - ▶ excess returns are a **martingale difference**:  $E[r_{t+1} - \mu \mid \text{past}] = 0$
- Log price  $p_t = \ln P_t$ , **random walk**:  $p_t = \mu + p_{t-1} + \varepsilon_t$ , so  $r_t = \mu + \varepsilon_t$  Campbell, Lo & MacKinlay (1997)
  - ▶ **RW1**:  $\varepsilon_t$  i.i.d. (independent, identically distributed); **RW2**: independent; **RW3**: uncorrelated
  - ▶  $\text{RW1} \Rightarrow \text{RW2} \Rightarrow \text{RW3}$ ; volatility clustering is allowed only by RW3 and by the martingale
- Under any of them,  $\text{Var}(r_t(q)) = q\sigma^2$  for the  $q$ -day return  $r_t(q) = r_t + \dots + r_{t-q+1}$ 
  - ▶ “square-root-of-time rule”:  $\text{sd}(r_t(q)) = \sigma\sqrt{q}$

## What You Need for Today (2/5): ACF and Portmanteau Tests

- Sample **ACF** (autocorrelation function), with  $e_t = r_t - \bar{r}$ :  $\hat{\rho}(k) = \sum_{t>k} e_t e_{t-k} / \sum_t e_t^2$ 
  - ▶ i.i.d. returns:  $\text{Var}(\hat{\rho}(k)) \approx 1/T$ , 95% band  $\pm 1.96/\sqrt{T}$
  - ▶ robust (volatility clustering):  $\text{Var}(\hat{\rho}(k)) \approx \hat{\delta}(k)/T$ ,  $\hat{\delta}(k) = T \sum_t e_t^2 e_{t-k}^2 / (\sum_t e_t^2)^2$  [Lo & MacKinlay \(1988\)](#)
- $H_0$ :  $\rho(1) = \dots = \rho(m) = 0$ ; three statistics, each  $\approx \chi^2(m)$  under its null
  - ▶ Box–Pierce  $Q_{BP} = T \sum_{k=1}^m \hat{\rho}(k)^2$  [Box & Pierce \(1970\)](#); **LB** (Ljung–Box)  
 $Q_{LB} = T(T+2) \sum_{k=1}^m \hat{\rho}(k)^2 / (T-k)$  [Ljung & Box \(1978\)](#)
  - ▶ robust  $\tilde{Q} = \sum_{k=1}^m T \hat{\rho}(k)^2 / \hat{\delta}(k)$  [Escanciano & Lobato \(2009\)](#): valid for a martingale with volatility clustering
- 5% critical values:  $\chi_{0.95}^2(3) = 7.81$ ,  $\chi_{0.95}^2(5) = 11.07$ ,  $\chi_{0.95}^2(10) = 18.31$

## What You Need for Today (3/5): the Runs Test

- ▣ A **run**: a maximal sequence of returns with the same sign [Wald & Wolfowitz \(1940\)](#); zero returns are dropped
  - ▶  $++---+-$ : 4 runs;  $n_1 = 3$  positive,  $n_2 = 4$  negative,  $n = n_1 + n_2 = 7$
- ▣ Under independence of the signs:
  - ▶  $E[R] = \frac{2n_1n_2}{n} + 1$ ,  $\text{Var}(R) = \frac{2n_1n_2(2n_1n_2 - n)}{n^2(n - 1)}$
  - ▶  $z = (R - E[R]) / \sqrt{\text{Var}(R)} \approx N(0, 1)$ ; reject at 5% if  $|z| > 1.96$
- ▣ Reading  $z$ 
  - ▶  $z < 0$ : too few runs, signs persist (positive dependence, trends)
  - ▶  $z > 0$ : too many runs, signs alternate (negative dependence, reversals)
- ▣ Signs only: the test ignores the size of the returns, so heavy tails and volatility do not affect it

## What You Need for Today (4/5): Unit Roots

- $I(0)$ : stationary (constant mean and variance, shocks fade);  $I(1)$ : the first difference is  $I(0)$ ; a random walk is  $I(1)$
- Dickey–Fuller regression [Dickey & Fuller \(1979\)](#):  $\Delta p_t = c [+ bt] + \gamma p_{t-1} [+ \sum_j \delta_j \Delta p_{t-j}] + \varepsilon_t$ 
  - ▶  $H_0: \gamma = 0$  (unit root);  $\tau = \hat{\gamma}/\text{SE}(\hat{\gamma})$ ; reject if  $\tau < \text{critical value}$
  - ▶ 5% critical values [MacKinnon \(1996\)](#):  $-2.86$  (constant),  $-3.41$  (constant and trend); not  $-1.645$
  - ▶ **ADF** (augmented Dickey–Fuller): with the lagged differences [Said & Dickey \(1984\)](#); **PP** (Phillips–Perron): without them, corrected  $\tau$  [Phillips & Perron \(1988\)](#)
- **KPSS** [Kwiatkowski et al. \(1992\)](#):  $H_0$  stationarity; reject if the statistic exceeds  $0.463$  (constant) or  $0.146$  (trend)
  - ▶ ADF does not reject + KPSS rejects:  $I(1)$ ; ADF rejects + KPSS does not:  $I(0)$ ; otherwise: inconclusive

## What You Need for Today (5/5): Variance Ratios

- **VR** (variance ratio):  $VR(q) = \frac{\text{Var}(r_t(q))}{q \text{Var}(r_t)} = 1 + 2 \sum_{k=1}^{q-1} \left(1 - \frac{k}{q}\right) \rho(k)$  [Lo & MacKinlay \(1988\)](#)
  - ▶ = 1: random walk; > 1: momentum (positive autocorrelation); < 1: mean reversion (negative)
- Tests with  $T$  returns
  - ▶ i.i.d.:  $Z(q) = (\widehat{VR}(q) - 1) / \sqrt{2(2q - 1)(q - 1)/(3qT)}$
  - ▶ robust:  $Z^*(q) = (\widehat{VR}(q) - 1) / \sqrt{\hat{\theta}(q)/T}$ ,  $\hat{\theta}(q) = \sum_{j=1}^{q-1} [2(q - j)/q]^2 \hat{\delta}(j)$
- Several horizons: **CD** (Chow–Denning) =  $\max_i |Z^*(q_i)|$  [Chow & Denning \(1993\)](#); 5% critical value 2.49 for  $m = 4$  horizons
  - ▶ the same rule for  $m$  markets: each test at level  $1 - 0.95^{1/m}$  (Šidák)
  - ▶ automatic VR [Choi \(1999\)](#): the horizon is chosen from the data; p-value by wild bootstrap [Kim \(2009\)](#)

## A1: Variance Ratios from Autocorrelations [Solved]

### Task

- ▣ Daily returns have a standard deviation  $\sigma = 1.2\%$ .
- ▣ 1. Under RW1, compute the standard deviation of the 5-day and of the 20-day return.
- ▣ 2. Suppose  $\rho(1) = 0.10$ ,  $\rho(2) = 0.05$  and  $\rho(k) = 0$  for  $k \geq 3$ ; compute VR(2) and VR(5).
- ▣ 3. Treat these as estimates from  $T = 2500$  i.i.d. returns and compute  $Z(2)$  and  $Z(5)$ .
- ▣ 4. Compute the 5-day standard deviation implied by VR(5).
- ▣ Report: two standard deviations, two VR, two  $Z$  and one sentence.

### Solution

- ▣ 1.  $1.2\sqrt{5} = 2.68\%$ ;  $1.2\sqrt{20} = 5.37\%$
- ▣ 2.  $VR(2) = 1 + 0.10 = 1.10$ ;  $VR(5) = 1 + 2(0.8 \times 0.10 + 0.6 \times 0.05) = 1.220$
- ▣ 3.  $\sqrt{2 \cdot 3 \cdot 1 / (6 \cdot 2500)} = 0.020$ ,  $Z(2) = 5.00$ ;  
 $\sqrt{2 \cdot 9 \cdot 4 / (15 \cdot 2500)} = 0.0438$ ,  $Z(5) = 5.02$
- ▣ 4.  $1.2\sqrt{5} \times 1.220 = 2.96\%$  instead of 2.68%: momentum makes weekly risk larger than the  $\sqrt{q}$  rule says.

## A2: Mean Reversion in a Volatile Asset [Proposed]

Solution discussed in class

### Task

- ▣ Daily returns:  $\sigma = 2\%$ ,  $\rho(1) = -0.08$ ,  $\rho(2) = 0.03$ ,  $\rho(k) = 0$  for  $k \geq 3$ ,  $T = 1000$ . Model: A1.
- ▣ 1. Compute  $VR(2)$  and  $VR(5)$ .
- ▣ 2. Compute  $Z(2)$  and  $Z(5)$  and decide at 5%.
- ▣ 3. Compare the 5-day standard deviation implied by  $VR(5)$  with  $2\sqrt{5}$ .
- ▣ Report: two  $VR$ , two  $Z$ , two standard deviations and one sentence.

## A3: Box–Pierce and Ljung–Box, Step by Step [Solved]

## Task

- ▣  $T = 500$  daily returns;  
 $\hat{\rho}(1), \dots, \hat{\rho}(5) = 0.12, -0.05, 0.04, 0.03, -0.06$ .
- ▣ 1. Compute  $Q_{BP}(5)$ .
- ▣ 2. Compute  $Q_{LB}(5)$ .
- ▣ 3. Compare with  $\chi^2_{0.95}(5) = 11.07$  and decide.
- ▣ 4. Say which lag drives the result.
- ▣ Report: two statistics, a decision and one sentence.

## Solution

- ▣ 1.  $T\hat{\rho}(k)^2$ : 7.20, 1.25, 0.80, 0.45, 1.80;  $Q_{BP}(5) = 11.50$
- ▣ 2.  $500 \cdot 502 \hat{\rho}(k)^2 / (500 - k)$ : 7.24, 1.26, 0.81, 0.46, 1.83;  
 $Q_{LB}(5) = 11.59$
- ▣ 3.  $11.59 > 11.07$ : reject  $H_0$  at 5% ( $p = 0.041$ )
- ▣ 4. Lag 1 alone gives 7.20 of 11.50: one autocorrelation of 0.12 drives the rejection.

## A4: Ljung–Box on Squared Returns [Proposed]

Solution discussed in class

### Task

- ▣  $T = 750$ ; the ACF of the squared returns: 0.25, 0.20, 0.18, 0.15, 0.12 at lags 1–5. Model: A3.
- ▣ 1. Compute  $Q_{LB}(5)$  and decide at 5%.
- ▣ 2. Say which of RW1, RW3 and the martingale this rejects.
- ▣ 3. Say whether weak-form efficiency is rejected.
- ▣ Report: one statistic and two sentences.

## A5: The Runs Test on 20 Days [Solved]

### Task

- ▣ Signs of 20 daily returns:  
++-++++--+- - -++-+- -++.
- ▣ 1. Count  $n_1$ ,  $n_2$  and the number of runs  $R$ .
- ▣ 2. Compute  $E[R]$  and  $\text{Var}(R)$ .
- ▣ 3. Compute  $z$  and decide at 5%.
- ▣ Report:  $R$ ,  $E[R]$ ,  $z$  and one sentence.

### Solution

- ▣ 1.  $n_1 = 11$ ,  $n_2 = 9$ ,  $R = 11$
- ▣ 2.  $E[R] = 2 \cdot 11 \cdot 9/20 + 1 = 10.90$ ;  
 $\text{Var}(R) = 198 \cdot 178/(400 \cdot 19) = 4.64$
- ▣ 3.  $z = (11 - 10.90)/2.15 = 0.05$  ( $p = 0.963$ ): do not reject independence.
- ▣ With 20 days the test has little power: only large departures can be detected.

## A6: A Sequence with Long Runs [Proposed]

Solution discussed in class

### Task

- ▣ Signs of 24 daily returns:  $+++--++--++--++--++--$ . Model: A5.
- ▣ 1. Count  $n_1$ ,  $n_2$  and  $R$ .
- ▣ 2. Compute  $E[R]$ ,  $\text{Var}(R)$  and  $z$ .
- ▣ 3. Decide at 5% and say what kind of dependence the signs show.
- ▣ Report:  $R$ ,  $E[R]$ ,  $z$  and one sentence.

## A7: ADF and KPSS from Regression Output [Solved]

### Task

- ▣ Log price,  $T = 2500$ :  
 $\Delta p_t = 0.021 - 0.0028 p_{t-1} + 0.05 \Delta p_{t-1} + e_t$ ,  
 $SE(\hat{\gamma}) = 0.0019$ ; KPSS (constant) = 2.10.
- ▣ Returns:  $\Delta r_t = 0.03 - 0.92 r_{t-1} + e_t$ ,  $SE(\hat{\gamma}) = 0.020$ ; KPSS (constant) = 0.09.
- ▣ 1. Compute  $\tau$  for the price and for the returns.
- ▣ 2. Decide with the 5% critical value for a constant.
- ▣ 3. Decide with KPSS and give the order of integration of each series.
- ▣ Report: two  $\tau$ , four decisions and one sentence.

### Solution

- ▣ 1. Price:  $\tau = -0.0028/0.0019 = -1.47$ ; returns:  
 $\tau = -0.92/0.020 = -46.0$
- ▣ 2. Price:  $-1.47 > -2.86$ : do not reject a unit root; returns:  
 $-46.0 < -2.86$ : reject
- ▣ 3. KPSS: price  $2.10 > 0.463$ : reject stationarity; returns  
 $0.09 < 0.463$ : do not reject
- ▣ The price is  $I(1)$ , the returns  $I(0)$ : the picture of a random walk, but not yet a proof of it.

## A8: When the Tests Disagree [Proposed]

Solution discussed in class


### Task

- ▣ Log price of a stock,  $T = 1500$ , constant and trend:  $\hat{\gamma} = -0.0105$ ,  $SE(\hat{\gamma}) = 0.0029$ ; KPSS (trend) = 0.21. Model: A7.
- ▣ 1. Compute  $\tau$  and decide with the 5% critical value for a constant and a trend.
- ▣ 2. Decide with KPSS.
- ▣ 3. Combine the two decisions and give two possible reasons for the conflict.
- ▣ Report:  $\tau$ , two decisions and two sentences.

## B1: Autocorrelation in the S&P 500 [Solved]

- **Question:** does the verdict on the autocorrelation of daily S&P 500 returns depend on whether we allow for volatility clustering?
- **Inputs:** S&P 500 closes since 1990, daily log returns in %
- **Tasks**
  1. Draw the ACF of the returns (lags 1–20) with the i.i.d. and the robust 95% bands, and the ACF of the squared returns.
  2. Report  $\hat{\rho}(1), \dots, \hat{\rho}(5)$  of the returns and of the squared returns.
  3. Compute  $Q_{LB}(10)$  and the robust  $\tilde{Q}(10)$ , with p-values, for the returns and for the squared returns.
  4. Say which random walk hypothesis each test rejects.
  5. Interpretation: is weak-form efficiency rejected?
- **Report:** the chart, ten autocorrelations, four statistics with p-values and two sentences
- **Notebook:** section B1

## B1: Solution [Solved]



- $\hat{\rho}(1..5)$ , returns:  $-0.079, -0.004, -0.006, -0.022, -0.016$ ; squared:  $0.311, 0.408, 0.250, 0.298, 0.308$ ;  $T = 9\,240$
- Returns:  $Q_{LB}(10) = 90.8$  ( $p < 0.001$ ),  $\tilde{Q}(10) = 18.8$  ( $p = 0.043$ ); squared:  $Q_{LB}(10) = 8015$ ,  $\tilde{Q}(10) = 189.4$  (both  $p < 0.001$ )
- Interpretation: squared returns reject RW1 and RW2 (volatility clustering); the robust test rejects RW3 only barely, because of a small negative  $\hat{\rho}(1)$ : a weak, short-lived reversal


## B2: BET, Bitcoin and Two BVB Stocks [Proposed]

- **Question:** for which of the BET, Bitcoin, Banca Transilvania and OMV Petrom does the verdict on autocorrelation change between the classic and the robust test? Model: B1.
- **Inputs:** BET since 1997, Bitcoin since 2014, TLV and SNP since 2010 (TLV without 30–31 May 2016), daily log returns in %
- **Tasks**
  1. Compute  $\hat{\rho}(1)$ ,  $Q_{LB}(10)$  and  $\tilde{Q}(10)$  with p-values for each series.
  2. Compute the runs test  $z$  and its p-value for each series.
  3. Mark the series for which the classic and the robust test disagree at 5%.
  4. Interpretation: why is the BET different from the other three?
- **Report:** one table and two sentences
- **Notebook:** section B2

## B3: Unit Roots in the DAX [Solved]

- **Question:** what is the order of integration of the DAX log price and of its returns?
- **Inputs:** DAX closes since 1990; log price and daily log returns
- **Tasks**
  1. Draw the log price with a fitted linear trend, and the returns.
  2. Run ADF (lag order by AIC), PP and KPSS on the log price with a constant and a trend.
  3. Run the same three tests on the returns with a constant.
  4. Interpretation: does a unit root in the DAX price prove that the DAX is weak-form efficient?
- **Report:** the chart, six statistics with p-values and two sentences
- **Notebook:** section B3

## B3: Solution [Solved]



|                    | ADF                    | PP                     | KPSS                |
|--------------------|------------------------|------------------------|---------------------|
| log price (trend)  | -2.69 ( $p = 0.239$ )  | -2.65 ( $p = 0.260$ )  | 0.94 ( $p < 0.01$ ) |
| returns (constant) | -23.30 ( $p < 0.001$ ) | -97.40 ( $p < 0.001$ ) | 0.04 ( $p > 0.10$ ) |

- ADF used 30 lags for the price (AIC); PP and KPSS use 38 Bartlett lags;  $T = 9288$
- Price: no rejection by ADF or PP, rejection by KPSS:  $I(1)$ ; returns: the opposite:  $I(0)$
- Interpretation: no; a unit root is necessary for a random walk but allows predictable increments; efficiency is tested on the returns (B1, B5)


## B4: BET, Bitcoin and TLV [Proposed]

- **Question:** do the three unit-root tests agree for the BET, Bitcoin and Banca Transilvania? Model: B3.
- **Inputs:** BET since 1997, Bitcoin since 2014, TLV since 2010 (without 30–31 May 2016)
- **Tasks**
  1. Run ADF, PP and KPSS on each log price (constant and trend) and on each return series (constant).
  2. Classify each series as  $I(1)$ ,  $I(0)$  or inconclusive.
  3. For an inconclusive case, propose one check that could settle it.
  4. Interpretation: would you trade a “mean-reverting” TLV price on this evidence?
- **Report:** one table, three classifications and two sentences
- **Notebook:** section B4

## B5: Variance Ratios of the S&P 500 [Solved]

- **Question:** is the S&P 500 mean-reverting over horizons of 2 to 20 days, once volatility clustering is taken into account?
- **Inputs:** S&P 500 closes since 1990, daily log returns in %
- **Tasks**
  1. Compute  $VR(q)$ ,  $Z(q)$  and  $Z^*(q)$  for  $q = 2, 5, 10, 20$ .
  2. Compute the Chow–Denning statistic and compare it with its 5% critical value.
  3. Draw  $VR(q)$  for  $q = 2, \dots, 40$  with the i.i.d. and the robust 95% bands.
  4. Interpretation: is the mean reversion large enough to matter for an investor?
- **Report:** a table of twelve numbers, one statistic, the chart and two sentences
- **Notebook:** section B5

## B5: Solution [Solved]



|          | $q = 2$ | $q = 5$ | $q = 10$ | $q = 20$ |
|----------|---------|---------|----------|----------|
| $VR(q)$  | 0.922   | 0.856   | 0.783    | 0.767    |
| $Z(q)$   | -7.54   | -6.33   | -6.17    | -4.50    |
| $Z^*(q)$ | -3.37   | -2.76   | -2.73    | -2.06    |

- ▣  $T = 9240$ ; Chow–Denning:  $\max |Z^*| = 3.37 > 2.49$  ( $p = 0.003$ ): reject the random walk
- ▣  $Z(2)$  is about 2.2 times  $Z^*(2)$ : ignoring volatility clustering exaggerates the evidence
- ▣ Interpretation: significant, but small: the weekly variance is about 86% of five daily variances; after trading costs this is hardly a profit

## B6: Five Emerging Markets [Proposed]

- **Question:** which of the WIG20, BUX, PX, BET and BIST 100 departs from a random walk over the last ten years? Model: B5.
- **Inputs:** daily log returns from 19 September 2016 to 18 September 2026
- **Tasks**
  1. Compute  $VR(5)$ ,  $Z^*(5)$  and its p-value for each market.
  2. Compute the Chow–Denning p-value ( $q = 2, 5, 10, 20$ ) and the automatic VR test with its wild-bootstrap p-value.
  3. Compute the Šidák critical value for five markets tested together at 5%.
  4. Interpretation: which markets would you call inefficient once you account for testing five markets?
- **Report:** one table, one critical value and two sentences
- **Notebook:** section B6

## C1: Has the BET Become More Efficient? [Proposed]

- **Question:** how has the predictability of the BET evolved since 1997, including around the FTSE Russell upgrade of September 2020?
- **Inputs:** BET daily log returns since 1997; models: B1, B5
- **Tasks**
  1. Compute  $VR(5)$  and  $Z^*(5)$  on rolling windows of 500 days, one every 21 days, and the share of windows with  $|Z^*(5)| > 1.96$  in the first and in the last five years.
  2. Compute  $\hat{\rho}(1)$ ,  $VR(5)$ ,  $Z^*(5)$  and the Chow–Denning p-value in 1997–2006, 2007–2016 and 2017–2026.
  3. Compare  $VR(5)$  in the five years before and after 21 September 2020.
  4. List what else changed in the Romanian market at the same time.
  5. Interpretation: how would you separate these changes from the effect of the upgrade?
- **Report:** a table, one chart and a plan for a project
- **Notebook:** section C1

## C2: Audit an AI Answer [Proposed]

- A student asked an AI assistant whether the DAX (daily, since 1990) is weak-form efficient. The answer:
- (a) ADF on the DAX log price gives  $\tau = -2.69$ ,  $p = 0.24$ : the unit root is not rejected, so the DAX is weak-form efficient.
- (b) Ljung-Box  $Q(10) = 21.83$ ,  $p = 0.016$ : DAX returns are predictable, so the DAX is not efficient.
- (c) For squared returns  $Q(10) = 3747$ : DAX returns are not a martingale.
- (d)  $VR(10) = 0.93 < 1$  shows positive autocorrelation (momentum) in the DAX.
- (e) Under RW1,  $VR(q) = 1$  at every horizon and the variance of  $q$ -day returns is  $q$  times the daily variance.
- **Tasks**
  - ▶ 1. For each statement, say whether it is correct; if not, give the correct statement and, where possible, the correct number from the notebook (section C2).
  - ▶ 2. Report: a list of five verdicts with one line of justification each.

## What You Should Take from Today

- ▣ A unit root in prices is the starting point, not the answer: efficiency is about the returns
- ▣ Test the martingale with robust statistics ( $\tilde{Q}$ ,  $Z^*$ ): classic tests mistake volatility clustering for predictability
- ▣  $VR > 1$ : momentum;  $VR < 1$ : mean reversion; the S&P 500 reverts slightly, the old BET trended
- ▣ Many horizons or many markets: use Chow–Denning or a Šidák correction
- ▣ An AI answer is a draft: check the null of each test, the sign of  $VR - 1$  and the robust version

## After the Seminar

- Lecture 7 develops each topic of today: efficiency, random walks, ACF tests, unit roots, variance ratios, adaptive markets
- Try the [Proposed] tasks in the notebook; the solutions are discussed in class
- C1 can grow into a team project: BET stocks, WIG20 and BUX as controls, a block bootstrap of the difference
- Reading: [Franke, Härdle & Hafner \(2019\)](#), Ch. 11; exercises in [Borak, Härdle & López-Cabrera \(2013\)](#), Ch. 11; [Campbell, Lo & MacKinlay \(1997\)](#), Ch. 2

## References

- Borak, S., Härdle, W.K., López-Cabrera, B. (2013). *Statistics of Financial Markets: Exercises and Solutions*, 2nd ed. Springer.
- Box, G.E.P., Pierce, D.A. (1970). Distribution of residual autocorrelations in autoregressive-integrated moving average time series models. *Journal of the American Statistical Association*, 65(332), 1509–1526.
- Campbell, J.Y., Lo, A.W., MacKinlay, A.C. (1997). *The Econometrics of Financial Markets*. Princeton University Press.
- Choi, I. (1999). Testing the random walk hypothesis for real exchange rates. *Journal of Applied Econometrics*, 14(3), 293–308.
- Chow, K.V., Denning, K.C. (1993). A simple multiple variance ratio test. *Journal of Econometrics*, 58(3), 385–401.
- Dickey, D.A., Fuller, W.A. (1979). Distribution of the estimators for autoregressive time series with a unit root. *Journal of the American Statistical Association*, 74(366), 427–431.
- Escanciano, J.C., Lobato, I.N. (2009). An automatic Portmanteau test for serial correlation. *Journal of Econometrics*, 151(2), 140–149.
- Fama, E.F. (1970). Efficient capital markets: a review of theory and empirical work. *Journal of Finance*, 25(2), 383–417.
- Franke, J., Härdle, W.K., Hafner, C.M. (2019). *Statistics of Financial Markets: An Introduction*, 5th ed. Springer.
- Kim, J.H. (2009). Automatic variance ratio test under conditional heteroskedasticity. *Finance Research Letters*, 6(3), 179–185.
- Kwiatkowski, D., Phillips, P.C.B., Schmidt, P., Shin, Y. (1992). Testing the null hypothesis of stationarity against the alternative of a unit root. *Journal of Econometrics*, 54(1–3), 159–178.
- Ljung, G.M., Box, G.E.P. (1978). On a measure of lack of fit in time series models. *Biometrika*, 65(2), 297–303.
- Lo, A.W., MacKinlay, A.C. (1988). Stock market prices do not follow random walks: evidence from a simple specification test. *Review of Financial Studies*, 1(1), 41–66.
- MacKinnon, J.G. (1996). Numerical distribution functions for unit root and cointegration tests. *Journal of Applied Econometrics*, 11(6), 601–618.
- Phillips, P.C.B., Perron, P. (1988). Testing for a unit root in time series regression. *Biometrika*, 75(2), 335–346.
- Said, S.E., Dickey, D.A. (1984). Testing for unit roots in autoregressive-moving average models of unknown order. *Biometrika*, 71(3), 599–607.
- Wald, A., Wolfowitz, J. (1940). On a test whether two samples are from the same population. *Annals of Mathematical Statistics*, 11(2), 147–162.