



Time Series Analysis and Forecasting

Chapter 7: Cointegration and VECM



Daniel Traian PELE

Bucharest University of Economic Studies

IDA Institute Digital Assets

Blockchain Research Center

AI4EFin Artificial Intelligence for Energy Finance

Romanian Academy, Institute for Economic Forecasting

MSCA Digital Finance

Outline

- Motivation
- Spurious Regression
- Cointegration Concept
- Engle-Granger Method
- Johansen Method
- VECM Estimation
- Practical Considerations
- Real-World Examples
- Case Study: Interest Rates
- Summary
- Quiz

Why Cointegration Matters

The Challenge

- ▣ Many economic/financial time series are **non-stationary** ($I(1)$)
- ▣ GDP, stock prices, exchange rates, interest rates all have unit roots
- ▣ Standard regression with $I(1)$ variables \Rightarrow **spurious results**
- ▣ Differencing removes non-stationarity but loses **long-run information**

The Solution: Cointegration

Some non-stationary series share a **common stochastic trend**—they move together in the long run.

Nobel Prize 2003

Granger & Engle received the Nobel Prize for “methods for analyzing economic time series with common trends.”

Real-World Applications

Finance

- ▣ **Pairs Trading:** Cointegrated stocks
- ▣ **Term Structure:** Interest rates
- ▣ **Spot-Futures:** Arbitrage

Macroeconomics

- ▣ **Consumption & Income**
- ▣ **Money & Prices**
- ▣ **PPP:** Exchange rates

Policy Analysis

- ▣ **Fiscal:** Spending & taxes
- ▣ **Monetary:** Rate pass-through
- ▣ **Labor:** Wages & productivity

The Spurious Regression Problem

Granger & Newbold (1974)

Regressing one random walk on another **independent** random walk: $Y_t = \alpha + \beta X_t + u_t$ where Y_t and X_t are independent $I(1)$ processes.


Symptoms of Spurious Regression

- ▣ High R^2 (often > 0.9) even though variables are **unrelated**
- ▣ Highly significant t -statistics (reject $H_0 : \beta = 0$)
- ▣ Very low Durbin-Watson statistic ($DW \approx 0$)
- ▣ Residuals are non-stationary (have unit root)

Rule of Thumb

If $R^2 > DW$, suspect spurious regression!

Spurious Regression: Visual Example



Definition of Cointegration

Definition 1 (Cointegration (Engle & Granger, 1987))

Variables $Y_{1t}, Y_{2t}, \dots, Y_{kt}$ are **cointegrated of order (d, b)** , written $CI(d, b)$, if:


1. All variables are integrated of order d : $Y_{it} \sim I(d)$
2. There exists a linear combination $\beta'Y_t = \beta_1 Y_{1t} + \dots + \beta_k Y_{kt}$ that is integrated of order $(d - b)$, where $b > 0$

Most Common Case: $CI(1, 1)$

- ▣ Variables are $I(1)$ (have unit roots)
- ▣ Linear combination is $I(0)$ (stationary)
- ▣ Vector $\beta = (\beta_1, \dots, \beta_k)'$ is the **cointegrating vector**

The cointegrating vector is unique only up to scalar multiplication. Usually normalized: $\beta_1 = 1$.

Cointegration: Visual Example



Intuition: Common Stochastic Trends

Why Does Cointegration Occur?

Cointegrated variables share **common stochastic trends**: $Y_{1t} = \gamma_1 \tau_t + S_{1t}$, $Y_{2t} = \gamma_2 \tau_t + S_{2t}$ where τ_t is a common random walk and S_{it} are stationary.

Linear Combination Eliminates the Trend

$$\gamma_2 Y_{1t} - \gamma_1 Y_{2t} = \gamma_2 S_{1t} - \gamma_1 S_{2t} \sim I(0)$$

Economic Interpretation

- Cointegration = **long-run equilibrium relationship**
- Variables may deviate in the short run, but are “pulled back”
- The cointegrating vector defines the equilibrium

Cointegrating Rank

How Many Cointegrating Relationships?

For k variables that are $I(1)$:

- Maximum possible cointegrating relationships: $r = k - 1$
- If $r = 0$: No cointegration (variables drift apart)
- If $r = k$: All variables are $I(0)$ (contradiction)

Example: 3 Variables

- $r = 0$: No cointegration
- $r = 1$: One cointegrating relationship
- $r = 2$: Two cointegrating relationships (only 1 common trend)

The number of common stochastic trends = $k - r$

Engle-Granger Two-Step Method

Step 1: Estimate Cointegrating Regression

Run OLS: $Y_t = \alpha + \beta X_t + e_t$. Save residuals: $\hat{e}_t = Y_t - \hat{\alpha} - \hat{\beta} X_t$

Step 2: Test Residuals for Stationarity

Test if \hat{e}_t is $I(0)$ using ADF: $\Delta \hat{e}_t = \rho \hat{e}_{t-1} + \sum_{j=1}^p \gamma_j \Delta \hat{e}_{t-j} + v_t$

- $H_0: \rho = 0$ (unit root \Rightarrow no cointegration)
- $H_1: \rho < 0$ (stationary \Rightarrow cointegration)

Important

Use **Engle-Granger critical values**, not standard ADF! (More negative because residuals are estimated)

Engle-Granger Critical Values

Critical Values for Cointegration Test

Variables	1%	5%	10%
2	-3.90	-3.34	-3.04
3	-4.29	-3.74	-3.45
4	-4.64	-4.10	-3.81
5	-4.96	-4.42	-4.13

MacKinnon (1991), $T = 100$

Limitations of Engle-Granger

- Only tests for **one** cointegrating relationship
- Results depend on choice of dependent variable
- Small sample bias; cannot test hypotheses on cointegrating vector

Johansen Cointegration Test

Advantages over Engle-Granger

- ▣ Tests for **multiple** cointegrating relationships
- ▣ Maximum likelihood estimation (more efficient)
- ▣ Can test restrictions on cointegrating vectors
- ▣ Does not require choosing a dependent variable

Starting Point: VAR in Levels

$$Y_t = c + A_1 Y_{t-1} + A_2 Y_{t-2} + \cdots + A_p Y_{t-p} + \varepsilon_t$$

Rewrite in **Vector Error Correction** form...

VECM Representation

Vector Error Correction Model

$$\Delta Y_t = c + \Pi Y_{t-1} + \sum_{j=1}^{p-1} \Gamma_j \Delta Y_{t-j} + \varepsilon_t$$

- $\Pi = \sum_i A_i - I$ (long-run impact); Γ_j (short-run dynamics)

Key Insight: Rank of Π

The **rank of Π** determines cointegration:

- $\text{rank}(\Pi) = 0$: No cointegration (VAR in differences)
- $\text{rank}(\Pi) = k$: All variables are $I(0)$ (VAR in levels)
- $0 < \text{rank}(\Pi) = r < k$: r cointegrating vectors

Decomposition of Π

When $\text{rank}(\Pi) = r < k$

$\Pi = \alpha\beta'$ where β ($k \times r$) = **cointegrating vectors**, α ($k \times r$) = **adjustment coefficients**

Interpretation

- ▣ $\beta'Y_{t-1}$ = deviations from equilibrium (error correction terms)
- ▣ α = speed of adjustment; rows show each variable's response

$$\text{VECM: } \Delta Y_t = c + \alpha(\beta'Y_{t-1}) + \sum_{j=1}^{p-1} \Gamma_j \Delta Y_{t-j} + \varepsilon_t$$

Johansen Test Statistics

Two Test Statistics

Based on eigenvalues $\hat{\lambda}_1 > \hat{\lambda}_2 > \dots > \hat{\lambda}_k$ of a certain matrix:

Trace Test:

$$\lambda_{\text{trace}}(r) = -T \sum_{i=r+1}^k \ln(1 - \hat{\lambda}_i)$$

Tests $H_0: \text{rank} \leq r$ vs $H_1: \text{rank} > r$

Maximum Eigenvalue Test:


$$\lambda_{\text{max}}(r, r+1) = -T \ln(1 - \hat{\lambda}_{r+1})$$

Tests $H_0: \text{rank} = r$ vs $H_1: \text{rank} = r+1$

Critical values from Johansen & Juselius (1990), depend on:

- Number of variables k
- Deterministic components (constant, trend)

Johansen Test: Visual Interpretation



charts/johansen_eigenvalues.pdf

Significant eigenvalues (above threshold) indicate cointegrating relationships. First eigenvalue significant $\Rightarrow r = 1$.

Testing Procedure

Sequential Testing (Trace Test)

1. Test $H_0: r = 0$. If rejected \Rightarrow continue
2. Test $H_0: r \leq 1$. If not rejected $\Rightarrow r = 1$
3. Continue until H_0 is not rejected

Deterministic Components

- ☐ No constant, no trend (rarely used)
- ☐ Constant in cointegrating relation only
- ☐ **Constant in both** (most common)
- ☐ Constant + trend in cointegrating relation

VECM Structure

Full VECM Specification

For $k = 2$ variables with $r = 1$ cointegrating relation:


$$\Delta Y_{1t} = c_1 + \alpha_1(Y_{1,t-1} - \beta Y_{2,t-1}) + \gamma_{11}\Delta Y_{1,t-1} + \gamma_{12}\Delta Y_{2,t-1} + \varepsilon_{1t}$$

$$\Delta Y_{2t} = c_2 + \alpha_2(Y_{1,t-1} - \beta Y_{2,t-1}) + \gamma_{21}\Delta Y_{1,t-1} + \gamma_{22}\Delta Y_{2,t-1} + \varepsilon_{2t}$$

Components

- $(Y_{1,t-1} - \beta Y_{2,t-1})$ = error correction term (deviation from equilibrium)
- α_1, α_2 = adjustment speeds (should have opposite signs)
- γ_{ij} = short-run dynamics
- ε_{it} = innovations

Error Correction Mechanism: Visual



Interpreting Adjustment Coefficients

The α Coefficients

If the cointegrating relation is $Y_1 - \beta Y_2 = 0$ (equilibrium):

- ▣ $\alpha_1 < 0$: Y_1 adjusts downward when above equilibrium
- ▣ $\alpha_2 > 0$: Y_2 adjusts upward when Y_1 is above equilibrium

Weak Exogeneity

If $\alpha_i = 0$, variable Y_i does **not** respond to disequilibrium.

- ▣ Y_i is **weakly exogenous** for the long-run parameters
- ▣ The other variable does all the adjusting
- ▣ Can simplify estimation (single-equation approach)

Test weak exogeneity: $H_0 : \alpha_i = 0$ using likelihood ratio test.

VECM vs VAR in Differences


When Variables are Cointegrated

	VAR in Differences	VECM
Long-run info	Lost	Preserved
Short-run dynamics	Yes	Yes
Error correction	No	Yes
Forecasting	Poor (long-run)	Better
IRF interpretation	Short-run only	Both

Granger Representation Theorem

If variables are cointegrated, there **must** exist an error correction representation. Ignoring cointegration = model misspecification!

VECM Impulse Response Functions



Practical Workflow

Step-by-Step Procedure

1. **Unit Root Tests:** Verify all variables are $I(1)$
 - ▶ ADF, KPSS on levels and first differences
2. **Lag Length Selection:** Choose p for VAR in levels
 - ▶ Use AIC, BIC, or sequential LR tests
3. **Cointegration Test:** Johansen trace/max-eigenvalue tests
 - ▶ Determine cointegrating rank r
4. **Estimate VECM:** If $0 < r < k$
 - ▶ Estimate α , β , Γ_j
5. **Diagnostics:** Check residuals for autocorrelation, normality
6. **Analysis:** IRF, FEVD, hypothesis tests

Common Pitfalls


Things to Watch Out For

- ▣ **Structural breaks:** Cause spurious unit roots or cointegration
- ▣ **Near-unit-root:** Tests have low power
- ▣ **Lag selection:** Too many/few lags bias results
- ▣ **Small samples:** Johansen test oversized


Recommendation

Always check: residual diagnostics, stability of cointegrating relationship, sensitivity to specification


Example 1: Term Structure of Interest Rates



Example 2: Pairs Trading in Finance



Example 3: Purchasing Power Parity (PPP)



Case Study: Cointegration of Interest Rates

Research Question

Are short-term and long-term interest rates cointegrated? Does the expectations hypothesis of the term structure hold?

Data

- US Monthly Data (1962-2023)
- 3-Month Treasury Bill Rate
- 10-Year Treasury Bond Yield
- Source: FRED Database

Methodology

- Unit root tests (ADF, PP)
- Engle-Granger cointegration test
- Johansen procedure
- VECM estimation
- Impulse response analysis

Step 1: Data Visualization



charts/ch7_case_raw_data.pdf

Step 2: Unit Root Tests

charts/ch7_case_unit_root.pdf

Step 3: Engle-Granger Cointegration Test

charts/ch7_case_cointegration.pdf

Step 4: VECM Estimation

charts/ch7_case_vecm.pdf

Step 5: Impulse Response Functions

charts/ch7_case_irf.pdf

Step 6: Forecasting

charts/ch7_case_forecast.pdf

Key Takeaways

Main Concepts

- ▣ **Cointegration:** $I(1)$ variables with stationary linear combination
- ▣ **Spurious regression:** High R^2 with unrelated $I(1)$ variables
- ▣ **VECM:** VAR with error correction for cointegrated systems

Testing Methods

- ▣ **Engle-Granger:** Simple, one vector only
- ▣ **Johansen:** Multiple vectors, MLE-based

Remember

Tests have low power in small samples. Theory should guide specification.

What's Next?

Extensions and Related Topics

- ▣ **Structural VECM:** Identifying structural shocks
- ▣ **Threshold cointegration:** Nonlinear adjustment
- ▣ **Panel cointegration:** Multiple cross-sections
- ▣ **Fractional cointegration:** Long memory
- ▣ **Time-varying cointegration:** Regime changes

Questions?

Quick Quiz

1. What does it mean for two $I(1)$ variables to be cointegrated?
2. What is the “spurious regression” problem?
3. In a VECM, what do the α coefficients represent?
4. What is the main advantage of Johansen over Engle-Granger?
5. If $\alpha_i = 0$ for variable Y_i , what does this imply?

Quiz Answers

1. **Cointegration:** A linear combination of the variables is $I(0)$ (stationary). They share a common stochastic trend.
2. **Spurious regression:** Regressing one $I(1)$ variable on another unrelated $I(1)$ variable gives high R^2 and significant coefficients even though there's no true relationship.
3. **α coefficients:** Speed of adjustment—how quickly each variable responds to deviations from long-run equilibrium.
4. **Johansen advantage:** Can test for multiple cointegrating relationships, uses MLE (more efficient), doesn't require choosing dependent variable.
5. **$\alpha_i = 0$:** Variable Y_i is weakly exogenous—it doesn't respond to disequilibrium. Other variables do all the adjusting.